



$$I(J^P) = \frac{1}{2}(0^-)$$

D^0 MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1864.80 ± 0.14				OUR FIT
1864.91 ± 0.17				OUR AVERAGE
1865.30 ± 0.33 ± 0.23	98 ± 13	ANASHIN	10A KEDR	e^+e^- at $\psi(3770)$
1864.847 ± 0.150 ± 0.095	319 ± 18	CAWLFIELD	07 CLEO	$D^0 \rightarrow K_S^0 \phi$
1864.6 ± 0.3 ± 1.0	641	BARLAG	90C ACCM	π^- Cu 230 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1852 ± 7	16	ADAMOVICH	87 EMUL	Photoproduction
1856 ± 36	22	ADAMOVICH	84B EMUL	Photoproduction
1861 ± 4		DERRICK	84 HRS	e^+e^- 29 GeV
1847 ± 7	1	FIORINO	81 EMUL	$\gamma N \rightarrow \bar{D}^0 +$
1863.8 ± 0.5		¹ SCHINDLER	81 MRK2	e^+e^- 3.77 GeV
1864.7 ± 0.6		¹ TRILLING	81 RVUE	e^+e^- 3.77 GeV
1863.0 ± 2.5	238	ASTON	80E OMEG	$\gamma p \rightarrow \bar{D}^0$
1860 ± 2	143	² AVERY	80 SPEC	$\gamma N \rightarrow D^{*+}$
1869 ± 4	35	² AVERY	80 SPEC	$\gamma N \rightarrow D^{*+}$
1854 ± 6	94	² ATIYA	79 SPEC	$\gamma N \rightarrow D^0 \bar{D}^0$
1850 ± 15	64	BALTAY	78C HBC	$\nu N \rightarrow K^0 \pi \pi$
1863 ± 3		GOLDHABER	77 MRK1	D^0, D^+ recoil spectra
1863.3 ± 0.9		¹ PERUZZI	77 LGW	e^+e^- 3.77 GeV
1868 ± 11		PICCOLO	77 MRK1	e^+e^- 4.03, 4.41 GeV
1865 ± 15	234	GOLDHABER	76 MRK1	$K\pi$ and $K3\pi$

¹PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision $J/\psi(1S)$ and $\psi(2S)$ measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted. TRILLING 81 enters the fit in the D^\pm mass, and PERUZZI 77 and SCHINDLER 81 enter in the $m_{D^\pm} - m_{D^0}$, below.

²Error does not include possible systematic mass scale shift, estimated to be less than 5 MeV.

$m_{D^\pm} - m_{D^0}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4.77 ± 0.10			OUR FIT Error includes scale factor of 1.1.
4.74 ± 0.28			OUR AVERAGE
4.7 ± 0.3	³ SCHINDLER	81 MRK2	e^+e^- 3.77 GeV
5.0 ± 0.8	³ PERUZZI	77 LGW	e^+e^- 3.77 GeV

³ See the footnote on TRILLING 81 in the D^0 and D^\pm sections on the mass.

D^0 MEAN LIFE

Measurements with an error $> 10 \times 10^{-15}$ s have been omitted from the average.

<u>VALUE (10^{-15} s)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
410.1 ± 1.5 OUR AVERAGE				
409.6 ± 1.1 ± 1.5	210k	LINK	02F FOCS	γ nucleus, ≈ 180 GeV
407.9 ± 6.0 ± 4.3	10k	KUSHNIR...	01 SELX	$K^- \pi^+$, $K^- \pi^+ \pi^+ \pi^-$
413 ± 3 ± 4	35k	AITALA	99E E791	$K^- \pi^+$
408.5 ± 4.1 ⁺ _{-3.4}	25k	BONVICINI	99 CLE2	$e^+ e^- \approx \Upsilon(4S)$
413 ± 4 ± 3	16k	FRABETTI	94D E687	$K^- \pi^+$, $K^- \pi^+ \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
424 ± 11 ± 7	5118	FRABETTI	91 E687	$K^- \pi^+$, $K^- \pi^+ \pi^+ \pi^-$
417 ± 18 ± 15	890	ALVAREZ	90 NA14	$K^- \pi^+$, $K^- \pi^+ \pi^+ \pi^-$
388 ⁺²³ ₋₂₁	641	⁴ BARLAG	90C ACCM	π^- Cu 230 GeV
480 ± 40 ± 30	776	ALBRECHT	88I ARG	$e^+ e^-$ 10 GeV
422 ± 8 ± 10	4212	RAAB	88 E691	Photoproduction
420 ± 50	90	BARLAG	87B ACCM	K^- and π^- 200 GeV

⁴ BARLAG 90C estimate systematic error to be negligible.

A REVIEW GOES HERE – Check our WWW List of Reviews

$$|m_{D_1^0} - m_{D_2^0}| = x \Gamma$$

The D_1^0 and D_2^0 are the mass eigenstates of the D^0 meson, as described in the note on “ D^0 - \bar{D}^0 Mixing,” above. The experiments usually present $x \equiv \Delta m / \Gamma$. Then $\Delta m = x \Gamma = x \hbar / \tau$.

We are leaving the D^0 mixing numbers unchanged from the 2010 Review, awaiting a comprehensive re-analysis by the Heavy Flavor Averaging Group. This would include as-yet-unpublished results from CLEO. Note, however, that there are herein new measurements of the D_1^0 - D_2^0 mass and width differences from BABAR.

<u>VALUE ($10^{10} \hbar s^{-1}$)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.39^{+0.59}_{-0.63} OUR EVALUATION HFAG fit; see the note on “ D^0 - \bar{D}^0 Mixing.”				
(1.0 ± 0.8) OUR AVERAGE Error includes scale factor of 1.5.				
0.39 ± 0.56 ± 0.35		⁵ DEL-AMO-SA..10D	BABR	$e^+ e^-$, 10.6 GeV
1.98 ± 0.73 ^{+0.32} _{-0.41}		⁶ ZHANG	07B BELL	$\Delta m < 3.9$, 95% CL

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.4	$\begin{matrix} +1.4 \\ -1.7 \end{matrix}$	± 1.0	7	AUBERT	09AN	BABR	e^+e^-	at 10.58 GeV
- 2	$\begin{matrix} +7 \\ -6 \end{matrix}$		8	LOWREY	09	CLEO	e^+e^-	at $\psi(3770)$
< 7		95	9	ZHANG	06	BELL	e^+e^-	
-11	to +22		6	ASNER	05	CLEO	e^+e^-	≈ 10 GeV
< 11		90		BITENC	05	BELL		
< 30		90		CAWLFIELD	05	CLEO		
< 7		95	9	LI	05A	BELL	See ZHANG	06
< 22		95	10	LINK	05H	FOCS	γ	nucleus
< 23		95		AUBERT	04Q	BABR		
< 11		95	9	AUBERT	03Z	BABR	e^+e^- ,	10.6 GeV
< 7		95	11	GODANG	00	CLE2	e^+e^-	
< 32		90	12,13	AITALA	98	E791	π^-	nucleus, 500 GeV
< 24		90	14	AITALA	96C	E791	π^-	nucleus, 500 GeV
< 21		90	13,15	ANJOS	88C	E691	Photoproduction	

⁵ DEL-AMO-SANCHEZ 10D uses $540,800 \pm 800 K_S^0 \pi^+ \pi^-$ and $79,900 \pm 300 K_S^0 K^+ K^-$ events in a time-dependent amplitude analysis of the D^0 and \bar{D}^0 Dalitz plots. No evidence was found for CP violation, and the values here assume no such violation.

⁶ The ASNER 05 and ZHANG 07B values are from the time-dependent Dalitz-plot analysis of $D^0 \rightarrow K_S^0 \pi^+ \pi^-$. Decay-time information and interference on the Dalitz plot are used to distinguish doubly Cabibbo-suppressed decays from mixing and to measure the relative phase between $D^0 \rightarrow K^{*+} \pi^-$ and $\bar{D}^0 \rightarrow K^{*+} \pi^-$. This value allows CP violation and is sensitive to the sign of Δm .

⁷ The AUBERT 09AN values are inferred from the branching ratio $\Gamma(D^0 \rightarrow K^+ \pi^- \pi^0 \text{ via } \bar{D}^0) / \Gamma(D^0 \rightarrow K^- \pi^+ \pi^0)$ given near the end of this Listings. Mixing is distinguished from DCS decays using decay-time information. Interference between mixing and DCS is allowed. The phase between $D^0 \rightarrow K^+ \pi^- \pi^0$ and $\bar{D}^0 \rightarrow K^+ \pi^- \pi^0$ is assumed to be small. The width difference here is y'' , which is not the same as y_{CP} in the note on D^0 - \bar{D}^0 mixing.

⁸ LOWREY 09 uses quantum correlations in $e^+e^- \rightarrow D^0 \bar{D}^0$ at the $\psi(3770)$. See below for coherence factors and average relative strong phases for both $D^0 \rightarrow K^- \pi^+ \pi^0$ and $D^0 \rightarrow K^- \pi^- 2\pi^+$. A fit that includes external measurements of charm mixing parameters gets $\Delta m = (2.34 \pm 0.61) \times 10^{10} \hbar s^{-1}$.

⁹ The AUBERT 03Z, LI 05A, and ZHANG 06 limits are inferred from the D^0 - \bar{D}^0 mixing ratio $\Gamma(K^+ \pi^- \text{ (via } \bar{D}^0)) / \Gamma(K^- \pi^+)$ given near the end of this D^0 Listings. Decay-time information is used to distinguish DCS decays from D^0 - \bar{D}^0 mixing. The limit allows interference between the DCS and mixing ratios, and also allows CP violation. AUBERT 03Z assumes the strong phase between $D^0 \rightarrow K^+ \pi^-$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ amplitudes is small; if an arbitrary phase is allowed, the limit degrades by 20%. The LI 05A and ZHANG 06 limits are valid for an arbitrary strong phase.

¹⁰ This LINK 05H limit is inferred from the D^0 - \bar{D}^0 mixing ratio $\Gamma(K^+ \pi^- \text{ (via } \bar{D}^0)) / \Gamma(K^- \pi^+)$ given near the end of this D^0 Listings. Decay-time information is used to distinguish DCS decays from D^0 - \bar{D}^0 mixing. The limit allows interference between the DCS and mixing ratios, and also allows CP violation. The strong phase between $D^0 \rightarrow K^+ \pi^-$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ is assumed to be small. If an arbitrary relative strong phase is allowed, the limit degrades by 25%.

¹¹ This GODANG 00 limit is inferred from the D^0 - \bar{D}^0 mixing ratio $\Gamma(K^+ \pi^- \text{ (via } \bar{D}^0)) / \Gamma(K^- \pi^+)$ given near the end of this D^0 Listings. Decay-time information is used to distinguish DCS decays from D^0 - \bar{D}^0 mixing. The limit allows interference between

the DCS and mixing ratios, and also allows CP violation. The strong phase between $D^0 \rightarrow K^+ \pi^-$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ is assumed to be small. If an arbitrary relative strong phase is allowed, the limit degrades by a factor of two.

- ¹² AITALA 98 allows interference between the doubly Cabibbo-suppressed and mixing amplitudes, and also allows CP violation in this term, but assumes that $A_D=A_R=0$. See the note on “ D^0 - \bar{D}^0 Mixing,” above.
- ¹³ This limit is inferred from R_M for $f = K^+ \pi^-$ and $f = K^+ \pi^- \pi^+ \pi^-$. See the note on “ D^0 - \bar{D}^0 Mixing,” above. Decay-time information is used to distinguish doubly Cabibbo-suppressed decays from D^0 - \bar{D}^0 mixing.
- ¹⁴ This limit is inferred from R_M for $f = K^+ \ell^- \bar{\nu}_\ell$. See the note on “ D^0 - \bar{D}^0 Mixing,” above.
- ¹⁵ ANJOS 88C assumes that $y = 0$. See the note on “ D^0 - \bar{D}^0 Mixing,” above. Without this assumption, the limit degrades by about a factor of two.

$$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2y$$

The D_1^0 and D_2^0 are the mass eigenstates of the D^0 meson, as described in the note on “ D^0 - \bar{D}^0 Mixing,” above.

Due to the strong phase difference between $D^0 \rightarrow K^+ \pi^-$ and $\bar{D}^0 \rightarrow K^+ \pi^-$, we exclude from the average those measurements of y' that are inferred from the D^0 - \bar{D}^0 mixing ratio $\Gamma(K^+ \pi^- \text{ via } \bar{D}^0) / \Gamma(K^+ \pi^-)$ given near the end of this D^0 Listings.

Some early results have been omitted. See our 2006 *Review* (Journal of Physics, G **33** 1 (2006)).

We are leaving the D^0 mixing numbers unchanged from the 2010 Review, awaiting a comprehensive re-analysis by the Heavy Flavor Averaging Group. This would include as-yet-unpublished results from CLEO. Note, however, that there are herein new measurements of the D_1^0 - D_2^0 mass and width differences from BABAR.

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	COMMENT
1.66 ± 0.32 OUR EVALUATION		HFAG fit; see the note on “ D^0 - \bar{D}^0 Mixing.”		
1.5 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
1.14 ± 0.40 ± 0.30		¹⁶ DEL-AMO-SA..10D	BABR	$e^+ e^-$, 10.6 GeV
2.32 ± 0.44 ± 0.36		¹⁷ AUBERT	09Al	BABR $e^+ e^- \approx \Upsilon(4S)$
0.22 ± 1.22 ± 1.04		¹⁸ ZUPANC	09	BELL $e^+ e^- \approx \Upsilon(4S)$
2.62 ± 0.64 ± 0.50	160k	¹⁹ STARIC	07	BELL $e^+ e^- \approx \Upsilon(4S)$
0.74 ± 0.50 ^{+0.20} _{-0.31}	534k	²⁰ ZHANG	07B	BELL $e^+ e^- \approx \Upsilon(4S)$
-1.0 ± 2.0 ^{+1.4} _{-1.6}	18k	²¹ ABE	02I	BELL $e^+ e^- \approx \Upsilon(4S)$
-2.4 ± 5.0 ± 2.8	3393	²² CSORNA	02	CLE2 $e^+ e^- \approx \Upsilon(4S)$
6.84 ± 2.78 ± 1.48	10k	²¹ LINK	00	FOCS γ nucleus
+1.6 ± 5.8 ± 2.1		²¹ AITALA	99E	E791 $K^- \pi^+, K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.12^{+1.10}_{-1.28} \pm 0.68$		²³ AUBERT	09AN BABR	e^+e^- at 10.58 GeV
$1.4^{+4.8}_{-5.4}$		²⁴ LOWREY	09 CLEO	e^+e^- at $\psi(3770)$
1.70 ± 1.52	$12.7 \pm 0.3k$	²⁵ AALTONEN	08E CDF	$p\bar{p}$, $\sqrt{s} = 1.96$ TeV
$2.06 \pm 0.66 \pm 0.38$		²⁶ AUBERT	08U BABR	See AUBERT 09AI
$1.94 \pm 0.88 \pm 0.62$	4030 ± 90	²⁵ AUBERT	07W BABR	$e^+e^- \approx 10.6$ GeV
-0.7 ± 4.9	$4k \pm 88$	^{25,27} ZHANG	06 BELL	e^+e^-
$-3.0^{+5.0}_{-4.8} +1.6_{-0.8}$		²⁰ ASNER	05 CLEO	$e^+e^- \approx 10$ GeV
-0.3 ± 5.7		^{25,27} LI	05A BELL	See ZHANG 06
$-5.2^{+18.4}_{-16.8}$		^{25,27} LINK	05H FOCS	γ nucleus
$1.6 \pm 0.8^{+1.0}_{-0.8}$	$450k$	²⁸ AUBERT	03P BABR	See AUBERT 08U
$1.6^{+6.2}_{-12.8}$		^{25,27} AUBERT	03Z BABR	e^+e^- , 10.6 GeV
$-5.0^{+2.8}_{-3.2} \pm 0.6$		²⁵ GODANG	00 CLE2	e^+e^-

¹⁶ DEL-AMO-SANCHEZ 10D uses $540,800 \pm 800 K_S^0 \pi^+ \pi^-$ and $79,900 \pm 300 K_S^0 K^+ K^-$ events in a time-dependent amplitude analyses of the D^0 and \bar{D}^0 Dalitz plots. No evidence was found for CP violation, and the values here assume no such violation.

¹⁷ This combines the $y_{CP} = (\tau_{K\pi}/\tau_{KK}) - 1$ using untagged $K^- \pi^+$ and $K^- K^+$ events of AUBERT 09AI with the disjoint y_{CP} using tagged $K^- \pi^+$, $K^- K^+$, and $\pi^- \pi^+$ events of AUBERT 08U.

¹⁸ ZUPANC 09 uses a method based on measuring the mean decay time of $D^0 \rightarrow K_S^0 K^+ K^-$ events for different $K^+ K^-$ mass intervals.

¹⁹ STARIC 07 compares the lifetimes of D^0 decay to the CP eigenstates $K^+ K^-$ and $\pi^+ \pi^-$ with D^0 decay to $K^- \pi^+$.

²⁰ The ASNER 05 and ZHANG 07B values are from the time-dependent Dalitz-plot analysis of $D^0 \rightarrow K_S^0 \pi^+ \pi^-$. Decay-time information and interference on the Dalitz plot are used to distinguish doubly Cabibbo-suppressed decays from mixing and to measure the relative phase between $D^0 \rightarrow K^{*+} \pi^-$ and $\bar{D}^0 \rightarrow K^{*+} \pi^-$. This limit allows CP violation.

²¹ LINK 00, AITALA 99E, and ABE 02I measure the lifetime difference between $D^0 \rightarrow K^- K^+$ (CP even) decays and $D^0 \rightarrow K^- \pi^+$ (CP mixed) decays, or $y_{CP} = [\Gamma(CP+) - \Gamma(CP-)] / [\Gamma(CP+) + \Gamma(CP-)]$. We list $2y_{CP} = \Delta\Gamma/\Gamma$.

²² CSORNA 02 measures the lifetime difference between $D^0 \rightarrow K^- K^+$ and $\pi^- \pi^+$ (CP even) decays and $D^0 \rightarrow K^- \pi^+$ (CP mixed) decays, or $y_{CP} = [\Gamma(CP+) - \Gamma(CP-)] / [\Gamma(CP+) + \Gamma(CP-)]$. We list $2y_{CP} = \Delta\Gamma/\Gamma$.

²³ The AUBERT 09AN values are inferred from the branching ratio $\Gamma(D^0 \rightarrow K^+ \pi^- \pi^0)$ via $\bar{D}^0) / \Gamma(D^0 \rightarrow K^- \pi^+ \pi^0)$ given near the end of this Listings. Mixing is distinguished from DCS decays using decay-time information. Interference between mixing and DCS is allowed. The phase between $D^0 \rightarrow K^+ \pi^- \pi^0$ and $\bar{D}^0 \rightarrow K^+ \pi^- \pi^0$ is assumed to be small. The width difference here is y'' , which is not the same as y_{CP} in the note on $D^0 - \bar{D}^0$ mixing.

²⁴ LOWREY 09 uses quantum correlations in $e^+ e^- \rightarrow D^0 \bar{D}^0$ at the $\psi(3770)$. See below for coherence factors and average relative strong phases for both $D^0 \rightarrow K^- \pi^+ \pi^0$ and $D^0 \rightarrow K^- \pi^- 2\pi^+$. A fit that includes external measurements of charm mixing parameters gets $2y = (1.62 \pm 0.32) \times 10^{-2}$.


²⁵ The GODANG 00, AUBERT 03Z, LINK 05H, LI 05A, ZHANG 06, AUBERT 07W, and AALTONEN 08E limits are inferred from the D^0 - \bar{D}^0 mixing ratio $\Gamma(K^+\pi^- \text{ (via } \bar{D}^0)) / \Gamma(K^-\pi^+)$ given near the end of this D^0 Listings. Decay-time information is used to distinguish DCS decays from D^0 - \bar{D}^0 mixing. The limits allow interference between the DCS and mixing ratios, and all except AUBERT 07W and AALTONEN 08E also allow CP violation. The phase between $D^0 \rightarrow K^+\pi^-$ and $\bar{D}^0 \rightarrow K^+\pi^-$ is assumed to be small. This is a measurement of y' and is not the same as the y_{CP} of our note above on " D^0 - \bar{D}^0 Mixing."

²⁶ This value combines the results of AUBERT 08U and AUBERT 03P.

²⁷ The ranges of AUBERT 03Z, LINK 05H, LI 05A, and ZHANG 06 measurements are for 95% confidence level.

²⁸ AUBERT 03P measures $Y \equiv 2\tau^0 / (\tau^+ + \tau^-) - 1$, where τ^0 is the $D^0 \rightarrow K^-\pi^+$ (and $\bar{D}^0 \rightarrow K^+\pi^-$) lifetime, and τ^+ and τ^- are the D^0 and \bar{D}^0 lifetimes to CP -even states (here K^-K^+ and $\pi^-\pi^+$). In the limit of CP conservation, $Y = y \equiv \Delta\Gamma / 2\Gamma$ (we list $2y = \Delta\Gamma/\Gamma$). AUBERT 03P also uses $\tau^+ - \tau^-$ to get $\Delta Y = -0.008 \pm 0.006 \pm 0.002$.

WEIGHTED AVERAGE
 1.5 ± 0.4 (Error scaled by 1.3)



|q/p|

The mass eigenstates D_1^0 and D_2^0 are related to the $C = \pm 1$ states by $|D_{1,2}^0\rangle = p|D^0\rangle + q|\bar{D}^0\rangle$. See the note on " D^0 - \bar{D}^0 Mixing" above.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.86^{+0.18}_{-0.15}$ OUR EVALUATION	HFAG fit;	see the note on " D^0 - \bar{D}^0 Mixing."	
$0.86^{+0.30}_{-0.29}$ +0.10 -0.08	²⁹ ZHANG	07B BELL	$e^+e^- \approx \tau(4S)$

²⁹ The phase of p/q is $(-14_{-18}^{+16} \pm 5)^\circ$. The ZHANG 07B value is from the time-dependent Dalitz-plot analysis of $D^0 \rightarrow K_S^0 \pi^+ \pi^-$. Decay-time information and interference on the Dalitz plot are used to distinguish doubly Cabibbo-suppressed decays from mixing and to measure the relative phase between $D^0 \rightarrow K^{*+} \pi^-$ and $\bar{D}^0 \rightarrow K^{*+} \pi^-$. This value allows CP violation.

A_Γ

A_Γ is the decay-rate asymmetry for CP -even final states $A_\Gamma = (\bar{\tau}_+ - \tau_+) / (\bar{\tau}_+ + \tau_+)$.

See the note on “ D^0 - \bar{D}^0 Mixing” above.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.4 ± 2.7 OUR AVERAGE			
$+2.6 \pm 3.6 \pm 0.8$	AUBERT	08U	BABR $e^+ e^- \approx \Upsilon(4S)$
$+0.1 \pm 3.0 \pm 2.5$	STARIC	07	BELL $e^+ e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$+8 \pm 6 \pm 2$	AUBERT	03P	BABR $e^+ e^- \approx \Upsilon(4S)$

$\cos \delta$

δ is the $D^0 \rightarrow K^+ \pi^-$ relative strong phase.

VALUE	DOCUMENT ID	TECN	COMMENT
$1.03_{-0.17}^{+0.31} \pm 0.06$	³⁰ ASNER	08	CLEO $e^+ e^- \rightarrow D^0 \bar{D}^0$, 3.77 GeV

³⁰ ASNER 08 uses quantum correlations in $e^+ e^- \rightarrow D^0 \bar{D}^0$ at the $\psi(3770)$, where decay rates of CP -tagged $K\pi$ final states depend on $\cos \delta$ because of interfering amplitudes. The above measurement implies $|\delta| < 75^\circ$ with a confidence level of 95%. A fit that includes external measurements of charm mixing parameters finds $\cos \delta = 1.10 \pm 0.35 \pm 0.07$. See also the note on “ D^0 - \bar{D}^0 Mixing” p. 783 in our 2008 Review (PDG 08).

$D^0 \rightarrow K^- \pi^+ \pi^0$ COHERENCE FACTOR $R_{K\pi\pi^0}$

See the note on ‘ D^0 - \bar{D}^0 Mixing’ for the definition. $R_{K\pi\pi^0}$ can have any value between 0 and 1. A value near 1 indicates the decay is dominated by a few intermediate states with limited interference.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.78_{-0.25}^{+0.11}$	³¹ LOWREY	09	CLEO $e^+ e^- \rightarrow D^0 \bar{D}^0$ at $\psi(3770)$

³¹ LOWREY 09 uses quantum correlations in $e^+ e^- \rightarrow D^0 \bar{D}^0$ at the $\psi(3770)$, where the decay rates of CP -tagged $K^- \pi^+ \pi^0$ final states depend on $R_{K\pi\pi^0}$ and $\delta^{K\pi\pi^0}$. A fit that includes external measurements of charm mixing parameters gets $R_{K\pi\pi^0} = 0.84 \pm 0.07$.

$D^0 \rightarrow K^- \pi^+ \pi^0$ AVERAGE RELATIVE STRONG PHASE $\delta^{K\pi\pi^0}$

VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
239_{-28}^{+32}	³² LOWREY	09	CLEO $e^+ e^- \rightarrow D^0 \bar{D}^0$ at $\psi(3770)$

³² LOWREY 09 uses quantum correlations in $e^+ e^- \rightarrow D^0 \bar{D}^0$ at the $\psi(3770)$, where the decay rates of CP -tagged $K^- \pi^+ \pi^0$ final states depend on $R_{K\pi\pi^0}$ and $\delta^{K\pi\pi^0}$. A fit that includes external measurements of charm mixing parameters gets $\delta^{K\pi\pi^0} = (227_{-17}^{+14})^\circ$.

$D^0 \rightarrow K^- \pi^- 2\pi^+$ COHERENCE FACTOR $R_{K3\pi}$

See the note on ' D^0 - \bar{D}^0 Mixing' for the definition. $R_{K3\pi}$ can have any value between 0 and 1. A value near 1 indicates the decay is dominated by a few intermediate states with limited interference.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.36^{+0.24}_{-0.30}$	³³ LOWREY	09	CLEO $e^+e^- \rightarrow D^0\bar{D}^0$ at $\psi(3770)$

³³ LOWREY 09 uses quantum correlations in $e^+e^- \rightarrow D^0\bar{D}^0$ at the $\psi(3770)$, where the decay rates of CP -tagged $K^- \pi^- 2\pi^+$ final states depend on $R_{K3\pi}$ and $\delta^{K3\pi}$. A fit that includes external measurements of charm mixing parameters gets $R_{K3\pi} = 0.33^{+0.26}_{-0.23}$.

$D^0 \rightarrow K^- \pi^- 2\pi^+$ AVERAGE RELATIVE STRONG PHASE $\delta^{K3\pi}$

VALUE (°)	DOCUMENT ID	TECN	COMMENT
118^{+62}_{-53}	³⁴ LOWREY	09	CLEO $e^+e^- \rightarrow D^0\bar{D}^0$ at $\psi(3770)$

³⁴ LOWREY 09 uses quantum correlations in $e^+e^- \rightarrow D^0\bar{D}^0$ at the $\psi(3770)$, where the decay rates of CP -tagged $K^- \pi^- 2\pi^+$ final states depend on $R_{K3\pi}$ and $\delta^{K3\pi}$. A fit that includes external measurements of charm mixing parameters gets $\delta^{K3\pi} = (114^{+26}_{-23})^\circ$.

D^0 DECAY MODES

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

Topological modes

Γ_1	0-prongs	[a] (17 ± 6) %	
Γ_2	2-prongs	(69 ± 6) %	
Γ_3	4-prongs	[b] (14.3 ± 0.5) %	
Γ_4	6-prongs	[c] (6.4 ± 1.3) × 10 ⁻⁴	

Inclusive modes

Γ_5	e^+ anything	[d] (6.49 ± 0.11) %	
Γ_6	μ^+ anything	(6.7 ± 0.6) %	
Γ_7	K^- anything	(54.7 ± 2.8) %	S=1.3
Γ_8	\bar{K}^0 anything + K^0 anything	(47 ± 4) %	
Γ_9	K^+ anything	(3.4 ± 0.4) %	
Γ_{10}	$K^*(892)^-$ anything	(15 ± 9) %	

Γ_{11}	$\bar{K}^*(892)^0$ anything	(9 ± 4) %	
Γ_{12}	$K^*(892)^+$ anything	< 3.6	CL=90%
Γ_{13}	$K^*(892)^0$ anything	(2.8 ± 1.3) %	
Γ_{14}	η anything	(9.5 ± 0.9) %	
Γ_{15}	η' anything	(2.48 ± 0.27) %	
Γ_{16}	ϕ anything	(1.05 ± 0.11) %	

Semileptonic modes

Γ_{17}	$K^- \ell^+ \nu_\ell$		
Γ_{18}	$K^- e^+ \nu_e$	(3.55 ± 0.04) %	S=1.1
Γ_{19}	$K^- \mu^+ \nu_\mu$	(3.30 ± 0.13) %	
Γ_{20}	$K^*(892)^- e^+ \nu_e$	(2.16 ± 0.16) %	
Γ_{21}	$K^*(892)^- \mu^+ \nu_\mu$	(1.90 ± 0.23) %	
Γ_{22}	$K^- \pi^0 e^+ \nu_e$	(1.6 + - 0.5) %	
Γ_{23}	$\bar{K}^0 \pi^- e^+ \nu_e$	(2.7 + - 0.7) %	
Γ_{24}	$K^- \pi^+ \pi^- e^+ \nu_e$	(2.8 + - 1.1) × 10 ⁻⁴	
Γ_{25}	$K_1(1270)^- e^+ \nu_e$	(7.6 + - 3.1) × 10 ⁻⁴	
Γ_{26}	$K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.2	CL=90%
Γ_{27}	$(\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu$	< 1.4	CL=90%
Γ_{28}	$\pi^- e^+ \nu_e$	(2.89 ± 0.08) × 10 ⁻³	S=1.1
Γ_{29}	$\pi^- \mu^+ \nu_\mu$	(2.37 ± 0.24) × 10 ⁻³	
Γ_{30}	$\rho^- e^+ \nu_e$	(1.9 ± 0.4) × 10 ⁻³	

Hadronic modes with one \bar{K}

Γ_{31}	$K^- \pi^+$	(3.87 ± 0.05) %	S=1.2
Γ_{32}	$K_S^0 \pi^0$	(1.19 ± 0.05) %	S=1.3
Γ_{33}	$K_L^0 \pi^0$	(10.0 ± 0.7) × 10 ⁻³	
Γ_{34}	$K_S^0 \pi^+ \pi^-$	[e] (2.81 ± 0.15) %	S=1.1
Γ_{35}	$K_S^0 \rho^0$	(6.3 + - 0.7) × 10 ⁻³	
Γ_{36}	$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-$	(2.0 ± 0.6) × 10 ⁻⁴	
Γ_{37}	$K_S^0 (\pi^+ \pi^-)_{S\text{-wave}}$	(3.3 ± 0.8) × 10 ⁻³	
Γ_{38}	$K_S^0 f_0(980),$ $f_0(980) \rightarrow \pi^+ \pi^-$	(1.21 + - 0.23) × 10 ⁻³	
Γ_{39}	$K_S^0 f_0(1370),$ $f_0(1370) \rightarrow \pi^+ \pi^-$	(2.8 + - 1.3) × 10 ⁻³	
Γ_{40}	$K_S^0 f_2(1270),$ $f_2(1270) \rightarrow \pi^+ \pi^-$	(9 + - 6) × 10 ⁻⁵	
Γ_{41}	$K^*(892)^- \pi^+,$ $K^*(892)^- \rightarrow K_S^0 \pi^-$	(1.65 + - 0.13) %	

Γ_{42}	$K_0^*(1430)^- \pi^+,$ $K_0^*(1430)^- \rightarrow K_S^0 \pi^-$	$(2.69 \pm 0.40) \times 10^{-3}$ $(- 0.31)$	
Γ_{43}	$K_2^*(1430)^- \pi^+,$ $K_2^*(1430)^- \rightarrow K_S^0 \pi^-$	$(3.4 \pm 1.9) \times 10^{-4}$ $(- 1.0)$	
Γ_{44}	$K^*(1680)^- \pi^+,$ $K^*(1680)^- \rightarrow K_S^0 \pi^-$	$(4 \pm 4) \times 10^{-4}$	
Γ_{45}	$K^*(892)^+ \pi^-,$ $K^*(892)^+ \rightarrow K_S^0 \pi^+$	[f] $(1.13 \pm 0.60) \times 10^{-4}$ $(- 0.34)$	
Γ_{46}	$K_0^*(1430)^+ \pi^-,$ $K_0^*(1430)^+ \rightarrow K_S^0 \pi^+$	[f] < 1.4	$\times 10^{-5}$ CL=95%
Γ_{47}	$K_2^*(1430)^+ \pi^-,$ $K_2^*(1430)^+ \rightarrow K_S^0 \pi^+$	[f] < 3.4	$\times 10^{-5}$ CL=95%
Γ_{48}	$K_S^0 \pi^+ \pi^-$ nonresonant	$(2.5 \pm 6.0) \times 10^{-4}$ $(- 1.6)$	
Γ_{49}	$K^- \pi^+ \pi^0$	[e] $(13.9 \pm 0.5) \%$	S=1.7
Γ_{50}	$K^- \rho^+$	$(10.8 \pm 0.7) \%$	
Γ_{51}	$K^- \rho(1700)^+,$ $\rho(1700)^+ \rightarrow \pi^+ \pi^0$	$(7.9 \pm 1.7) \times 10^{-3}$	
Γ_{52}	$K^*(892)^- \pi^+,$ $K^*(892)^- \rightarrow K^- \pi^0$	$(2.21 \pm 0.40) \%$ $(- 0.19)$	
Γ_{53}	$\bar{K}^*(892)^0 \pi^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(1.88 \pm 0.23) \%$	
Γ_{54}	$K_0^*(1430)^- \pi^+,$ $K_0^*(1430)^- \rightarrow K^- \pi^0$	$(4.6 \pm 2.1) \times 10^{-3}$	
Γ_{55}	$\bar{K}_0^*(1430)^0 \pi^0,$ $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	$(5.7 \pm 5.0) \times 10^{-3}$ $(- 1.5)$	
Γ_{56}	$K^*(1680)^- \pi^+,$ $K^*(1680)^- \rightarrow K^- \pi^0$	$(1.8 \pm 0.7) \times 10^{-3}$	
Γ_{57}	$K^- \pi^+ \pi^0$ nonresonant	$(1.11 \pm 0.50) \%$ $(- 0.19)$	
Γ_{58}	$K_S^0 2\pi^0$	$(8.3 \pm 0.6) \times 10^{-3}$	
Γ_{59}	$\bar{K}^*(892)^0 \pi^0,$ $\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	$(6.6 \pm 1.8) \times 10^{-3}$ $(- 1.5)$	
Γ_{60}	$K_S^0 2\pi^0$ nonresonant	$(4.4 \pm 1.1) \times 10^{-3}$	
Γ_{61}	$K^- 2\pi^+ \pi^-$	[e] $(8.07 \pm 0.21) \%$ $(- 0.19)$	S=1.4
Γ_{62}	$K^- \pi^+ \rho^0$ total	$(6.74 \pm 0.33) \%$	
Γ_{63}	$K^- \pi^+ \rho^0$ 3-body	$(5.1 \pm 2.3) \times 10^{-3}$	
Γ_{64}	$\bar{K}^*(892)^0 \rho^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(1.05 \pm 0.23) \%$	
Γ_{65}	$K^- a_1(1260)^+,$ $a_1(1260)^+ \rightarrow 2\pi^+ \pi^-$	$(3.6 \pm 0.6) \%$	

Γ ₆₆	$\bar{K}^*(892)^0 \pi^+ \pi^-$ total, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.6 ± 0.4) %	
Γ ₆₇	$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(9.9 ± 2.3) × 10 ⁻³	
Γ ₆₈	$K_1(1270)^- \pi^+$, $K_1(1270)^- \rightarrow K^- \pi^+ \pi^-$	[g] (2.9 ± 0.3) × 10 ⁻³	
Γ ₆₉	$K^- 2\pi^+ \pi^-$ nonresonant	(1.88 ± 0.26) %	
Γ ₇₀	$K_S^0 \pi^+ \pi^- \pi^0$	[h] (5.2 ± 0.6) %	
Γ ₇₁	$K_S^0 \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$	(1.02 ± 0.09) × 10 ⁻³	
Γ ₇₂	$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	(9.9 ± 0.5) × 10 ⁻³	
Γ ₇₃	$K^- \pi^+ 2\pi^0$		
Γ ₇₄	$K^- 2\pi^+ \pi^- \pi^0$	(4.2 ± 0.4) %	
Γ ₇₅	$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.3 ± 0.6) %	
Γ ₇₆	$K^- \pi^+ \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	(2.7 ± 0.5) %	
Γ ₇₇	$\bar{K}^*(892)^0 \omega$, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$, $\omega \rightarrow \pi^+ \pi^- \pi^0$	(6.5 ± 3.0) × 10 ⁻³	
Γ ₇₈	$K_S^0 \eta \pi^0$	(5.5 ± 1.1) × 10 ⁻³	
Γ ₇₉	$K_S^0 a_0(980), a_0(980) \rightarrow \eta \pi^0$	(6.5 ± 2.0) × 10 ⁻³	
Γ ₈₀	$\bar{K}^*(892)^0 \eta$, $\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	(1.6 ± 0.5) × 10 ⁻³	
Γ ₈₁	$K_S^0 2\pi^+ 2\pi^-$	(2.67 ± 0.28) × 10 ⁻³	
Γ ₈₂	$K_S^0 \rho^0 \pi^+ \pi^-$, no $K^*(892)^-$	(1.1 ± 0.7) × 10 ⁻³	
Γ ₈₃	$K^*(892)^- 2\pi^+ \pi^-$, $K^*(892)^- \rightarrow K_S^0 \pi^-$, no ρ^0	(5 ± 8) × 10 ⁻⁴	
Γ ₈₄	$K^*(892)^- \rho^0 \pi^+$, $K^*(892)^- \rightarrow K_S^0 \pi^-$	(1.6 ± 0.6) × 10 ⁻³	
Γ ₈₅	$K_S^0 2\pi^+ 2\pi^-$ nonresonant	< 1.2 × 10 ⁻³	CL=90%
Γ ₈₆	$\bar{K}^0 \pi^+ \pi^- 2\pi^0 (\pi^0)$		
Γ ₈₇	$K^- 3\pi^+ 2\pi^-$	(2.2 ± 0.6) × 10 ⁻⁴	

Fractions of many of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. (Modes for which there are only upper limits and $\bar{K}^*(892)\rho$ submodes only appear below.)

Γ ₈₈	$K_S^0 \eta$	(4.5 ± 0.4) × 10 ⁻³	S=1.5
Γ ₈₉	$K_S^0 \omega$	(1.11 ± 0.06) %	
Γ ₉₀	$K_S^0 \eta'(958)$	(9.4 ± 0.5) × 10 ⁻³	
Γ ₉₁	$K^- a_1(1260)^+$	(7.8 ± 1.1) %	
Γ ₉₂	$K^- a_2(1320)^+$	< 2 × 10 ⁻³	CL=90%
Γ ₉₃	$\bar{K}^*(892)^0 \pi^+ \pi^-$ total	(2.4 ± 0.5) %	

Γ_{94}	$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body	(1.48 ± 0.34) %	
Γ_{95}	$\bar{K}^*(892)^0 \rho^0$	(1.57 ± 0.34) %	
Γ_{96}	$\bar{K}^*(892)^0 \rho^0$ transverse	(1.7 ± 0.6) %	
Γ_{97}	$\bar{K}^*(892)^0 \rho^0$ S-wave	(3.0 ± 0.6) %	
Γ_{98}	$\bar{K}^*(892)^0 \rho^0$ S-wave long.	< 3	$\times 10^{-3}$ CL=90%
Γ_{99}	$\bar{K}^*(892)^0 \rho^0$ P-wave	< 3	$\times 10^{-3}$ CL=90%
Γ_{100}	$\bar{K}^*(892)^0 \rho^0$ D-wave	(2.1 ± 0.6) %	
Γ_{101}	$K^- \pi^+ f_0(980)$		
Γ_{102}	$\bar{K}^*(892)^0 f_0(980)$		
Γ_{103}	$K_1(1270)^- \pi^+$	[g] (1.6 ± 0.8) %	
Γ_{104}	$K_1(1400)^- \pi^+$	< 1.2	% CL=90%
Γ_{105}	$K^*(1410)^- \pi^+$		
Γ_{106}	$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$	(1.9 ± 0.9) %	
Γ_{107}	$\bar{K}^*(892)^0 \eta$		
Γ_{108}	$K^- \pi^+ \omega$	(3.0 ± 0.6) %	
Γ_{109}	$\bar{K}^*(892)^0 \omega$	(1.1 ± 0.5) %	
Γ_{110}	$K^- \pi^+ \eta'(958)$	(7.5 ± 1.9) $\times 10^{-3}$	
Γ_{111}	$\bar{K}^*(892)^0 \eta'(958)$	< 1.1	$\times 10^{-3}$ CL=90%

Hadronic modes with three K's

Γ_{112}	$K_S^0 K^+ K^-$	(4.45 ± 0.28) $\times 10^{-3}$	
Γ_{113}	$K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-$	(3.0 ± 0.4) $\times 10^{-3}$	
Γ_{114}	$K^- a_0(980)^+, a_0^+ \rightarrow K^+ K_S^0$	(6.0 ± 1.8) $\times 10^{-4}$	
Γ_{115}	$K^+ a_0(980)^-, a_0^- \rightarrow K^- K_S^0$	< 1.1	$\times 10^{-4}$ CL=95%
Γ_{116}	$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$	< 9	$\times 10^{-5}$ CL=95%
Γ_{117}	$K_S^0 \phi, \phi \rightarrow K^+ K^-$	(2.04 ± 0.14) $\times 10^{-3}$	
Γ_{118}	$K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-$	(1.7 ± 1.1) $\times 10^{-4}$	
Γ_{119}	$3K_S^0$	(9.1 ± 1.3) $\times 10^{-4}$	
Γ_{120}	$K^+ 2K^- \pi^+$	(2.21 ± 0.31) $\times 10^{-4}$	
Γ_{121}	$K^+ K^- \bar{K}^*(892)^0, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(4.4 ± 1.7) $\times 10^{-5}$	
Γ_{122}	$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$	(4.0 ± 1.7) $\times 10^{-5}$	
Γ_{123}	$\phi \bar{K}^*(892)^0, \phi \rightarrow K^+ K^-, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.06 ± 0.20) $\times 10^{-4}$	
Γ_{124}	$K^+ 2K^- \pi^+$ nonresonant	(3.3 ± 1.5) $\times 10^{-5}$	
Γ_{125}	$2K_S^0 K^\pm \pi^\mp$	(6.0 ± 1.3) $\times 10^{-4}$	

Pionic modes

Γ_{126}	$\pi^+ \pi^-$	(1.400 ± 0.026) $\times 10^{-3}$	S=1.1
Γ_{127}	$2\pi^0$	(8.0 ± 0.5) $\times 10^{-4}$	
Γ_{128}	$\pi^+ \pi^- \pi^0$	(1.43 ± 0.06) %	S=1.9
Γ_{129}	$\rho^+ \pi^-$	(9.8 ± 0.4) $\times 10^{-3}$	
Γ_{130}	$\rho^0 \pi^0$	(3.72 ± 0.22) $\times 10^{-3}$	

Γ_{131}	$\rho^- \pi^+$	$(4.96 \pm 0.24) \times 10^{-3}$	
Γ_{132}	$\rho(1450)^+ \pi^-, \rho(1450)^+ \rightarrow$ $\pi^+ \pi^0$	$(1.6 \pm 2.0) \times 10^{-5}$	
Γ_{133}	$\rho(1450)^0 \pi^0, \rho(1450)^0 \rightarrow$ $\pi^+ \pi^-$	$(4.3 \pm 1.9) \times 10^{-5}$	
Γ_{134}	$\rho(1450)^- \pi^+, \rho(1450)^- \rightarrow$ $\pi^- \pi^0$	$(2.6 \pm 0.4) \times 10^{-4}$	
Γ_{135}	$\rho(1700)^+ \pi^-, \rho(1700)^+ \rightarrow$ $\pi^+ \pi^0$	$(5.9 \pm 1.4) \times 10^{-4}$	
Γ_{136}	$\rho(1700)^0 \pi^0, \rho(1700)^0 \rightarrow$ $\pi^+ \pi^-$	$(7.2 \pm 1.7) \times 10^{-4}$	
Γ_{137}	$\rho(1700)^- \pi^+, \rho(1700)^- \rightarrow$ $\pi^- \pi^0$	$(4.6 \pm 1.1) \times 10^{-4}$	
Γ_{138}	$f_0(980) \pi^0, f_0(980) \rightarrow$	$(3.6 \pm 0.8) \times 10^{-5}$	
Γ_{139}	$f_0(600) \pi^0, f_0(600) \rightarrow$	$(1.17 \pm 0.21) \times 10^{-4}$	
Γ_{140}	$(\pi^+ \pi^-)_{S\text{-wave}} \pi^0$		
Γ_{141}	$f_0(1370) \pi^0, f_0(1370) \rightarrow$	$(5.3 \pm 2.0) \times 10^{-5}$	
Γ_{142}	$f_0(1500) \pi^0, f_0(1500) \rightarrow$ $\pi^+ \pi^-$	$(5.6 \pm 1.5) \times 10^{-5}$	
Γ_{143}	$f_0(1710) \pi^0, f_0(1710) \rightarrow$ $\pi^+ \pi^-$	$(4.4 \pm 1.5) \times 10^{-5}$	
Γ_{144}	$f_2(1270) \pi^0, f_2(1270) \rightarrow$ $\pi^+ \pi^-$	$(1.89 \pm 0.20) \times 10^{-4}$	
Γ_{145}	$\pi^+ \pi^- \pi^0$ nonresonant	$(1.20 \pm 0.35) \times 10^{-4}$	
Γ_{146}	$3\pi^0$	$< 3.5 \times 10^{-4}$	CL=90%
Γ_{147}	$2\pi^+ 2\pi^-$	$(7.42 \pm 0.21) \times 10^{-3}$	S=1.1
Γ_{148}	$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$ $2\pi^+ \pi^-$ total	$(4.45 \pm 0.31) \times 10^{-3}$	
Γ_{149}	$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$ $\rho^0 \pi^+$ S-wave	$(3.21 \pm 0.25) \times 10^{-3}$	
Γ_{150}	$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$ $\rho^0 \pi^+$ D-wave	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{151}	$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$	$(6.2 \pm 0.7) \times 10^{-4}$	
Γ_{152}	$2\rho^0$ total	$(1.82 \pm 0.13) \times 10^{-3}$	
Γ_{153}	$2\rho^0$, parallel helicities	$(8.2 \pm 3.2) \times 10^{-5}$	
Γ_{154}	$2\rho^0$, perpendicular helicities	$(4.7 \pm 0.6) \times 10^{-4}$	
Γ_{155}	$2\rho^0$, longitudinal helicities	$(1.25 \pm 0.10) \times 10^{-3}$	
Γ_{156}	Resonant $(\pi^+ \pi^-) \pi^+ \pi^-$ 3-body total	$(1.48 \pm 0.12) \times 10^{-3}$	
Γ_{157}	$\sigma \pi^+ \pi^-$	$(6.1 \pm 0.9) \times 10^{-4}$	
Γ_{158}	$f_0(980) \pi^+ \pi^-, f_0 \rightarrow$ $\pi^+ \pi^-$	$(1.8 \pm 0.5) \times 10^{-4}$	
Γ_{159}	$f_2(1270) \pi^+ \pi^-, f_2 \rightarrow$ $\pi^+ \pi^-$	$(3.6 \pm 0.6) \times 10^{-4}$	
Γ_{160}	$\pi^+ \pi^- 2\pi^0$	$(10.0 \pm 0.9) \times 10^{-3}$	

Γ_{161}	$\eta\pi^0$	$[i]$	$(6.8 \pm 0.7) \times 10^{-4}$	
Γ_{162}	$\omega\pi^0$	$[i]$	$< 2.6 \times 10^{-4}$	CL=90%
Γ_{163}	$2\pi^+2\pi^-\pi^0$		$(4.1 \pm 0.5) \times 10^{-3}$	
Γ_{164}	$\eta\pi^+\pi^-$	$[i]$	$(1.09 \pm 0.16) \times 10^{-3}$	
Γ_{165}	$\omega\pi^+\pi^-$	$[i]$	$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{166}	$3\pi^+3\pi^-$		$(4.2 \pm 1.2) \times 10^{-4}$	
Γ_{167}	$\eta'(958)\pi^0$		$(8.9 \pm 1.4) \times 10^{-4}$	
Γ_{168}	$\eta'(958)\pi^+\pi^-$		$(4.5 \pm 1.7) \times 10^{-4}$	
Γ_{169}	2η		$(1.67 \pm 0.20) \times 10^{-3}$	
Γ_{170}	$\eta\eta'(958)$		$(1.05 \pm 0.26) \times 10^{-3}$	

Hadronic modes with a $K\bar{K}$ pair

Γ_{171}	K^+K^-		$(3.96 \pm 0.08) \times 10^{-3}$	S=1.4
Γ_{172}	$2K_S^0$		$(1.73 \pm 0.29) \times 10^{-4}$	S=1.7
Γ_{173}	$K_S^0K^-\pi^+$		$(3.3 \pm 0.5) \times 10^{-3}$	S=1.1
Γ_{174}	$\bar{K}^*(892)^0 K_S^0,$ $\bar{K}^*(892)^0 \rightarrow K^-\pi^+$		$< 5 \times 10^{-4}$	CL=90%
Γ_{175}	$K_S^0K^+\pi^-$		$(2.6 \pm 0.5) \times 10^{-3}$	
Γ_{176}	$K^*(892)^0 K_S^0,$ $K^*(892)^0 \rightarrow K^+\pi^-$		$< 2.8 \times 10^{-4}$	CL=90%
Γ_{177}	$K^+K^-\pi^0$		$(3.28 \pm 0.14) \times 10^{-3}$	
Γ_{178}	$K^*(892)^+K^-,$ $K^*(892)^+ \rightarrow K^+\pi^0$		$(1.46 \pm 0.07) \times 10^{-3}$	
Γ_{179}	$K^*(892)^-K^+,$ $K^*(892)^- \rightarrow K^-\pi^0$		$(5.2 \pm 0.4) \times 10^{-4}$	
Γ_{180}	$(K^+\pi^0)_{S-wave}K^-$		$(2.33 \pm 0.17) \times 10^{-3}$	
Γ_{181}	$(K^-\pi^0)_{S-wave}K^+$		$(1.3 \pm 0.4) \times 10^{-4}$	
Γ_{182}	$f_0(980)\pi^0, f_0 \rightarrow K^+K^-$		$(3.4 \pm 0.6) \times 10^{-4}$	
Γ_{183}	$\phi\pi^0, \phi \rightarrow K^+K^-$		$(6.4 \pm 0.4) \times 10^{-4}$	
Γ_{184}	$K^+K^-\pi^0$ nonresonant			
Γ_{185}	$2K_S^0\pi^0$		$< 5.9 \times 10^{-4}$	
Γ_{186}	$K^+K^-\pi^+\pi^-$	$[j]$	$(2.42 \pm 0.12) \times 10^{-3}$	
Γ_{187}	$\phi\pi^+\pi^-$ 3-body, $\phi \rightarrow K^+K^-$		$(2.4 \pm 2.4) \times 10^{-5}$	
Γ_{188}	$\phi\rho^0, \phi \rightarrow K^+K^-$		$(7.0 \pm 0.6) \times 10^{-4}$	
Γ_{189}	$K^+K^-\rho^0$ 3-body		$(5 \pm 7) \times 10^{-5}$	
Γ_{190}	$f_0(980)\pi^+\pi^-, f_0 \rightarrow K^+K^-$		$(3.6 \pm 0.9) \times 10^{-4}$	
Γ_{191}	$K^*(892)^0 K^\mp\pi^\pm$ 3-body, $K^{*0} \rightarrow K^\pm\pi^\mp$	$[k]$	$(2.7 \pm 0.6) \times 10^{-4}$	
Γ_{192}	$K^*(892)^0 \bar{K}^*(892)^0, K^{*0} \rightarrow$ $K^\pm\pi^\mp$		$(7 \pm 5) \times 10^{-5}$	
Γ_{193}	$K_1(1270)^\pm K^\mp,$ $K_1(1270)^\pm \rightarrow K^\pm\pi^+\pi^-$		$(8.0 \pm 1.8) \times 10^{-4}$	
Γ_{194}	$K_1(1400)^\pm K^\mp,$ $K_1(1400)^\pm \rightarrow K^\pm\pi^+\pi^-$		$(5.3 \pm 1.2) \times 10^{-4}$	

Γ_{195}	$2K_S^0 \pi^+ \pi^-$		$(1.22 \pm 0.23) \times 10^{-3}$	
Γ_{196}	$K_S^0 K^- 2\pi^+ \pi^-$		$< 1.4 \times 10^{-4}$	CL=90%
Γ_{197}	$K^+ K^- \pi^+ \pi^- \pi^0$		$(3.1 \pm 2.0) \times 10^{-3}$	

Other $K\bar{K}X$ modes. They include all decay modes of the ϕ , η , and ω .

Γ_{198}	$\phi \pi^0$			
Γ_{199}	$\phi \eta$		$(1.4 \pm 0.5) \times 10^{-4}$	
Γ_{200}	$\phi \omega$		$< 2.1 \times 10^{-3}$	CL=90%

Radiative modes

Γ_{201}	$\rho^0 \gamma$		$< 2.4 \times 10^{-4}$	CL=90%
Γ_{202}	$\omega \gamma$		$< 2.4 \times 10^{-4}$	CL=90%
Γ_{203}	$\phi \gamma$		$(2.69 \pm 0.35) \times 10^{-5}$	
Γ_{204}	$\bar{K}^*(892)^0 \gamma$		$(3.27 \pm 0.34) \times 10^{-4}$	

Doubly Cabibbo suppressed (DC) modes or $\Delta C = 2$ forbidden via mixing (C2M) modes

Γ_{205}	$K^+ \ell^- \bar{\nu}_\ell$ via \bar{D}^0		$< 2.2 \times 10^{-5}$	CL=90%
Γ_{206}	K^+ or $K^*(892)^+$ $e^- \bar{\nu}_e$ via \bar{D}^0		$< 6 \times 10^{-5}$	CL=90%
Γ_{207}	$K^+ \pi^-$	DC	$(1.47 \pm 0.07) \times 10^{-4}$	S=2.8
Γ_{208}	$K^+ \pi^-$ via DCS		$(1.31 \pm 0.08) \times 10^{-4}$	
Γ_{209}	$K^+ \pi^-$ via \bar{D}^0		$< 1.5 \times 10^{-5}$	CL=95%
Γ_{210}	$K_S^0 \pi^+ \pi^-$ in $D^0 \rightarrow \bar{D}^0$		$< 1.8 \times 10^{-4}$	CL=95%
Γ_{211}	$K^*(892)^+ \pi^-$, $K^*(892)^+ \rightarrow K_S^0 \pi^+$	DC	$(1.13 \pm_{-0.34}^{0.60}) \times 10^{-4}$	
Γ_{212}	$K_0^*(1430)^+ \pi^-$, $K_0^*(1430)^+ \rightarrow K_S^0 \pi^+$	DC	$< 1.4 \times 10^{-5}$	
Γ_{213}	$K_2^*(1430)^+ \pi^-$, $K_2^*(1430)^+ \rightarrow K_S^0 \pi^+$	DC	$< 3.4 \times 10^{-5}$	
Γ_{214}	$K^+ \pi^- \pi^0$	DC	$(3.04 \pm 0.17) \times 10^{-4}$	
Γ_{215}	$K^+ \pi^- \pi^0$ via \bar{D}^0		$(7.3 \pm 0.5) \times 10^{-4}$	
Γ_{216}	$K^+ \pi^+ 2\pi^-$	DC	$(2.61 \pm_{-0.19}^{0.21}) \times 10^{-4}$	
Γ_{217}	$K^+ \pi^+ 2\pi^-$ via \bar{D}^0		$< 4 \times 10^{-4}$	CL=90%
Γ_{218}	$K^+ \pi^-$ or $K^+ \pi^+ 2\pi^-$ via \bar{D}^0			
Γ_{219}	μ^- anything via \bar{D}^0		$< 4 \times 10^{-4}$	CL=90%

$\Delta C = 1$ weak neutral current (C1) modes, Lepton Family number (LF) violating modes, Lepton (L) or Baryon (B) number violating modes

Γ_{220}	$\gamma \gamma$	C1	$< 2.6 \times 10^{-5}$	CL=90%
Γ_{221}	$e^+ e^-$	C1	$< 7.9 \times 10^{-8}$	CL=90%
Γ_{222}	$\mu^+ \mu^-$	C1	$< 1.4 \times 10^{-7}$	CL=90%

Γ ₂₂₃	$\pi^0 e^+ e^-$	<i>C1</i>	< 4.5	$\times 10^{-5}$	CL=90%
Γ ₂₂₄	$\pi^0 \mu^+ \mu^-$	<i>C1</i>	< 1.8	$\times 10^{-4}$	CL=90%
Γ ₂₂₅	$\eta e^+ e^-$	<i>C1</i>	< 1.1	$\times 10^{-4}$	CL=90%
Γ ₂₂₆	$\eta \mu^+ \mu^-$	<i>C1</i>	< 5.3	$\times 10^{-4}$	CL=90%
Γ ₂₂₇	$\pi^+ \pi^- e^+ e^-$	<i>C1</i>	< 3.73	$\times 10^{-4}$	CL=90%
Γ ₂₂₈	$\rho^0 e^+ e^-$	<i>C1</i>	< 1.0	$\times 10^{-4}$	CL=90%
Γ ₂₂₉	$\pi^+ \pi^- \mu^+ \mu^-$	<i>C1</i>	< 3.0	$\times 10^{-5}$	CL=90%
Γ ₂₃₀	$\rho^0 \mu^+ \mu^-$	<i>C1</i>	< 2.2	$\times 10^{-5}$	CL=90%
Γ ₂₃₁	$\omega e^+ e^-$	<i>C1</i>	< 1.8	$\times 10^{-4}$	CL=90%
Γ ₂₃₂	$\omega \mu^+ \mu^-$	<i>C1</i>	< 8.3	$\times 10^{-4}$	CL=90%
Γ ₂₃₃	$K^- K^+ e^+ e^-$	<i>C1</i>	< 3.15	$\times 10^{-4}$	CL=90%
Γ ₂₃₄	$\phi e^+ e^-$	<i>C1</i>	< 5.2	$\times 10^{-5}$	CL=90%
Γ ₂₃₅	$K^- K^+ \mu^+ \mu^-$	<i>C1</i>	< 3.3	$\times 10^{-5}$	CL=90%
Γ ₂₃₆	$\phi \mu^+ \mu^-$	<i>C1</i>	< 3.1	$\times 10^{-5}$	CL=90%
Γ ₂₃₇	$\bar{K}^0 e^+ e^-$		[<i>l</i>] < 1.1	$\times 10^{-4}$	CL=90%
Γ ₂₃₈	$\bar{K}^0 \mu^+ \mu^-$		[<i>l</i>] < 2.6	$\times 10^{-4}$	CL=90%
Γ ₂₃₉	$K^- \pi^+ e^+ e^-$	<i>C1</i>	< 3.85	$\times 10^{-4}$	CL=90%
Γ ₂₄₀	$\bar{K}^*(892)^0 e^+ e^-$		[<i>l</i>] < 4.7	$\times 10^{-5}$	CL=90%
Γ ₂₄₁	$K^- \pi^+ \mu^+ \mu^-$	<i>C1</i>	< 3.59	$\times 10^{-4}$	CL=90%
Γ ₂₄₂	$\bar{K}^*(892)^0 \mu^+ \mu^-$		[<i>l</i>] < 2.4	$\times 10^{-5}$	CL=90%
Γ ₂₄₃	$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	<i>C1</i>	< 8.1	$\times 10^{-4}$	CL=90%
Γ ₂₄₄	$\mu^\pm e^\mp$	<i>LF</i>	[<i>m</i>] < 2.6	$\times 10^{-7}$	CL=90%
Γ ₂₄₅	$\pi^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 8.6	$\times 10^{-5}$	CL=90%
Γ ₂₄₆	$\eta e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 1.0	$\times 10^{-4}$	CL=90%
Γ ₂₄₇	$\pi^+ \pi^- e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 1.5	$\times 10^{-5}$	CL=90%
Γ ₂₄₈	$\rho^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 4.9	$\times 10^{-5}$	CL=90%
Γ ₂₄₉	$\omega e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 1.2	$\times 10^{-4}$	CL=90%
Γ ₂₅₀	$K^- K^+ e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 1.8	$\times 10^{-4}$	CL=90%
Γ ₂₅₁	$\phi e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 3.4	$\times 10^{-5}$	CL=90%
Γ ₂₅₂	$\bar{K}^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 1.0	$\times 10^{-4}$	CL=90%
Γ ₂₅₃	$K^- \pi^+ e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 5.53	$\times 10^{-4}$	CL=90%
Γ ₂₅₄	$\bar{K}^*(892)^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>m</i>] < 8.3	$\times 10^{-5}$	CL=90%
Γ ₂₅₅	$2\pi^- 2e^+ + \text{c.c.}$	<i>L</i>	< 1.12	$\times 10^{-4}$	CL=90%
Γ ₂₅₆	$2\pi^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 2.9	$\times 10^{-5}$	CL=90%
Γ ₂₅₇	$K^- \pi^- 2e^+ + \text{c.c.}$	<i>L</i>	< 2.06	$\times 10^{-4}$	CL=90%
Γ ₂₅₈	$K^- \pi^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 3.9	$\times 10^{-4}$	CL=90%
Γ ₂₅₉	$2K^- 2e^+ + \text{c.c.}$	<i>L</i>	< 1.52	$\times 10^{-4}$	CL=90%
Γ ₂₆₀	$2K^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 9.4	$\times 10^{-5}$	CL=90%
Γ ₂₆₁	$\pi^- \pi^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 7.9	$\times 10^{-5}$	CL=90%
Γ ₂₆₂	$K^- \pi^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 2.18	$\times 10^{-4}$	CL=90%
Γ ₂₆₃	$2K^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 5.7	$\times 10^{-5}$	CL=90%
Γ ₂₆₄	$p e^-$	<i>L,B</i>	[<i>n</i>] < 1.0	$\times 10^{-5}$	CL=90%
Γ ₂₆₅	$\bar{p} e^+$	<i>L,B</i>	[<i>o</i>] < 1.1	$\times 10^{-5}$	CL=90%

Γ_{266} A dummy mode used by the fit. (38.2 ± 1.3) % $S=1.1$

- [a] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
 - [b] This is the sum of our $K^- 2\pi^+ \pi^-$, $K^- 2\pi^+ \pi^- \pi^0$, $\bar{K}^0 2\pi^+ 2\pi^-$, $K^+ 2K^- \pi^+$, $2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^- \pi^0$, $K^+ K^- \pi^+ \pi^-$, and $K^+ K^- \pi^+ \pi^- \pi^0$, branching fractions.
 - [c] This is the sum of our $K^- 3\pi^+ 2\pi^-$ and $3\pi^+ 3\pi^-$ branching fractions.
 - [d] The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to 6.20 ± 0.17 %.
 - [e] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.
 - [f] This is a doubly Cabibbo-suppressed mode.
 - [g] The two experiments measuring this fraction are in serious disagreement. See the Particle Listings.
 - [h] Submodes of the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ mode with a K^* and/or ρ were studied by COFFMAN 92B, but with only 140 events. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
 - [i] This branching fraction includes all the decay modes of the resonance in the final state.
 - [j] The experiments on the division of this charge mode amongst its submodes disagree, and the submode branching fractions here add up to considerably more than the charged-mode fraction.
 - [k] However, these upper limits are in serious disagreement with values obtained in another experiment.
 - [l] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
 - [m] The value is for the sum of the charge states or particle/antiparticle states indicated.
 - [n] This limit is for either D^0 or \bar{D}^0 to $p e^-$.
 - [o] This limit is for either D^0 or \bar{D}^0 to $\bar{p} e^+$.
-

CONSTRAINED FIT INFORMATION

An overall fit to 55 branching ratios uses 111 measurements and one constraint to determine 31 parameters. The overall fit has a $\chi^2 = 104.8$ for 81 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{18}	2										
x_{19}	20	9									
x_{20}	0	1	0								
x_{28}	0	0	0	0							
x_{29}	3	2	17	0	0						
x_{31}	4	48	19	2	0	3					
x_{32}	1	14	5	2	0	1	28				
x_{34}	0	6	2	12	0	0	12	16			
x_{49}	0	-2	-1	0	0	0	-4	-1	0		
x_{61}	1	10	4	0	0	1	22	6	3	55	
x_{70}	0	2	1	4	0	0	4	5	33	0	
x_{74}	0	4	2	0	0	0	8	2	1	8	
x_{88}	0	6	2	2	0	0	13	14	16	0	
x_{89}	0	0	0	0	0	0	0	1	4	0	
x_{90}	1	10	4	2	0	1	20	8	17	-1	
x_{126}	2	30	12	1	0	2	63	18	8	-2	
x_{127}	1	9	4	0	0	1	19	6	2	-1	
x_{128}	0	-1	0	0	0	0	-1	0	0	82	
x_{147}	1	13	5	0	0	1	27	8	3	29	
x_{161}	0	5	2	0	0	0	11	3	1	0	
x_{167}	0	4	1	0	0	0	7	2	1	0	
x_{169}	0	5	2	0	0	0	10	3	1	0	
x_{170}	0	2	1	0	0	0	5	1	1	0	
x_{171}	2	30	11	1	0	2	61	17	8	-2	
x_{172}	0	3	1	0	0	0	7	2	1	0	
x_{173}	0	2	1	4	0	0	5	5	30	0	
x_{175}	0	2	1	3	0	0	4	4	21	0	
x_{203}	0	4	2	0	0	0	9	3	1	0	
x_{207}	1	12	5	0	0	1	24	7	3	-1	
x_{266}	-49	-13	-23	-16	-1	-6	-21	-13	-33	-53	
	x_6	x_{18}	x_{19}	x_{20}	x_{28}	x_{29}	x_{31}	x_{32}	x_{34}	x_{49}	

x70	1										
x74	16	0									
x88	3	5	1								
x89	0	12	0	1							
x90	4	6	2	5	1						
x126	14	3	5	8	0	13					
x127	4	1	2	2	0	4	12				
x128	46	0	7	0	0	0	-1	0			
x147	58	1	10	3	0	5	17	5	25		
x161	2	0	1	1	0	2	7	2	0	3	
x167	2	0	1	1	0	2	5	1	0	2	
x169	2	0	1	1	0	2	6	2	0	3	
x170	1	0	0	1	0	1	3	1	0	1	
x171	13	2	5	8	0	12	38	12	-1	16	
x172	1	0	1	1	0	1	4	1	0	2	
x173	1	10	0	5	1	5	3	1	0	1	
x175	1	7	0	3	1	4	3	1	0	1	
x203	2	0	1	1	0	2	6	2	0	2	
x207	5	1	2	3	0	5	15	5	0	7	
x266	-48	-51	-38	-10	-11	-12	-13	-4	-46	-30	
	x61	x70	x74	x88	x89	x90	x126	x127	x128	x147	
x167	1										
x169	1	1									
x170	1	0	0								
x171	7	5	6	3							
x172	1	0	1	0	4						
x173	1	0	1	0	3	0					
x175	0	0	0	0	3	0	6				
x203	1	1	1	0	8	1	0	0			
x207	3	2	3	1	15	2	1	1	2		
x266	-3	-3	-4	-3	-13	-2	-14	-11	-2	-5	
	x161	x167	x169	x170	x171	x172	x173	x175	x203	x207	

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 3 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 0.0$ for 0 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100		
x_3	-47	40	
x_4	0	0	0
	x_1	x_2	x_3

D^0 BRANCHING RATIOS

Some older now obsolete results have been omitted from these Listings.

Topological modes

$\Gamma(0\text{-prongs})/\Gamma_{\text{total}}$

Γ_1/Γ

This value is obtained by subtracting the branching fractions for 2-, 4-, and 6-prongs from unity.

VALUE	DOCUMENT ID
0.17±0.06 OUR FIT	

$\Gamma(4\text{-prongs})/\Gamma_{\text{total}}$

Γ_3/Γ

This is the sum of our $K^- 2\pi^+ \pi^-$, $K^- 2\pi^+ \pi^- \pi^0$, $\bar{K}^0 2\pi^+ 2\pi^-$, $K^+ 2K^- \pi^+$, $2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^- \pi^0$, $K^+ K^- \pi^+ \pi^-$, and $K^+ K^- \pi^+ \pi^- \pi^0$ branching fractions.

VALUE	DOCUMENT ID
0.143±0.005 OUR FIT	
0.143±0.005	PDG 10

$\Gamma(4\text{-prongs})/\Gamma(2\text{-prongs})$

Γ_3/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.207±0.016 OUR FIT				
0.207±0.016±0.004	226	ONENGUT	05	CHRS ν_μ emulsion, $\bar{E}_\nu \approx 27$ GeV

$\Gamma(6\text{-prongs})/\Gamma_{\text{total}}$

Γ_4/Γ

This is the sum of our $K^- 3\pi^+ 2\pi^-$ and $3\pi^+ 3\pi^-$ branching fractions.

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(6.4±1.3) OUR FIT				
6.4± 1.3		PDG	10	

• • • We do not use the following data for averages, fits, limits, etc. • • •

12 $\begin{smallmatrix} +13 \\ -9 \end{smallmatrix}$ ±2	3	ONENGUT	05	CHRS ν_μ emulsion, $\bar{E}_\nu \approx 27$ GeV
---	---	---------	----	---

————— Inclusive modes —————

$\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_5/Γ

The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to 6.20 ± 0.17 %.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.49±0.11 OUR AVERAGE				
6.46±0.09±0.11	6584 ± 96	³⁵ ASNER	10 CLEO	$e^+ e^-$ at 3774 MeV
6.3 ±0.7 ±0.4	290 ± 32	ABLIKIM	07G BES2	$e^+ e^- \approx \psi(3770)$
6.46±0.17±0.13	2246 ± 57	ADAM	06A CLEO	See ASNER 10
6.9 ±0.3 ±0.5	1670	ALBRECHT	96C ARG	$e^+ e^- \approx 10$ GeV
6.64±0.18±0.29	4609	KUBOTA	96B CLE2	$e^+ e^- \approx \Upsilon(4S)$

³⁵ Using the D^+ and D^0 lifetimes, ASNER 10 finds that the ratio of the D^+ and D^0 semileptonic widths is $0.985 \pm 0.015 \pm 0.024$.

$\Gamma(\mu^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_6/Γ

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.7±0.6 OUR FIT				
6.4±0.8 OUR AVERAGE				
6.8±1.5±0.8	79 ± 10	³⁶ ABLIKIM	08L BES2	$e^+ e^- \approx \psi(3772)$
6.5±1.2±0.3	36	KAYIS-TOPAK.05	CHRS	ν_μ emulsion
6.0±0.7±1.2	310	ALBRECHT	96C ARG	$e^+ e^- \approx 10$ GeV


³⁶ ABLIKIM 08L finds the ratio of $D^+ \rightarrow \mu^+ X$ and $D^0 \rightarrow \mu^+ X$ branching fractions to be $2.59 \pm 0.70 \pm 0.25$, in accord with the ratio of D^+ and D^0 lifetimes, 2.54 ± 0.02 .

$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$

Γ_7/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.547±0.028 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.				
0.578±0.016±0.032	2098 ± 59	ABLIKIM	07G BES2	$e^+ e^- \approx \psi(3770)$
0.546 ^{+0.039} _{-0.038}		³⁷ BARLAG	92C ACCM	π^- Cu 230 GeV
0.609±0.032±0.052		COFFMAN	91 MRK3	$e^+ e^-$ 3.77 GeV
0.42 ±0.08		AGUILAR-...	87E HYBR	$\pi p, pp$ 360, 400 GeV
0.55 ±0.11	121	SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV
0.35 ±0.10	19	VUILLEMIN	78 LGW	$e^+ e^-$ 3.772 GeV

³⁷ BARLAG 92C computes the branching fraction using topological normalization.



$[\Gamma(\bar{K}^0 \text{ anything}) + \Gamma(K^0 \text{ anything})]/\Gamma_{\text{total}}$ **Γ_8/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.47 ± 0.04	OUR AVERAGE			
$0.476 \pm 0.048 \pm 0.030$	250 ± 25	ABLIKIM	06U BES2	e^+e^- at 3773 MeV
$0.455 \pm 0.050 \pm 0.032$		COFFMAN	91 MRK3	e^+e^- 3.77 GeV

$\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$ **Γ_9/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.034 ± 0.004	OUR AVERAGE			
$0.035 \pm 0.007 \pm 0.003$	119 ± 23	ABLIKIM	07G BES2	$e^+e^- \approx \psi(3770)$
$0.034^{+0.007}_{-0.005}$	³⁸	BARLAG	92C ACCM	π^- Cu 230 GeV
$0.028 \pm 0.009 \pm 0.004$		COFFMAN	91 MRK3	e^+e^- 3.77 GeV
$0.03^{+0.05}_{-0.02}$		AGUILAR-...	87E HYBR	$\pi p, pp$ 360, 400 GeV
0.08 ± 0.03	25	SCHINDLER	81 MRK2	e^+e^- 3.771 GeV

³⁸ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^*(892)^- \text{ anything})/\Gamma_{\text{total}}$ **Γ_{10}/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.153 \pm 0.083 \pm 0.019$	28 ± 15	ABLIKIM	06U BES2	e^+e^- at 3773 MeV

$\Gamma(\bar{K}^*(892)^0 \text{ anything})/\Gamma_{\text{total}}$ **Γ_{11}/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.087 \pm 0.040 \pm 0.012$	96 ± 44	ABLIKIM	05P BES	$e^+e^- \approx 3773$ MeV

$\Gamma(K^*(892)^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.036	90	ABLIKIM	06U BES2	e^+e^- at 3773 MeV

$\Gamma(K^*(892)^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.028 \pm 0.012 \pm 0.004$	31 ± 12	ABLIKIM	05P BES	$e^+e^- \approx 3773$ MeV

$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$ Γ_{14}/Γ

This ratio includes η' particles from η' decays.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.5 \pm 0.4 \pm 0.8$	4463 ± 197	HUANG	06B CLEO	e^+e^- at $\psi(3770)$

$\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.48 \pm 0.17 \pm 0.21$	299 ± 21	HUANG	06B CLEO	e^+e^- at $\psi(3770)$

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.05 \pm 0.08 \pm 0.07$	368 ± 24	HUANG	06B CLEO	e^+e^- at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.71^{+0.76}_{-0.71} \pm 0.17$	9	BAI	00C BES	$e^+e^- \rightarrow D\bar{D}^*, D^*\bar{D}^*$
---------------------------------	---	-----	---------	---

————— Semileptonic modes —————

$\Gamma(K^- e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.55 ± 0.04 OUR FIT Error includes scale factor of 1.1.

3.50 ± 0.05 OUR AVERAGE

$3.50 \pm 0.03 \pm 0.04$	14.1k	³⁹ BESSON	09 CLEO	e^+e^- at $\psi(3770)$
$3.45 \pm 0.10 \pm 0.19$	1318 ± 38	⁴⁰ WIDHALM	06 BELL	$e^+e^- \approx \Upsilon(4S)$
$3.82 \pm 0.40 \pm 0.27$	104 ± 11	ABLIKIM	04C BES	e^+e^- , 3.773 GeV
$3.4 \pm 0.5 \pm 0.4$	55	ADLER	89 MRK3	e^+e^- 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.56 \pm 0.03 \pm 0.09$		⁴¹ DOBBS	08 CLEO	See BESSON 09
$3.44 \pm 0.10 \pm 0.10$	1311 ± 37	COAN	05 CLEO	See DOBBS 08

³⁹ See the form-factor parameters near the end of this D^0 Listing.

⁴⁰ The $\pi^- e^+ \nu_e$ and $K^- e^+ \nu_e$ results of WIDHALM 06 give $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}|^2 = 0.042 \pm 0.003 \pm 0.003$.

⁴¹ DOBBS 08 establishes $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}| = 0.188 \pm 0.008 \pm 0.002$ from the D^+ and D^0 decays to $\bar{K} e^+ \nu_e$ and $\pi e^+ \nu_e$.

$\Gamma(K^- e^+ \nu_e)/\Gamma(K^- \pi^+)$					Γ_{18}/Γ_{31}
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.915±0.011 OUR FIT					Error includes scale factor of 1.1.
0.930±0.013 OUR AVERAGE					
0.927±0.007±0.012	76k±323	42 AUBERT	07BG BABR	$e^+e^- \approx \Upsilon(4S)$	
0.978±0.027±0.044	2510	43 BEAN	93C CLE2	$e^+e^- \approx \Upsilon(4S)$	
0.90 ±0.06 ±0.06	584	44 CRAWFORD	91B CLEO	$e^+e^- \approx 10.5$ GeV	
0.91 ±0.07 ±0.11	250	45 ANJOS	89F E691	Photoproduction	

⁴²The event samples in this AUBERT 07BG result include radiative photons. The $D^0 \rightarrow K^- e^+ \nu_e$ form factor at $q^2 = 0$ is $f_+(0) = 0.727 \pm 0.007 \pm 0.005 \pm 0.007$.

⁴³BEAN 93C uses $K^- \mu^+ \nu_\mu$ as well as $K^- e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events. A pole mass of $2.00 \pm 0.12 \pm 0.18$ GeV/ c^2 is obtained from the q^2 dependence of the decay rate.

⁴⁴CRAWFORD 91B uses $K^- e^+ \nu_e$ and $K^- \mu^+ \nu_\mu$ candidates to measure a pole mass of $2.1_{-0.2}^{+0.4+0.3}$ GeV/ c^2 from the q^2 dependence of the decay rate.

⁴⁵ANJOS 89F measures a pole mass of $2.1_{-0.2}^{+0.4} \pm 0.2$ GeV/ c^2 from the q^2 dependence of the decay rate.

$\Gamma(K^- \mu^+ \nu_\mu)/\Gamma_{\text{total}}$					Γ_{19}/Γ
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.30±0.13 OUR FIT					
3.45±0.10±0.21	1249 ± 43	WIDHALM	06 BELL	$e^+e^- \approx \Upsilon(4S)$	

$\Gamma(K^- \mu^+ \nu_\mu)/\Gamma(K^- \pi^+)$					Γ_{19}/Γ_{31}
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.853±0.033 OUR FIT					
0.84 ±0.04 OUR AVERAGE					
0.852±0.034±0.028	1897	46 FRABETTI	95G E687	γ Be $\bar{E}_\gamma = 220$ GeV	
0.82 ±0.13 ±0.13	338	47 FRABETTI	93I E687	γ Be $\bar{E}_\gamma = 221$ GeV	
0.79 ±0.08 ±0.09	231	48 CRAWFORD	91B CLEO	$e^+e^- \approx 10.5$ GeV	

⁴⁶FRABETTI 95G extracts the ratio of form factors $f_-(0)/f_+(0) = -1.3_{-3.4}^{+3.6} \pm 0.6$, and measures a pole mass of $1.87_{-0.08}^{+0.11+0.07}$ GeV/ c^2 from the q^2 dependence of the decay rate.

⁴⁷FRABETTI 93I measures a pole mass of $2.1_{-0.3}^{+0.7+0.7}$ GeV/ c^2 from the q^2 dependence of the decay rate.

⁴⁸CRAWFORD 91B measures a pole mass of $2.00 \pm 0.12 \pm 0.18$ GeV/ c^2 from the q^2 dependence of the decay rate.

$\Gamma(K^- \mu^+ \nu_\mu)/\Gamma(\mu^+ \text{anything})$					Γ_{19}/Γ_6
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.50 ±0.05 OUR FIT					
0.472±0.051±0.040	232	KODAMA	94 E653	π^- emulsion 600 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.32 ±0.05 ±0.05	124	KODAMA	91 EMUL	pA 800 GeV	

$\Gamma(K^- \pi^0 e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{22} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.016^{+0.013}_{-0.005} \pm 0.002$	4	49 BAI	91	MRK3 $e^+ e^- \approx 3.77$ GeV

⁴⁹BAI 91 finds that a fraction $0.79^{+0.15+0.09}_{-0.17-0.03}$ of combined D^+ and D^0 decays to $\bar{K} \pi e^+ \nu_e$ (24 events) are $\bar{K}^*(892) e^+ \nu_e$. BAI 91 uses 56 $K^- e^+ \nu_e$ events to measure a pole mass of $1.8 \pm 0.3 \pm 0.2$ GeV/ c^2 from the q^2 dependence of the decay rate.

$\Gamma(\bar{K}^0 \pi^- e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{23} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.7 $^{+0.9}_{-0.7}$ OUR AVERAGE				

2.61 $\pm 1.04 \pm 0.28$ 9 ± 3 ABLIKIM 06O BES2 $e^+ e^-$ at 3773 MeV

2.8 $^{+1.7}_{-0.8} \pm 0.3$ 6 50 BAI 91 MRK3 $e^+ e^- \approx 3.77$ GeV

⁵⁰BAI 91 finds that a fraction $0.79^{+0.15+0.09}_{-0.17-0.03}$ of combined D^+ and D^0 decays to $\bar{K} \pi e^+ \nu_e$ (24 events) are $\bar{K}^*(892) e^+ \nu_e$.

$\Gamma(K^*(892)^- e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{20} / Γ

Both decay modes of the $K^*(892)^-$ are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.16 ± 0.16 OUR FIT				

2.16 $\pm 0.15 \pm 0.08$ 219 ± 16 ⁵¹COAN 05 CLEO $e^+ e^-$ at $\psi(3770)$

⁵¹COAN 05 uses both $K^- \pi^0$ and $K_S^0 \pi^-$ events.

$\Gamma(K^*(892)^- e^+ \nu_e) / \Gamma(K_S^0 \pi^+ \pi^-)$ $\Gamma_{20} / \Gamma_{34}$

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.77 ± 0.06 OUR FIT				

0.76 $\pm 0.12 \pm 0.06$ 152 ⁵²BEAN 93C CLE2 $e^+ e^- \approx \Upsilon(4S)$

⁵²BEAN 93C uses $K^{*-} \mu^+ \nu_\mu$ as well as $K^{*-} e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

$\Gamma(K^*(892)^- \mu^+ \nu_\mu) / \Gamma(K_S^0 \pi^+ \pi^-)$ $\Gamma_{21} / \Gamma_{34}$

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.674 $\pm 0.068 \pm 0.026$	175 ± 17	⁵³ LINK	05B	FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV

⁵³LINK 05B finds that in $D^0 \rightarrow \bar{K}^0 \pi^- \mu^+ \nu_\mu$ the $\bar{K}^0 \pi^-$ system is 6% in S-wave.

$\Gamma(K^- \pi^+ \pi^- e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{24} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8 $^{+1.4}_{-1.1} \pm 0.3$	8	ARTUSO	07A	CLEO $e^+ e^-$ at $\Upsilon(3770)$

$\Gamma(K_1(1270)^- e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{25} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.6^{+4.1}_{-3.0} \pm 0.9$	8	⁵⁴ ARTUSO	07A	CLEO $e^+ e^-$ at $\Upsilon(3770)$

⁵⁴ This ARTUSO 07A result is corrected for all decay modes of the $K_1(1270)^-$.

$\Gamma(K^- \pi^+ \pi^- \mu^+ \nu_\mu) / \Gamma(K^- \mu^+ \nu_\mu)$ $\Gamma_{26} / \Gamma_{19}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.037	90	KODAMA	93B	E653 π^- emulsion 600 GeV

$\Gamma((\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu) / \Gamma(K^- \mu^+ \nu_\mu)$ $\Gamma_{27} / \Gamma_{19}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.043	90	⁵⁵ KODAMA	93B	E653 π^- emulsion 600 GeV

⁵⁵ KODAMA 93B searched in $K^- \pi^+ \pi^- \mu^+ \nu_\mu$, but the limit includes other $(\bar{K}^*(892)\pi)^-$ charge states.

$\Gamma(\pi^- e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{28} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.289 ± 0.008 OUR FIT Error includes scale factor of 1.1.

0.287 ± 0.008 OUR AVERAGE

0.288 ± 0.008 ± 0.003 1374 ⁵⁶ BESSON 09 CLEO $e^+ e^-$ at $\psi(3770)$

0.279 ± 0.027 ± 0.016 126 ± 12 ⁵⁷ WIDHALM 06 BELL $e^+ e^- \approx \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.299 ± 0.011 ± 0.009 ⁵⁸ DOBBS 08 CLEO See BESSON 09

0.262 ± 0.025 ± 0.008 117 ± 11 COAN 05 CLEO See DOBBS 08

⁵⁶ See the form-factor parameters near the end of this D^0 Listing.

⁵⁷ The $\pi^- e^+ \nu_e$ and $K^- e^+ \nu_e$ results of WIDHALM 06 give $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}|^2 = 0.042 \pm 0.003 \pm 0.003$.

⁵⁸ DOBBS 08 establishes $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}| = 0.188 \pm 0.008 \pm 0.002$ from the D^+ and D^0 decays to $\bar{K} e^+ \nu_e$ and $\pi e^+ \nu_e$.

$\Gamma(\pi^- e^+ \nu_e) / \Gamma(K^- e^+ \nu_e)$ $\Gamma_{28} / \Gamma_{18}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0814 ± 0.0025 OUR FIT Error includes scale factor of 1.1.

0.085 ± 0.007 OUR AVERAGE

0.082 ± 0.006 ± 0.005 ⁵⁹ HUANG 05 CLEO $e^+ e^- \approx \Upsilon(4S)$

0.101 ± 0.020 ± 0.003 91 ⁶⁰ FRABETTI 96B E687 γ Be, $\bar{E}_\gamma \approx 200$ GeV

0.103 ± 0.039 ± 0.013 87 ⁶¹ BUTLER 95 CLE2 < 0.156 (90% CL)

⁵⁹ HUANG 05 uses both e and μ events, and makes a small correction to the μ events to make them effectively e events. This result gives $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}|^2 = 0.038^{+0.006+0.005}_{-0.007-0.003}$.

⁶⁰ FRABETTI 96B uses both e and μ events, and makes a small correction to the μ events to make them effectively e events. This result gives $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}|^2 = 0.050 \pm 0.011 \pm 0.002$.

⁶¹ BUTLER 95 has $87 \pm 33 \pi^- e^+ \nu_e$ events. The result gives $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}|^2 = 0.052 \pm 0.020 \pm 0.007$.

$\Gamma(\pi^- \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.237 ± 0.024 OUR FIT				
0.231 ± 0.026 ± 0.019	106 ± 13	WIDHALM	06 BELL	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\pi^- \mu^+ \nu_\mu)/\Gamma(K^- \mu^+ \nu_\mu)$ Γ_{29}/Γ_{19}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.072 ± 0.007 OUR FIT				
0.074 ± 0.008 ± 0.007	288 ± 29	⁶² LINK	05 FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$

⁶² LINK 05 finds the form-factor ratio $|f_0^\pi(0)/f_0^K(0)|$ to be $0.85 \pm 0.04 \pm 0.04 \pm 0.01$.

$\Gamma(\rho^- e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.194 ± 0.039 ± 0.013	31 ± 6	COAN	05 CLEO	$e^+ e^-$ at $\psi(3770)$

————— Hadronic modes with a single \bar{K} —————

$\Gamma(K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.87 ± 0.05 OUR FIT				Error includes scale factor of 1.2.
3.91 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.1.

4.007 ± 0.037 ± 0.072	33.8 ± 0.3k	AUBERT	08L BABR	$e^+ e^-$ at $\Upsilon(4S)$
3.891 ± 0.035 ± 0.069		⁶³ DOBBS	07 CLEO	$e^+ e^-$ at $\psi(3770)$
3.82 ± 0.07 ± 0.12		⁶⁴ ARTUSO	98 CLE2	CLEO average
3.90 ± 0.09 ± 0.12	5392	⁶⁵ BARATE	97C ALEP	From Z decays
3.41 ± 0.12 ± 0.28	1173 ± 37	⁶⁵ ALBRECHT	94F ARG	$e^+ e^- \approx \Upsilon(4S)$
3.62 ± 0.34 ± 0.44		⁶⁵ DECAMP	91J ALEP	From Z decays

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.91 ± 0.08 ± 0.09	10.3k ± 100	⁶³ HE	05 CLEO	See DOBBS 07
3.81 ± 0.15 ± 0.16	1165	⁶⁶ ARTUSO	98 CLE2	$e^+ e^-$ at $\Upsilon(4S)$
3.69 ± 0.11 ± 0.16		⁶⁷ COAN	98 CLE2	See ARTUSO 98
4.5 ± 0.6 ± 0.4		⁶⁸ ALBRECHT	94 ARG	$e^+ e^- \approx \Upsilon(4S)$
3.95 ± 0.08 ± 0.17	4208	^{65,69} AKERIB	93 CLE2	See ARTUSO 98
4.5 ± 0.8 ± 0.5	56	⁶⁵ ABACHI	88 HRS	$e^+ e^-$ 29 GeV
4.2 ± 0.4 ± 0.4	930	ADLER	88C MRK3	$e^+ e^-$ 3.77 GeV
4.1 ± 0.6	263 ± 17	⁷⁰ SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV
4.3 ± 1.0	130	⁷¹ PERUZZI	77 LGW	$e^+ e^-$ 3.77 GeV

⁶³ DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

⁶⁴ This combines the CLEO results of ARTUSO 98, COAN 98, and AKERIB 93.

⁶⁵ ABACHI 88, DECAMP 91J, AKERIB 93, ALBRECHT 94F, and BARATE 97C use $D^*(2010)^+ \rightarrow D^0 \pi^+$ decays. The π^+ is both slow and of low p_T with respect to the event thrust axis or nearest jet ($\approx D^{*+}$ direction). The excess number of such π^+ 's over background gives the number of $D^*(2010)^+ \rightarrow D^0 \pi^+$ events, and the fraction with $D^0 \rightarrow K^- \pi^+$ gives the $D^0 \rightarrow K^- \pi^+$ branching fraction.

⁶⁶ ARTUSO 98, following ALBRECHT 94, uses D^0 mesons from $\bar{B}^0 \rightarrow D^{*+}(2010)^+ X \ell^- \bar{\nu}_\ell$ decays. Our average uses the CLEO average of this value with the values of COAN 98 and AKERIB 93.

⁶⁷ COAN 98 assumes that $\Gamma(B \rightarrow \bar{D} X \ell^+ \nu)/\Gamma(B \rightarrow X \ell^+ \nu) = 1.0 - 3|V_{ub}/V_{cb}|^2 - 0.010 \pm 0.005$, the last term accounting for $\bar{B} \rightarrow D_s^+ K X \ell^- \bar{\nu}$. COAN 98 is included in the CLEO average in ARTUSO 98.

⁶⁸ ALBRECHT 94 uses D^0 mesons from $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}_\ell$ decays. This is a different set of events than used by ALBRECHT 94F.

⁶⁹ This AKERIB 93 value includes radiative corrections; without them, the value is $0.0391 \pm 0.0008 \pm 0.0017$. AKERIB 93 is included in the CLEO average in ARTUSO 98.

⁷⁰ SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.24 ± 0.02 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

⁷¹ PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.25 ± 0.05 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

$\Gamma(K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1.240 \pm 0.017 \pm 0.056$	614	HE	08	CLEO See MENDEZ 10

$\Gamma(K_S^0 \pi^0)/\Gamma(K^- \pi^+)$ Γ_{32}/Γ_{31}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.68 \pm 0.12 \pm 0.11$	119	ANJOS	92B	E691 γ Be 80–240 GeV

$\Gamma(K_S^0 \pi^0)/[\Gamma(K^- \pi^+) + \Gamma(K^+ \pi^-)]$ $\Gamma_{32}/(\Gamma_{31} + \Gamma_{207})$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
30.7 ± 1.2 OUR FIT	Error includes scale factor of 1.3.			
$30.4 \pm 0.3 \pm 0.9$	20k	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV

$\Gamma(K_S^0 \pi^0)/\Gamma(K_S^0 \pi^+ \pi^-)$ Γ_{32}/Γ_{34}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.425 ± 0.026 OUR FIT Error includes scale factor of 1.1.				
$0.44 \pm 0.02 \pm 0.05$	1942 ± 64	PROCARIO	93B	CLE2 $e^+ e^-$ 10.36–10.7 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.34 \pm 0.04 \pm 0.02$	92	⁷² ALBRECHT	92P	ARG $e^+ e^- \approx 10$ GeV
$0.36 \pm 0.04 \pm 0.08$	104	KINOSHITA	91	CLEO $e^+ e^- \sim 10.7$ GeV

⁷² This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K_L^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.998 \pm 0.049 \pm 0.048$	1116	⁷³ HE	08	CLEO $e^+ e^-$ at $\psi(3770)$

⁷³ The difference of HE 08 $D^0 \rightarrow K_S^0 \pi^0$ and $K_L^0 \pi^0$ branching fractions over the sum is $0.108 \pm 0.025 \pm 0.024$. This is consistent with U-spin symmetry and the Cabibbo angle.

$\Gamma(K_S^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{34} / Γ

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

2.81 ± 0.15 OUR FIT Error includes scale factor of 1.1.

2.68 ± 0.29 OUR AVERAGE

2.52 ± 0.20 ± 0.25 284 ± 22 ⁷⁴ ALBRECHT 94F ARG $e^+ e^- \approx \Upsilon(4S)$

3.2 ± 0.3 ± 0.5 ADLER 87 MRK3 $e^+ e^-$ 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.8 32 ± 8 ⁷⁵ SCHINDLER 81 MRK2 $e^+ e^-$ 3.771 GeV

4.0 ± 1.2 28 ⁷⁶ PERUZZI 77 LGW $e^+ e^-$ 3.77 GeV

⁷⁴ See the footnote on the ALBRECHT 94F measurement of $\Gamma(K^- \pi^+) / \Gamma_{\text{total}}$ for the method used.

⁷⁵ SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.30 ± 0.08 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

⁷⁶ PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.46 ± 0.12 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

$\Gamma(K_S^0 \pi^+ \pi^-) / \Gamma(K^- \pi^+)$ $\Gamma_{34} / \Gamma_{31}$

VALUE EVTS DOCUMENT ID TECN COMMENT

0.73 ± 0.04 OUR FIT Error includes scale factor of 1.1.

0.81 ± 0.05 ± 0.08 856 ± 35 FRABETTI 94J E687 γ Be $\bar{E}_\gamma = 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.85 ± 0.40 35 AVERY 80 SPEC $\gamma N \rightarrow D^{*+}$

1.4 ± 0.5 116 PICCOLO 77 MRK1 $e^+ e^-$ 4.03, 4.41 GeV

$\Gamma(K_S^0 \rho^0) / \Gamma(K_S^0 \pi^+ \pi^-)$ $\Gamma_{35} / \Gamma_{34}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE DOCUMENT ID TECN COMMENT

0.224^{+0.017}_{-0.023} OUR AVERAGE Error includes scale factor of 1.7.

0.210 ± 0.016 ⁷⁷ AUBERT 08AL BABR Dalitz fit, ≈ 487 k evts

0.264 ± 0.009^{+0.010}_{-0.026} MURAMATSU 02 CLE2 Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.267 ± 0.011^{+0.009}_{-0.028} ASNER 04A CLEO See MURAMATSU 02

0.350 ± 0.028 ± 0.067 FRABETTI 94G E687 Dalitz fit, 597 evts

0.227 ± 0.032 ± 0.009 ALBRECHT 93D ARG Dalitz fit, 440 evts

0.215 ± 0.051 ± 0.037 ANJOS 93 E691 γ Be 90–260 GeV

0.20 ± 0.06 ± 0.03 FRABETTI 92B E687 γ Be, $\bar{E}_\gamma = 221$ GeV

0.12 ± 0.01 ± 0.07 ADLER 87 MRK3 $e^+ e^-$ 3.77 GeV

⁷⁷ The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$\Gamma(K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-) / \Gamma(K_S^0 \pi^+ \pi^-)$ $\Gamma_{36} / \Gamma_{34}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE DOCUMENT ID TECN COMMENT

0.0073 ± 0.0020 OUR AVERAGE

0.009 ± 0.010 ⁷⁸ AUBERT 08AL BABR Dalitz fit, ≈ 487 k evts

0.0072 ± 0.0018^{+0.0010}_{-0.0009} MURAMATSU 02 CLE2 Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0081 ± 0.0019^{+0.0018}_{-0.0010} ASNER 04A CLEO See MURAMATSU 02

⁷⁸The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$$\Gamma(K_S^0(\pi^+\pi^-)_{S\text{-wave}})/\Gamma(K_S^0\pi^+\pi^-) \quad \Gamma_{37}/\Gamma_{34}$$

This is the “fit fraction” from the Dalitz-plot analysis. The $(\pi^+\pi^-)_{S\text{-wave}}$ includes what in isobar models are the $f_0(980)$ and $f_0(1370)$; see the following two data blocks.

VALUE	DOCUMENT ID	TECN	COMMENT
0.119±0.026	⁷⁹ AUBERT	08AL BABR	Dalitz fit, ≈ 487 k evts

⁷⁹The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$$\Gamma(K_S^0 f_0(980), f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(K_S^0\pi^+\pi^-) \quad \Gamma_{38}/\Gamma_{34}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.043±0.005^{+0.012}_{-0.006}	MURAMATSU 02	CLE2	Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.042±0.005 ^{+0.011} _{-0.005}	ASNER	04A CLEO	See MURAMATSU 02
0.068±0.016±0.018	FRABETTI	94G E687	Dalitz fit, 597 evts
0.046±0.018±0.006	ALBRECHT	93D ARG	Dalitz fit, 440 evts

$$\Gamma(K_S^0 f_0(1370), f_0(1370) \rightarrow \pi^+\pi^-)/\Gamma(K_S^0\pi^+\pi^-) \quad \Gamma_{39}/\Gamma_{34}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.099±0.011^{+0.028}_{-0.044}	MURAMATSU 02	CLE2	Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.098±0.014 ^{+0.026} _{-0.036}	ASNER	04A CLEO	See MURAMATSU 02
0.077±0.022±0.031	FRABETTI	94G E687	Dalitz fit, 597 evts
0.082±0.028±0.013	ALBRECHT	93D ARG	Dalitz fit, 440 evts

$$\Gamma(K_S^0 f_2(1270), f_2(1270) \rightarrow \pi^+\pi^-)/\Gamma(K_S^0\pi^+\pi^-) \quad \Gamma_{40}/\Gamma_{34}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0032^{+0.0035}_{-0.0022} OUR AVERAGE			

0.006 ±0.007	⁸⁰ AUBERT	08AL BABR	Dalitz fit, ≈ 487 k evts
0.0027±0.0015 ^{+0.0037} _{-0.0017}	MURAMATSU 02	CLE2	Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0036±0.0022 ^{+0.0032} _{-0.0019}	ASNER	04A CLEO	See MURAMATSU 02
0.037 ±0.014 ±0.017	FRABETTI	94G E687	Dalitz fit, 597 evts
0.050 ±0.021 ±0.008	ALBRECHT	93D ARG	Dalitz fit, 440 evts

⁸⁰The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$\Gamma(K^*(892)^-\pi^+, K^*(892)^-\rightarrow K_S^0\pi^-)/\Gamma(K_S^0\pi^+\pi^-)$ Γ_{41}/Γ_{34}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.588^{+0.034}_{-0.050} OUR AVERAGE Error includes scale factor of 2.0.

0.557 ± 0.028	⁸¹ AUBERT	08AL BABR	Dalitz fit, ≈ 487 k evts
0.657 ± 0.013 ^{+0.018} _{-0.040}	MURAMATSU 02	CLE2	Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.663 ± 0.013 ^{+0.024} _{-0.043}	ASNER	04A CLEO	See MURAMATSU 02
0.625 ± 0.036 ± 0.026	FRABETTI	94G E687	Dalitz fit, 597 evts
0.718 ± 0.042 ± 0.030	ALBRECHT	93D ARG	Dalitz fit, 440 evts
0.480 ± 0.097	ANJOS	93 E691	γ Be 90–260 GeV
0.56 ± 0.04 ± 0.05	ADLER	87 MRK3	e ⁺ e ⁻ 3.77 GeV

⁸¹ The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$\Gamma(K_0^*(1430)^-\pi^+, K_0^*(1430)^-\rightarrow K_S^0\pi^-)/\Gamma(K_S^0\pi^+\pi^-)$ Γ_{42}/Γ_{34}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.095^{+0.014}_{-0.010} OUR AVERAGE

0.102 ± 0.015	⁸² AUBERT	08AL BABR	Dalitz fit, ≈ 487 k evts
0.073 ± 0.007 ^{+0.031} _{-0.011}	MURAMATSU 02	CLE2	Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.072 ± 0.007 ^{+0.014} _{-0.013}	ASNER	04A CLEO	See MURAMATSU 02
0.109 ± 0.027 ± 0.029	FRABETTI	94G E687	Dalitz fit, 597 evts
0.129 ± 0.034 ± 0.021	ALBRECHT	93D ARG	Dalitz fit, 440 evts

⁸² The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$\Gamma(K_2^*(1430)^-\pi^+, K_2^*(1430)^-\rightarrow K_S^0\pi^-)/\Gamma(K_S^0\pi^+\pi^-)$ Γ_{43}/Γ_{34}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0120^{+0.0070}_{-0.0035} OUR AVERAGE

0.022 ± 0.016	⁸³ AUBERT	08AL BABR	Dalitz fit, ≈ 487 k evts
0.011 ± 0.002 ^{+0.007} _{-0.003}	MURAMATSU 02	CLE2	Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.011 ± 0.002 ^{+0.005} _{-0.003}	ASNER	04A CLEO	See MURAMATSU 02
---	-------	----------	------------------

⁸³ The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$\Gamma(K^*(1680)^-\pi^+, K^*(1680)^-\pi^0)/\Gamma(K_S^0\pi^+\pi^-)$ Γ_{44}/Γ_{34}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.016±0.013 OUR AVERAGE

0.007±0.019 ⁸⁴ AUBERT 08AL BABR Dalitz fit, ≈ 487 k evts

0.022±0.004^{+0.018}_{-0.015} MURAMATSU 02 CLE2 Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.023±0.005^{+0.007}_{-0.014} ASNER 04A CLEO See MURAMATSU 02

⁸⁴ The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$\Gamma(K^*(892)^+\pi^-, K^*(892)^+\pi^0)/\Gamma(K_S^0\pi^+\pi^-)$ Γ_{45}/Γ_{34}

This is the "fit fraction" from the Dalitz-plot analysis. This is a doubly Cabibbo-suppressed mode.

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

4.0^{+2.0}_{-1.2} OUR AVERAGE

4.6±2.3 ⁸⁵ AUBERT 08AL BABR Dalitz fit, ≈ 487 k evts

3.4±1.3^{+4.1}_{-0.4} MURAMATSU 02 CLE2 Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.4±1.3^{+3.6}_{-0.5} ASNER 04A CLEO See MURAMATSU 02

⁸⁵ The error on this AUBERT 08AL value includes both statistical and systematic uncertainties; the latter dominates.

$\Gamma(K_0^*(1430)^+\pi^-, K_0^*(1430)^+\pi^0)/\Gamma(K_S^0\pi^+\pi^-)$ Γ_{46}/Γ_{34}

This is the "fit fraction" from the Dalitz-plot analysis. This is a doubly Cabibbo-suppressed mode.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<5 × 10⁻⁴ 95 AUBERT 08AL BABR Dalitz fit, ≈ 487 k evts

$\Gamma(K_2^*(1430)^+\pi^-, K_2^*(1430)^+\pi^0)/\Gamma(K_S^0\pi^+\pi^-)$ Γ_{47}/Γ_{34}

This is the "fit fraction" from the Dalitz-plot analysis. This is a doubly Cabibbo-suppressed mode.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.2 × 10⁻³ 95 AUBERT 08AL BABR Dalitz fit, ≈ 487 k evts

$\Gamma(K_S^0\pi^+\pi^- \text{ nonresonant})/\Gamma(K_S^0\pi^+\pi^-)$ Γ_{48}/Γ_{34}

This is the "fit fraction" from the Dalitz-plot analysis. Neither FRABETTI 94G nor ALBRECHT 93D (quoted in many of the earlier submodes of $K_S^0\pi^+\pi^-$) sees evidence for a nonresonant component.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.009±0.004^{+0.020}_{-0.004} MURAMATSU 02 CLE2 Dalitz fit, 5299 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.007±0.007^{+0.021}_{-0.006} ASNER 04A CLEO See MURAMATSU 02

0.263±0.024±0.041 ANJOS 93 E691 γ Be 90–260 GeV

0.26 ±0.08 ±0.05 FRABETTI 92B E687 γ Be, $\bar{E}_\gamma = 221$ GeV

0.33 ±0.05 ±0.10 ADLER 87 MRK3 e^+e^- 3.77 GeV

$\Gamma(K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$ **Γ_{49}/Γ**

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

13.9 ± 0.5 OUR FIT Error includes scale factor of 1.7.
14.57 ± 0.12 ± 0.38 ⁸⁶ DOBBS 07 CLEO $e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.9 ± 0.3 ± 0.5 19k ± 150 ⁸⁶ HE 05 CLEO See DOBBS 07
 13.3 ± 1.2 ± 1.3 931 ADLER 88C MRK3 $e^+ e^-$ 3.77 GeV
 11.7 ± 4.3 37 ⁸⁷ SCHINDLER 81 MRK2 $e^+ e^-$ 3.771 GeV

⁸⁶ DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

⁸⁷ SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.68 ± 0.23 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.


$\Gamma(K^- \pi^+ \pi^0)/\Gamma(K^- \pi^+)$ **Γ_{49}/Γ_{31}**

VALUE EVTS DOCUMENT ID TECN COMMENT

3.58 ± 0.14 OUR FIT Error includes scale factor of 1.9.
3.44 ± 0.30 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

3.81 ± 0.07 ± 0.26 10k BARISH 96 CLE2 $e^+ e^- \approx \Upsilon(4S)$
 3.04 ± 0.16 ± 0.34 931 ⁸⁸ ALBRECHT 92P ARG $e^+ e^- \approx 10$ GeV
 2.8 ± 0.14 ± 0.52 1050 KINOSHITA 91 CLEO $e^+ e^- \sim 10.7$ GeV

⁸⁸ This value is calculated from numbers in Table 1 of ALBRECHT 92P.



$\Gamma(K^- \rho^+)/\Gamma(K^- \pi^+ \pi^0)$ Γ_{50}/Γ_{49}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.78 ± 0.04 OUR AVERAGE			
0.788 ± 0.019 ± 0.048	KOPP	01 CLE2	Dalitz fit, ≈ 7,000 evts
0.765 ± 0.041 ± 0.054	FRABETTI	94G E687	Dalitz fit, 530 evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.647 ± 0.039 ± 0.150	ANJOS	93 E691	γBe 90–260 GeV
0.81 ± 0.03 ± 0.06	ADLER	87 MRK3	e ⁺ e ⁻ 3.77 GeV

$\Gamma(K^- \rho(1700)^+, \rho(1700)^+ \rightarrow \pi^+ \pi^0)/\Gamma(K^- \pi^+ \pi^0)$ Γ_{51}/Γ_{49}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.057 ± 0.008 ± 0.009	KOPP	01 CLE2	Dalitz fit, ≈ 7,000 evts

$\Gamma(K^*(892)^- \pi^+, K^*(892)^- \rightarrow K^- \pi^0)/\Gamma(K^- \pi^+ \pi^0)$ Γ_{52}/Γ_{49}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.160^{+0.025}_{-0.013} OUR AVERAGE			
0.161 ± 0.007 ^{+0.027} _{-0.011}	KOPP	01 CLE2	Dalitz fit, ≈ 7,000 evts
0.148 ± 0.028 ± 0.049	FRABETTI	94G E687	Dalitz fit, 530 evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.084 ± 0.011 ± 0.012	ANJOS	93 E691	γBe 90–260 GeV
0.12 ± 0.02 ± 0.03	ADLER	87 MRK3	e ⁺ e ⁻ 3.77 GeV

$\Gamma(\bar{K}^*(892)^0 \pi^0, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- \pi^+ \pi^0)$ Γ_{53}/Γ_{49}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.135 ± 0.016 OUR AVERAGE			
0.127 ± 0.009 ± 0.016	KOPP	01 CLE2	Dalitz fit, ≈ 7,000 evts
0.165 ± 0.031 ± 0.015	FRABETTI	94G E687	Dalitz fit, 530 evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.142 ± 0.018 ± 0.024	ANJOS	93 E691	γBe 90–260 GeV
0.13 ± 0.02 ± 0.03	ADLER	87 MRK3	e ⁺ e ⁻ 3.77 GeV

$\Gamma(K_0^*(1430)^- \pi^+, K_0^*(1430)^- \rightarrow K^- \pi^0)/\Gamma(K^- \pi^+ \pi^0)$ Γ_{54}/Γ_{49}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.033 ± 0.006 ± 0.014	KOPP	01 CLE2	Dalitz fit, ≈ 7,000 evts

$\Gamma(\bar{K}_0^*(1430)^0 \pi^0, \bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^- \pi^+ \pi^0)$ Γ_{55}/Γ_{49}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.041 ± 0.006^{+0.032}_{-0.009}	KOPP	01 CLE2	Dalitz fit, ≈ 7,000 evts

$\Gamma(K^*(1680)^- \pi^+, K^*(1680)^- \rightarrow K^- \pi^0)/\Gamma(K^- \pi^+ \pi^0)$ Γ_{56}/Γ_{49}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.013 ± 0.003 ± 0.004	KOPP	01 CLE2	Dalitz fit, ≈ 7,000 evts

$\Gamma(K^- \pi^+ \pi^0 \text{ nonresonant})/\Gamma(K^- \pi^+ \pi^0)$ Γ_{57}/Γ_{49}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.080^{+0.040}_{-0.014} OUR AVERAGE

0.075 ± 0.009 ^{+0.056} _{-0.011}		KOPP	01 CLE2	Dalitz fit, ≈ 7,000 evts
---	--	------	---------	--------------------------

0.101 ± 0.033 ± 0.040		FRABETTI	94G E687	Dalitz fit, 530 evts
-----------------------	--	----------	----------	----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.036 ± 0.004 ± 0.018		ANJOS	93 E691	γBe 90–260 GeV
-----------------------	--	-------	---------	----------------

0.09 ± 0.02 ± 0.04		ADLER	87 MRK3	e ⁺ e ⁻ 3.77 GeV
--------------------	--	-------	---------	--

0.51 ± 0.22	21	SUMMERS	84 E691	Photoproduction
-------------	----	---------	---------	-----------------

$\Gamma(K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE (units 10⁻³)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	--------------------	-------------	----------------

8.34 ± 0.45 ± 0.42	ASNER	08 CLEO	e ⁺ e ⁻ → D ⁰ \bar{D}^0 , 3.77 GeV
---------------------------	-------	---------	---

$\Gamma(\bar{K}^*(892)^0 \pi^0, \bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0)/\Gamma(K_S^0 \pi^0)$ Γ_{59}/Γ_{32}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.55^{+0.13}_{-0.10} ± 0.07	PROCARIO	93B CLE2	Dalitz plot fit, 122 evts
--	----------	----------	---------------------------

$\Gamma(K_S^0 2\pi^0 \text{ nonresonant})/\Gamma(K_S^0 \pi^0)$ Γ_{60}/Γ_{32}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.37 ± 0.08 ± 0.04	PROCARIO	93B CLE2	Dalitz plot fit, 122 evts
---------------------------	----------	----------	---------------------------

$\Gamma(K^- 2\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ

<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	-------------	--------------------	-------------	----------------

8.07^{+0.21}_{-0.19} OUR FIT Error includes scale factor of 1.4.

8.17 ± 0.33 OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.

8.30 ± 0.07 ± 0.20		⁸⁹ DOBBS	07 CLEO	e ⁺ e ⁻ at ψ(3770)
--------------------	--	---------------------	---------	--

7.9 ± 1.5 ± 0.9		⁹⁰ ALBRECHT	94 ARG	e ⁺ e ⁻ ≈ Υ(4S)
-----------------	--	------------------------	--------	---------------------------------------

6.80 ± 0.27 ± 0.57	1430 ± 52	⁹¹ ALBRECHT	94F ARG	e ⁺ e ⁻ ≈ Υ(4S)
--------------------	-----------	------------------------	---------	---------------------------------------

9.1 ± 0.8 ± 0.8	992	ADLER	88C MRK3	e ⁺ e ⁻ 3.77 GeV
-----------------	-----	-------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.3 ± 0.2 ± 0.3	15k ± 130	⁸⁹ HE	05 CLEO	See DOBBS 07
-----------------	-----------	------------------	---------	--------------

11.7 ± 2.5	185	⁹² SCHINDLER	81 MRK2	e ⁺ e ⁻ 3.771 GeV
------------	-----	-------------------------	---------	---

6.2 ± 1.9	44	⁹³ PERUZZI	77 LGW	e ⁺ e ⁻ 3.77 GeV
-----------	----	-----------------------	--------	--

⁸⁹ DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.


⁹⁰ ALBRECHT 94 uses D⁰ mesons from $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}_\ell$ decays. This is a different set of events than used by ALBRECHT 94F.

⁹¹ See the footnote on the ALBRECHT 94F measurement of $\Gamma(K^- \pi^+)/\Gamma_{\text{total}}$ for the method used.

⁹² SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.68 ± 0.11 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

⁹³ PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.36 ± 0.10 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

WEIGHTED AVERAGE
8.17±0.33 (Error scaled by 1.7)



$\Gamma(K^- 2\pi^+ \pi^-) / \Gamma(K^- \pi^+)$

$\Gamma_{61} / \Gamma_{31}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
2.08±0.05 OUR FIT				Error includes scale factor of 1.6.
1.97±0.09 OUR AVERAGE				
1.94±0.07 ^{+0.09} _{-0.11}		JUN 00	SELX	Σ^- nucleus, 600 GeV
1.7 ±0.2 ±0.2	1745	ANJOS 92C	E691	γ Be 90–260 GeV
1.90±0.25±0.20	337	ALVAREZ 91B	NA14	Photoproduction
2.12±0.16±0.09		BORTOLETTO88	CLEO	$e^+ e^-$ 10.55 GeV
2.17±0.28±0.23		ALBRECHT 85F	ARG	$e^+ e^-$ 10 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.0 ±0.9	48	BAILEY 86	ACCM	π^- Be fixed target
2.0 ±1.0	10	BAILEY 83B	SPEC	π^- Be $\rightarrow D^0$
2.2 ±0.8	214	PICCOLO 77	MRK1	$e^+ e^-$ 4.03, 4.41 GeV

$\Gamma(K^- \pi^+ \rho^0 \text{ total}) / \Gamma(K^- 2\pi^+ \pi^-)$

$\Gamma_{62} / \Gamma_{61}$

This includes $K^- a_1(1260)^+$, $\bar{K}^*(892)^0 \rho^0$, etc. The next entry gives the specifically 3-body fraction. We rely on the MARK III and E691 full amplitude analyses of the $K^- \pi^+ \pi^+ \pi^-$ channel for values of the resonant substructure.

VALUE	DOCUMENT ID	TECN	COMMENT
0.835±0.035 OUR AVERAGE			
0.80 ±0.03 ±0.05	ANJOS 92C	E691	1745 $K^- 2\pi^+ \pi^-$ evts
0.855±0.032±0.030	COFFMAN 92B	MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.98 ±0.12 ±0.10	ALVAREZ 91B	NA14	Photoproduction

$\Gamma(K^- \pi^+ \rho^0 \text{ 3-body})/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{63}/Γ_{61}

We rely on the MARK III and E691 full amplitude analyses of the $K^- \pi^+ \pi^+ \pi^-$ channel for values of the resonant substructure.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.063±0.028 OUR AVERAGE				
0.05 ±0.03 ±0.02		ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts
0.084±0.022±0.04		COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.77 ±0.06 ±0.06		⁹⁴ ALVAREZ	91B NA14	Photoproduction
0.85 ^{+0.11} _{-0.22}	180	PICCOLO	77 MRK1	$e^+ e^-$ 4.03, 4.41 GeV

⁹⁴This value is for ρ^0 ($K^- \pi^+$)-nonresonant. ALVAREZ 91B cannot determine what fraction of this is $K^- a_1(1260)^+$.

$\Gamma(\bar{K}^*(892)^0 \rho^0)/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{95}/Γ_{61}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included. We rely on the MARK III and E691 full amplitude analyses of the $K^- \pi^+ \pi^+ \pi^-$ channel for values of the resonant substructure.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.195±0.03±0.03				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.34 ±0.09±0.09		ALVAREZ	91B NA14	Photoproduction
0.75 ±0.3	5	BAILEY	83B SPEC	$\pi \text{Be} \rightarrow D^0$
0.15 ^{+0.16} _{-0.15}	20	PICCOLO	77 MRK1	$e^+ e^-$ 4.03, 4.41 GeV

$\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ transverse})/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{96}/Γ_{61}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.213±0.024±0.075	COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ S-wave})/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{97}/Γ_{61}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.375±0.045±0.06	ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ S-wave long.})/\Gamma_{\text{total}}$ Γ_{98}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.003	90	COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ P-wave})/\Gamma_{\text{total}}$ Γ_{99}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.003	90	COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.009	90	ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(\bar{K}^*(892)^0 \rho^0 D\text{-wave})/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{100}/Γ_{61}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.255 ± 0.045 ± 0.06		ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(K^- \pi^+ f_0(980))/\Gamma_{\text{total}}$ Γ_{101}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.011	90	ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts
--------	----	-------	----------	------------------------------

$\Gamma(\bar{K}^*(892)^0 f_0(980))/\Gamma_{\text{total}}$ Γ_{102}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.007	90	ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts
--------	----	-------	----------	------------------------------

$\Gamma(K^- a_1(1260)^+)/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{91}/Γ_{61}

Unseen decay modes of the $a_1(1260)^+$ are included, assuming that the $a_1(1260)^+$ decays entirely to $\rho\pi$ [or at least to $(\pi\pi)_{J=1} \pi$].

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

0.97 ± 0.14 OUR AVERAGE

0.94 ± 0.13 ± 0.20		ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts
0.984 ± 0.048 ± 0.16		COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(K^- a_2(1320)^+)/\Gamma_{\text{total}}$ Γ_{92}/Γ

Unseen decay modes of the $a_2(1320)^+$ are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.002 90 ANJOS 92C E691 1745 $K^- 2\pi^+ \pi^-$ evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.006	90	COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts
--------	----	---------	----------	-----------------------------------

$\Gamma(K_1(1270)^- \pi^+)/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{103}/Γ_{61}

Unseen decay modes of the $K_1(1270)^-$ are included. The MARK3 and E691 experiments disagree considerably here.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

0.194 ± 0.056 ± 0.088 COFFMAN 92B MRK3 1281 ± 45 $K^- 2\pi^+ \pi^-$ evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.013	90	ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts
--------	----	-------	----------	------------------------------

$\Gamma(K_1(1400)^- \pi^+)/\Gamma_{\text{total}}$ Γ_{104}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.012 90 COFFMAN 92B MRK3 1281 ± 45 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(K^*(1410)^- \pi^+)/\Gamma_{\text{total}}$ Γ_{105}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.012	90	COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts
--------	----	---------	----------	-----------------------------------

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^- \text{ total}) / \Gamma(K^- 2\pi^+ \pi^-)$ $\Gamma_{93} / \Gamma_{61}$

This includes $\bar{K}^*(892)^0 \rho^0$, etc. The next entry gives the specifically 3-body fraction.
Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.30 ± 0.06 ± 0.03	ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^- \text{ 3-body}) / \Gamma(K^- 2\pi^+ \pi^-)$ $\Gamma_{94} / \Gamma_{61}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.18 ± 0.04 OUR AVERAGE			
0.165 ± 0.03 ± 0.045	ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts
0.210 ± 0.027 ± 0.06	COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(K^- 2\pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^- 2\pi^+ \pi^-)$ $\Gamma_{69} / \Gamma_{61}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.233 ± 0.032 OUR AVERAGE			
0.23 ± 0.02 ± 0.03	ANJOS	92C E691	1745 $K^- 2\pi^+ \pi^-$ evts
0.242 ± 0.025 ± 0.06	COFFMAN	92B MRK3	1281 ± 45 $K^- 2\pi^+ \pi^-$ evts

$\Gamma(K_S^0 \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{70} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.2 ± 0.6 OUR FIT				
5.2 ± 1.1 ± 1.2	140	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.7 ^{+1.6} _{-1.7}		⁹⁵ BARLAG	92C ACCM	π^- Cu 230 GeV
-------------------------------------	--	----------------------	----------	--------------------

⁹⁵ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K_S^0 \pi^+ \pi^- \pi^0) / \Gamma(K_S^0 \pi^+ \pi^-)$ $\Gamma_{70} / \Gamma_{34}$

Branching fractions for submodes of this mode with narrow resonances (the η , ω , η') are fairly well determined (see below). COFFMAN 92B gives fractions of K^* and ρ submodes, but with only 140 ± 28 events above background could not determine them with much accuracy. We omit those measurements here; they are in our 2008 Review (Physics Letters **B667** 1 (2008)).

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.84 ± 0.20 OUR FIT				
1.86 ± 0.23 OUR AVERAGE				
1.80 ± 0.20 ± 0.21	190	⁹⁶ ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV
2.8 ± 0.8 ± 0.8	46	ANJOS	92C E691	γ Be 90–260 GeV
1.85 ± 0.26 ± 0.30	158	KINOSHITA	91 CLEO	$e^+ e^- \sim 10.7$ GeV

⁹⁶ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K_S^0 \eta) / \Gamma_{\text{total}}$ Γ_{88} / Γ

Unseen decay modes of the η are included.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.42 ± 0.15 ± 0.28	ASNER	08 CLEO	See MENDEZ 10

$$\Gamma(K_S^0 \eta) / [\Gamma(K^- \pi^+) + \Gamma(K^+ \pi^-)] \quad \Gamma_{88} / (\Gamma_{31} + \Gamma_{207})$$

Unseen decay modes of the η are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
11.6 ± 1.0 OUR FIT				Error includes scale factor of 1.5.
12.3 ± 0.3 ± 0.7	2864 ± 65	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV

$$\Gamma(K_S^0 \eta) / \Gamma(K_S^0 \pi^0) \quad \Gamma_{88} / \Gamma_{32}$$

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.379 ± 0.033 OUR FIT				Error includes scale factor of 1.4.
0.32 ± 0.04 ± 0.03	225 ± 30	PROCARIO	93B	CLE2 $\eta \rightarrow \gamma\gamma$

$$\Gamma(K_S^0 \eta) / \Gamma(K_S^0 \pi^+ \pi^-) \quad \Gamma_{88} / \Gamma_{34}$$

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.161 ± 0.014 OUR FIT				Error includes scale factor of 1.2.
0.14 ± 0.02 ± 0.02	80 ± 12	PROCARIO	93B	CLE2 $\eta \rightarrow \pi^+ \pi^- \pi^0$

$$\Gamma(K_S^0 \omega) / \Gamma_{\text{total}} \quad \Gamma_{89} / \Gamma$$

Unseen decay modes of the ω are included.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
1.11 ± 0.06 OUR FIT			
1.12 ± 0.04 ± 0.05	ASNER	08	CLEO $e^+ e^- \rightarrow D^0 \bar{D}^0$, 3.77 GeV

$$\Gamma(K_S^0 \omega) / \Gamma(K^- \pi^+) \quad \Gamma_{89} / \Gamma_{31}$$

Unseen decay modes of the ω are included.

VALUE	DOCUMENT ID	TECN	COMMENT
• • •			We do not use the following data for averages, fits, limits, etc. • • •
0.50 ± 0.18 ± 0.10	ALBRECHT	89D	ARG $e^+ e^-$ 10 GeV

$$\Gamma(K_S^0 \omega) / \Gamma(K_S^0 \pi^+ \pi^-) \quad \Gamma_{89} / \Gamma_{34}$$

Unseen decay modes of the ω are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.394 ± 0.030 OUR FIT				Error includes scale factor of 1.1.
0.33 ± 0.09 OUR AVERAGE				Error includes scale factor of 1.1.
0.29 ± 0.08 ± 0.05	16	⁹⁷ ALBRECHT	92P	ARG $e^+ e^- \approx 10$ GeV
0.54 ± 0.14 ± 0.16	40	KINOSHITA	91	CLEO $e^+ e^- \sim 10.7$ GeV

⁹⁷ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$$\Gamma(K_S^0 \omega) / \Gamma(K_S^0 \pi^+ \pi^- \pi^0) \quad \Gamma_{89} / \Gamma_{70}$$

Unseen decay modes of the ω are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.214 ± 0.026 OUR FIT			
0.220 ± 0.048 ± 0.0116	COFFMAN	92B	MRK3 1281 ± 45 $K^- 2\pi^+ \pi^-$ evts

$$\Gamma(K_S^0 \eta'(958)) / [\Gamma(K^- \pi^+) + \Gamma(K^+ \pi^-)] \quad \Gamma_{90} / (\Gamma_{31} + \Gamma_{207})$$

Unseen decay modes of the $\eta'(958)$ are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
24.1 ± 1.3 OUR FIT				
24.3 ± 0.8 ± 1.1	1321 ± 42	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV

$\Gamma(K_S^0 \eta'(958))/\Gamma(K_S^0 \pi^+ \pi^-)$ Γ_{90}/Γ_{34}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.332±0.023 OUR FIT

0.32 ±0.04 OUR AVERAGE

0.31 ±0.02 ±0.04	594	PROCARIO	93B CLE2	$\eta' \rightarrow \eta \pi^+ \pi^-, \rho^0 \gamma$
0.37 ±0.13 ±0.06	18	⁹⁸ ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV

⁹⁸ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^- \pi^+ 2\pi^0)/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.177±0.029		⁹⁹ BARLAG	92C ACCM	π^- Cu 230 GeV
0.149±0.037 ±0.030	24	¹⁰⁰ ADLER	88C MRK3	$e^+ e^-$ 3.77 GeV
0.209 ^{+0.074} _{-0.043} ±0.012	9	⁹⁹ AGUILAR-...	87F HYBR	$\pi p, p p$ 360, 400 GeV

⁹⁹ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction using topological normalization. They do not distinguish the presence of a third π^0 , and thus are not included in the average.

¹⁰⁰ ADLER 88C uses an absolute normalization method finding this decay channel opposite a detected $\bar{D}^0 \rightarrow K^+ \pi^-$ in pure $D\bar{D}$ events.

$\Gamma(K^- 2\pi^+ \pi^- \pi^0)/\Gamma(K^- \pi^+)$ Γ_{74}/Γ_{31}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

1.09±0.10 OUR FIT

0.98±0.11±0.11 225 ¹⁰¹ ALBRECHT 92P ARG $e^+ e^- \approx 10$ GeV

¹⁰¹ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^- 2\pi^+ \pi^- \pi^0)/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{74}/Γ_{61}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.52±0.05 OUR FIT

0.56±0.07 OUR AVERAGE

0.55±0.07 ^{+0.12} _{-0.09}	167	KINOSHITA	91 CLEO	$e^+ e^- \sim 10.7$ GeV
0.57±0.06 ±0.05	180	ANJOS	90D E691	Photoproduction

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0)/\Gamma(K^- 2\pi^+ \pi^- \pi^0)$ Γ_{106}/Γ_{74}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.45±0.15±0.15 ANJOS 90D E691 Photoproduction

$\Gamma(\bar{K}^*(892)^0 \eta)/\Gamma(K^- \pi^+)$ Γ_{107}/Γ_{31}

Unseen decay modes of the $\bar{K}^*(892)^0$ and η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.58±0.19 ^{+0.24} _{-0.28}	46	KINOSHITA	91 CLEO	$e^+ e^- \sim 10.7$ GeV
---	----	-----------	---------	-------------------------

$\Gamma(\bar{K}^*(892)^0 \eta) / \Gamma(K^- \pi^+ \pi^0)$ $\Gamma_{107} / \Gamma_{49}$

Unseen decay modes of the $\bar{K}^*(892)^0$ and η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13 ± 0.02 ± 0.03	214	PROCARIO	93B CLE2	$\bar{K}^{*0} \eta \rightarrow K^- \pi^+ / \gamma \gamma$
--------------------	-----	----------	----------	---

$\Gamma(K_S^0 \eta \pi^0) / \Gamma(K_S^0 \pi^0)$ $\Gamma_{78} / \Gamma_{32}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.46 ± 0.07 ± 0.06	155 ± 22	102 RUBIN	04 CLEO	$e^+ e^- \approx 10$ GeV
--------------------	----------	-----------	---------	--------------------------

¹⁰² The η here is detected in its $\gamma \gamma$ mode, but other η modes are included in the value given.

$\Gamma(K_S^0 a_0(980), a_0(980) \rightarrow \eta \pi^0) / \Gamma(K_S^0 \eta \pi^0)$ $\Gamma_{79} / \Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis, with interference.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.19 ± 0.09 ± 0.26	103 RUBIN	04 CLEO	Dalitz fit, 155 evts
--------------------	-----------	---------	----------------------

¹⁰³ In addition to $K_S^0 a_0(980)$ and $\bar{K}^*(892)^0 \eta$ modes, RUBIN 04 finds a fit fraction of $0.246 \pm 0.092 \pm 0.091$ for other, undetermined modes.

$\Gamma(\bar{K}^*(892)^0 \eta, \bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0) / \Gamma(K_S^0 \eta \pi^0)$ $\Gamma_{80} / \Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis, with interference.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.293 ± 0.062 ± 0.035	104 RUBIN	04 CLEO	Dalitz fit, 155 evts
-----------------------	-----------	---------	----------------------

¹⁰⁴ See the note on RUBIN 04 in the preceding data block.

$\Gamma(K^- \pi^+ \omega) / \Gamma(K^- \pi^+)$ $\Gamma_{108} / \Gamma_{31}$

Unseen decay modes of the ω are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.78 ± 0.12 ± 0.10	99	105 ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV
--------------------	----	--------------	---------	--------------------------

¹⁰⁵ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\bar{K}^*(892)^0 \omega) / \Gamma(K^- \pi^+)$ $\Gamma_{109} / \Gamma_{31}$

Unseen decay modes of the $\bar{K}^*(892)^0$ and ω are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.28 ± 0.11 ± 0.04	17	106 ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV
--------------------	----	--------------	---------	--------------------------

¹⁰⁶ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^- \pi^+ \eta'(958)) / \Gamma(K^- 2\pi^+ \pi^-)$ $\Gamma_{110} / \Gamma_{61}$

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.093 ± 0.014 ± 0.019	286	PROCARIO	93B CLE2	$\eta' \rightarrow \eta \pi^+ \pi^-, \rho^0 \gamma$
-----------------------	-----	----------	----------	---

$\Gamma(\bar{K}^*(892)^0 \eta'(958)) / \Gamma(K^- \pi^+ \eta'(958))$ $\Gamma_{111} / \Gamma_{110}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN
-------	-----	-------------	------

<0.15	90	PROCARIO	93B CLE2
-------	----	----------	----------

$\Gamma(K_S^0 2\pi^+ 2\pi^-)/\Gamma(K_S^0 \pi^+ \pi^-)$ Γ_{81}/Γ_{34}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.095±0.005±0.007	1283 ± 57	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 ± 0.02 ± 0.01	11	¹⁰⁷ ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV
0.149 ± 0.026	56	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV
0.18 ± 0.07 ± 0.04	6	ANJOS	90D E691	Photoproduction

¹⁰⁷ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K_S^0 \rho^0 \pi^+ \pi^-, \text{no } K^*(892)^-)/\Gamma(K_S^0 2\pi^+ 2\pi^-)$ Γ_{82}/Γ_{81}

VALUE	DOCUMENT ID	TECN	COMMENT
0.40±0.24±0.07	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^*(892)^- 2\pi^+ \pi^-, K^*(892)^- \rightarrow K_S^0 \pi^-, \text{no } \rho^0)/\Gamma(K_S^0 2\pi^+ 2\pi^-)$ Γ_{83}/Γ_{81}

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.28±0.02	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^*(892)^- \rho^0 \pi^+, K^*(892)^- \rightarrow K_S^0 \pi^-)/\Gamma(K_S^0 2\pi^+ 2\pi^-)$ Γ_{84}/Γ_{81}

VALUE	DOCUMENT ID	TECN	COMMENT
0.60±0.21±0.09	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K_S^0 2\pi^+ 2\pi^- \text{ nonresonant})/\Gamma(K_S^0 2\pi^+ 2\pi^-)$ Γ_{85}/Γ_{81}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.46	90	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^- 3\pi^+ 2\pi^-)/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{87}/Γ_{61}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.70±0.58±0.38	48 ± 10	LINK	04B FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

————— Hadronic modes with three *K*'s —————

$\Gamma(K_S^0 K^+ K^-)/\Gamma(K_S^0 \pi^+ \pi^-)$ Γ_{112}/Γ_{34}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.158±0.001±0.005	14k ± 116	AUBERT,B	05J BABR	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.20 ± 0.05 ± 0.04	47	FRABETTI	92B E687	γ Be, $\bar{E}_\gamma = 221$ GeV
0.170 ± 0.022	136	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV
0.24 ± 0.08		BEBEK	86 CLEO	$e^+ e^-$ near $\gamma(4S)$
0.185 ± 0.055	52	ALBRECHT	85B ARG	$e^+ e^- 10$ GeV

$\Gamma(K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-)/\Gamma(K_S^0 K^+ K^-)$ $\Gamma_{113}/\Gamma_{112}$

This is the "fit fraction" from the Dalitz-plot analysis, with interference.

VALUE	DOCUMENT ID	TECN	COMMENT
0.664±0.016±0.070	AUBERT,B	05J BABR	Dalitz fit, 12540 ± 112 evts

$\Gamma(K^- a_0(980)^+, a_0^+ \rightarrow K^+ K_S^0)/\Gamma(K_S^0 K^+ K^-)$ $\Gamma_{114}/\Gamma_{112}$

This is the "fit fraction" from the Dalitz-plot analysis, with interference.

VALUE	DOCUMENT ID	TECN	COMMENT
0.134±0.011±0.037	AUBERT,B	05J BABR	Dalitz fit, 12540 ± 112 evts

$$\Gamma(K^+ a_0(980)^-, a_0^- \rightarrow K^- K_S^0) / \Gamma(K_S^0 K^+ K^-) \quad \Gamma_{115} / \Gamma_{112}$$

This is a doubly Cabibbo-suppressed mode.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.025	95	AUBERT,B	05J BABR	Dalitz fit, 12540 ± 112 evts

$$\Gamma(K_S^0 f_0(980), f_0 \rightarrow K^+ K^-) / \Gamma(K_S^0 K^+ K^-) \quad \Gamma_{116} / \Gamma_{112}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.021	95	AUBERT,B	05J BABR	Dalitz fit, 12540 ± 112 evts

$$\Gamma(K_S^0 \phi, \phi \rightarrow K^+ K^-) / \Gamma(K_S^0 K^+ K^-) \quad \Gamma_{117} / \Gamma_{112}$$

This is the "fit fraction" from the Dalitz-plot analysis, with interference.

VALUE	DOCUMENT ID	TECN	COMMENT
0.459 ± 0.007 ± 0.007	AUBERT,B	05J BABR	Dalitz fit, 12540 ± 112 evts

$$\Gamma(K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-) / \Gamma(K_S^0 K^+ K^-) \quad \Gamma_{118} / \Gamma_{112}$$

This is the "fit fraction" from the Dalitz-plot analysis, with interference.

VALUE	DOCUMENT ID	TECN	COMMENT
0.038 ± 0.007 ± 0.023	¹⁰⁸ AUBERT,B	05J BABR	Dalitz fit, 12540 ± 112 evts

¹⁰⁸ AUBERT,B 05J calls the mode $K_S^0 f_0(1400)$, but insofar as it is seen here at all, it is certainly the same as $f_0(1370)$.

$$\Gamma(3K_S^0) / \Gamma(K_S^0 \pi^+ \pi^-) \quad \Gamma_{119} / \Gamma_{34}$$

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.4 OUR AVERAGE				
3.58 ± 0.54 ± 0.52	170 ± 26	LINK	05A FOCS	γ Be, $\bar{E}_\gamma \approx 180$ GeV
2.78 ± 0.38 ± 0.48	61	ASNER	96B CLE2	$e^+ e^- \approx \Upsilon(4S)$
7.0 ± 2.4 ± 1.2	10 ± 3	FRABETTI	94J E687	γ Be, $\bar{E}_\gamma = 220$ GeV
3.2 ± 1.0	22	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV
3.4 ± 1.4 ± 1.0	5	ALBRECHT	90C ARG	$e^+ e^- \approx 10$ GeV

$$\Gamma(K^+ 2K^- \pi^+) / \Gamma(K^- 2\pi^+ \pi^-) \quad \Gamma_{120} / \Gamma_{61}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0027 ± 0.0004 OUR AVERAGE				Error includes scale factor of 1.1.
0.00257 ± 0.00034 ± 0.00024	143	LINK	03G FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
0.0054 ± 0.0016 ± 0.0008	18	AITALA	01D E791	π^- A, 500 GeV
0.0028 ± 0.0007 ± 0.0001	20	FRABETTI	95C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(\phi \bar{K}^*(892)^0, \phi \rightarrow K^+ K^-, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^+ 2K^- \pi^+) \quad \Gamma_{123} / \Gamma_{120}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.48 ± 0.06 ± 0.01	LINK	03G FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$$\Gamma(K^- \pi^+ \phi, \phi \rightarrow K^+ K^-) / \Gamma(K^+ 2K^- \pi^+) \quad \Gamma_{122} / \Gamma_{120}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.18 ± 0.06 ± 0.04	LINK	03G FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ K^- \bar{K}^*(892)^0, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^+ 2K^- \pi^+)$ $\Gamma_{121} / \Gamma_{120}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.20 ± 0.07 ± 0.02	LINK	03G FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ 2K^- \pi^+ \text{ nonresonant}) / \Gamma(K^+ 2K^- \pi^+)$ $\Gamma_{124} / \Gamma_{120}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.15 ± 0.06 ± 0.02	LINK	03G FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(2K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 \pi^+ \pi^-)$ $\Gamma_{125} / \Gamma_{34}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.12 ± 0.38 ± 0.20	57 ± 10	LINK	05A FOCS	γ Be, $\bar{E}_\gamma \approx 180$ GeV

———— Pionic modes ————

$\Gamma(\pi^+ \pi^-) / \Gamma(K^- \pi^+)$ $\Gamma_{126} / \Gamma_{31}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.62 ± 0.05 OUR FIT				
3.59 ± 0.06 OUR AVERAGE				
3.594 ± 0.054 ± 0.040	7334 ± 97	ACOSTA	05C CDF	$p\bar{p}$, $\sqrt{s} = 1.96$ TeV
3.53 ± 0.12 ± 0.06	3453	LINK	03 FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
3.51 ± 0.16 ± 0.17	710	CSORNA	02 CLE2	$e^+ e^- \approx \Upsilon(4S)$
4.0 ± 0.2 ± 0.3	2043	AITALA	98C E791	π^- A, 500 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3.62 ± 0.10 ± 0.08	2085 ± 54	RUBIN	06 CLEO	See MENDEZ 10
3.4 ± 0.7 ± 0.1	76 ± 15	ABLIKIM	05F BES	$e^+ e^- \approx \psi(3770)$
4.3 ± 0.7 ± 0.3	177	FRABETTI	94C E687	γ Be $\bar{E}_\gamma = 220$ GeV
3.48 ± 0.30 ± 0.23	227	SELEN	93 CLE2	$e^+ e^- \approx \Upsilon(4S)$
5.5 ± 0.8 ± 0.5	120	ANJOS	91D E691	Photoproduction
5.0 ± 0.7 ± 0.5	110	ALEXANDER	90 CLEO	$e^+ e^-$ 10.5–11 GeV

$\Gamma(\pi^+ \pi^-) / [\Gamma(K^- \pi^+) + \Gamma(K^+ \pi^-)]$ $\Gamma_{126} / (\Gamma_{31} + \Gamma_{207})$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.60 ± 0.05 OUR FIT				
3.70 ± 0.06 ± 0.09	6210 ± 93	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV

$\Gamma(2\pi^0) / \Gamma(K^- \pi^+)$ $\Gamma_{127} / \Gamma_{31}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.05 ± 0.13 ± 0.16	499 ± 32	RUBIN	06 CLEO	See MENDEZ 10
2.2 ± 0.4 ± 0.4	40	SELEN	93 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(2\pi^0) / [\Gamma(K^- \pi^+) + \Gamma(K^+ \pi^-)]$ $\Gamma_{127} / (\Gamma_{31} + \Gamma_{207})$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.06 ± 0.12 OUR FIT				
2.06 ± 0.07 ± 0.10	1567 ± 54	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+)$ Γ_{128}/Γ_{31}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
37.0±1.6 OUR FIT				Error includes scale factor of 2.1.
34.4±0.5±1.2	11k±164	RUBIN	06	CLEO e^+e^- at $\psi(3770)$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+\pi^0)$ Γ_{128}/Γ_{49}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.34±0.24 OUR FIT				Error includes scale factor of 2.2.
10.41±0.23 OUR AVERAGE				Error includes scale factor of 2.0.
10.12±0.04±0.18	123k±490	ARINSTEIN	08	BELL $e^+e^- \approx \Upsilon(4S)$
10.59±0.06±0.13	60k±343	AUBERT,B	06X	BABR $e^+e^- \approx \Upsilon(4S)$

$\Gamma(\rho^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{129}/\Gamma_{128}$

This is the "fit fraction" from the Dalitz-plot analysis, with interference. See GASPERO 08 and BHATTACHARYA 10A for isospin decompositions of the $D^0 \rightarrow \pi^+\pi^0\pi^-$ Dalitz plot, both based on the amplitudes of AUBERT 07BJ. They quantify the conclusion that the final state is dominantly isospin 0.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
68.1±0.6 OUR AVERAGE			
67.8±0.0±0.6	AUBERT	07BJ	BABR Dalitz fit, 45k events
76.3±1.9±2.5	CRONIN-HEN..05	CLEO	$e^+e^- \approx 10$ GeV

$\Gamma(\rho^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{130}/\Gamma_{128}$

This is the "fit fraction" from the Dalitz-plot analysis, with interference.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
25.9±1.1 OUR AVERAGE			
26.2±0.5±1.1	AUBERT	07BJ	BABR Dalitz fit, 45k events
24.4±2.0±2.1	CRONIN-HEN..05	CLEO	$e^+e^- \approx 10$ GeV

$\Gamma(\rho^-\pi^+)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{131}/\Gamma_{128}$

This is the "fit fraction" from the Dalitz-plot analysis, with interference.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
34.6±0.8 OUR AVERAGE			
34.6±0.8±0.3	AUBERT	07BJ	BABR Dalitz fit, 45k events
34.5±2.4±1.3	CRONIN-HEN..05	CLEO	$e^+e^- \approx 10$ GeV

$\Gamma(\rho(1450)^+\pi^-, \rho(1450)^+ \rightarrow \pi^+\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{132}/\Gamma_{128}$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.11±0.07±0.12	AUBERT	07BJ	BABR Dalitz fit, 45k events

$\Gamma(\rho(1450)^0\pi^0, \rho(1450)^0 \rightarrow \pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{133}/\Gamma_{128}$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.30±0.11±0.07	AUBERT	07BJ	BABR Dalitz fit, 45k events

$\Gamma(\rho(1450)^-\pi^+, \rho(1450)^- \rightarrow \pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{134}/\Gamma_{128}$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.79±0.22±0.12	AUBERT	07BJ	BABR Dalitz fit, 45k events

$$\Gamma(\rho(1700)^+ \pi^-, \rho(1700)^+ \rightarrow \pi^+ \pi^0) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{135} / \Gamma_{128}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.1 ± 0.7 ± 0.7	AUBERT	07BJ BABR	Dalitz fit, 45k events

$$\Gamma(\rho(1700)^0 \pi^0, \rho(1700)^0 \rightarrow \pi^+ \pi^-) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{136} / \Gamma_{128}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.0 ± 0.6 ± 1.0	AUBERT	07BJ BABR	Dalitz fit, 45k events

$$\Gamma(\rho(1700)^- \pi^+, \rho(1700)^- \rightarrow \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{137} / \Gamma_{128}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.2 ± 0.4 ± 0.6	AUBERT	07BJ BABR	Dalitz fit, 45k events

$$\Gamma(f_0(980) \pi^0, f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{138} / \Gamma_{128}$$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.25 ± 0.04 ± 0.04		AUBERT	07BJ BABR	Dalitz fit, 45k events

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.026 95 ¹⁰⁹ CRONIN-HEN..05 CLEO $e^+ e^- \approx 10$ GeV

¹⁰⁹ The CRONIN-HENNESSY 05 fit here includes, in addition to the three $\rho\pi$ charged states, only the $f_0(980)\pi^0$ mode. See also the next entries for limits obtained in the same way for the $f_0(600)\pi^0$ mode and for an S -wave $\pi^+ \pi^-$ parametrized using a K -matrix. Our $\rho\pi$ branching ratios, given above, use the fit with the K -matrix S wave.

$$\Gamma(f_0(600) \pi^0, f_0(600) \rightarrow \pi^+ \pi^-) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{139} / \Gamma_{128}$$

The $f_0(600)$ is the σ .

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.82 ± 0.10 ± 0.10		AUBERT	07BJ BABR	Dalitz fit, 45k events

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.21 95 ¹¹⁰ CRONIN-HEN..05 CLEO $e^+ e^- \approx 10$ GeV

¹¹⁰ See the note on CRONIN-HENNESSY 05 in the proceeding data block.

$$\Gamma((\pi^+ \pi^-)_{S\text{-wave}} \pi^0) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{140} / \Gamma_{128}$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.019		95 ¹¹¹ CRONIN-HEN..05	CLEO	$e^+ e^- \approx 10$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.019 95 ¹¹¹ CRONIN-HEN..05 CLEO $e^+ e^- \approx 10$ GeV

¹¹¹ See the note on CRONIN-HENNESSY 05 two data blocks up.

$$\Gamma(f_0(1370) \pi^0, f_0(1370) \rightarrow \pi^+ \pi^-) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{141} / \Gamma_{128}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.37 ± 0.11 ± 0.09	AUBERT	07BJ BABR	Dalitz fit, 45k events

$$\Gamma(f_0(1500) \pi^0, f_0(1500) \rightarrow \pi^+ \pi^-) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{142} / \Gamma_{128}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.39 ± 0.08 ± 0.07	AUBERT	07BJ BABR	Dalitz fit, 45k events

$$\Gamma(f_0(1710) \pi^0, f_0(1710) \rightarrow \pi^+ \pi^-) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{143} / \Gamma_{128}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.31 ± 0.07 ± 0.08	AUBERT	07BJ BABR	Dalitz fit, 45k events

$\Gamma(f_2(1270)\pi^0, f_2(1270) \rightarrow \pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{144}/\Gamma_{128}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.32±0.08±0.10	AUBERT	07BJ BABR	Dalitz fit, 45k events

$\Gamma(\pi^+\pi^-\pi^0 \text{ nonresonant})/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{145}/\Gamma_{128}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
0.84±0.21±0.12	AUBERT	07BJ BABR	Dalitz fit, 45k events

$\Gamma(3\pi^0)/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.5 × 10⁻⁴	90	RUBIN	06 CLEO	e^+e^- at $\psi(3770)$

$\Gamma(2\pi^+2\pi^-)/\Gamma(K^-\pi^+)$ Γ_{147}/Γ_{31}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
19.1±0.5 OUR FIT	Error includes scale factor of 1.1.			
19.1±0.4±0.6	7331 ± 130	RUBIN	06 CLEO	e^+e^- at $\psi(3770)$

$\Gamma(2\pi^+2\pi^-)/\Gamma(K^-2\pi^+\pi^-)$ Γ_{147}/Γ_{61}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.19±0.23 OUR FIT	Error includes scale factor of 1.1.			
9.20±0.26 OUR AVERAGE				
9.14±0.18±0.22	6360 ± 115	LINK	07A FOCS	$\gamma\text{Be}, \bar{E}_\gamma \approx 180 \text{ GeV}$
7.9 ± 1.8 ± 0.5	162	ABLIKIM	05F BES	$e^+e^- \approx \psi(3770)$
9.5 ± 0.7 ± 0.2	814	FRABETTI	95C E687	$\gamma\text{Be}, \bar{E}_\gamma \approx 200 \text{ GeV}$
10.2 ± 1.3	345	AMMAR	91 CLEO	$e^+e^- \approx 10.5 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
11.5 ± 2.3 ± 1.6	64	ADAMOVICH	92 OMEG	π^- 340 GeV
10.8 ± 2.4 ± 0.8	79	FRABETTI	92 E687	γBe
9.6 ± 1.8 ± 0.7	66	ANJOS	91 E691	γBe 80–240 GeV

$\Gamma(a_1(1260)^+\pi^-, a_1^+ \rightarrow 2\pi^+\pi^- \text{ total})/\Gamma(2\pi^+2\pi^-)$ $\Gamma_{148}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
60.0±3.0±2.4	LINK	07A FOCS	4-body fit, $\approx 5.7\text{k}$ evts

$\Gamma(a_1(1260)^+\pi^-, a_1^+ \rightarrow \rho^0\pi^+ \text{ S-wave})/\Gamma(2\pi^+2\pi^-)$ $\Gamma_{149}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
43.3±2.5±1.9	LINK	07A FOCS	4-body fit, $\approx 5.7\text{k}$ evts

$\Gamma(a_1(1260)^+\pi^-, a_1^+ \rightarrow \rho^0\pi^+ \text{ D-wave})/\Gamma(2\pi^+2\pi^-)$ $\Gamma_{150}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
2.5±0.5±0.4	LINK	07A FOCS	4-body fit, $\approx 5.7\text{k}$ evts

$\Gamma(a_1(1260)^+\pi^-, a_1^+ \rightarrow \sigma\pi^+)/\Gamma(2\pi^+2\pi^-)$ $\Gamma_{151}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
8.3±0.7±0.6	LINK	07A FOCS	4-body fit, $\approx 5.7\text{k}$ evts

$\Gamma(2\rho^0 \text{ total})/\Gamma(2\pi^+ 2\pi^-)$ $\Gamma_{152}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
24.5±1.3±1.0	LINK	07A	FOCS 4-body fit, \approx 5.7k evts

$\Gamma(2\rho^0, \text{ parallel helicities})/\Gamma(2\pi^+ 2\pi^-)$ $\Gamma_{153}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.1±0.3±0.3	LINK	07A	FOCS 4-body fit, \approx 5.7k evts

$\Gamma(2\rho^0, \text{ perpendicular helicities})/\Gamma(2\pi^+ 2\pi^-)$ $\Gamma_{154}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.4±0.6±0.5	LINK	07A	FOCS 4-body fit, \approx 5.7k evts

$\Gamma(2\rho^0, \text{ longitudinal helicities})/\Gamma(2\pi^+ 2\pi^-)$ $\Gamma_{155}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16.8±1.0±0.8	LINK	07A	FOCS 4-body fit, \approx 5.7k evts

$\Gamma(\text{Resonant } (\pi^+ \pi^-) \pi^+ \pi^- \text{ 3-body total})/\Gamma(2\pi^+ 2\pi^-)$ $\Gamma_{156}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
20.0±1.2±1.0	LINK	07A	FOCS 4-body fit, \approx 5.7k evts

$\Gamma(\sigma \pi^+ \pi^-)/\Gamma(2\pi^+ 2\pi^-)$ $\Gamma_{157}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.2±0.9±0.7	LINK	07A	FOCS 4-body fit, \approx 5.7k evts

$\Gamma(f_0(980) \pi^+ \pi^-, f_0 \rightarrow \pi^+ \pi^-)/\Gamma(2\pi^+ 2\pi^-)$ $\Gamma_{158}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.4±0.5±0.4	LINK	07A	FOCS 4-body fit, \approx 5.7k evts

$\Gamma(f_2(1270) \pi^+ \pi^-, f_2 \rightarrow \pi^+ \pi^-)/\Gamma(2\pi^+ 2\pi^-)$ $\Gamma_{159}/\Gamma_{147}$

This is the fit fraction from the coherent amplitude analysis.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.9±0.6±0.5	LINK	07A	FOCS 4-body fit, \approx 5.7k evts

$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(K^- \pi^+)$ Γ_{160}/Γ_{31}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
25.8±1.5±1.8	2724 ± 166	RUBIN	06	CLEO $e^+ e^-$ at $\psi(3770)$

$\Gamma(\eta \pi^0)/\Gamma_{\text{total}}$ Γ_{161}/Γ

Unseen decay modes of the η are included.

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.4±1.0±0.4	156 ± 24	ARTUSO	08	CLEO See MENDEZ 10

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.4±1.0±0.4	156 ± 24	ARTUSO	08	CLEO See MENDEZ 10
-------------	----------	--------	----	--------------------

$\Gamma(\eta\pi^0)/\Gamma(K^-\pi^+)$ Γ_{161}/Γ_{31}

Unseen decay modes of the η are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.47 \pm 0.34 \pm 0.11$	62 ± 14	RUBIN	06	CLEO See ARTUSO 08

$\Gamma(\eta\pi^0)/[\Gamma(K^-\pi^+) + \Gamma(K^+\pi^-)]$ $\Gamma_{161}/(\Gamma_{31} + \Gamma_{207})$

Unseen decay modes of the η are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ± 0.19 OUR FIT				
$1.74 \pm 0.15 \pm 0.11$	481 ± 40	MENDEZ	10	CLEO e^+e^- at 3774 MeV

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{162}/Γ

Unseen decay modes of the ω are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.6 \times 10^{-4}$	90	RUBIN	06	CLEO e^+e^- at $\psi(3770)$

$\Gamma(2\pi^+2\pi^-\pi^0)/\Gamma(K^-\pi^+)$ Γ_{163}/Γ_{31}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$10.7 \pm 1.2 \pm 0.5$	1614 ± 171	RUBIN	06	CLEO e^+e^- at $\psi(3770)$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{164}/Γ

Unseen decay modes of the η are included.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$10.9 \pm 1.3 \pm 0.9$		257 ± 32	ARTUSO	08	CLEO e^+e^- at $\psi(3770)$
< 19	90		RUBIN	06	CLEO e^+e^- at $\psi(3770)$

$\Gamma(\omega\pi^+\pi^-)/\Gamma(K^-\pi^+)$ Γ_{165}/Γ_{31}

Unseen decay modes of the ω are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.1 \pm 1.2 \pm 0.4$	472 ± 132	RUBIN	06	CLEO e^+e^- at $\psi(3770)$

$\Gamma(3\pi^+3\pi^-)/\Gamma(K^-\pi^+)$ Γ_{166}/Γ_{61}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.23 \pm 0.59 \pm 1.35$	149 ± 17	LINK	04B	FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV

$\Gamma(3\pi^+3\pi^-)/\Gamma(K^-\pi^+2\pi^-)$ Γ_{166}/Γ_{87}

VALUE	DOCUMENT ID	TECN	COMMENT
$1.93 \pm 0.47 \pm 0.48$	¹¹² LINK	04B	FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV

¹¹² This LINK 04B result is not independent of other results in these Listings.

$\Gamma(\eta'(958)\pi^0)/\Gamma_{\text{total}}$ Γ_{167}/Γ

Unseen decay modes of the $\eta'(958)$ are included.

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.1 \pm 1.5 \pm 0.6$	50 ± 9	ARTUSO	08	CLEO See MENDEZ 10

$$\Gamma(\eta'(958)\pi^0)/[\Gamma(K^-\pi^+) + \Gamma(K^+\pi^-)] \quad \Gamma_{167}/(\Gamma_{31}+\Gamma_{207})$$

Unseen decay modes of the $\eta'(958)$ are included.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.3±0.4 OUR FIT				
2.3±0.3±0.2	159 ± 19	MENDEZ	10	CLEO e^+e^- at 3774 MeV

$$\Gamma(\eta'(958)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{168}/\Gamma$$

Unseen decay modes of the $\eta'(958)$ are included.

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.5±1.6±0.5	21 ± 8	ARTUSO	08	CLEO e^+e^- at $\psi(3770)$

$$\Gamma(2\eta)/\Gamma_{\text{total}} \quad \Gamma_{169}/\Gamma$$

Unseen decay modes of the η are included.

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16.7±1.4±1.3	255 ± 22	ARTUSO	08	CLEO See MENDEZ 10

• • • We do not use the following data for averages, fits, limits, etc. • • •

16.7±1.4±1.3 255 ± 22 ARTUSO 08 CLEO See MENDEZ 10

$$\Gamma(2\eta)/[\Gamma(K^-\pi^+) + \Gamma(K^+\pi^-)] \quad \Gamma_{169}/(\Gamma_{31}+\Gamma_{207})$$

Unseen decay modes of the η are included.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.3±0.5 OUR FIT				
4.3±0.3±0.4	430 ± 29	MENDEZ	10	CLEO e^+e^- at 3774 MeV

$$\Gamma(\eta\eta'(958))/\Gamma_{\text{total}} \quad \Gamma_{170}/\Gamma$$

Unseen decay modes of the η and $\eta'(958)$ are included.

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.6±2.5±1.1	46 ± 9	ARTUSO	08	CLEO See MENDEZ 10

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.6±2.5±1.1 46 ± 9 ARTUSO 08 CLEO See MENDEZ 10

$$\Gamma(\eta\eta'(958))/[\Gamma(K^-\pi^+) + \Gamma(K^+\pi^-)] \quad \Gamma_{170}/(\Gamma_{31}+\Gamma_{207})$$

Unseen decay modes of the η and $\eta'(958)$ are included.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.7±0.7 OUR FIT				
2.7±0.6±0.3	66 ± 15	MENDEZ	10	CLEO e^+e^- at 3774 MeV

————— **Hadronic modes with a $K\bar{K}$ pair** —————

$$\Gamma(K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_{171}/\Gamma$$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.96±0.08 OUR FIT	Error includes scale factor of 1.4.			
4.08±0.08±0.09	4746 ± 74	BONVICINI	08	CLEO See MENDEZ 10

• • • We do not use the following data for averages, fits, limits, etc. • • •


4.08±0.08±0.09 4746 ± 74 BONVICINI 08 CLEO See MENDEZ 10

$\Gamma(K^+ K^-)/\Gamma(K^- \pi^+)$

Γ_{171}/Γ_{31}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1021±0.0015 OUR FIT	Error includes scale factor of 1.7.			
0.1010±0.0016 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			
0.122 ±0.011 ±0.004	242 ± 20	ABLIKIM	05F BES	$e^+ e^- \approx \psi(3770)$
0.0992±0.0011±0.0012	16k±200	ACOSTA	05C CDF	$p\bar{p}, \sqrt{s}=1.96$ TeV
0.0993±0.0014±0.0014	11k	LINK	03 FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.1040±0.0033±0.0027	1900	CSORNA	02 CLE2	$e^+ e^- \approx \Upsilon(4S)$
0.109 ±0.003 ±0.003	3317	AITALA	98C E791	π^- nucleus, 500 GeV
0.116 ±0.007 ±0.007	1102	ASNER	96B CLE2	$e^+ e^- \approx \Upsilon(4S)$
0.109 ±0.007 ±0.009	581	FRABETTI	94C E687	γ Be $\bar{E}_\gamma=220$ GeV
0.107 ±0.010 ±0.009	193	ANJOS	91D E691	Photoproduction
0.117 ±0.010 ±0.007	249	ALEXANDER	90 CLEO	$e^+ e^-$ 10.5–11 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.107 ±0.029 ±0.015	103	ADAMOVICH	92 OMEG	π^- 340 GeV
0.138 ±0.027 ±0.010	155	FRABETTI	92 E687	γ Be
0.16 ±0.05	34	ALVAREZ	91B NA14	Photoproduction
0.10 ±0.02 ±0.01	131	ALBRECHT	90C ARG	$e^+ e^- \approx 10$ GeV
0.122 ±0.018 ±0.012	118	BALTRUSAIT	85E MRK3	$e^+ e^-$ 3.77 GeV
0.113 ±0.030		ABRAMS	79D MRK2	$e^+ e^-$ 3.77 GeV

WEIGHTED AVERAGE
0.1010±0.0016 (Error scaled by 1.4)



$\Gamma(K^+ K^-) / [\Gamma(K^- \pi^+) + \Gamma(K^+ \pi^-)]$ $\Gamma_{171} / (\Gamma_{31} + \Gamma_{207})$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
10.18 ± 0.15 OUR FIT	Error includes scale factor of 1.7.			
10.41 ± 0.11 ± 0.12	13.8k	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV

$\Gamma(K^+ K^-) / \Gamma(\pi^+ \pi^-)$ $\Gamma_{171} / \Gamma_{126}$

The unused results here are redundant with $\Gamma(K^+ K^-) / \Gamma(K^- \pi^+)$ and $\Gamma(\pi^+ \pi^-) / \Gamma(K^- \pi^+)$ measurements by the same experiments.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.760 ± 0.040 ± 0.034	7334	ACOSTA	05C	CDF $p\bar{p}$, $\sqrt{s}=1.96$ TeV
2.81 ± 0.10 ± 0.06		LINK	03	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
2.96 ± 0.16 ± 0.15	710	CSORNA	02	CLE2 $e^+ e^- \approx \Upsilon(4S)$
2.75 ± 0.15 ± 0.16		AITALA	98C	E791 π^- nucleus, 500 GeV
2.53 ± 0.46 ± 0.19		FRABETTI	94C	E687 γ Be $\bar{E}_\gamma = 220$ GeV
2.23 ± 0.81 ± 0.46		ADAMOVICH	92	OMEG π^- 340 GeV
1.95 ± 0.34 ± 0.22		ANJOS	91D	E691 Photoproduction
2.5 ± 0.7		ALBRECHT	90C	ARG $e^+ e^- \approx 10$ GeV
2.35 ± 0.37 ± 0.28		ALEXANDER	90	CLEO $e^+ e^-$ 10.5–11 GeV

$\Gamma(2K_S^0) / \Gamma_{\text{total}}$ Γ_{172} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.46 ± 0.32 ± 0.09	68 ± 15	BONVICINI	08	CLEO See MENDEZ 10

$\Gamma(2K_S^0) / [\Gamma(K^- \pi^+) + \Gamma(K^+ \pi^-)]$ $\Gamma_{172} / (\Gamma_{31} + \Gamma_{207})$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.07 OUR FIT	Error includes scale factor of 1.7.			
0.41 ± 0.04 ± 0.02	215 ± 23	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV

$\Gamma(2K_S^0) / \Gamma(K_S^0 \pi^+ \pi^-)$ $\Gamma_{172} / \Gamma_{34}$

This is the same as $\Gamma(K^0 \bar{K}^0) / \Gamma(\bar{K}^0 \pi^+ \pi^-)$ because $D^0 \rightarrow K_S^0 K_L^0$ is forbidden by CP conservation.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0062 ± 0.0011 OUR FIT	Error includes scale factor of 1.6.			
0.0120 ± 0.0022 OUR AVERAGE				
0.0144 ± 0.0032 ± 0.0016	79 ± 17	LINK	05A	FOCS γ Be, $\bar{E}_\gamma \approx 180$ GeV
0.0101 ± 0.0022 ± 0.0016	26	ASNER	96B	CLE2 $e^+ e^- \approx \Upsilon(4S)$
0.039 ± 0.013 ± 0.013	20 ± 7	FRABETTI	94J	E687 γ Be $\bar{E}_\gamma = 220$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.021 $^{+0.011}_{-0.008}$ ± 0.002	5	ALEXANDER	90	CLEO $e^+ e^-$ 10.5–11 GeV

$\Gamma(K_S^0 K^- \pi^+) / \Gamma(K^- \pi^+)$ $\Gamma_{173} / \Gamma_{31}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.086 ± 0.013 OUR FIT	Error includes scale factor of 1.1.		
0.08 ± 0.03	¹¹³ ANJOS	91	E691 γ Be 80–240 GeV
¹¹³ The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.			

$\Gamma(K_S^0 K^- \pi^+)/\Gamma(K_S^0 \pi^+ \pi^-)$ Γ_{173}/Γ_{34}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.118±0.017 OUR FIT				Error includes scale factor of 1.1.
0.119±0.021 OUR AVERAGE				Error includes scale factor of 1.3.
0.108±0.019	61	AMMAR	91	CLEO $e^+ e^- \approx 10.5$ GeV
0.16 ±0.03 ±0.02	39	ALBRECHT	90C	ARG $e^+ e^- \approx 10$ GeV

$\Gamma(\bar{K}^*(892)^0 K_S^0, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K_S^0 \pi^+ \pi^-)$ Γ_{174}/Γ_{34}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.019	90	AMMAR	91	CLEO $e^+ e^- \approx 10.5$ GeV
• • •				We do not use the following data for averages, fits, limits, etc. • • •
<0.02	90	ALBRECHT	90C	ARG $e^+ e^- \approx 10$ GeV

$\Gamma(K_S^0 K^+ \pi^-)/\Gamma(K^- \pi^+)$ Γ_{175}/Γ_{31}

VALUE	DOCUMENT ID	TECN	COMMENT
0.066±0.013 OUR FIT			
0.05 ±0.025	¹¹⁴ ANJOS	91	E691 γ Be 80–240 GeV

¹¹⁴ The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(K_S^0 K^+ \pi^-)/\Gamma(K_S^0 \pi^+ \pi^-)$ Γ_{175}/Γ_{34}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.091±0.017 OUR FIT				
0.098±0.020	55	AMMAR	91	CLEO $e^+ e^- \approx 10.5$ GeV

$\Gamma(K^*(892)^0 K_S^0, K^*(892)^0 \rightarrow K^+ \pi^-)/\Gamma(K_S^0 \pi^+ \pi^-)$ Γ_{176}/Γ_{34}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	90	AMMAR	91	CLEO $e^+ e^- \approx 10.5$ GeV

$\Gamma(K^+ K^- \pi^0)/\Gamma(K^- \pi^+ \pi^0)$ Γ_{177}/Γ_{49}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.37±0.03±0.04	11k±122	AUBERT,B	06X BABR	$e^+ e^- \approx \Upsilon(4S)$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
0.95±0.26	151	ASNER	96B CLE2	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(K^*(892)^+ K^-, K^*(892)^+ \rightarrow K^+ \pi^0)/\Gamma(K^+ K^- \pi^0)$ $\Gamma_{178}/\Gamma_{177}$

This is the “fit fraction” from the Dalitz-plot analysis with interference.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
44.4±0.8±0.6	AUBERT	07T	BABR Dalitz fit II, 11k evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

46.1±3.1	¹¹⁵ CAWLFIELD	06A	CLEO Dalitz fit, 627 ± 30 evts
----------	--------------------------	-----	--------------------------------

¹¹⁵ The error on this CAWLFIELD 06A result is statistical only.

$$\Gamma(K^*(892)^- K^+),$$

$$K^*(892)^- \rightarrow K^- \pi^0 / \Gamma(K^+ K^- \pi^0)$$

 $\Gamma_{179} / \Gamma_{177}$

This is the “fit fraction” from the Dalitz-plot analysis with interference.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
15.9 ± 0.7 ± 0.6	AUBERT	07T	BABR Dalitz fit II, 11k evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.3 ± 2.2 ¹¹⁶ CAWLFIELD 06A CLEO Dalitz fit, 627 ± 30 evts

¹¹⁶ The error on this CAWLFIELD 06A result is statistical only.

$$\Gamma((K^+ \pi^0)_{S\text{-wave}} K^-) / \Gamma(K^+ K^- \pi^0)$$

 $\Gamma_{180} / \Gamma_{177}$

This is the “fit fraction” from the Dalitz-plot analysis with interference.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
71.1 ± 3.7 ± 1.9	¹¹⁷ AUBERT	07T	BABR Dalitz fit II, 11k evts

¹¹⁷ The only major difference between fits I and II in the AUBERT 07T analysis is in this mode, where the fit-I fraction is $(16.3 \pm 3.4 \pm 2.1)\%$.

$$\Gamma((K^- \pi^0)_{S\text{-wave}} K^+) / \Gamma(K^+ K^- \pi^0)$$

 $\Gamma_{181} / \Gamma_{177}$

This is the “fit fraction” from the Dalitz-plot analysis with interference.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
3.9 ± 0.9 ± 1.0	AUBERT	07T	BABR Dalitz fit II, 11k evts

$$\Gamma(f_0(980) \pi^0, f_0 \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^0)$$

 $\Gamma_{182} / \Gamma_{177}$

This is the “fit fraction” from the Dalitz-plot analysis with interference.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
10.5 ± 1.1 ± 1.2	¹¹⁸ AUBERT	07T	BABR Dalitz fit II, 11k evts

¹¹⁸ When AUBERT 07T replace the $f_0(980) \pi^0$ mode with $a_0(980) \pi^0$, the fit fraction is a negligibly different $(11.0 \pm 1.5 \pm 1.2)\%$.

$$\Gamma(\phi \pi^0, \phi \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^0)$$

 $\Gamma_{183} / \Gamma_{177}$

This is the “fit fraction” from the Dalitz-plot analysis with interference.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
19.4 ± 0.6 ± 0.5	AUBERT	07T	BABR Dalitz fit II, 11k evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.9 ± 1.6 ¹¹⁹ CAWLFIELD 06A CLEO Dalitz fit, 627 ± 30 evts

¹¹⁹ The error on this CAWLFIELD 06A result is statistical only.

$$\Gamma(K^+ K^- \pi^0 \text{ nonresonant}) / \Gamma(K^+ K^- \pi^0)$$

 $\Gamma_{184} / \Gamma_{177}$

This is the “fit fraction” from the Dalitz-plot analysis with interference.

VALUE	DOCUMENT ID	TECN	COMMENT
0.360 ± 0.037	¹²⁰ CAWLFIELD	06A	CLEO Dalitz fit, 627 ± 30 evts

¹²⁰ The error is statistical only. CAWLFIELD 06A also fits the Dalitz plot replacing this flat nonresonant background with broad S -wave $\kappa^\pm \rightarrow K^\pm \pi^0$ resonances. There is no significant improvement in the fit, and $K^{*\pm} K^\mp$ and $\phi \pi^0$ results are not much changed.

$$\Gamma(2K_S^0 \pi^0) / \Gamma_{\text{total}}$$

 Γ_{185} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
< 0.00059	ASNER	96B	CLE2 $e^+ e^- \approx \gamma(4S)$

$\Gamma(\phi\pi^0)/\Gamma(K^+K^-)$ $\Gamma_{198}/\Gamma_{171}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.194 \pm 0.006 \pm 0.009$	1254	TAJIMA	04	BELL e^+e^- at $\Upsilon(4S)$

$\Gamma(\phi\eta)/\Gamma(K^+K^-)$ $\Gamma_{199}/\Gamma_{171}$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.59 \pm 1.14 \pm 0.18$	31	TAJIMA	04	BELL e^+e^- at $\Upsilon(4S)$

$\Gamma(\phi\omega)/\Gamma_{\text{total}}$ Γ_{200}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0021	90	ALBRECHT	94I	ARG $e^+e^- \approx 10$ GeV

$\Gamma(K^+K^-\pi^+\pi^-)/\Gamma(K^-2\pi^+\pi^-)$ Γ_{186}/Γ_{61}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.00 ± 0.13 OUR AVERAGE				
$2.95 \pm 0.11 \pm 0.08$	2669 ± 101	¹²¹ LINK	05G FOCS	γBe , $\bar{E}_\gamma \approx 180$ GeV
$3.13 \pm 0.37 \pm 0.36$	136 ± 15	AITALA	98D E791	π^- nucleus, 500 GeV
$3.5 \pm 0.4 \pm 0.2$	244 ± 26	FRABETTI	95C E687	γBe , $\bar{E}_\gamma \approx 200$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.4 \pm 1.8 \pm 0.5$	19 ± 8	ABLIKIM	05F BES	$e^+e^- \approx \psi(3770)$
$4.1 \pm 0.7 \pm 0.5$	114 ± 20	ALBRECHT	94I ARG	$e^+e^- \approx 10$ GeV
3.14 ± 1.0	89 ± 29	AMMAR	91 CLEO	$e^+e^- \approx 10.5$ GeV
$2.8^{+0.8}_{-0.7}$		ANJOS	91 E691	γBe 80–240 GeV

¹²¹LINK 05G uses a smaller, cleaner subset of 1279 ± 48 events for the amplitude analysis that gives the results in the next data blocks.

$\Gamma(\phi\pi^+\pi^- \text{ 3-body}, \phi \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+\pi^-)$ $\Gamma_{187}/\Gamma_{186}$

This is the fraction from a coherent amplitude analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.01 ± 0.01	LINK	05G FOCS	1279 ± 48 $K^+K^-\pi^+\pi^-$ evts.

$\Gamma(\phi\rho^0, \phi \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+\pi^-)$ $\Gamma_{188}/\Gamma_{186}$

This is the fraction from a coherent amplitude analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.29 \pm 0.02 \pm 0.01$	LINK	05G FOCS	1279 ± 48 $K^+K^-\pi^+\pi^-$ evts.

$\Gamma(K^+K^-\rho^0 \text{ 3-body})/\Gamma(K^+K^-\pi^+\pi^-)$ $\Gamma_{189}/\Gamma_{186}$

This is the fraction from a coherent amplitude analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.02 \pm 0.02 \pm 0.02$	LINK	05G FOCS	1279 ± 48 $K^+K^-\pi^+\pi^-$ evts.

$\Gamma(f_0(980)\pi^+\pi^-, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+\pi^-)$ $\Gamma_{190}/\Gamma_{186}$

This is the fraction from a coherent amplitude analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.15 \pm 0.03 \pm 0.02$	LINK	05G FOCS	1279 ± 48 $K^+K^-\pi^+\pi^-$ evts.

$$\Gamma(K^*(892)^0 K^\mp \pi^\pm \text{ 3-body}, K^{*0} \rightarrow K^\pm \pi^\mp) / \Gamma(K^+ K^- \pi^+ \pi^-) \quad \Gamma_{191} / \Gamma_{186}$$

This is the fraction from a coherent amplitude analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.11 ± 0.02 ± 0.01	LINK	05G FOCS	1279 ± 48 $K^+ K^- \pi^+ \pi^-$ evts.

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0, K^{*0} \rightarrow K^\pm \pi^\mp) / \Gamma(K^+ K^- \pi^+ \pi^-) \quad \Gamma_{192} / \Gamma_{186}$$

This is the fraction from a coherent amplitude analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.03 ± 0.02 ± 0.01	LINK	05G FOCS	1279 ± 48 $K^+ K^- \pi^+ \pi^-$ evts.

$$\Gamma(K_1(1270)^\pm K^\mp, K_1(1270)^\pm \rightarrow K^\pm \pi^+ \pi^-) / \Gamma(K^+ K^- \pi^+ \pi^-) \quad \Gamma_{193} / \Gamma_{186}$$

This is the fraction from a coherent amplitude analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.33 ± 0.06 ± 0.04	122 LINK	05G FOCS	1279 ± 48 $K^+ K^- \pi^+ \pi^-$ evts.

122 This LINK 05G value includes $K_1(1270)^\pm \rightarrow \rho^0 K^\pm, \rightarrow K_0^*(1430)^0 \pi^\pm$, and $K^*(892)^0 \pi^\pm$.

$$\Gamma(K_1(1400)^\pm K^\mp, K_1(1400)^\pm \rightarrow K^\pm \pi^+ \pi^-) / \Gamma(K^+ K^- \pi^+ \pi^-) \quad \Gamma_{194} / \Gamma_{186}$$

This is the fraction from a coherent amplitude analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.22 ± 0.03 ± 0.04	LINK	05G FOCS	1279 ± 48 $K^+ K^- \pi^+ \pi^-$ evts.

$$\Gamma(2K_S^0 \pi^+ \pi^-) / \Gamma(K_S^0 \pi^+ \pi^-) \quad \Gamma_{195} / \Gamma_{34}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3 ± 0.8 OUR AVERAGE				
4.16 ± 0.70 ± 0.42	113 ± 21	LINK	05A FOCS	γ Be, $\bar{E}_\gamma \approx 180$ GeV
6.2 ± 2.0 ± 1.6	25	ALBRECHT	94I ARG	$e^+ e^- \approx 10$ GeV

$$\Gamma(K_S^0 K^- 2\pi^+ \pi^-) / \Gamma(K_S^0 2\pi^+ 2\pi^-) \quad \Gamma_{196} / \Gamma_{81}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.054	90	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{197} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0031 ± 0.0020	123 BARLAG	92C ACCM	π^- Cu 230 GeV

123 BARLAG 92C computes the branching fraction using topological normalization.

Radiative modes

$$\Gamma(\rho^0 \gamma) / \Gamma_{\text{total}} \quad \Gamma_{201} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN
<2.4 × 10⁻⁴	90	ASNER	98 CLE2

$$\Gamma(\omega \gamma) / \Gamma_{\text{total}} \quad \Gamma_{202} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN
<2.4 × 10⁻⁴	90	ASNER	98 CLE2

$\Gamma(\phi\gamma)/\Gamma(K^+K^-)$						$\Gamma_{203}/\Gamma_{171}$
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
6.8 ± 0.9				OUR FIT		
6.31^{+1.70+0.30}_{-1.48-0.36}	28	TAJIMA	04	BELL	e^+e^- at $\Upsilon(4S)$	

$\Gamma(\phi\gamma)/\Gamma(K^-\pi^+)$						Γ_{203}/Γ_{31}
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
(7.0±0.9)				OUR FIT		
7.15±0.78±0.69	243 ± 25	AUBERT	08AZ	BABR	$e^+e^- \approx 10.6$ GeV	

$\Gamma(\bar{K}^*(892)^0\gamma)/\Gamma(K^-\pi^+)$						Γ_{204}/Γ_{31}
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
8.43±0.51±0.70	2286 ± 113	AUBERT	08AZ	BABR	$e^+e^- \approx 10.6$ GeV	

———— Doubly Cabibbo-suppressed / Mixing modes ————

$\Gamma(K^+\ell^-\bar{\nu}_\ell \text{ via } \bar{D}^0)/\Gamma(K^-\ell^+\nu_\ell)$						Γ_{205}/Γ_{17}
This is a limit on R_M without the complications of possible doubly Cabibbo-suppressed decays that occur when using hadronic modes. For the limits on $ m_1 - m_2 $ and $(\Gamma_1 - \Gamma_2)/\Gamma$ that come from the best mixing limit, see near the beginning of these D^0 Listings.						

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 6.1 × 10⁻⁴	90	¹²⁴ BITENC	08	BELL	e^+e^- , 10.58 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<50 × 10 ⁻⁴	90	¹²⁵ AITALA	96C	E791	π^- nucleus, 500 GeV
¹²⁴ The BITENC 08 right-sign sample includes about 15% of $D^0 \rightarrow K^-\pi^0\ell^+\nu_\ell$ and other decays.					
¹²⁵ AITALA 96C uses $D^{*+} \rightarrow D^0\pi^+$ (and charge conjugate) decays to identify the charm at production and $D^0 \rightarrow K^-\ell^+\nu_\ell$ (and charge conjugate) decays to identify the charm at decay.					

$\Gamma(K^+ \text{ or } K^*(892)^+ e^-\bar{\nu}_e \text{ via } \bar{D}^0)/[\Gamma(K^-\ell^+\nu_\ell) + \Gamma(K^*(892)^-\ell^+\nu_\ell)]$						$\Gamma_{206}/(\Gamma_{18}+\Gamma_{20})$
This is a limit on R_M without the complications of possible doubly Cabibbo-suppressed decays that occur when using hadronic modes. The experiments use $D^{*+} \rightarrow D^0\pi^+$ (and charge conjugate) decays to identify the charm at production and the charge of the e to identify the charm at decay. These limits do not allow CP violation. For the limits on $ m_1 - m_2 $ and $(\Gamma_1 - \Gamma_2)/\Gamma$ that come from the best mixing limit, see near the beginning of these D^0 Listings.						

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.001	90	BITENC	05	BELL	$e^+e^- \approx 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
-0.0013 < R < +0.0012	90	AUBERT	07AB	BABR	$e^+e^- \approx 10.58$ GeV
<0.0078	90	CAWLFIELD	05	CLEO	$e^+e^- \approx 10.6$ GeV
<0.0042	90	AUBERT,B	04Q	BABR	See AUBERT 07AB

$\Gamma(K^+\pi^-)/\Gamma(K^-\pi^+)$

Γ_{207}/Γ_{31}

This is R , the time-integrated wrong-sign rate compared to the right-sign rate. See the note on “ D^0 - \bar{D}^0 Mixing,” near the start of the D^0 Listings.

The experiments here use the charge of the pion in $D^*(2010)^\pm \rightarrow (D^0 \text{ or } \bar{D}^0) \pi^\pm$ decay to tell whether a D^0 or a \bar{D}^0 was born. The $D^0 \rightarrow K^+\pi^-$ decay can occur directly by doubly Cabibbo-suppressed (DCS) decay, or indirectly by $D^0 \rightarrow \bar{D}^0$ mixing followed by $\bar{D}^0 \rightarrow K^+\pi^-$ decay. Some of the experiments can use the decay-time information to disentangle the two mechanisms. Here, we list the experimental branching ratio, which if there is no mixing is the DCS ratio. See the next data block for values of the DCS ratio R_D , and the following data block for limits on the mixing ratio R_M . See the section on CP -violating asymmetries near the end of this D^0 Listing for values of A_D , and the note on “ D^0 - \bar{D}^0 Mixing” for limits on x' and y' .

Some early limits have been omitted from this Listing; see our 1998 edition (The European Physical Journal **C3** 1 (1998)) and our 2006 edition (Journal of Physics, G **33** 1 (2006)).

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.79±0.18 OUR FIT	Error includes scale factor of 3.3.			
3.80±0.18 OUR AVERAGE	Error includes scale factor of 3.3. See the ideogram below.			
4.15±0.10	12.7 ± 0.3k	¹²⁶ AALTONEN	08E CDF	$p\bar{p}$, $\sqrt{s} = 1.96$ TeV
3.53±0.08±0.04	4030 ± 90	¹²⁷ AUBERT	07W BABR	$e^+e^- \approx 10.6$ GeV
3.77±0.08±0.05	4024 ± 88	¹²⁶ ZHANG	06 BELL	e^+e^-
4.29 ^{+0.63} _{-0.61} ± 0.27	234	¹²⁸ LINK	05H FOCS	γ nucleus
3.32 ^{+0.63} _{-0.65} ± 0.40	45	¹²⁶ GODANG	00 CLE2	e^+e^-
6.8 ^{+3.4} _{-3.3} ± 0.7	34	¹²⁷ AITALA	98 E791	π^- nucl., 500 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.05±0.21±0.11	2.0 ± 0.1k	¹²⁹ ABULENCIA	06X CDF	See AALTONEN 08E
3.81±0.17 ^{+0.08} _{-0.16}	845 ± 40	¹²⁷ LI	05A BELL	See ZHANG 06
3.57±0.22±0.27		¹³⁰ AUBERT	03Z BABR	See AUBERT 07W
4.04±0.85±0.25	149	¹³¹ LINK	01 FOCS	γ nucleus

¹²⁶ GODANG 00, ZHANG 06, and AALTONEN 08E allow CP violation.

¹²⁷ AITALA 98, LI 05A, and AUBERT 07W assume no CP violation.


¹²⁸ This LINK 05H result assumes no mixing but allows CP violation. If neither mixing nor CP violation is allowed, $R = (4.29 \pm 0.63 \pm 0.28) \times 10^{-3}$.

¹²⁹ This ABULENCIA 06X result assumes no mixing.

¹³⁰ This AUBERT 03Z result allows CP violation. If CP violation is not allowed, $R = 0.00359 \pm 0.00020 \pm 0.00027$.

¹³¹ This LINK 01 result assumes no mixing or CP violation.

WEIGHTED AVERAGE
 3.80 ± 0.18 (Error scaled by 3.3)




$\Gamma(K^+ \pi^- \text{ via DCS}) / \Gamma(K^- \pi^+)$

$\Gamma_{208} / \Gamma_{31}$

This is R_D , the doubly Cabibbo-suppressed ratio when mixing is allowed.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.37 ± 0.21 OUR AVERAGE					Error includes scale factor of 1.8. See the ideogram below.
3.04 ± 0.55		$12.7 \pm 0.3k$	AALTONEN	08E CDF	$p\bar{p}$, $\sqrt{s} = 1.96$ TeV
$3.03 \pm 0.16 \pm 0.10$		4030 ± 90	¹³² AUBERT	07W BABR	$e^+ e^- \approx 10.6$ GeV
3.64 ± 0.17		4024 ± 88	¹³³ ZHANG	06 BELL	$e^+ e^-$
$5.17^{+1.47}_{-1.58} \pm 0.76$		234	¹³⁴ LINK	05H FOCS	γ nucleus
$4.8 \pm 1.2 \pm 0.4$		45	¹³⁵ GODANG	00 CLE2	$e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2.87 ± 0.37		845 ± 40	LI	05A BELL	See ZHANG 06
$2.3 < R_D < 5.2$	95		¹³⁶ AUBERT	03Z BABR	See AUBERT 07W
$9.0^{+12.0}_{-10.9} \pm 4.4$		34	¹³⁷ AITALA	98 E791	π^- nucl., 500 GeV

¹³² This AUBERT 07W result is the same whether or not CP violation is allowed.
¹³³ This ZHANG 06 assumes no CP violation.
¹³⁴ This LINK 05H result allows CP violation. Allowing mixing but not CP violation, $R_D = (3.81^{+1.67}_{-1.63} \pm 0.92) \times 10^{-3}$.
¹³⁵ This GODANG 00 result allows CP violation.
¹³⁶ This AUBERT 03Z result allows CP violation. If only mixing is allowed, the 95% confidence level interval is $(2.4 < R_D < 4.9) \times 10^{-3}$.
¹³⁷ This AITALA 98 result assumes no CP violation.



$\Gamma(K^+ \pi^- \text{ via } \bar{D}^0)/\Gamma(K^- \pi^+)$

Γ_{209}/Γ_{31}

This is R_M in the note on “ D^0 - \bar{D}^0 Mixing” near the start of the D^0 Listings. The experiments here (1) use the charge of the pion in $D^*(2010)^\pm \rightarrow (D^0 \text{ or } \bar{D}^0) \pi^\pm$ decay to tell whether a D^0 or a \bar{D}^0 was born; and (2) use the decay-time distribution to disentangle doubly Cabibbo-suppressed decay and mixing. For the limits on $|m_1 - m_2|$ and $(\Gamma_1 - \Gamma_2)/\Gamma$ that come from the best mixing limit, see near the beginning of these D^0 Listings.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.00040	95	138	ZHANG	06	BELL $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.00046	95	139	LI	05A	BELL See ZHANG 06
<0.0063	95	140	LINK	05H	FOCS γ nucleus
<0.0013	95	141	AUBERT	03Z	BABR $e^+ e^-$, 10.6 GeV
<0.00041	95	142	GODANG	00	CLE2 $e^+ e^-$
<0.0092	95	143	BARATE	98W	ALEP $e^+ e^-$ at Z^0
<0.005	90	1 ± 4	144 ANJOS	88C	E691 Photoproduction

138 This ZHANG 06 result allows CP violation, but the result does not change if CP violation is not allowed.
 139 This LI 05A result allows CP violation. The limit becomes < 0.00042 (95% CL) if CP violation is not allowed.
 140 LINK 05H obtains the same result whether or not CP violation is allowed.
 141 This AUBERT 03Z result allows CP violation and assumes that the strong phase between $D^0 \rightarrow K^+ \pi^-$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ is small, and limits only $D^0 \rightarrow \bar{D}^0$ transitions via off-shell intermediate states. The limit on transitions via on-shell intermediate states is 0.0016.

142 This GODANG 00 result allows CP violation and assumes that the strong phase between $D^0 \rightarrow K^+ \pi^-$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ is small, and limits only $D^0 \rightarrow \bar{D}^0$ transitions via off-shell intermediate states. The limit on transitions via on-shell intermediate states is 0.0017.

143 This BARATE 98W result assumes no interference between the DCS and mixing amplitudes ($y' = 0$ in the note on " D^0 - \bar{D}^0 Mixing" near the start of the D^0 Listings). When interference is allowed, the limit degrades to 0.036 (95%CL).

144 This ANJOS 88C result assumes no interference between the DCS and mixing amplitudes ($y' = 0$ in the note on " D^0 - \bar{D}^0 Mixing" near the start of the D^0 Listings). When interference is allowed, the limit degrades to 0.019.

$\Gamma(K_S^0 \pi^+ \pi^- \text{ in } D^0 \rightarrow \bar{D}^0) / \Gamma(K_S^0 \pi^+ \pi^-)$ $\Gamma_{210} / \Gamma_{34}$

This is R_M in the note on " D^0 - \bar{D}^0 Mixing" near the start of the D^0 Listings. The experiments here (1) use the charge of the pion in $D^*(2010)^\pm \rightarrow (D^0 \text{ or } \bar{D}^0) \pi^\pm$ decay to tell whether a D^0 or a \bar{D}^0 was born; and (2) use the decay-time distribution to disentangle doubly Cabibbo-suppressed decay and mixing. For the limits on $|m_1 - m_2|$ and $(\Gamma_1 - \Gamma_2) / \Gamma$ that come from the best mixing limit, see near the beginning of these D^0 Listings.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0063	95	145 ASNER	05	CLEO $e^+ e^- \approx 10$ GeV

145 This ASNER 05 limit allows CP violation. If CP violation is not allowed, the limit is 0.0042 at 95% CL.

$\Gamma(K^+ \pi^- \pi^0) / \Gamma(K^- \pi^+ \pi^0)$ $\Gamma_{214} / \Gamma_{49}$

The experiments here use the charge of the pion in $D^*(2010)^\pm \rightarrow (D^0 \text{ or } \bar{D}^0) \pi^\pm$ decay to tell whether a D^0 or a \bar{D}^0 was born. The $D^0 \rightarrow K^+ \pi^- \pi^0$ decay can occur directly by doubly Cabibbo-suppressed (DCS) decay, or indirectly by $D^0 \rightarrow \bar{D}^0$ mixing followed by $\bar{D}^0 \rightarrow K^+ \pi^- \pi^0$ decay.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20 ± 0.10 OUR AVERAGE				
2.14 ± 0.08 ± 0.08	763 ± 51	146 AUBERT,B	06N BABR	$e^+ e^- \approx \Upsilon(4S)$
2.29 ± 0.15 $^{+0.13}_{-0.09}$	1978 ± 104	TIAN	05 BELL	$e^+ e^- \approx \Upsilon(4S)$
4.3 $^{+1.1}_{-1.0} \pm 0.7$	38	BRANDENB...	01 CLE2	$e^+ e^- \approx \Upsilon(4S)$

146 This AUBERT,B 06N result assumes no mixing.

$\Gamma(K^+ \pi^- \pi^0 \text{ via } \bar{D}^0) / \Gamma(K^- \pi^+ \pi^0)$ $\Gamma_{215} / \Gamma_{49}$

This is R_M in the note on " D^0 - \bar{D}^0 Mixing" near the start of the D^0 Listings. The experiments here (1) use the charge of the pion in $D^*(2010)^\pm \rightarrow (D^0 \text{ or } \bar{D}^0) \pi^\pm$ decay to tell whether a D^0 or a \bar{D}^0 was born; and (2) use the decay-time distribution to disentangle doubly Cabibbo-suppressed decay and mixing. For the limits on $|m_1 - m_2|$ and $(\Gamma_1 - \Gamma_2) / \Gamma$ that come from the best mixing limit, see near the beginning of these D^0 Listings.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
5.25 $^{+0.25}_{-0.31} \pm 0.12$		AUBERT	09AN BABR	$e^+ e^-$ at 10.58 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.54	95	147 AUBERT,B	06N BABR	$e^+ e^- \approx \Upsilon(4S)$
-------	----	--------------	----------	--------------------------------

¹⁴⁷ This AUBERT,B 06N limit assumes no CP violation. The measured value corresponding to the limit is $(2.3_{-1.4}^{+1.8} \pm 0.4) \times 10^{-4}$. If CP violation is allowed, this becomes $(1.0_{-0.7}^{+2.2} \pm 0.3) \times 10^{-4}$.

$\Gamma(K^+ \pi^+ 2\pi^-)/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{216}/Γ_{61}

The experiments here use the charge of the pion in $D^*(2010)^\pm \rightarrow (D^0 \text{ or } \bar{D}^0) \pi^\pm$ decay to tell whether a D^0 or a \bar{D}^0 was born. The $D^0 \rightarrow K^+ \pi^- \pi^+ \pi^-$ decay can occur directly by doubly Cabibbo-suppressed (DCS) decay, or indirectly by $D^0 \rightarrow \bar{D}^0$ mixing followed by $\bar{D}^0 \rightarrow K^+ \pi^- \pi^+ \pi^-$ decay. Some of the experiments can use the decay-time information to disentangle the two mechanisms. Here, we list the experimental branching ratio, which if there is no mixing is the DCS ratio; in the next data block we give the limits on the mixing ratio.

Some early limits have been omitted from this Listing; see our 1998 edition (EPJ **C3** 1).

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$3.24_{-0.22}^{+0.25}$ OUR AVERAGE

$3.20 \pm 0.18_{-0.13}^{+0.18}$	1721 ± 75	¹⁴⁸ TIAN	05	BELL	$e^+ e^- \approx \Upsilon(4S)$
$4.4_{-1.2}^{+1.3} \pm 0.4$	54	¹⁴⁸ DYTMAN	01	CLE2	$e^+ e^- \approx \Upsilon(4S)$
$2.5_{-3.4}^{+3.6} \pm 0.3$		¹⁴⁹ AITALA	98	E791	π^- nucl., 500 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18	90	¹⁴⁸ AMMAR	91	CLEO	$e^+ e^- \approx 10.5$ GeV
<18	90	5 ± 12	¹⁵⁰ ANJOS	88C E691	Photoproduction

¹⁴⁸ AMMAR 91 cannot and DYTMAN 01 and TIAN 05 do not distinguish between doubly Cabibbo-suppressed decay and D^0 - \bar{D}^0 mixing.

¹⁴⁹ This AITALA 98 result assumes no D^0 - \bar{D}^0 mixing (R_M in the note on “ D^0 - \bar{D}^0 Mixing”). It becomes $-0.0020_{-0.0106}^{+0.0117} \pm 0.0035$ when mixing is allowed and decay-time information is used to distinguish doubly Cabibbo-suppressed decays from mixing.

¹⁵⁰ ANJOS 88C uses decay-time information to distinguish doubly Cabibbo-suppressed (DCS) decays from D^0 - \bar{D}^0 mixing. However, the result assumes no interference between the DCS and mixing amplitudes ($y' = 0$ in the note on “ D^0 - \bar{D}^0 Mixing” near the start of the D^0 Listings). When interference is allowed, the limit degrades to 0.033.

$\Gamma(K^+ \pi^+ 2\pi^- \text{ via } \bar{D}^0)/\Gamma(K^- 2\pi^+ \pi^-)$ Γ_{217}/Γ_{61}

This is a D^0 - \bar{D}^0 mixing limit. The experiments here (1) use the charge of the pion in $D^*(2010)^\pm \rightarrow (D^0 \text{ or } \bar{D}^0) \pi^\pm$ decay to tell whether a D^0 or a \bar{D}^0 was born; and (2) use the decay-time distribution to disentangle doubly Cabibbo-suppressed decay and mixing. For the limits on $|m_{D_1^0} - m_{D_2^0}|$ and $(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma_{D^0}$ that come from the best mixing limit, see near the beginning of these D^0 Listings.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

<0.005	90	0 ± 4	¹⁵¹ ANJOS	88C E691	Photoproduction
------------------	----	-----------	----------------------	----------	-----------------

¹⁵¹ ANJOS 88C uses decay-time information to distinguish doubly Cabibbo-suppressed (DCS) decays from D^0 - \bar{D}^0 mixing. However, the result assumes no interference between the DCS and mixing amplitudes ($y' = 0$ in the note on “ D^0 - \bar{D}^0 Mixing” near the start of the D^0 Listings). When interference is allowed, the limit degrades to 0.007.

$\Gamma(K^+\pi^- \text{ or } K^+\pi^+2\pi^- \text{ via } \bar{D}^0)/\Gamma(K^-\pi^+ \text{ or } K^-2\pi^+\pi^-)$ **Γ_{218}/Γ_0**

This is a $D^0-\bar{D}^0$ mixing limit. For the limits on $|m_{D_1^0} - m_{D_2^0}|$ and $(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma_{D^0}$

that come from the best mixing limit, see near the beginning of these D^0 Listings.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0085	90	152 AITALA	98 E791	π^- nucleus, 500 GeV
<0.0037	90	153 ANJOS	88C E691	Photoproduction

152 AITALA 98 uses decay-time information to distinguish doubly Cabibbo-suppressed decays from $D^0-\bar{D}^0$ mixing. The fit allows interference between the two amplitudes, and also allows CP violation in this term. The central value obtained is $0.0039^{+0.0036}_{-0.0032} \pm 0.0016$.

When interference is disallowed, the result becomes $0.0021 \pm 0.0009 \pm 0.0002$.

153 This combines results of ANJOS 88C on $K^+\pi^-$ and $K^+\pi^-\pi^+\pi^-$ (via \bar{D}^0) reported in the data block above (see footnotes there). It assumes no interference.

$\Gamma(\mu^- \text{ anything via } \bar{D}^0)/\Gamma(\mu^+ \text{ anything})$ **Γ_{219}/Γ_6**

This is a $D^0-\bar{D}^0$ mixing limit. See the somewhat better limits above.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.0056	90	LOUIS	86 SPEC	π^- W 225 GeV
-------------------	----	-------	---------	-------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.012	90	BENVENUTI	85 CNTR	μC , 200 GeV
<0.044	90	BODEK	82 SPEC	π^- , $p\text{Fe} \rightarrow D^0$

————— **Rare or forbidden modes** —————

$\Gamma(\gamma\gamma)/\Gamma(2\pi^0)$ **$\Gamma_{220}/\Gamma_{127}$**

$D^0 \rightarrow \gamma\gamma$ is a flavor-changing neutral-current decay, forbidden in the Standard Model at the tree level.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.033	90	COAN	03 CLE2	$e^+e^- \approx \Upsilon(4S)$
------------------	----	------	---------	-------------------------------

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ **Γ_{221}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	-------------	--------------------	-------------	----------------

<7.9 $\times 10^{-8}$	90		PETRIC	10 BELL	$e^+e^- \approx \Upsilon(4S)$
--	----	--	--------	---------	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.2 $\times 10^{-6}$	90	3	AUBERT,B	04Y BABR	$e^+e^- \approx \Upsilon(4S)$
<8.19 $\times 10^{-6}$	90		PRIPSTEIN	00 E789	p nucleus, 800 GeV
<6.2 $\times 10^{-6}$	90		AITALA	99G E791	$\pi^- N$ 500 GeV
<1.3 $\times 10^{-5}$	90	0	FREYBERGER	96 CLE2	$e^+e^- \approx \Upsilon(4S)$
<1.3 $\times 10^{-4}$	90		ADLER	88 MRK3	e^+e^- 3.77 GeV
<1.7 $\times 10^{-4}$	90	7	ALBRECHT	88G ARG	e^+e^- 10 GeV
<2.2 $\times 10^{-4}$	90	8	HAAS	88 CLEO	e^+e^- 10 GeV

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{222}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-7}$	90		PETRIC	10	BELL $e^+ e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.1 \times 10^{-7}$	90	4	AALTONEN	10X	CDF $p\bar{p}, \sqrt{s} = 1.96$ TeV
$<2.0 \times 10^{-6}$	90		ABT	04	HERB $pA, 920$ GeV
$<1.3 \times 10^{-6}$	90	1	AUBERT,B	04Y	BABR $e^+ e^- \approx \Upsilon(4S)$
$<2.5 \times 10^{-6}$	90		ACOSTA	03F	CDF See AALTONEN 10X
$<1.56 \times 10^{-5}$	90		PRIPSTEIN	00	E789 p nucleus, 800 GeV
$<5.2 \times 10^{-6}$	90		AITALA	99G	E791 $\pi^- N$ 500 GeV
$<4.1 \times 10^{-6}$	90		ADAMOVICH	97	BEAT π^- Cu, W 350 GeV
$<4.2 \times 10^{-6}$	90		ALEXOPOU...	96	E771 p Si, 800 GeV
$<3.4 \times 10^{-5}$	90	1	FREYBERGER	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$
$<7.6 \times 10^{-6}$	90	0	ADAMOVICH	95	BEAT See ADAMOVICH 97
$<4.4 \times 10^{-5}$	90	0	KODAMA	95	E653 π^- emulsion 600 GeV
$<3.1 \times 10^{-5}$	90	154	MISHRA	94	E789 -4.1 ± 4.8 events
$<7.0 \times 10^{-5}$	90	3	ALBRECHT	88G	ARG $e^+ e^-$ 10 GeV
$<1.1 \times 10^{-5}$	90		LOUIS	86	SPEC π^- W 225 GeV
$<3.4 \times 10^{-4}$	90		AUBERT	85	EMC Deep inelast. $\mu^- N$

¹⁵⁴ Here MISHRA 94 uses "the statistical approach advocated by the PDG." For an alternate approach, giving a limit of 9×10^{-6} at 90% confidence level, see the paper.

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{223}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<4.5 \times 10^{-5}$	90	0	FREYBERGER	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{224}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-4}$	90	2	KODAMA	95	E653 π^- emulsion 600 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<5.4 \times 10^{-4}$	90	3	FREYBERGER	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{225}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-4}$	90	0	FREYBERGER	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\eta \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{226}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<5.3 \times 10^{-4}$	90	0	FREYBERGER	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\pi^+ \pi^- e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{227}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<3.73 \times 10^{-4}$	90	9	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\rho^0 e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{228}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-4}$	90	2	¹⁵⁵ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.24 \times 10^{-4}$	90	1	AITALA	01C E791	π^- nucleus, 500 GeV
$<4.5 \times 10^{-4}$	90	2	HAAS	88 CLEO	$e^+ e^-$ 10 GeV

¹⁵⁵ This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 1.8 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(\pi^+ \pi^- \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{229}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-5}$	90	2	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\rho^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{230}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	0	AITALA	01C E791	π^- nucleus, 500 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4.9 \times 10^{-4}$	90	1	¹⁵⁶ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$
$<2.3 \times 10^{-4}$	90	0	KODAMA 95	E653	π^- emulsion 600 GeV
$<8.1 \times 10^{-4}$	90	5	HAAS 88	CLEO	$e^+ e^-$ 10 GeV

¹⁵⁶ This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 4.5 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(\omega e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{231}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-4}$	90	1	¹⁵⁷ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

¹⁵⁷ This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 2.7 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(\omega \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{232}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<8.3 \times 10^{-4}$	90	0	¹⁵⁸ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

¹⁵⁸ This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 6.5 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(K^- K^+ e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{233}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.15 \times 10^{-4}$	90	9	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\phi e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{234}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.2 \times 10^{-5}$	90	2	¹⁵⁹ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<5.9 \times 10^{-5}$	90	0	AITALA	01C E791	π^- nucleus, 500 GeV
-----------------------	----	---	--------	----------	--------------------------

¹⁵⁹ This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 7.6 \times 10^{-5}$ using a photon pole amplitude model.

$\Gamma(K^- K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{235}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.3 \times 10^{-5}$	90	0	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\phi \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{236}/Γ**

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.1 \times 10^{-5}$	90	0	AITALA	01C E791	π^- nucleus, 500 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4.1 \times 10^{-4}$	90	0	¹⁶⁰ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$
-----------------------	----	---	------------------------------	------	------------------------------

¹⁶⁰ This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 2.4 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(\bar{K}^0 e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{237}/Γ**

Not a useful test for $\Delta C = 1$ weak neutral current because both quarks must change flavor.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.1 \times 10^{-4}$	90	0	FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.7 \times 10^{-3}$	90		ADLER	89C MRK3	$e^+ e^-$ 3.77 GeV
-----------------------	----	--	-------	----------	--------------------

$\Gamma(\bar{K}^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{238}/Γ**

Not a useful test for $\Delta C = 1$ weak neutral current because both quarks must change flavor.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.6 \times 10^{-4}$	90	2	KODAMA 95	E653	π^- emulsion 600 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6.7 \times 10^{-4}$	90	1	FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$
-----------------------	----	---	---------------	------	------------------------------

$\Gamma(K^- \pi^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{239}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 3.85 \times 10^{-4}$	90	6	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\bar{K}^*(892)^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{240}/Γ

Not a useful test for $\Delta C = 1$ weak neutral current because both quarks must change flavor.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 4.7 \times 10^{-5}$	90	2	AITALA	01C E791	π^- nucleus, 500 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.4 \times 10^{-4}$	90	1	¹⁶¹ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$
------------------------	----	---	------------------------------	------	------------------------------

¹⁶¹This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 2.0 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(K^- \pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{241}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 3.59 \times 10^{-4}$	90	12	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\bar{K}^*(892)^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{242}/Γ

Not a useful test for $\Delta C = 1$ weak neutral current because both quarks must change flavor.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 2.4 \times 10^{-5}$	90	3	AITALA	01C E791	π^- nucleus, 500 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.18 \times 10^{-3}$	90	1	¹⁶² FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$
-------------------------	----	---	------------------------------	------	------------------------------

¹⁶²This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 1.0 \times 10^{-3}$ using a photon pole amplitude model.

$\Gamma(\pi^+ \pi^- \pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{243}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 8.1 \times 10^{-4}$	90	1	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(\mu^\pm e^\mp)/\Gamma_{\text{total}}$ Γ_{244}/Γ

A test of lepton family number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 2.6 \times 10^{-7}$	90		PETRIC	10 BELL	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 8.1 \times 10^{-7}$	90	0	AUBERT,B	04Y BABR	$e^+ e^- \approx \gamma(4S)$
$< 1.72 \times 10^{-5}$	90		PRIPSTEIN	00 E789	p nucleus, 800 GeV
$< 8.1 \times 10^{-6}$	90		AITALA	99G E791	$\pi^- N$ 500 GeV
$< 1.9 \times 10^{-5}$	90	2	¹⁶³ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$
$< 1.0 \times 10^{-4}$	90	4	ALBRECHT	88G ARG	$e^+ e^-$ 10 GeV
$< 2.7 \times 10^{-4}$	90	9	HAAS	88 CLEO	$e^+ e^-$ 10 GeV
$< 1.2 \times 10^{-4}$	90		BECKER	87C MRK3	$e^+ e^-$ 3.77 GeV
$< 9 \times 10^{-4}$	90		PALKA	87 SILI	200 GeV πp
$< 21 \times 10^{-4}$	90	0	¹⁶⁴ RILES	87 MRK2	$e^+ e^-$ 29 GeV

¹⁶³ This is the corrected result given in the erratum to FREYBERGER 96.

¹⁶⁴ RILES 87 assumes $B(D \rightarrow K\pi) = 3.0\%$ and has production model dependency.

$\Gamma(\pi^0 e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{245}/Γ**

A test of lepton family number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.6 \times 10^{-5}$	90	2	FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(\eta e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{246}/Γ**

A test of lepton family number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.0 \times 10^{-4}$	90	0	FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(\pi^+ \pi^- e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{247}/Γ**

A test of lepton family-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.5 \times 10^{-5}$	90	1	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\rho^0 e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{248}/Γ**

A test of lepton family number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.9 \times 10^{-5}$	90	0	¹⁶⁵ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6.6 \times 10^{-5}$	90	1	AITALA	01C E791	π^- nucleus, 500 GeV
-----------------------	----	---	--------	----------	--------------------------

¹⁶⁵ This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 5.0 \times 10^{-5}$ using a photon pole amplitude model.

$\Gamma(\omega e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{249}/Γ**

A test of lepton family number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.2 \times 10^{-4}$	90	0	¹⁶⁶ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

¹⁶⁶ This FREYBERGER 96 limit is obtained using a phase-space model. The same limit is obtained using a photon pole amplitude model.

$\Gamma(K^- K^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{250}/Γ**

A test of lepton family-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.8 \times 10^{-4}$	90	5	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\phi e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{251}/Γ**

A test of lepton family number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.4 \times 10^{-5}$	90	0	¹⁶⁷ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4.7 \times 10^{-5}$	90	0	AITALA	01C E791	π^- nucleus, 500 GeV
-----------------------	----	---	--------	----------	--------------------------

¹⁶⁷ This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 3.3 \times 10^{-5}$ using a photon pole amplitude model.

$\Gamma(\bar{K}^0 e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{252}/Γ**

A test of lepton family number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.0 \times 10^{-4}$	90	0	FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(K^- \pi^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{253}/Γ**

A test of lepton family-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.53 \times 10^{-4}$	90	15	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\bar{K}^*(892)^0 e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{254}/Γ**

A test of lepton family number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.3 \times 10^{-5}$	90	9	AITALA	01C E791	π^- nucleus, 500 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.0 \times 10^{-4}$	90	0	¹⁶⁸ FREYBERGER 96	CLE2	$e^+ e^- \approx \gamma(4S)$
-----------------------	----	---	------------------------------	------	------------------------------

¹⁶⁸ This FREYBERGER 96 limit is obtained using a phase-space model. The same limit is obtained using a photon pole amplitude model.

$\Gamma(2\pi^- 2e^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{255}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.12 \times 10^{-4}$	90	1	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(2\pi^- 2\mu^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{256}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.9 \times 10^{-5}$	90	1	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(K^- \pi^- 2e^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{257}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.06 \times 10^{-4}$	90	2	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(K^- \pi^- 2\mu^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{258}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.9 \times 10^{-4}$	90	14	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(2K^- 2e^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{259}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.52 \times 10^{-4}$	90	2	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(2K^- 2\mu^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{260}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.4 \times 10^{-5}$	90	1	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(\pi^- \pi^- e^+ \mu^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{261}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.9 \times 10^{-5}$	90	4	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(K^- \pi^- e^+ \mu^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{262}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.18 \times 10^{-4}$	90	7	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(2K^- e^+ \mu^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{263}/Γ**

A test of lepton-number conservation. The value is for the sum of the two charge states.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.7 \times 10^{-5}$	90	0	AITALA	01C E791	π^- nucleus, 500 GeV

$\Gamma(pe^-)/\Gamma_{\text{total}}$ **Γ_{264}/Γ**

A test of baryon- and lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.0 \times 10^{-5}$	90	169	RUBIN	09 CLEO	$e^+ e^-$ at $\psi(3770)$

¹⁶⁹ This RUBIN 09 limit is for either $D^0 \rightarrow pe^-$ or $\bar{D}^0 \rightarrow pe^-$ decay.

$\Gamma(\bar{p}e^+)/\Gamma_{\text{total}}$ **Γ_{265}/Γ**

A test of baryon- and lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.1 \times 10^{-5}$	90	170	RUBIN	09 CLEO	$e^+ e^-$ at $\psi(3770)$

¹⁷⁰ This RUBIN 09 limit is for either $D^0 \rightarrow \bar{p}e^+$ or $\bar{D}^0 \rightarrow \bar{p}e^+$ decay.

D^0 CP-VIOLATING DECAY-RATE ASYMMETRIES

This is the difference between D^0 and \bar{D}^0 partial widths for these modes divided by the sum of the widths. The D^0 and \bar{D}^0 are distinguished by the charge of the parent D^* : $D^{*+} \rightarrow D^0 \pi^+$ and $D^{*-} \rightarrow \bar{D}^0 \pi^-$.


$A_{CP}(K^+ K^-)$ in $D^0, \bar{D}^0 \rightarrow K^+ K^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.17 ± 0.31 OUR AVERAGE		Error includes scale factor of 1.3.		See the ideogram below.
$0.00 \pm 0.34 \pm 0.13$	129k	171 AUBERT	08M BABR	$e^+ e^- \approx 10.6$ GeV
$-0.43 \pm 0.30 \pm 0.11$	120k	172 STARIC	08 BELL	$e^+ e^- \approx \Upsilon(4S)$
$+2.0 \pm 1.2 \pm 0.6$		173 ACOSTA	05C CDF	$p\bar{p}, \sqrt{s}=1.96$ TeV
$0.0 \pm 2.2 \pm 0.8$	3023	173 CSORNA	02 CLE2	$e^+ e^- \approx \Upsilon(4S)$
$-0.1 \pm 2.2 \pm 1.5$	3330	173 LINK	00B FOCS	
$-1.0 \pm 4.9 \pm 1.2$	609	173 AITALA	98C E791	$-0.093 < A_{CP} < +0.073$ (90% CL)

171 AUBERT 08M uses corrected numbers of events directly, not ratios with $K^\mp \pi^\pm$ events.

172 STARIC 08 uses $D^0 \rightarrow K^- \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ decays to correct for detector-induced asymmetries.

173 AITALA 98C, LINK 00B, CSORNA 02, and ACOSTA 05C measure $N(D^0 \rightarrow K^+ K^-)/N(D^0 \rightarrow K^- \pi^+)$, the ratio of numbers of events observed, and similarly for the \bar{D}^0 .



$A_{CP}(K^+ K^-)$ in $D^0, \bar{D}^0 \rightarrow K^+ K^-$ (%)

$A_{CP}(K_S^0 K_S^0)$ in $D^0, \bar{D}^0 \rightarrow K_S^0 K_S^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-23 ± 19	65	BONVICINI 01	CLE2	$e^+ e^- \approx 10.6$ GeV

$A_{CP}(\pi^+\pi^-)$ in $D^0, \bar{D}^0 \rightarrow \pi^+\pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.2 ± 0.4 OUR AVERAGE				
-0.24 ± 0.52 ± 0.22	63.7k	174 AUBERT	08M BABR	$e^+e^- \approx 10.6$ GeV
+0.43 ± 0.52 ± 0.12	51k	175 STARIC	08 BELL	$e^+e^- \approx \Upsilon(4S)$
+1.0 ± 1.3 ± 0.6		176 ACOSTA	05C CDF	$p\bar{p}, \sqrt{s}=1.96$ TeV
+1.9 ± 3.2 ± 0.8	1136	176 CSORNA	02 CLE2	$e^+e^- \approx \Upsilon(4S)$
+4.8 ± 3.9 ± 2.5	1177	176 LINK	00B FOCS	
-4.9 ± 7.8 ± 3.0	343	176 AITALA	98C E791	$-0.186 < A_{CP} < +0.088$ (90% CL)

174 AUBERT 08M uses corrected numbers of events directly, not ratios with $K^\mp\pi^\pm$ events.

175 STARIC 08 uses $D^0 \rightarrow K^-\pi^+$ and $\bar{D}^0 \rightarrow K^+\pi^-$ decays to correct for detector-induced asymmetries.

176 AITALA 98C, LINK 00B, CSORNA 02, and ACOSTA 05C measure $N(D^0 \rightarrow \pi^+\pi^-)/N(D^0 \rightarrow K^-\pi^+)$, the ratio of numbers of events observed, and similarly for the \bar{D}^0 .

$A_{CP}(\pi^0\pi^0)$ in $D^0, \bar{D}^0 \rightarrow \pi^0\pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
+0.1 ± 4.8	810	BONVICINI	01 CLE2	$e^+e^- \approx 10.6$ GeV

$A_{CP}(\pi^+\pi^-\pi^0)$ in $D^0, \bar{D}^0 \rightarrow \pi^+\pi^-\pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.3 ± 0.4 OUR AVERAGE				
+0.43 ± 1.30	123k ± 490	ARINSTEIN	08 BELL	$e^+e^- \approx \Upsilon(4S)$
+0.31 ± 0.41 ± 0.17	80 ± .3k	AUBERT	08AO BABR	$e^+e^- \approx 10.6$ GeV
+1 ⁺⁹ / ₋₇ ± 5		CRONIN-HEN..05	CLEO	$e^+e^- \approx 10$ GeV

$A_{CP}(\rho(770)^+\pi^- \rightarrow \pi^+\pi^-\pi^0)$ in $D^0 \rightarrow \rho^+\pi^-, \bar{D}^0 \rightarrow \rho^-\pi^+$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+1.2 ± 0.8 ± 0.3	AUBERT	08AO BABR	Table 1, -Col.5/2 × Col.2

$A_{CP}(\rho(770)^0\pi^0 \rightarrow \pi^+\pi^-\pi^0)$ in $D^0, \bar{D}^0 \rightarrow \rho^0\pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-3.1 ± 2.7 ± 1.2	AUBERT	08AO BABR	Table 1, -Col.5/2 × Col.2

$A_{CP}(\rho(770)^-\pi^+ \rightarrow \pi^+\pi^-\pi^0)$ in $D^0 \rightarrow \rho^-\pi^+, \bar{D}^0 \rightarrow \rho^+\pi^-$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-1.0 ± 1.6 ± 0.7	AUBERT	08AO BABR	Table 1, -Col.5/2 × Col.2

$A_{CP}(\rho(1450)^+\pi^- \rightarrow \pi^+\pi^-\pi^0)$ in $D^0 \rightarrow \rho(1450)^+\pi^-, \bar{D}^0 \rightarrow \rho(1450)^-\pi^+$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0 ± 50 ± 50	AUBERT	08AO BABR	Table 1, -Col.5/2 × Col.2

$A_{CP}(\rho(1450)^0\pi^0 \rightarrow \pi^+\pi^-\pi^0)$ in $D^0, \bar{D}^0 \rightarrow \rho(1450)^0\pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-17 ± 33 ± 17	AUBERT	08AO BABR	Table 1, -Col.5/2 × Col.2

$A_{CP}(\rho(1450)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0 \rightarrow \rho(1450)^- \pi^+, \bar{D}^0 \rightarrow \rho(1450)^+ \pi^-$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+6±8±3	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(\rho(1700)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0 \rightarrow \rho(1700)^+ \pi^-, \bar{D}^0 \rightarrow \rho(1700)^- \pi^+$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-5±13±5	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(\rho(1700)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow \rho(1700)^0 \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+13±8±3	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(\rho(1700)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0 \rightarrow \rho(1700)^- \pi^+, \bar{D}^0 \rightarrow \rho(1700)^+ \pi^-$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+8±10±5	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(f_0(980) \pi^0 \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow f_0(980) \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0±25±25	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(f_0(1370) \pi^0 \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow f_0(1370) \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+25±13±13	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(f_0(1500) \pi^0 \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow f_0(1500) \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0±13±13	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(f_0(1710) \pi^0 \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow f_0(1710) \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0±17±17	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(f_2(1270) \pi^0 \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow f_2(1270) \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-4±4±4	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(\sigma(400) \pi^0 \rightarrow \pi^+ \pi^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow \sigma(400) \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+6±6±6	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(\text{nonresonant } \pi^+ \pi^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow \text{nonresonant } \pi^+ \pi^- \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-13±19±13	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(K^+ K^- \pi^0)$ in $D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-1.00±1.67±0.25	11 ± 0.11k	AUBERT	08AO BABR	$e^+ e^- \approx 10.6 \text{ GeV}$

$$A_{CP}(K^*(892)^+ K^- \rightarrow K^+ K^- \pi^0) \text{ in } D^0 \rightarrow K^*(892)^+ K^-, \bar{D}^0 \rightarrow K^*(892)^- K^+$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-0.9 \pm 1.2 \pm 0.4$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$$A_{CP}(K^*(1410)^+ K^- \rightarrow K^+ K^- \pi^0) \text{ in } D^0 \rightarrow K^*(1410)^+ K^-, \bar{D}^0 \rightarrow K^*(1410)^- K^+$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-21 \pm 23 \pm 8$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$$A_{CP}((K^+ \pi^0)_S K^- \rightarrow K^+ K^- \pi^0) \text{ in } D^0 \rightarrow (K^+ \pi^0)_S K^-, \bar{D}^0 \rightarrow (K^- \pi^0)_S K^+$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$+7 \pm 15 \pm 3$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$$A_{CP}(\phi(1020) \pi^0 \rightarrow K^+ K^- \pi^0) \text{ in } D^0, \bar{D}^0 \rightarrow \phi(1020) \pi^0$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$+1.1 \pm 2.1 \pm 0.5$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$$A_{CP}(f_0(980) \pi^0 \rightarrow K^+ K^- \pi^0) \text{ in } D^0, \bar{D}^0 \rightarrow f_0(980) \pi^0$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-3 \pm 19 \pm 1$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$$A_{CP}(a_0(980)^0 \pi^0 \rightarrow K^+ K^- \pi^0) \text{ in } D^0, \bar{D}^0 \rightarrow a_0(980)^0 \pi^0$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-5 \pm 16 \pm 2$	¹⁷⁷ AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

¹⁷⁷ This AUBERT 08AO value is obtained when the $a_0(980)^0$ replaces the $f_0(980)$ in the fit.

$$A_{CP}(f_2'(1525) \pi^0 \rightarrow K^+ K^- \pi^0) \text{ in } D^0, \bar{D}^0 \rightarrow f_2'(1525) \pi^0$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$0 \pm 50 \pm 150$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$$A_{CP}(K^*(892)^- K^+ \rightarrow K^+ K^- \pi^0) \text{ in } D^0 \rightarrow K^*(892)^- K^+, \bar{D}^0 \rightarrow K^*(892)^+ K^-$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-5 \pm 4 \pm 1$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$$A_{CP}(K^*(1410)^- K^+ \rightarrow K^+ K^- \pi^0) \text{ in } D^0 \rightarrow K^*(1410)^- K^+, \bar{D}^0 \rightarrow K^*(1410)^+ K^-$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-17 \pm 28 \pm 7$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$$A_{CP}((K^- \pi^0)_S K^+ \rightarrow K^+ K^- \pi^0) \text{ in } D^0 \rightarrow (K^- \pi^0)_S K^+, \bar{D}^0 \rightarrow (K^+ \pi^0)_S K^-$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-7 \pm 40 \pm 8$	AUBERT	08AO BABR	Table 1, -Col.5/2×Col.2

$A_{CP}(K_S^0 \phi)$ in $D^0, \bar{D}^0 \rightarrow K_S^0 \phi$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-2.8 ± 9.4	BARTELT 95	CLE2	$-0.182 < A_{CP} < +0.126$ (90%CL)

$A_{CP}(K_S^0 \pi^0)$ in $D^0, \bar{D}^0 \rightarrow K_S^0 \pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$+0.1 \pm 1.3$	9099	BONVICINI 01	CLE2	$e^+ e^- \approx 10.6$ GeV
-1.8 ± 3.0		BARTELT 95	CLE2	See BONVICINI 01

$A_{CP}(K^\mp \pi^\pm)$ in $D^0 \rightarrow K^- \pi^+, \bar{D}^0 \rightarrow K^+ \pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.1 ± 0.7 OUR AVERAGE				
$+0.5 \pm 0.4 \pm 0.9$	150k	MENDEZ 10	CLEO	$e^+ e^-$ at 3774 MeV
$-0.4 \pm 0.5 \pm 0.9$		DOBBS 07	CLEO	$e^+ e^-$ at $\psi(3770)$

$A_{CP}(K^\pm \pi^\mp)$ in $D^0 \rightarrow K^+ \pi^-, \bar{D}^0 \rightarrow K^- \pi^+$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 3.2 OUR AVERAGE				
$-2.1 \pm 5.2 \pm 1.5$	4030 ± 90	AUBERT 07W	BABR	$e^+ e^- \approx 10.6$ GeV
$+2.3 \pm 4.7$	4024 ± 88	178 ZHANG 06	BELL	$e^+ e^-$
$+18 \pm 14 \pm 4$		179 LINK 05H	FOCS	γ nucleus
$+9.5 \pm 6.1 \pm 8.3$		180 AUBERT 03Z	BABR	$e^+ e^-$, 10.6 GeV
$+2 \pm 19 \pm 20$	45	181 GODANG 00	CLE2	$-0.43 < A_{CP} < +0.34$ (95%CL)

• • • We do not use the following data for averages, fits, limits, etc. • • •

-8.0 ± 7.7 845 ± 40 182 LI 05A BELL See ZHANG 06

178 This ZHANG 06 result allows mixing.

179 This LINK 05H result assumes no mixing. If mixing is allowed, it becomes $0.13^{+0.33}_{-0.25} \pm 0.10$.

180 This AUBERT 03Z limit assumes no mixing. If mixing is allowed, the 95% confidence-level interval is $(-2.8 < A_D < 4.9) \times 10^{-3}$.

181 This GODANG 00 result assumes no D^0 - \bar{D}^0 mixing; it becomes $-0.01^{+0.16}_{-0.17} \pm 0.01$ when mixing is allowed.

182 This LI 05A result allows mixing.

$A_{CP}(K^\mp \pi^\pm \pi^0)$ in $D^0 \rightarrow K^- \pi^+ \pi^0, \bar{D}^0 \rightarrow K^+ \pi^- \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.2 ± 0.9 OUR AVERAGE			
$+0.2 \pm 0.4 \pm 0.8$	DOBBS 07	CLEO	$e^+ e^-$ at $\psi(3770)$
-3.1 ± 8.6	183 KOPP 01	CLE2	$e^+ e^- \approx 10.6$ GeV

183 KOPP 01 fits separately the D^0 and \bar{D}^0 Dalitz plots and then calculates the integrated difference of normalized densities divided by the integrated sum.

$A_{CP}(K^\pm \pi^\mp \pi^0)$ in $D^0 \rightarrow K^+ \pi^- \pi^0, \bar{D}^0 \rightarrow K^- \pi^+ \pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0 ± 5 OUR AVERAGE				
-0.6 ± 5.3	1978 ± 104	TIAN 05	BELL	$e^+ e^- \approx \Upsilon(4S)$
$+9 \pm 25 \pm 22$	38	BRANDENB... 01	CLE2	$e^+ e^- \approx \Upsilon(4S)$

$A_{CP}(K_S^0 \pi^+ \pi^-)$ in $D^0, \bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.9 \pm 2.1_{-5.7}^{+1.6}$	4854	184 ASNER	04A CLEO	$e^+ e^- \approx 10$ GeV

¹⁸⁴ This is the overall result of ASNER 04A; CP -violating limits are also given below for each of the 10 resonant submodes found in an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots. These limits range from $< 3.5 \times 10^{-4}$ to 28.4×10^{-4} at 95% CL.

$A_{CP}(K^*(892)^{\mp} \pi^{\pm} \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow K^{*-} \pi^+, \bar{D}^0 \rightarrow K^{*+} \pi^-$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.5	95	185 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁸⁵ This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K^*(892)^{\pm} \pi^{\mp} \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow K^{*+} \pi^-, \bar{D}^0 \rightarrow K^{*-} \pi^+$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 7.8	95	186 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁸⁶ This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K_S^0 \rho^0 \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow \bar{K}^0 \rho^0, \bar{D}^0 \rightarrow K^0 \rho^0$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.8	95	187 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁸⁷ This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K_S^0 \omega \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow \bar{K}^0 \omega, \bar{D}^0 \rightarrow K^0 \omega$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 9.2	95	188 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁸⁸ This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K_S^0 f_0(980) \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow \bar{K}^0 f_0(980), \bar{D}^0 \rightarrow K^0 f_0(980)$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 6.8	95	189 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁸⁹ This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K_S^0 f_2(1270) \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow \bar{K}^0 f_2(1270), \bar{D}^0 \rightarrow K^0 f_2(1270)$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 13.5	95	190 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁹⁰ This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K_S^0 f_0(1370) \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow \bar{K}^0 f_0(1370), \bar{D}^0 \rightarrow K^0 f_0(1370)$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<25.5	95	191 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁹¹This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K_0^*(1430)^{\mp} \pi^{\pm} \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow K_0^*(1430)^- \pi^+, \bar{D}^0 \rightarrow K_0^*(1430)^+ \pi^-$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<9.0	95	192 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁹²This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K_2^*(1430)^{\mp} \pi^{\pm} \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow K_2^*(1430)^- \pi^+, \bar{D}^0 \rightarrow K_2^*(1430)^+ \pi^-$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6.5	95	193 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁹³This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K^*(1680)^{\mp} \pi^{\pm} \rightarrow K_S^0 \pi^+ \pi^-)$ in $D^0 \rightarrow K^*(1680)^- \pi^+, \bar{D}^0 \rightarrow K^*(1680)^+ \pi^-$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<28.4	95	194 ASNER	04A CLEO	Dalitz fit, 4854 $D^0 + \bar{D}^0$ evts

¹⁹⁴This ASNER 04A limit comes from an amplitude analysis of the D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ Dalitz plots.

$A_{CP}(K^- \pi^+ \pi^+ \pi^-)$ in $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-, \bar{D}^0 \rightarrow K^+ \pi^- \pi^- \pi^+$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+0.7±0.5±0.9	DOBBS	07 CLEO	$e^+ e^-$ at $\psi(3770)$

$A_{CP}(K^{\pm} \pi^{\mp} \pi^+ \pi^-)$ in $D^0 \rightarrow K^+ \pi^- \pi^+ \pi^-, \bar{D}^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-1.8±4.4	1721 ± 75	TIAN	05 BELL	$e^+ e^- \approx \Upsilon(4S)$

$A_{CP}(K^+ K^- \pi^+ \pi^-)$ in $D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^+ \pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-8.2±5.6±4.7	828 ± 46	LINK	05E FOCS	$\gamma A, \bar{E}_{\gamma} \approx 180$ GeV

D^0 - \bar{D}^0 T -VIOLATING DECAY-RATE ASYMMETRIES

D^0 and \bar{D}^0 are distinguished by the charge of the parent D^* : $D^{*+} \rightarrow D^0 \pi^+$ and $D^{*-} \rightarrow \bar{D}^0 \pi^-$. Assuming CPT is good, T violation implies CP violation.

$A_{Tviol}(K^+ K^- \pi^+ \pi^-)$ in $D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^+ \pi^-$

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$ is a T -odd correlation of the K^+ , π^+ , and π^- momenta (evaluated in the D^0 rest frame) for the D^0 . $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$ is the corresponding quantity for the \bar{D}^0 .

$$A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)]$$

would, in the absence of strong phases, test for T violation in D^0 decays (the Γ 's are partial widths). With

$$\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)],$$

the asymmetry $A_{Tviol} \equiv \frac{1}{2}(A_T - \bar{A}_T)$ tests for T violation even with nonzero strong phases.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1 ± 7 OUR AVERAGE				
+ 1.0 ± 5.1 ± 4.4	47k	DEL-AMO-SA..10	BABR	$e^+ e^- \approx 10.6$ GeV
+10 ± 57 ± 37	828 ± 46	LINK	05E FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

D^0 CPT -VIOLATING DECAY-RATE ASYMMETRIES

$A_{CPT}(K^\mp \pi^\pm)$ in $D^0 \rightarrow K^- \pi^+, \bar{D}^0 \rightarrow K^+ \pi^-$

$A_{CPT}(t)$ is defined in terms of the time-dependent decay probabilities $P(D^0 \rightarrow K^- \pi^+)$ and $\bar{P}(\bar{D}^0 \rightarrow K^+ \pi^-)$ by $A_{CPT}(t) = (\bar{P} - P)/(\bar{P} + P)$. For small mixing parameters $x \equiv \Delta m/\Gamma$ and $y \equiv \Delta\Gamma/2\Gamma$ (as is the case), and times t , $A_{CPT}(t)$ reduces to $[y \operatorname{Re} \xi - x \operatorname{Im} \xi] \Gamma t$, where ξ is the CPT -violating parameter.

The following is actually $y \operatorname{Re} \xi - x \operatorname{Im} \xi$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0083 ± 0.0065 ± 0.0041	LINK	03B	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

$D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell$ FORM FACTORS

$r_V \equiv V(0)/A_1(0)$ in $D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell$

VALUE	DOCUMENT ID	TECN	COMMENT
1.71 ± 0.68 ± 0.34	LINK	05B	FOCS $K^*(892)^- \mu^+ \nu_\mu$

$r_2 \equiv A_2(0)/A_1(0)$ in $D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell$

VALUE	DOCUMENT ID	TECN	COMMENT
0.91 ± 0.37 ± 0.10	LINK	05B	FOCS $K^*(892)^- \mu^+ \nu_\mu$

$D^0 \rightarrow K^- / \pi^- \ell^+ \nu_\ell$ FORM FACTORS

$f_+(0)|V_{cs}|$ in $D^0 \rightarrow K^- \ell^+ \nu_\ell$

VALUE	DOCUMENT ID	TECN	COMMENT
0.726 ± 0.008 ± 0.004	BESSION	09	CLEO $K^- e^+ \nu_e$ 3-parameter fit

$$r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell$$

VALUE	DOCUMENT ID	TECN	COMMENT
$-2.65 \pm 0.34 \pm 0.08$	BESSON 09	CLEO	$K^- e^+ \nu_e$ 3-parameter fit

$$r_2 \equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell$$

VALUE	DOCUMENT ID	TECN	COMMENT
$13 \pm 9 \pm 1$	BESSON 09	CLEO	$K^- e^+ \nu_e$ 3-parameter fit

$$f_+(0)|V_{cd}| \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.152 \pm 0.005 \pm 0.001$	BESSON 09	CLEO	$\pi^- e^+ \nu_e$ 3-parameter fit

$$r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell$$

VALUE	DOCUMENT ID	TECN	COMMENT
$-2.80 \pm 0.49 \pm 0.04$	BESSON 09	CLEO	$\pi^- e^+ \nu_e$ 3-parameter fit

$$r_2 \equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell$$

VALUE	DOCUMENT ID	TECN	COMMENT
$6 \pm 3 \pm 0$	BESSON 09	CLEO	$\pi^- e^+ \nu_e$ 3-parameter fit

D⁰ REFERENCES

AALTONEN 10X	PR D82 091105R	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ANASHIN 10A	PL B686 84	V.V. Anashin <i>et al.</i>	(VEPP-4M KEDR Collab.)
ASNER 10	PR D81 052007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BHATTACHAR.10A	PR D81 096008	B. Bhattacharya, C.-W. Chiang, J.L. Rosner	(CHIC+)
DEL-AMO-SA... 10	PR D81 111103R	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
DEL-AMO-SA... 10D	PRL 105 081803	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MENDEZ 10	PR D81 052013	H. Mendez <i>et al.</i>	(CLEO Collab.)
PDG 10	JPG 37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)
PETRIC 10	PR D81 091102R	M. Petric <i>et al.</i>	(BELLE Collab.)
AUBERT 09AI	PR D80 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 09AN	PRL 103 211801	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON 09	PR D80 032005	D. Besson <i>et al.</i>	(CLEO Collab.)
Also	PR D79 052010	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
LOWREY 09	PR D80 031105R	N. Lowrey <i>et al.</i>	(CLEO Collab.)
RUBIN 09	PR D79 097101	P. Rubin <i>et al.</i>	(CLEO Collab.)
ZUPANC 09	PR D80 052006	A. Zupanc <i>et al.</i>	(BELLE Collab.)
AALTONEN 08E	PRL 100 121802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABLIKIM 08L	PL B665 16	M. Ablikim <i>et al.</i>	(BES Collab.)
ARINSTEIN 08	PL B662 102	K. Arinstein <i>et al.</i>	(BELLE Collab.)
ARTUSO 08	PR D77 092003	M. Artuso <i>et al.</i>	(CLEO Collab.)
ASNER 08	PR D78 012001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT 08AL	PR D78 034023	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 08AO	PR D78 051102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 08AZ	PR D78 071101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 08L	PRL 100 051802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 08M	PRL 100 061803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 08U	PR D78 011105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
BITENC 08	PR D77 112003	U. Bitenc <i>et al.</i>	(BELLE Collab.)
BONVICINI 08	PR D77 091106R	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
DOBBS 08	PR D77 112005	S. Dobbs <i>et al.</i>	(CLEO Collab.)
Also	PRL 100 251802	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
GASPERO 08	PR D78 014015	M. Gaspero <i>et al.</i>	(ROMA, CINN, TELA)
HE 08	PRL 100 091801	Q. He <i>et al.</i>	(CLEO Collab.)
PDG 08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
STARIC 08	PL B670 190	M. Staric <i>et al.</i>	(BELLE Collab.)
ABLIKIM 07G	PL B658 1	M. Ablikim <i>et al.</i>	(BES Collab.)
ARTUSO 07A	PRL 99 191801	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT 07AB	PR D76 014018	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 07BG	PR D76 052005	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT	07BJ	PRL 99 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07T	PR D76 011102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07W	PRL 98 211802	B. Aubert <i>et al.</i>	(BABAR Collab.)
CAWLFIELD	07	PRL 98 092002	C. Cawfield <i>et al.</i>	(CLEO Collab.)
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)
LINK	07A	PR D75 052003	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
STARIC	07	PRL 98 211803	M. Staric <i>et al.</i>	(BELLE Collab.)
ZHANG	07B	PRL 99 131803	L.M. Zhang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06O	EPJ C47 31	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06U	PL B643 246	M. Ablikim <i>et al.</i>	(BES Collab.)
ABULENCIA	06X	PR D74 031109R	A. Abulencia <i>et al.</i>	(CDF Collab.)
ADAM	06A	PRL 97 251801	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT,B	06N	PRL 97 221803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06X	PR D74 091102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
CAWLFIELD	06A	PR D74 031108R	C. Cawfield <i>et al.</i>	(CLEO Collab.)
HUANG	06B	PR D74 112005	G.S. Huang <i>et al.</i>	(CLEO Collab.)
PDG	06	JPG 33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
RUBIN	06	PRL 96 081802	P. Rubin <i>et al.</i>	(CLEO Collab.)
WIDHALM	06	PRL 97 061804	L. Widhalm <i>et al.</i>	(BELLE Collab.)
ZHANG	06	PRL 96 151801	L.M. Zhang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05F	PL B622 6	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05P	PL B625 196	M. Ablikim <i>et al.</i>	(BES Collab.)
ACOSTA	05C	PRL 94 122001	D. Acosta <i>et al.</i>	(FNAL CDF Collab.)
ASNER	05	PR D72 012001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT,B	05J	PR D72 052008	B. Aubert <i>et al.</i>	(BABAR Collab.)
BITENC	05	PR D72 071101R	U. Bitenc <i>et al.</i>	(BELLE Collab.)
CAWLFIELD	05	PR D71 077101	C. Cawfield <i>et al.</i>	(CLEO Collab.)
COAN	05	PRL 95 181802	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... HE	05	PR D72 031102R PRL 95 121801	D. Cronin-Hennessy <i>et al.</i> Q. He <i>et al.</i>	(CLEO Collab.) (CLEO Collab.)
Also		PRL 96 199903 (errata.)	Q. He <i>et al.</i>	(CLEO Collab.)
HUANG	05	PRL 94 011802	G.S. Huang <i>et al.</i>	(CLEO Collab.)
KAYIS-TOPAK... LI	05A	PL B626 24 PRL 94 071801	A. Kayis-Topaksu <i>et al.</i> J. Li <i>et al.</i>	(CERN CHORUS Collab.) (BELLE Collab.)
LINK	05	PL B607 51	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05A	PL B607 59	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05B	PL B607 67	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05E	PL B622 239	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05G	PL B610 225	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05H	PL B618 23	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ONENGUT	05	PL B613 105	G. Onengut <i>et al.</i>	(CERN CHORUS Collab.)
TIAN	05	PRL 95 231801	X.C. Tian <i>et al.</i>	(BELLE Collab.)
ABLIKIM	04C	PL B597 39	M. Ablikim <i>et al.</i>	(BEPC BES Collab.)
ABT	04	PL B596 173	I. Abt <i>et al.</i>	(HERA B Collab.)
ASNER	04A	PR D70 091101R	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04Q	PR D69 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04Q	PR D70 091102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04Y	PRL 93 191801	B. Aubert <i>et al.</i>	(BaBar Collab.)
LINK	04B	PL B586 21	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04D	PL B586 191	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
RUBIN	04	PRL 93 111801	P. Rubin <i>et al.</i>	(CLEO Collab.)
TAJIMA	04	PRL 92 101803	O. Tajima <i>et al.</i>	(BELLE Collab.)
ACOSTA	03F	PR D68 091101R	D. Acosta <i>et al.</i>	(CDF Collab.)
AUBERT	03P	PRL 91 121801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03Z	PRL 91 171801	B. Aubert <i>et al.</i>	(BaBar Collab.)
COAN	03	PRL 90 101801	T.E. Coan <i>et al.</i>	(CLEO Collab.)
LINK	03	PL B555 167	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03B	PL B556 7	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03G	PL B575 190	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABE	02I	PRL 88 162001	K. Abe <i>et al.</i>	(KEK BELLE Collab.)
CSORNA	02	PR D65 092001	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
LINK	02F	PL B537 192	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
MURAMATSU	02	PRL 89 251802	H. Muramatsu <i>et al.</i>	(CLEO Collab.)
Also		PRL 90 059901 (erratum)	H. Muramatsu <i>et al.</i>	(CLEO Collab.)
AITALA	01C	PRL 86 3969	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	01D	PR D64 112003	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BONVICINI	01	PR D63 071101R	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BRANDENB... DYTMAN	01	PRL 87 071802 PR D64 111101R	G. Brandenburg <i>et al.</i> S.A. Dytman <i>et al.</i>	(CLEO Collab.) (CLEO Collab.)
KOPP	01	PR D63 092001	S. Kopp <i>et al.</i>	(CLEO Collab.)
KUSHNIR... KUSHNIR...	01	PRL 86 5243	A. Kushnirenko <i>et al.</i>	(FNAL SELEX Collab.)

LINK	01	PRL 86 2955	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
BAI	00C	PR D62 052001	J.Z. Bai <i>et al.</i>	(BEP C BES Collab.)
GODANG	00	PRL 84 5038	R. Godang <i>et al.</i>	(CLEO Collab.)
JUN	00	PRL 84 1857	S.Y. Jun <i>et al.</i>	(FNAL SELEX Collab.)
LINK	00	PL B485 62	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	00B	PL B491 232	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
Also		PL B495 443 (erratum)	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
PRIPSTEIN	00	PR D61 032005	D. Pripstein <i>et al.</i>	(FNAL E789 Collab.)
AITALA	99E	PRL 83 32	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99G	PL B462 401	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AITALA	98	PR D57 13	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	98C	PL B421 405	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	98D	PL B423 185	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ARTUSO	98	PRL 80 3193	M. Artuso <i>et al.</i>	(CLEO Collab.)
ASNER	98	PR D58 092001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARATE	98W	PL B436 211	R. Barate <i>et al.</i>	(ALEPH Collab.)
COAN	98	PRL 80 1150	T.E. Coan <i>et al.</i>	(CLEO Collab.)
PDG	98	EPJ C3 1	C. Caso <i>et al.</i>	
ADAMOVICH	97	PL B408 469	M.I. Adamovich <i>et al.</i>	(CERN BEATRICE Collab.)
BARATE	97C	PL B403 367	R. Barate <i>et al.</i>	(ALEPH Collab.)
AITALA	96C	PRL 77 2384	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ALBRECHT	96C	PL B374 249	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXOPOU...	96	PRL 77 2380	T. Alexopoulos <i>et al.</i>	(FNAL E771 Collab.)
ASNER	96B	PR D54 4211	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARISH	96	PL B373 334	B.C. Barish <i>et al.</i>	(CLEO Collab.)
FRABETTI	96B	PL B382 312	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FREYBERGER	96	PRL 76 3065	A. Freyberger <i>et al.</i>	(CLEO Collab.)
Also		PRL 77 2147 (erratum)	A. Freyberger <i>et al.</i>	(CLEO Collab.)
KUBOTA	96B	PR D54 2994	Y. Kubota <i>et al.</i>	(CLEO Collab.)
ADAMOVICH	95	PL B353 563	M.I. Adamovich <i>et al.</i>	(CERN BEATRICE Collab.)
BARTELT	95	PR D52 4860	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
BUTLER	95	PR D52 2656	F. Butler <i>et al.</i>	(CLEO Collab.)
FRABETTI	95C	PL B354 486	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	95G	PL B364 127	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	94	PL B324 249	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	94F	PL B340 125	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	94I	ZPHY C64 375	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
FRABETTI	94C	PL B321 295	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	94D	PL B323 459	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	94G	PL B331 217	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	94J	PL B340 254	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	94	PL B336 605	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
MISHRA	94	PR D50 R9	C.S. Mishra <i>et al.</i>	(FNAL E789 Collab.)
AKERIB	93	PRL 71 3070	D.S. Akerib <i>et al.</i>	(CLEO Collab.)
ALBRECHT	93D	PL B308 435	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	93	PR D48 56	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BEAN	93C	PL B317 647	A. Bean <i>et al.</i>	(CLEO Collab.)
FRABETTI	93I	PL B315 203	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	93B	PL B313 260	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
PROCARIO	93B	PR D48 4007	M. Procaro <i>et al.</i>	(CLEO Collab.)
SELEN	93	PRL 71 1973	M.A. Selen <i>et al.</i>	(CLEO Collab.)
ADAMOVICH	92	PL B280 163	M.I. Adamovich <i>et al.</i>	(CERN WA82 Collab.)
ALBRECHT	92P	ZPHY C56 7	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	92B	PR D46 R1	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	92C	PR D46 1941	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
Also		ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
COFFMAN	92B	PR D45 2196	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
Also		PRL 64 2615	J. Adler <i>et al.</i>	(Mark III Collab.)
FRABETTI	92	PL B281 167	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	92B	PL B286 195	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALVAREZ	91B	ZPHY C50 11	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
AMMAR	91	PR D44 3383	R. Ammar <i>et al.</i>	(CLEO Collab.)
ANJOS	91	PR D43 R635	J.C. Anjos <i>et al.</i>	(FNAL-TPS Collab.)
ANJOS	91D	PR D44 R3371	J.C. Anjos <i>et al.</i>	(FNAL-TPS Collab.)
BAI	91	PRL 66 1011	Z. Bai <i>et al.</i>	(Mark III Collab.)
COFFMAN	91	PL B263 135	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
CRAWFORD	91B	PR D44 3394	G. Crawford <i>et al.</i>	(CLEO Collab.)
DECAMP	91J	PL B266 218	D. Decamp <i>et al.</i>	(ALEPH Collab.)

FRABETTI	91	PL B263 584	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KINOSHITA	91	PR D43 2836	K. Kinoshita <i>et al.</i>	(CLEO Collab.)
KODAMA	91	PRL 66 1819	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	90C	ZPHY C46 9	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	90	PRL 65 1184	J. Alexander <i>et al.</i>	(CLEO Collab.)
ALVAREZ	90	ZPHY C47 539	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ANJOS	90D	PR D42 2414	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
ADLER	89	PRL 62 1821	J. Adler <i>et al.</i>	(Mark III Collab.)
ADLER	89C	PR D40 906	J. Adler <i>et al.</i>	(Mark III Collab.)
ALBRECHT	89D	ZPHY C43 181	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	89F	PRL 62 1587	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ABACHI	88	PL B205 411	S. Abachi <i>et al.</i>	(HRS Collab.)
ADLER	88	PR D37 2023	J. Adler <i>et al.</i>	(Mark III Collab.)
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)
ALBRECHT	88G	PL B209 380	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88I	PL B210 267	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	88C	PRL 60 1239	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BORTOLETTO	88	PR D37 1719	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also		PR D39 1471 (erratum)	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
HAAS	88	PRL 60 1614	S. Haas <i>et al.</i>	(CLEO Collab.)
RAAB	88	PR D37 2391	J.R. Raab <i>et al.</i>	(FNAL E691 Collab.)
ADAMOVICH	87	EPL 4 887	M.I. Adamovich <i>et al.</i>	(Photon Emulsion Collab.)
ADLER	87	PL B196 107	J. Adler <i>et al.</i>	(Mark III Collab.)
AGUILAR-...	87E	ZPHY C36 551	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		ZPHY C40 321	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
AGUILAR-...	87F	ZPHY C36 559	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		ZPHY C38 520 (erratum)	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
BARLAG	87B	ZPHY C37 17	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
BECKER	87C	PL B193 147	J.J. Becker <i>et al.</i>	(Mark III Collab.)
Also		PL B198 590 (erratum)	J.J. Becker <i>et al.</i>	(Mark III Collab.)
PALKA	87	PL B189 238	H. Palka <i>et al.</i>	(ACCMOR Collab.)
RILES	87	PR D35 2914	K. Riles <i>et al.</i>	(Mark II Collab.)
BAILEY	86	ZPHY C30 51	R. Bailey <i>et al.</i>	(ACCMOR Collab.)
BEBEK	86	PRL 56 1893	C. Bebek <i>et al.</i>	(CLEO Collab.)
LOUIS	86	PRL 56 1027	W.C. Louis <i>et al.</i>	(PRIN, CHIC, ISU)
ALBRECHT	85B	PL 158B 525	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85F	PL 150B 235	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AUBERT	85	PL 155B 461	J.J. Aubert <i>et al.</i>	(EMC Collab.)
BALTRUSAITIS...	85E	PRL 55 150	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BENVENUTI	85	PL 158B 531	A.C. Benvenuti <i>et al.</i>	(BCDMS Collab.)
ADAMOVICH	84B	PL 140B 123	M.I. Adamovich <i>et al.</i>	(CERN WA58 Collab.)
DERRICK	84	PRL 53 1971	M. Derrick <i>et al.</i>	(HRS Collab.)
SUMMERS	84	PRL 52 410	D.J. Summers <i>et al.</i>	(UCSB, CARL, COLO+)
BAILEY	83B	PL 132B 237	R. Bailey <i>et al.</i>	(ACCMOR Collab.)
BODEK	82	PL 113B 82	A. Bodek <i>et al.</i>	(ROCH, CIT, CHIC, FNAL+)
FIORINO	81	LNC 30 166	A. Fiorino <i>et al.</i>	(Photon-Emul/Omega-Photon)
SCHINDLER	81	PR D24 78	R.H. Schindler <i>et al.</i>	(Mark II Collab.)
TRILLING	81	PRPL 75 57	G.H. Trilling	(LBL, UCB) J
ASTON	80E	PL 94B 113	D. Aston <i>et al.</i>	(BONN, CERN, EPOL, GLAS+)
VERY	80	PRL 44 1309	P. Avery <i>et al.</i>	(ILL, FNAL, COLU)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		
ABRAMS	79D	PRL 43 481	G.S. Abrams <i>et al.</i>	(Mark II Collab.)
ATIYA	79	PRL 43 414	M.S. Atiya <i>et al.</i>	(COLU, ILL, FNAL)
BALTAY	78C	PRL 41 73	C. Baltay <i>et al.</i>	(COLU, BNL)
VUILLEMIN	78	PRL 41 1149	V. Vuillemin <i>et al.</i>	(LGW Collab.)
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)
PERUZZI	77	PRL 39 1301	I. Peruzzi <i>et al.</i>	(LGW Collab.)
PICCOLO	77	PL 70B 260	M. Piccolo <i>et al.</i>	(Mark I Collab.)
GOLDHABER	76	PRL 37 255	G. Goldhaber <i>et al.</i>	(Mark I Collab.)

OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	J.D. Richman, P.R. Burchat	(UCSB, STAN)
ROSNER	95	CNPP 21 369	J. Rosner	(CHIC)