

$\psi(4160)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

### $\psi(4160)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4153 <math>\pm</math> 3 OUR ESTIMATE</b>			
<b>4191.7 <math>\pm</math> 6.5</b>	<sup>1</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4193 $\pm$ 7	<sup>2</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4151 $\pm$ 4	<sup>3</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4155 $\pm$ 5	<sup>4</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4159 $\pm$ 20	BRANDELIK	78C DASP	$e^+e^-$
<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$ , $\psi(4040)$ , $\psi(4160)$ , and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$ .			
<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$ , $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.			
<sup>4</sup> From a fit to BES (BAI 02C) data.			

### $\psi(4160)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>103 <math>\pm</math> 8 OUR ESTIMATE</b>			
<b>71.8 <math>\pm</math> 12.3</b>	<sup>5</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
79 $\pm$ 14	<sup>6</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
107 $\pm$ 10	<sup>7</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
107 $\pm$ 16	<sup>8</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
78 $\pm$ 20	BRANDELIK	78C DASP	$e^+e^-$
<sup>5</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$ , $\psi(4040)$ , $\psi(4160)$ , and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$ .			
<sup>6</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$ , $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
<sup>7</sup> From a fit to Crystal Ball (OSTERHELD 86) data.			
<sup>8</sup> From a fit to BES (BAI 02C) data.			

**$\psi(4160)$  DECAY MODES**

Due to the complexity of the  $c\bar{c}$  threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective  $\sqrt{s}$  near this particle’s central mass value, more (less) than  $2\sigma$  above zero, without regard to any peaking behavior in  $\sqrt{s}$  or absence thereof. See mode listing(s) for details and references.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $e^+e^-$	$(8.1 \pm 0.9) \times 10^{-6}$	
$\Gamma_2$ $D\bar{D}$	seen	
$\Gamma_3$ $D^0\bar{D}^0$	seen	
$\Gamma_4$ $D^+D^-$	seen	
$\Gamma_5$ $D^*\bar{D} + \text{c.c.}$	seen	
$\Gamma_6$ $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_7$ $D^*(2010)^+D^- + \text{c.c.}$	seen	
$\Gamma_8$ $D^*\bar{D}^*$	seen	
$\Gamma_9$ $D^*(2007)^0\bar{D}^*(2007)^0$	seen	
$\Gamma_{10}$ $D^*(2010)^+D^*(2010)^-$	seen	
$\Gamma_{11}$ $D^0D^-\pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0\bar{D}^0 + \text{c.c.}$ , $D^*(2010)^+D^- + \text{c.c.}$ )	not seen	
$\Gamma_{12}$ $D\bar{D}^*\pi + \text{c.c.}$ (excl. $D^*\bar{D}^*$ )	seen	
$\Gamma_{13}$ $D^0D^{*-}\pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+D^*(2010)^-$ )	not seen	
$\Gamma_{14}$ $D_s^+D_s^-$	not seen	
$\Gamma_{15}$ $D_s^{*+}D_s^- + \text{c.c.}$	seen	
$\Gamma_{16}$ $J/\psi\pi^+\pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{17}$ $J/\psi\pi^0\pi^0$	$< 3 \times 10^{-3}$	90%
$\Gamma_{18}$ $J/\psi K^+K^-$	$< 2 \times 10^{-3}$	90%
$\Gamma_{19}$ $J/\psi\eta$	$< 8 \times 10^{-3}$	90%
$\Gamma_{20}$ $J/\psi\pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{21}$ $J/\psi\eta'$	$< 5 \times 10^{-3}$	90%
$\Gamma_{22}$ $J/\psi\pi^+\pi^-\pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{23}$ $\psi(2S)\pi^+\pi^-$	$< 4 \times 10^{-3}$	90%
$\Gamma_{24}$ $\chi_{c1}\gamma$	$< 7 \times 10^{-3}$	90%
$\Gamma_{25}$ $\chi_{c2}\gamma$	$< 1.3 \%$	90%
$\Gamma_{26}$ $\chi_{c1}\pi^+\pi^-\pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{27}$ $\chi_{c2}\pi^+\pi^-\pi^0$	$< 8 \times 10^{-3}$	90%
$\Gamma_{28}$ $\phi\pi^+\pi^-$	$< 2 \times 10^{-3}$	90%

## $\psi(4160)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$					$\Gamma_1$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>0.83±0.07 OUR ESTIMATE</b>					
<b>0.48±0.22</b>	<sup>9</sup> ABLIKIM	08D	BES2	$e^+e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.4 to 1.1	<sup>10</sup> MO	10	RVUE	$e^+e^- \rightarrow$ hadrons	
0.83±0.08	<sup>11</sup> SETH	05A	RVUE	$e^+e^- \rightarrow$ hadrons	
0.84±0.13	<sup>12</sup> SETH	05A	RVUE	$e^+e^- \rightarrow$ hadrons	
0.77±0.23	BRANDELIK	78C	DASP	$e^+e^-$	
<sup>9</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$ , $\psi(4040)$ , $\psi(4160)$ , and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$ .					
<sup>10</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$ , $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+e^-$ partial widths. We quote only the range of values.					
<sup>11</sup> From a fit to Crystal Ball (OSTERHELD 86) data.					
<sup>12</sup> From a fit to BES (BAI 02C) data.					

## $\psi(4160)$ BRANCHING RATIOS

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$					$\Gamma_2/\Gamma_8$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>0.02±0.03±0.02</b>	AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$	
$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$					$\Gamma_3/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^0\bar{D}^0$		
<b>seen</b>	PAKHLOVA	08	BELL	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$	
$\Gamma(D^+D^-)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^+D^-$		
<b>seen</b>	PAKHLOVA	08	BELL	$e^+e^- \rightarrow D^+D^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^+D^-\gamma$	
$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_6/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>seen</b>	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$	
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^0$		

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^-$
<b>seen</b>	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$

$\Gamma(D^* \bar{D} + \text{c.c.})/\Gamma(D^* \bar{D}^*)$   $\Gamma_5/\Gamma_8$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.34 ± 0.14 ± 0.05</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$

$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$

$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$
<b>seen</b>	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.))}/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*))/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$

$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
<b>not seen</b>	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
<b>not seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
<b>seen</b>	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$

$\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$				$\Gamma_{16}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$				$\Gamma_{17}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(J/\psi K^+ K^-)/\Gamma_{\text{total}}$				$\Gamma_{18}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$				$\Gamma_{19}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(J/\psi \pi^0)/\Gamma_{\text{total}}$				$\Gamma_{20}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(J/\psi \eta')/\Gamma_{\text{total}}$				$\Gamma_{21}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(J/\psi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$				$\Gamma_{22}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(\psi(2S) \pi^+ \pi^-)/\Gamma_{\text{total}}$				$\Gamma_{23}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(\chi_{c1} \gamma)/\Gamma_{\text{total}}$				$\Gamma_{24}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<7	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(\chi_{c2} \gamma)/\Gamma_{\text{total}}$				$\Gamma_{25}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<13	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons
$\Gamma(\chi_{c1} \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$				$\Gamma_{26}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN 06	CLEO	4.12-4.2 $e^+ e^- \rightarrow$ hadrons

$\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{27}/\Gamma$
VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<8	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{28}/\Gamma$
VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<2	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons

### $\psi(4160)$ REFERENCES

PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101R	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103R	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)