D^0 – THIS IS PART 2 OF 3

To reduce the size of this section's PostScript file, we have divided it into three PostScript files. We present the following index:

PART 1

Page #	Section name
1	Mass
1	Mass differences
2	Mean life
3	Mean life difference/average
3	Decay modes
11	Branching ratios
11	Inclusive modes
13	Semileptonic modes

PART 2

Page #	Section name
	Branching ratios (continued)
16	Hadronic modes with a $\overline{K}$ or $\overline{K}K\overline{K}$

PART 3

Page #	Section name
	Branching ratios (continued)
30	Pionic modes
31	Hadronic modes with a $K\overline{K}$ pair
37	Rare or forbidden modes
43	CP-violating decay-rate asymmetries
43	Production cross section at $\psi(3770)$
44	References

Hadronic modes with a $\overline{K}$ or $\overline{K}K\overline{K}$

$\Gamma(K^-\pi^+)/\Gamma_{total}$ Γ_{20}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0382±0.0012 OUR FIT Error includes scale factor of 1.1.					
0.0386±0.0014 OUR AVERAGE					
0.045 ±0.006 ±0.004		[29] ALBRECHT	94	ARG	$e^+e^- \approx \Upsilon(4S)$
0.0341±0.0012±0.0028	1173	[30] ALBRECHT	94F	ARG	$e^+e^- \approx \Upsilon(4S)$
0.0391±0.0008±0.0017	4208	[30,31] AKERIB	93	CLEO	$e^+e^- \approx \Upsilon(4S)$
0.0362±0.0034±0.0044		[30] DECAMP	91J	ALEP	From Z decays
0.045 ±0.008 ±0.005	56	[30] ABACHI	88	HRS	e^+e^- 29 GeV
0.042 ±0.004 ±0.004	930	ADLER	88C	MRK3	e^+e^- 3.77 GeV
0.041 ±0.006	263	[32] SCHINDLER	81	MRK2	e^+e^- 3.771 GeV
0.043 ±0.010	130	[33] PERUZZI	77	MRK1	e^+e^- 3.77 GeV

[29] ALBRECHT 94 uses D^0 mesons from $\overline{B}^0 \to D^{*+}\ell^-\overline{\nu}_\ell$ decays. This is a different set of events than used by ALBRECHT 94F.

[30] ABACHI 88, DECAMP 91J, AKERIB 93, and ALBRECHT 94F use $D^*(2010)^+ \to D^0\pi^+$ decays. The π^+ is both slow and of low p_T with respect to the event thrust axis ($\approx D^{*+}$ direction). The excess number of such π^+'s over background gives the number of $D^*(2010)^+ \to D^0\pi^+$ events, and the fraction with $D^0 \to K^-\pi^+$ gives the $D^0 \to K^-\pi^+$ branching fraction.

[31] Radiative corrections increase this AKERIB 93 value to 0.0395 ± 0.0008 ± 0.0017.

[32] SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.24 ± 0.02 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

[33] PERUZZI 77 (MARK-1) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.25 ± 0.05 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

$\Gamma(\overline{K}^0\pi^0)/\Gamma(K^-\pi^+)$ Γ_{21}/Γ_{20}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.55±0.06 OUR FIT Error includes scale factor of 1.1.				
1.36±0.23±0.22	119	ANJOS	92B E691	γ Be 80–240 GeV

$\Gamma(\overline{K}^0\pi^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{21}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.390±0.031 OUR FIT					
0.378±0.033 OUR AVERAGE					
0.44 ±0.02 ±0.05	1942	PROCARIO	93B	CLEO	e^+e^- 10.36–10.7 GeV
0.34 ±0.04 ±0.02	92	[34] ALBRECHT	92P	ARG	$e^+e^- \approx$ 10 GeV
0.36 ±0.04 ±0.08	104	KINOSHITA	91	CLEO	$e^+e^- \sim$ 10.7 GeV

[34] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^0\pi^+\pi^-)/\Gamma_{total}$ Γ_{22}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.054 ±0.004 OUR FIT Error includes scale factor of 1.2.					
0.055 ±0.005 OUR AVERAGE					
0.0503±0.0039±0.0049	284	[35] ALBRECHT	94F	ARG	$e^+e^- \approx \Upsilon(4S)$
0.064 ±0.005 ±0.010		ADLER	87	MRK3	e^+e^- 3.77 GeV
0.052 ±0.016	32	[36] SCHINDLER	81	MRK2	e^+e^- 3.771 GeV
0.079 ±0.023	28	[37] PERUZZI	77	MRK1	e^+e^- 3.77 GeV

[35] See the footnote on the ALBRECHT 94F measurement of $\Gamma(K^-\pi^+)/\Gamma_{\text{total}}$ for the method used.

[36] SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.30 ± 0.08 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

[37] PERUZZI 77 (MARK-1) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.46 ± 0.12 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

$\Gamma(\overline{K}^0\pi^+\pi^-)/\Gamma(K^-\pi^+)$ Γ_{22}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.42±0.11 OUR FIT Error includes scale factor of 1.2.					
1.65±0.17 OUR AVERAGE					
1.61±0.10±0.15	856	FRABETTI	94J	E687	γ Be $\overline{E}_\gamma$=220 GeV
1.7 ±0.8	35	AVERY	80	SPEC	$\gamma N \to D^{*+}$
2.8 ±1.0	116	PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV

$\Gamma(\overline{K}^0\rho^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{23}/Γ_{22}

VALUE	DOCUMENT ID		TECN	COMMENT
0.223±0.027 OUR AVERAGE Error includes scale factor of 1.2.				
0.350±0.028±0.067	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.227±0.032±0.009	ALBRECHT	93D	ARG	$e^+e^- \approx$ 10 GeV
0.215±0.051±0.037	ANJOS	93	E691	γ Be 90–260 GeV
0.20 ±0.06 ±0.03	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.12 ±0.01 ±0.07	ADLER	87	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0 f_0(980))/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{74}/Γ_{22}

Unseen decay modes of the $f_0(980)$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.105±0.029 OUR AVERAGE				
0.131±0.031±0.034	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.088±0.035±0.012	ALBRECHT	93D	ARG	$e^+e^- \approx$ 10 GeV

$\Gamma(\overline{K}^0 f_2(1270))/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{78}/Γ_{22}

Unseen decay modes of the $f_2(1270)$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.076±0.028 OUR AVERAGE				
0.065±0.025±0.030	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.088±0.037±0.014	ALBRECHT	93D	ARG	$e^+e^- \approx$ 10 GeV

$\Gamma(\overline{K}^0 f_0(1370))/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{80}/Γ_{22}

Unseen decay modes of the $f_0(1370)$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.13 ±0.04 OUR AVERAGE				
0.123±0.035±0.049	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.131±0.045±0.021	ALBRECHT	93D	ARG	$e^+e^- \approx$ 10 GeV

$\Gamma(K^*(892)^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{81}/Γ_{22}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.93 ±0.04 OUR FIT	Error includes scale factor of 1.1.				
0.96 ±0.04 OUR AVERAGE					
0.938±0.054±0.038		FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
1.08 ±0.063±0.045		ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV
0.720±0.145±0.185		ANJOS	93	E691	γ Be 90–260 GeV
0.96 ±0.12 ±0.075		FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.84 ±0.06 ±0.08		ADLER	87	MRK3	e^+e^- 3.77 GeV
$1.05\ ^{+0.23}_{-0.26}\ ^{+0.07}_{-0.09}$	25	SCHINDLER	81	MRK2	e^+e^- 3.771 GeV

$\Gamma(K_0^*(1430)^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{103}/Γ_{22}

Unseen decay modes of the $\overline{K}_0^*(1430)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.19 ±0.05 OUR AVERAGE				
0.176±0.044±0.047	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.208±0.055±0.034	ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(K_2^*(1430)^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{104}/Γ_{22}

Unseen decay modes of the $\overline{K}_2^*(1430)^-$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.15	90	ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(\overline{K}^0\pi^+\pi^-$ **nonresonant**$)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{29}/Γ_{22}

VALUE	DOCUMENT ID		TECN	COMMENT
0.27 ±0.04 OUR AVERAGE				
0.263±0.024±0.041	ANJOS	93	E691	γ Be 90–260 GeV
0.26 ±0.08 ±0.05	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.33 ±0.05 ±0.10	ADLER	87	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.139±0.009 OUR FIT	Error includes scale factor of 1.3.				
0.131±0.016 OUR AVERAGE					
0.133±0.012±0.013	931	ADLER	88C	MRK3	e^+e^- 3.77 GeV
0.117±0.043	37	[38] SCHINDLER	81	MRK2	e^+e^- 3.771 GeV

[38] SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.68 ± 0.23 nb. We use the MARK-3 (ADLER 88C) value of σ = 5.8 ± 0.5 ± 0.6 nb.

$\Gamma(K^-\pi^+\pi^0)/\Gamma(K^-\pi^+)$ Γ_{30}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
3.63±0.24 OUR FIT	Error includes scale factor of 1.4.				
3.47±0.30 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.				
3.81±0.07±0.26	10k	BARISH	96	CLEO	$e^+e^- \approx \Upsilon(4S)$
3.04±0.16±0.34	931	[39] ALBRECHT	92P	ARG	$e^+e^- \approx$ 10 GeV
4.0 ±0.9 ±1.0	69	ALVAREZ	91B	NA14	Photoproduction
2.8 ±0.14±0.52	1050	KINOSHITA	91	CLEO	$e^+e^- \sim$ 10.7 GeV
4.2 ±1.4	41	SUMMERS	84	E691	Photoproduction

[39] This value is calculated from numbers in Table 1 of ALBRECHT 92P.



$\Gamma\left(K^-\pi^+\pi^0\right)/\Gamma\left(K^-\pi^+\right)$

$\Gamma\left(K^-\rho^+\right)/\Gamma\left(K^-\pi^+\pi^0\right)$ Γ_{31}/Γ_{30}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.78 ±0.05 OUR AVERAGE					
0.765±0.041±0.054		FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.647±0.039±0.150		ANJOS	93	E691	γ Be 90–260 GeV
0.81 ±0.03 ±0.06		ADLER	87	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.31\ ^{+0.20}_{-0.14}$	13	SUMMERS	84	E691	Photoproduction
$0.85\ ^{+0.11}_{-0.15}\ ^{+0.09}_{-0.10}$	31	SCHINDLER	81	MRK2	e^+e^- 3.771 GeV

$\Gamma\left(K^*(892)^-\pi^+\right)/\Gamma\left(K^-\pi^+\pi^0\right)$ Γ_{81}/Γ_{30}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.36 ±0.04 OUR FIT Error includes scale factor of 1.3.				
0.28 ±0.04 OUR AVERAGE				
0.444±0.084±0.147	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.252±0.033±0.035	ANJOS	93	E691	γ Be 90–260 GeV
0.36 ±0.06 ±0.09	ADLER	87	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\pi^0)/\Gamma(K^-\pi^+\pi^0)$ Γ_{82}/Γ_{30}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.227±0.027 OUR FIT				
0.221±0.029 OUR AVERAGE				
0.248±0.047±0.023	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx 220$ GeV
0.213±0.027±0.035	ANJOS	93	E691	γ Be 90–260 GeV
0.20 ±0.03 ±0.05	ADLER	87	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-\pi^+\pi^0$ **nonresonant**$)/\Gamma(K^-\pi^+\pi^0)$ Γ_{34}/Γ_{30}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.049±0.018 OUR AVERAGE		Error includes scale factor of 1.1.			
0.101±0.033±0.040		FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx 220$ GeV
0.036±0.004±0.018		ANJOS	93	E691	γ Be 90–260 GeV
0.09 ±0.02 ±0.04		ADLER	87	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.51 ±0.22	21	SUMMERS	84	E691	Photoproduction

$\Gamma(\overline{K}^*(892)^0\pi^0)/\Gamma(\overline{K}^0\pi^0)$ Γ_{82}/Γ_{21}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.49±0.23 OUR FIT		Error includes scale factor of 1.1.			
$\mathbf{1.65^{+0.39}_{-0.31}\pm0.20}$	122	PROCARIO	93B	CLEO	$\overline{K}^0\pi^0\pi^0$ Dalitz plot

$\Gamma(\overline{K}_2^*(1430)^0\pi^0)/\Gamma(\overline{K}^*(892)^0\pi^0)$ Γ_{105}/Γ_{82}

Unseen decay modes of the $\overline{K}_2^*(1430)^0$ and $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.12	90	PROCARIO	93B	CLEO	$\overline{K}^0\pi^0\pi^0$ Dalitz plot

$\Gamma(\overline{K}^0\pi^0\pi^0$ **nonresonant**$)/\Gamma(\overline{K}^0\pi^0)$ Γ_{37}/Γ_{21}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.37±0.08±0.04	76	PROCARIO	93B	CLEO	$\overline{K}^0\pi^0\pi^0$ Dalitz plot

$\Gamma(K^-\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.075 ±0.004 OUR FIT		Error includes scale factor of 1.1.			
0.075 ±0.006 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
0.079 ±0.015 ±0.009		[40] ALBRECHT	94	ARG	$e^+e^- \approx \Upsilon(4S)$
0.0680±0.0027±0.0057	1430	[41] ALBRECHT	94F	ARG	$e^+e^- \approx \Upsilon(4S)$
0.091 ±0.008 ±0.008	992	ADLER	88C	MRK3	e^+e^- 3.77 GeV
0.117 ±0.025	185	[42] SCHINDLER	81	MRK2	e^+e^- 3.771 GeV
0.062 ±0.019	44	[43] PERUZZI	77	MRK1	e^+e^- 3.77 GeV

[40] ALBRECHT 94 uses D^0 mesons from $\overline{B}^0 \to D^{*+}\ell^-\overline{\nu}_\ell$ decays. This is a different set of events than used by ALBRECHT 94F.

[41] See the footnote on the ALBRECHT 94F measurement of $\Gamma(K^-\pi^+)/\Gamma_{\text{total}}$ for the method used.

[42] SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.68 ± 0.11 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

[43] PERUZZI 77 (MARK-1) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.36 ± 0.10 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.



$\Gamma\left(K^-\pi^+\pi^+\pi^-\right)/\Gamma_{\text{total}}$

$\Gamma(K^-\pi^+\pi^+\pi^-)/\Gamma(K^-\pi^+)$ Γ_{38}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.97±0.10 OUR FIT					
2.01±0.13 OUR AVERAGE					
1.7 ±0.2 ±0.2	1745	ANJOS	92C	E691	γ Be 90–260 GeV
1.90±0.25±0.20	337	ALVAREZ	91B	NA14	Photoproduction
2.12±0.16±0.09		BORTOLETTO	88	CLEO	e^+e^- 10.55 GeV
2.0 ±0.9	48	BAILEY	86	ACCM	π^- Be fixed target
2.17±0.28±0.23		ALBRECHT	85F	ARG	e^+e^- 10 GeV
2.0 ±1.0	10	BAILEY	83B	SPEC	π^- Be $\to D^0$
2.2 ±0.8	214	PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV

$\Gamma(K^-\pi^+\rho^0\,\text{total})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{39}/Γ_{38}

This includes $K^-a_1(1260)^+$, $\overline{K}^*(892)^0\rho^0$, etc. The next entry gives the specifically 3-body fraction. We rely on the MARK III and E691 full amplitude analyses of the $K^-\pi^+\pi^+\pi^-$ channel for values of the resonant substructure.

VALUE	DOCUMENT ID		TECN	COMMENT
0.835±0.035 OUR AVERAGE				
0.80 ±0.03 ±0.05	ANJOS	92C	E691	γ Be 90–260 GeV
0.855±0.032±0.030	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.98 ±0.12 ±0.10	ALVAREZ	91B	NA14	Photoproduction

$\Gamma(K^-\pi^+\rho^0\,3\text{-body})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{40}/Γ_{38}

We rely on the MARK III and E691 full amplitude analyses of the $K^-\pi^+\pi^+\pi^-$ channel for values of the resonant substructure.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.063±0.028 OUR AVERAGE					
0.05 ±0.03 ±0.02		ANJOS	92C	E691	γ Be 90–260 GeV
0.084±0.022±0.04		COFFMAN	92B	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.77 ±0.06 ±0.06		[44] ALVAREZ	91B	NA14	Photoproduction
0.85 $^{+0.11}_{-0.22}$	180	PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV

[44] This value is for ρ^0 $(K^-\pi^+)$-nonresonant. ALVAREZ 91B cannot determine what fraction of this is $K^-a_1(1260)^+$.

$\Gamma(\overline{K}^*(892)^0\rho^0)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{87}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included. We rely on the MARK III and E691 full amplitude analyses of the $K^-\pi^+\pi^+\pi^-$ channel for values of the resonant substructure.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.195±0.03±0.03		ANJOS	92C	E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.34 ±0.09±0.09		ALVAREZ	91B	NA14	Photoproduction
0.75 ±0.3	5	BAILEY	83B	SPEC	π Be $\rightarrow$ D^0
0.15 $^{+0.16}_{-0.15}$	20	PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,\text{transverse})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{88}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.20 ±0.07 OUR FIT				
0.213±0.024±0.075	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,S\text{-wave})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{89}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.375±0.045±0.06	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,S\text{-wave long.})/\Gamma_{\text{total}}$ Γ_{90}/Γ

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.003	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,P\text{-wave})/\Gamma_{\text{total}}$ Γ_{91}/Γ

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.003	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.009	90	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(\overline{K}^*(892)^0 \rho^0 D\text{-wave})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{92}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.255±0.045±0.06	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(K^-\pi^+ f_0(980))/\Gamma_{total}$ Γ_{97}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.011	90	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(\overline{K}^*(892)^0 f_0(980))/\Gamma_{total}$ Γ_{98}/Γ

Unseen decay modes of the $\overline{K}^*(892)^0$ and $f_0(980)$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.007	90	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(K^- a_1(1260)^+)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{76}/Γ_{38}

Unseen decay modes of the $a_1(1260)^+$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.97 ±0.14 OUR AVERAGE				
0.94 ±0.13 ±0.20	ANJOS	92C	E691	γ Be 90–260 GeV
0.984±0.048±0.16	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^- a_2(1320)^+)/\Gamma_{total}$ Γ_{79}/Γ

Unseen decay modes of the $a_2(1320)^+$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.002	90	ANJOS	92C	E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.006	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K_1(1270)^-\pi^+)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{99}/Γ_{38}

Unseen decay modes of the $K_1(1270)^-$ are included. The MARK3 and E691 experiments disagree considerably here.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
0.14 ±0.04 OUR FIT					
0.194±0.056±0.088		COFFMAN	92B	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.013	90	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(K_1(1400)^-\pi^+)/\Gamma_{total}$ Γ_{100}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.012	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(1410)^-\pi^+)/\Gamma_{total}$ Γ_{102}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.012	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\pi^+\pi^-\text{total})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{83}/Γ_{38}

This includes $\overline{K}^*(892)^0\rho^0$, *etc.* The next entry gives the specifically 3-body fraction.

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.30±0.06±0.03	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(\overline{K}^*(892)^0\pi^+\pi^-$ 3-body$)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{84}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.19 ±0.04 OUR FIT			
0.18 ±0.04 OUR AVERAGE			
0.165±0.03 ±0.045	ANJOS	92C E691	γ Be 90–260 GeV
0.210±0.027±0.06	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-\pi^+\pi^+\pi^-$ nonresonant$)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{46}/Γ_{38}

VALUE	DOCUMENT ID	TECN	COMMENT
0.233±0.032 OUR AVERAGE			
0.23 ±0.02 ±0.03	ANJOS	92C E691	γ Be 90–260 GeV
0.242±0.025±0.06	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.100±0.012 OUR FIT				
0.103±0.022±0.025	140	COFFMAN	92B MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.134^{+0.032}_{-0.033}$		[45] BARLAG	92C ACCM	π^- Cu 230 GeV

[45] BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{47}/Γ_{22}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.84±0.20 OUR FIT				
1.86±0.23 OUR AVERAGE				
1.80±0.20±0.21	190	[46] ALBRECHT	92P ARG	$e^+e^- \approx$ 10 GeV
2.8 ±0.8 ±0.8	46	ANJOS	92C E691	γ Be 90–260 GeV
1.85±0.26±0.30	158	KINOSHITA	91 CLEO	$e^+e^- \sim$ 10.7 GeV

[46] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^0\eta)/\Gamma(K^-\pi^+)$ Γ_{69}/Γ_{20}

Unseen decay modes of the η are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.64	90	ALBRECHT	89D ARG	e^+e^- 10 GeV

$\Gamma(\overline{K}^0\eta)/\Gamma(\overline{K}^0\pi^0)$ Γ_{69}/Γ_{21}

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.33±0.04 OUR FIT				
0.32±0.04±0.03	225	PROCARIO	93B CLEO	$\eta \to \gamma\gamma$

$\Gamma(\overline{K}^0\eta)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{69}/Γ_{22}

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.130±0.017 OUR FIT				
0.14 ±0.02 ±0.02	80	PROCARIO	93B CLEO	$\eta \to \pi^+\pi^-\pi^0$

$\Gamma(\overline{K}^0\omega)/\Gamma(K^-\pi^+)$ Γ_{72}/Γ_{20}

Unseen decay modes of the ω are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.54±0.10 OUR FIT				
1.00±0.36±0.20	ALBRECHT	89D	ARG	e^+e^- 10 GeV

$\Gamma(\overline{K}^0\omega)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{72}/Γ_{22}

Unseen decay modes of the ω are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.38±0.07 OUR FIT					
0.33±0.09 OUR AVERAGE Error includes scale factor of 1.1.					
0.29±0.08±0.05	16	[47] ALBRECHT	92P	ARG	$e^+e^-\approx$ 10 GeV
0.54±0.14±0.16	40	KINOSHITA	91	CLEO	$e^+e^-\sim$ 10.7 GeV

[47] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^0\omega)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{72}/Γ_{47}

Unseen decay modes of the ω are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.21 ±0.04 OUR FIT				
0.220±0.048±0.0116	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0\eta'(958))/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{73}/Γ_{22}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.32±0.04 OUR AVERAGE					
0.31±0.02±0.04	594	PROCARIO	93B	CLEO	$\eta' \to \eta\pi^+\pi^-$, $\rho^0\gamma$
0.37±0.13±0.06	18	[48] ALBRECHT	92P	ARG	$e^+e^-\approx$ 10 GeV

[48] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^*(892)^-\rho^+)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{93}/Γ_{47}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.606±0.188±0.126	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(892)^-\rho^+$ longitudinal$)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{94}/Γ_{47}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.290±0.111	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(892)^-\rho^+$ transverse$)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{95}/Γ_{47}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.317±0.180	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(892)^-\rho^+$ P-wave$)/\Gamma_{\text{total}}$ Γ_{96}/Γ

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.015	90	[49] COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

[49] Obtained using other $\overline{K}^*(892)\rho$ P-wave limits and isospin relations.

$\Gamma(\overline{K}^*(892)^0\rho^0\,\text{transverse})/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{88}/Γ_{47}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.15 ±0.06 OUR FIT			
0.126±0.111	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0 a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{77}/Γ

Unseen decay modes of the $a_1(1260)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.019	90	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(K_1(1270)^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{99}/Γ_{47}

Unseen decay modes of the $K_1(1270)^-$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.106±0.028 OUR FIT			
0.10 ±0.03	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}_1(1400)^0\pi^0)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.037	90	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\pi^+\pi^-\text{3-body})/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{84}/Γ_{47}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.14 ±0.04 OUR FIT Error includes scale factor of 1.1.			
0.191±0.105	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0\pi^+\pi^-\pi^0\,\text{nonresonant})/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{54}/Γ_{47}

VALUE	DOCUMENT ID	TECN	COMMENT
0.210±0.147±0.150	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-\pi^+\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.149±0.037±0.030	24	[50] ADLER	88C MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.177 ± 0.029		[51] BARLAG	92C ACCM	π^- Cu 230 GeV
$0.209^{+0.074}_{-0.043}\pm0.012$	9	[51] AGUILAR-...	87F HYBR	πp, pp 360, 400 GeV

[50] ADLER 88C uses an absolute normalization method finding this decay channel opposite a detected $\overline{D}^0 \to K^+\pi^-$ in pure $D\overline{D}$ events.

[51] AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction using topological normalization. They do not distinguish the presence of a third π^0, and thus are not included in the average.

$\Gamma(K^-\pi^+\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+)$ Γ_{56}/Γ_{20}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.06±0.10 OUR FIT				
0.98±0.11±0.11	225	[52] ALBRECHT	92P ARG	$e^+e^- \approx$ 10 GeV

[52] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^-\pi^+\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{56}/Γ_{38}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.54±0.05 OUR FIT					
0.56±0.07 OUR AVERAGE					
$0.55\pm0.07^{+0.12}_{-0.09}$	167	KINOSHITA	91	CLEO	$e^+e^-\sim 10.7$ GeV
0.57±0.06±0.05	180	ANJOS	90D	E691	Photoproduction

$\Gamma(\overline{K}^*(892)^0\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+\pi^+\pi^-\pi^0)$ Γ_{106}/Γ_{56}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.45±0.15±0.15	ANJOS	90D	E691	Photoproduction

$\Gamma(\overline{K}^*(892)^0\eta)/\Gamma(K^-\pi^+)$ Γ_{107}/Γ_{20}

Unseen decay modes of the $\overline{K}^*(892)^0$ and η are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.49±0.12 OUR FIT					
$\mathbf{0.58\pm0.19^{+0.24}_{-0.28}}$	46	KINOSHITA	91	CLEO	$e^+e^-\sim 10.7$ GeV

$\Gamma(\overline{K}^*(892)^0\eta)/\Gamma(K^-\pi^+\pi^0)$ Γ_{107}/Γ_{30}

Unseen decay modes of the $\overline{K}^*(892)^0$ and η are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.134±0.034 OUR FIT					
0.13 ±0.02 ±0.03	214	PROCARIO	93B	CLEO	$\overline{K}^{*0}\eta \to K^-\pi^+/\gamma\gamma$

$\Gamma(K^-\pi^+\omega)/\Gamma(K^-\pi^+)$ Γ_{108}/Γ_{20}

Unseen decay modes of the ω are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.78±0.12±0.10	99	[53] ALBRECHT	92P	ARG	$e^+e^-\approx$ 10 GeV

[53] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^*(892)^0\omega)/\Gamma(K^-\pi^+)$ Γ_{109}/Γ_{20}

Unseen decay modes of the $\overline{K}^*(892)^0$ and ω are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.28±0.11±0.04	17	[54] ALBRECHT	92P	ARG	$e^+e^-\approx$ 10 GeV

[54] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^*(892)^0\omega)/\Gamma(K^-\pi^+\pi^+\pi^-\pi^0)$ Γ_{109}/Γ_{56}

Unseen decay modes of the $\overline{K}^*(892)^0$ and ω are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.44	90	[55] ANJOS	90D	E691	Photoproduction

[55] Recovered from the published limit, $\Gamma(\overline{K}^*(892)^0\omega)/\Gamma_{\text{total}}$, in order to make our normalization consistent.

$\Gamma(K^-\pi^+\eta'(958))/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{110}/Γ_{38}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.093±0.014±0.019	286	PROCARIO	93B	CLEO	$\eta' \to \eta\pi^+\pi^-, \rho^0\gamma$

$\Gamma(\overline{K}^*(892)^0\eta'(958))/\Gamma(K^-\pi^+\eta'(958))$ $\Gamma_{111}/\Gamma_{110}$

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN
<0.15	90	PROCARIO	93B	CLEO

$\Gamma(\overline{K}^0\pi^+\pi^+\pi^-\pi^-)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{61}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.107±0.029 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.			
0.07 ±0.02 ±0.01	11	[56] ALBRECHT	92P	ARG	$e^+e^-\approx$ 10 GeV
0.149±0.026	56	AMMAR	91	CLEO	$e^+e^-\approx$ 10.5 GeV
0.18 ±0.07 ±0.04	6	ANJOS	90D	E691	Photoproduction

[56] This value is calculated from numbers in Table 1 of ALBRECHT 92P.



$\Gamma(\overline{K}^0\pi^+\pi^+\pi^-\pi^-)/\Gamma(\overline{K}^0\pi^+\pi^-)$

$\Gamma(\overline{K}^0\pi^+\pi^-\pi^0\pi^0(\pi^0))/\Gamma_{total}$ Γ_{62}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.106^{+0.073}_{-0.029}\pm0.006}$	4	[57] AGUILAR-...	87F	HYBR	πp, pp 360, 400 GeV

[57] AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization, and does not distinguish the presence of a third π^0.

$\Gamma(\overline{K}^0 K^+ K^-)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ $\Gamma_{63}/\Gamma_{22} = (\Gamma_{65}+\frac{1}{2}\Gamma_{75})/\Gamma_{22}$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.172±0.014 OUR FIT					
0.178±0.019 OUR AVERAGE					
0.20 ±0.05 ±0.04	47	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.170±0.022	136	AMMAR	91	CLEO	e^+e^- ≈ 10.5 GeV
0.24 ±0.08		BEBEK	86	CLEO	e^+e^- near $\Upsilon(4S)$
0.185±0.055	52	ALBRECHT	85B	ARG	e^+e^- 10 GeV

$\Gamma(\overline{K}^0 \phi)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ Γ_{75}/Γ_{22}

Unseen decay modes of the ϕ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.158±0.016 OUR FIT					
0.156±0.017 OUR AVERAGE					
0.13 ±0.06 ±0.02	13	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.163±0.023	63	AMMAR	91	CLEO	e^+e^- ≈ 10.5 GeV
0.155±0.033	56	ALBRECHT	87E	ARG	e^+e^- 10 GeV
0.14 ±0.05	29	BEBEK	86	CLEO	e^+e^- near $\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.186±0.052	26	ALBRECHT	85B	ARG	See ALBRECHT 87E

$\Gamma(\overline{K}^0 K^+ K^-$ non-$\phi)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ Γ_{65}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.093±0.014 OUR FIT					
0.088±0.019 OUR AVERAGE					
0.11 ±0.04 ±0.03	20	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.084±0.020		ALBRECHT	87E	ARG	e^+e^- 10 GeV

$\Gamma(K^0_S K^0_S K^0_S)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ Γ_{66}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0154±0.0025 OUR AVERAGE					
0.0139±0.0019±0.0024	61	ASNER	96B	CLEO	e^+e^- ≈ $\Upsilon(4S)$
0.035 ±0.012 ±0.006	10	FRABETTI	94J	E687	γ Be $\overline{E}_\gamma$=220 GeV
0.016 ±0.005	22	AMMAR	91	CLEO	e^+e^- ≈ 10.5 GeV
0.017 ±0.007 ±0.005	5	ALBRECHT	90C	ARG	e^+e^- ≈ 10 GeV

$\Gamma(K^+ K^- K^- \pi^+)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{67}/Γ_{38}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0028±0.0007±0.0001	20	FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma$ ≈ 200 GeV

$\Gamma(K^+ K^- \overline{K}^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE		DOCUMENT ID		TECN	COMMENT
$\mathbf{0.0072^{+0.0048}_{-0.0035}}$	[58]	BARLAG	92C	ACCM	π^- Cu 230 GeV

[58] BARLAG 92C computes the branching fraction using topological normalization.