

$N(1650) S_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-) \text{ Status: } ****$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $N(1650)$ BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1640 to 1680 (≈ 1650) OUR ESTIMATE			
1659 \pm 9	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
1650 \pm 30	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1670 \pm 8	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1677 \pm 8	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
1667	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1712	¹ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1669 \pm 17	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
1713 \pm 27	² BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
1674	LI	93	IPWA $\gamma N \rightarrow \pi N$
1688	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
1672	MUSETTE	80	IPWA $\pi^- p \rightarrow \Lambda K^0$
1680	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
1680	BAKER	78	DPWA $\pi^- p \rightarrow \Lambda K^0$
1694	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
1700 \pm 5	³ BAKER	77	IPWA $\pi^- p \rightarrow \Lambda K^0$
1680	³ BAKER	77	DPWA $\pi^- p \rightarrow \Lambda K^0$
1700	⁴ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
1675	KNASEL	75	DPWA $\pi^- p \rightarrow \Lambda K^0$
1660	⁵ LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

 $N(1650)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
145 to 190 (≈ 150) OUR ESTIMATE			
167.9 \pm 9.4	GREEN	97	DPWA $\pi N \rightarrow \pi N, \eta N$
173 \pm 12	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
150 \pm 40	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
180 \pm 20	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
160 \pm 12	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
90	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
184	¹ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
215 \pm 32	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
279 \pm 54	² BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
225	LI	93	IPWA $\gamma N \rightarrow \pi N$
183	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
179	MUSETTE	80	IPWA $\pi^- p \rightarrow \Lambda K^0$

120		SAXON	80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
90		BAKER	78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
193		BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$
130 ± 10		³ BAKER	77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
90		³ BAKER	77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
170		⁴ LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
170		KNASEL	75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
130		⁵ LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1650) POLE POSITION

REAL PART

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1640 to 1680 (≈ 1660) OUR ESTIMATE				
1673		ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1689		¹ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1670		⁶ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
1640 ± 20		CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1657		ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
1648 or 1651		⁷ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
1699 or 1698		⁴ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

– 2×IMAGINARY PART

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
150 to 170 (≈ 160) OUR ESTIMATE				
82		ARNDT	95	DPWA $\pi N \rightarrow N\pi$
192		¹ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
163		⁶ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
150 ± 30		CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
160		ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
117 or 119		⁷ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
174 or 173		⁴ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

N(1650) ELASTIC POLE RESIDUE

MODULUS $|r|$

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
22		ARNDT	95	DPWA $\pi N \rightarrow N\pi$
72		¹ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
39		HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
60 ± 10		CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
54		ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

PHASE θ

<u>VALUE ($^{\circ}$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
29	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
-85	¹ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
-37	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
-75 \pm 25	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-38	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

 $N(1650)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	55–90 %
Γ_2 $N\eta$	3–10 %
Γ_3 ΛK	3–11 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	10–20 %
Γ_6 $\Delta\pi$	1–7 %
Γ_7 $\Delta(1232)\pi$, D -wave	
Γ_8 $N\rho$	4–12 %
Γ_9 $N\rho$, $S=1/2$, S -wave	
Γ_{10} $N\rho$, $S=3/2$, D -wave	
Γ_{11} $N(\pi\pi)_{S\text{-wave}}^{I=0}$	<4 %
Γ_{12} $N(1440)\pi$	<5 %
Γ_{13} $p\gamma$	0.04–0.18 %
Γ_{14} $p\gamma$, helicity=1/2	0.04–0.18 %
Γ_{15} $n\gamma$	0.003–0.17 %
Γ_{16} $n\gamma$, helicity=1/2	0.003–0.17 %

 $N(1650)$ BRANCHING RATIOS

<u>$\Gamma(N\pi)/\Gamma_{\text{total}}$</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_1/Γ</u>
0.55 to 0.90 OUR ESTIMATE				
0.735 \pm 0.011	GREEN	97	DPWA $\pi N \rightarrow \pi N, \eta N$	
0.89 \pm 0.07	MANLEY	92	IPWA $\pi N \rightarrow \pi N \& N\pi\pi$	
0.65 \pm 0.10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.61 \pm 0.04	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.99	ARNDT	95	DPWA $\pi N \rightarrow N\pi$	
0.27	¹ ARNDT	95	DPWA $\pi N \rightarrow N\pi$	
0.94 \pm 0.07	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$	
0.49 \pm 0.21	² BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$	

$$\Gamma(N\eta)/\Gamma_{\text{total}} \qquad \qquad \qquad \Gamma_2/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.06 ± 0.05	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
0.02 ± 0.03	² BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$

$$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1650) \rightarrow N\eta \qquad \qquad \qquad (\Gamma_1\Gamma_2)^{1/2}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.09	⁸ BAKER	79	DPWA $\pi^- p \rightarrow n\eta$

$$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1650) \rightarrow \Lambda K \qquad \qquad \qquad (\Gamma_1\Gamma_3)^{1/2}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.27 to -0.17 OUR ESTIMATE			
-0.22	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$
-0.22	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.25	⁹ BAKER	78	DPWA See SAXON 80
-0.23 ± 0.01	³ BAKER	77	IPWA $\pi^- p \rightarrow \Lambda K^0$
-0.25	³ BAKER	77	DPWA $\pi^- p \rightarrow \Lambda K^0$
0.12	KNASEL	75	DPWA $\pi^- p \rightarrow \Lambda K^0$

$$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1650) \rightarrow \Sigma K \qquad \qquad \qquad (\Gamma_1\Gamma_4)^{1/2}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.254	LIVANOS	80	DPWA $\pi p \rightarrow \Sigma K$
0.066 to 0.137	¹⁰ DEANS	75	DPWA $\pi N \rightarrow \Sigma K$
0.20	KNASEL	75	DPWA

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1650) \rightarrow \Delta(1232)\pi, D\text{-wave} \qquad \qquad \qquad (\Gamma_1\Gamma_7)^{1/2}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+0.15 to 0.23 OUR ESTIMATE			
+0.12 ± 0.04	MANLEY	92	IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
+0.29	^{4,11} LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
+0.15	⁵ LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1650) \rightarrow N\rho, S=1/2, S\text{-wave} \qquad \qquad \qquad (\Gamma_1\Gamma_9)^{1/2}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
±0.03 to ±0.19 OUR ESTIMATE			
-0.01 ± 0.09	MANLEY	92	IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
+0.17	^{4,11} LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
-0.16	⁵ LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1650) \rightarrow N\rho, S=3/2, D\text{-wave}$ $(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.17 to +0.29 OUR ESTIMATE

+0.16 ± 0.06	MANLEY	92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
+0.29	4,11 LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1650) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0}$ $(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.04 to +0.18 OUR ESTIMATE

+0.12 ± 0.08	MANLEY	92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
0.00	4,11 LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.25	5 LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1650) \rightarrow N(1440)\pi$ $(\Gamma_1 \Gamma_{12})^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.11 ± 0.06	MANLEY	92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
--------------	--------	----	------	---------------------------------------

N(1650) PHOTON DECAY AMPLITUDES

$N(1650) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV^{-1/2}) DOCUMENT ID TECN COMMENT

+0.053 ± 0.016 OUR ESTIMATE

0.069 ± 0.005	ARNDT	96	IPWA	$\gamma N \rightarrow \pi N$
0.033 ± 0.015	CRAWFORD	83	IPWA	$\gamma N \rightarrow \pi N$
0.050 ± 0.010	AWAJI	81	DPWA	$\gamma N \rightarrow \pi N$
0.065 ± 0.005	ARAI	80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.061 ± 0.005	ARAI	80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.031 ± 0.017	CRAWFORD	80	DPWA	$\gamma N \rightarrow \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.068 ± 0.003	LI	93	IPWA	$\gamma N \rightarrow \pi N$
0.091	WADA	84	DPWA	Compton scattering
+0.048 ± 0.017	BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$
+0.068 ± 0.009	FELLER	76	DPWA	$\gamma N \rightarrow \pi N$

$N(1650) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV^{-1/2}) DOCUMENT ID TECN COMMENT

-0.015 ± 0.021 OUR ESTIMATE

-0.015 ± 0.005	ARNDT	96	IPWA	$\gamma N \rightarrow \pi N$
-0.008 ± 0.004	AWAJI	81	DPWA	$\gamma N \rightarrow \pi N$
0.004 ± 0.004	FUJII	81	DPWA	$\gamma N \rightarrow \pi N$
0.010 ± 0.020	ARAI	80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.008 ± 0.019	ARAI	80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.068 ± 0.040	CRAWFORD	80	DPWA	$\gamma N \rightarrow \pi N$
-0.011 ± 0.011	TAKEDA	80	DPWA	$\gamma N \rightarrow \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.002 ± 0.002	LI	93	IPWA	$\gamma N \rightarrow \pi N$
-0.045 ± 0.024	BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$

$N(1650) \quad \gamma p \rightarrow \Lambda K^+$ AMPLITUDES **$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1650) \rightarrow \Lambda K^+$ (E_{0+} amplitude)**VALUE (units 10^{-3}) DOCUMENT ID TECN

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.8 \pm 0.3	WORKMAN	90	DPWA
8.13	TANABE	89	DPWA

 $p\gamma \rightarrow N(1650) \rightarrow \Lambda K^+$ phase angle θ (E_{0+} amplitude)VALUE (degrees) DOCUMENT ID TECN

• • • We do not use the following data for averages, fits, limits, etc. • • •

-107 \pm 3	WORKMAN	90	DPWA
-107.8	TANABE	89	DPWA

 $N(1650)$ FOOTNOTES¹ ARNDT 95 finds two distinct states.² BATINIC 95 finds two distinct states. This second resonance was associated with the $N(2090) S_{11}$.³ The two BAKER 77 entries are from an IPWA using the Barrelet-zero method and from a conventional energy-dependent analysis.⁴ LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.⁵ From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.⁶ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.⁷ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.⁸ BAKER 79 fixed this coupling during fitting, but the negative sign relative to the $N(1535)$ is well determined.⁹ The overall phase of BAKER 78 couplings has been changed to agree with previous conventions. Superseded by SAXON 80.¹⁰ The range given for DEANS 75 is from the four best solutions.¹¹ LONGACRE 77 considers this coupling to be well determined.

N(1650) REFERENCESFor early references, see Physics Letters **111B** 70 (1982).

GREEN	97	PR C55 R2167	+Wycech	(HELS, WINR)
ARNDT	96	PR C53 430	+Strakovsky, Workman	(VPI)
ARNDT	95	PR C52 2120	+Strakovsky, Workman, Pavan	(VPI, BRCO)
BATINIC	95	PR C51 2310	+Slaus, Svarc, Nefkens	(BOSK, UCLA)
HOEHLER	93	π N Newsletter 9 1		(KARL)
LI	93	PR C47 2759	+Arndt, Roper, Workman	(VPI)
MANLEY	92	PR D45 4002	+Saleski	(KENT) IJP
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
ARNDT	91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
WORKMAN	90	PR C42 781		(VPI)
TANABE	89	PR C39 741	+Kohno, Bennhold	(MANZ)
Also	89	NC 102A 193	Kohno, Tanabe, Bennhold	(MANZ)
WADA	84	NP B247 313	+Egawa, Imanishi, Ishii, Kato, Ukai+	(INUS)
BELL	83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+	(RL) IJP
CRAWFORD	83	NP B211 1	+Morton	(GLAS)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
FUJII	81	NP B187 53	+Hayashii, Iwata, Kajikawa+	(NAGO, OSAK)
ARAI	80	Toronto Conf. 93		(INUS)
Also	82	NP B194 251	Arai, Fujii	(INUS)
CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	+Baton, Coutures, Kochowski, Neveu	(SACL) IJP
MUSETTE	80	NC 57A 37		(BRUX) IJP
SAXON	80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
TAKEDA	80	NP B168 17	+Arai, Fujii, Ikeda, Iwasaki+	(TOKY, INUS)
BAKER	79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
BAKER	78	NP B141 29	+Blissett, Bloodworth, Broome+	(RL, CAVE) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons	(GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
BAKER	77	NP B126 365	+Blissett, Bloodworth, Broome, Hart+	(RHEL) IJP
LONGACRE	77	NP B122 493	+Dolbeau	(SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet	(SACL) IJP
FELLER	76	NP B104 219	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
DEANS	75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
KNASEL	75	PR D11 1	+Lindquist, Nelson+	(CHIC, WUSL, OSU, ANL) IJP
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP