

**$\Sigma(1560)$  Bumps**

$$I(J^P) = 1(?^?) \quad \text{Status: } **$$

## OMITTED FROM SUMMARY TABLE

This entry lists peaks reported in mass spectra around 1560 MeV without implying that they are necessarily related.

DIONISI 78B observes a 6 standard-deviation enhancement at 1553 MeV in the charged  $\Lambda/\Sigma\pi$  mass spectra from  $K^- p \rightarrow (\Lambda/\Sigma)\pi K\bar{K}$  at 4.2 GeV/c. In a CERN ISR experiment, LOCKMAN 78 reports a narrow 6 standard-deviation enhancement at 1572 MeV in  $\Lambda\pi^\pm$  from the reaction  $pp \rightarrow \Lambda\pi^+\pi^- X$ . These enhancements are unlikely to be associated with the  $\Sigma(1580)$  (which has not been confirmed by several recent experiments – see the next entry in the Listings).

CARROLL 76 observes a bump at 1550 MeV (as well as one at 1580 MeV) in the isospin-1  $\bar{K}N$  total cross section, but uncertainties in cross section measurements outside the mass range of the experiment preclude estimating its significance.

See also MEADOWS 80 for a review of this state.

---

 **$\Sigma(1560)$  MASS  
(PRODUCTION EXPERIMENTS)**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b><math>\approx 1560</math> OUR ESTIMATE</b>					
$1553 \pm 7$	121	DIONISI	78B HBC	$\pm$	$K^- p \rightarrow (Y\pi)K\bar{K}$
$1572 \pm 4$	40	LOCKMAN	78 SPEC	$\pm$	$pp \rightarrow \Lambda\pi^+\pi^- X$

---

 **$\Sigma(1560)$  WIDTH  
(PRODUCTION EXPERIMENTS)**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
$79 \pm 30$	121	DIONISI	78B HBC	$\pm$	$K^- p \rightarrow (Y\pi)K\bar{K}$
$15 \pm 6$	40	<sup>1</sup> LOCKMAN	78 SPEC	$\pm$	$pp \rightarrow \Lambda\pi^+\pi^- X$

---

 **$\Sigma(1560)$  DECAY MODES  
(PRODUCTION EXPERIMENTS)**

	<u>Mode</u>	<u>Fraction (<math>\Gamma_j/\Gamma</math>)</u>
$\Gamma_1$	$\Lambda\pi$	seen
$\Gamma_2$	$\Sigma\pi$	

---

**$\Sigma(1560)$  BRANCHING RATIOS  
(PRODUCTION EXPERIMENTS)**

$\Gamma(\Sigma\pi)/[\Gamma(\Lambda\pi) + \Gamma(\Sigma\pi)]$	$\Gamma_2/(\Gamma_1+\Gamma_2)$			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
$0.35 \pm 0.12$	DIONISI	78B HBC	$\pm$	$K^- p \rightarrow$ $(Y\pi)K\bar{K}$

  

$\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b>seen</b>	LOCKMAN	78 SPEC	$\pm$	$pp \rightarrow \Lambda\pi^+\pi^-X$

**$\Sigma(1560)$  FOOTNOTES  
(PRODUCTION EXPERIMENTS)**

<sup>1</sup> The width observed by LOCKMAN 78 is consistent with experimental resolution.

**$\Sigma(1560)$  REFERENCES  
(PRODUCTION EXPERIMENTS)**

MEADOWS	80	Toronto Conf. 283		(CINC)
DIONISI	78B	PL 78B 154	+Armenteros, Diaz	(CERN, AMST, NIJM, OXF)
LOCKMAN	78	Saclay DPHPE 78-01	+Meyer, Rander, Poster, Schlein+	(UCLA, SACL)
CARROLL	76	PRL 37 806	+Chiang, Kycia, Li, Mazur, Michael+	(BNL)