

$\Xi_c(2645)$

 $I(J^P) = ?(?^?)$  Status: \*\*\*

A narrow peak seen in the  $\Xi_c \pi$  mass spectrum. The natural assignment is that this is the  $J^P = 3/2^+$  excitation of the  $\Xi_c$  in the same SU(4) multiplet as the  $\Delta(1232)$ .

### $\Xi_c(2645)$ MASSES

The masses are obtained from the mass-difference measurements that follow.

#### $\Xi_c(2645)^+$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
<b>2644.6 ± 2.1 OUR FIT</b>	Error includes scale factor of 1.2.

#### $\Xi_c(2645)^0$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
<b>2643.8 ± 1.8 OUR FIT</b>	

### $m_{\Xi_c(2645)} - m_{\Xi_c}$

#### $m_{\Xi_c(2645)^+} - m_{\Xi_c^0}$

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>174.3 ± 1.1 OUR FIT</b>				
<b>174.3 ± 0.5 ± 1.0</b>	34	GIBBONS	96 CLE2	$e^+ e^- \approx \Upsilon(4S)$

#### $m_{\Xi_c(2645)^0} - m_{\Xi_c^+}$

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>178.2 ± 1.1 OUR FIT</b>				
<b>178.2 ± 0.5 ± 1.0</b>	55	AVERY	95 CLE2	$e^+ e^- \approx \Upsilon(4S)$

### $\Xi_c(2645)$ WIDTHS

#### $\Xi_c(2645)^+$ WIDTH

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 3.1</b>	90	GIBBONS	96 CLE2	$e^+ e^- \approx \Upsilon(4S)$

#### $\Xi_c(2645)^0$ WIDTH

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 5.5</b>	90	55	AVERY	95 CLE2	$e^+ e^- \approx \Upsilon(4S)$

### $\Xi_c(2645)$ DECAY MODES

$\Xi_c \pi$  is the only strong decay allowed to a  $\Xi_c$  resonance having this mass.

	Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$	$\Xi_c^0 \pi^+$	seen
$\Gamma_2$	$\Xi_c^+ \pi^-$	seen

### $\Xi_c(2645)$ REFERENCES

GIBBONS	96	PRL 77 810	+Johnson, Kwon+	(CLEO Collab.)
AVERY	95	PRL 75 4364	+Freyberger, Lingel+	(CLEO Collab.)