1. PHYSICAL CONSTANTS

Table 1.1. Reviewed 1998 by B.N. Taylor (NIST). Based mainly on the "1986 Adjustment of the Fundamental Physical Constants" by E.R. Cohen and B.N. Taylor, Rev. Mod. Phys. **59**, 1121 (1987). The last group of constants (beginning with the Fermi coupling constant) comes from the Particle Data Group. The figures in parentheses after the values give the 1-standard-deviation uncertainties in the last digits; the corresponding uncertainties in parts per million (ppm) are given in the last column. This set of constants (aside from the last group) is recommended for international use by CODATA (the Committee on Data for Science and Technology).

Since the 1986 adjustment, new experiments have yielded improved values for a number of constants, including the Rydberg constant R_∞, the Planck constant h, the fine-structure constant α, and the molar gas constant R, and hence also for constants directly derived from these, such as the Boltzmann constant k and Stefan-Boltzmann constant σ. The new results and their impact on the 1986 recommended values are discussed extensively in "Recommended Values of the Fundamental Physical Constants: A Status Report," B.N. Taylor and E.R. Cohen, J. Res. Natl. Inst. Stand. Technol. **95**, 497 (1990); see also E.R. Cohen and B.N. Taylor, "The Fundamental Physical Constants," Phys. Today, August 1997 Part 2, BG7. In general, the new results give uncertainties for the affected constants that are 5 to 7 times smaller than the 1986 uncertainties, but the changes in the values themselves are smaller than twice the 1986 uncertainties. Because the output values of a least-squares adjustment are correlated, the new results cannot readily be incorporated with the 1986 values. Until the next complete adjustment of the constants (expected by the end of 1998), the 1986 CODATA set, given (in part) below, remains the set of choice. The full 1986 set (to be replaced by the new set, when available) may be found at `http://physics.nist.gov/cuu`.

c **speed of light in vacuum**
Value: 299 792 458 m s^{-1}
Uncert. (ppm): exact*

* The meter is the length of the path traveled by light in vacuum during a time interval of 1/299 792 458 of a second.

h **Planck constant**
Value: $6.626\,075\,5(40)\times10^{-34}$ J s
Uncert. (ppm): 0.60

$\hbar \equiv h/2\pi$ **Planck constant, reduced**
Value: $1.054\,572\,66(63)\times10^{-34}$ J s
Uncert. (ppm): 0.60

Value: $= 6.582\,122\,0(20)\times10^{-22}$ MeV s
Uncert. (ppm): 0.30

e **electron charge magnitude**
Value: $1.602\,177\,33(49)\times10^{-19}$ C $= 4.803\,206\,8(15)\times10^{-10}$ esu
Uncert. (ppm): 0.30, 0.30

$\hbar c$ **conversion constant**
Value: 197.327 053(59) MeV fm
Uncert. (ppm): 0.30

$(\hbar c)^2$ **conversion constant**
Value: 0.389 379 66(23) GeV^2 mbarn
Uncert. (ppm): 0.59

---

m_e **electron mass**
Value: 0.510 999 06(15) MeV/c^2 $= 9.109\,389\,7(54)\times10^{-31}$ kg
Uncert. (ppm): 0.30, 0.59

m_p **proton mass**
Value: 938.272 31(28) MeV/c^2 $= 1.672\,623\,1(10)\times10^{-27}$ kg
Uncert. (ppm): 0.30, 0.59

Value: $= 1.007\,276\,470(12)$ u $= 1836.152\,701(37)\ m_e$
Uncert. (ppm):0.012, 0.020

m_d **deuteron mass**
Value: 1875.613 39(57) MeV/c^2
Uncert. (ppm): 0.30

(mass ^{12}C atom)/12 = (1 g)/(N_A mol) **unified atomic mass unit (u)**
Value: 931.494 32(28) MeV/c^2 = 1.660 540 2(10)$\times 10^{-27}$ kg
Uncert. (ppm): 0.30, 0.59

ϵ_0 $\epsilon_0\mu_0 = 1/c^2$ **permittivity of free space**
Value: 8.854 187 817 ... $\times 10^{-12}$ F m^{-1}
Uncert. (ppm): exact

μ_0 $\epsilon_0\mu_0 = 1/c^2$ **permeability of free space**
Value: $4\pi \times 10^{-7}$ N A^{-2} = 12.566 370 614 ... $\times 10^{-7}$ N A^{-2}
Uncert. (ppm): exact

$\alpha = e^2/4\pi\epsilon_0\hbar c$ **fine-structure constant**
Value: 1/137.035 989 5(61)†
Uncert. (ppm): 0.045

† At $Q^2 = 0$. At $Q^2 \approx m_W^2$ the value is approximately 1/128.

$r_e = e^2/4\pi\epsilon_0 m_e c^2$ **classical electron radius**
Value: 2.817 940 92(38)$\times 10^{-15}$ m
Uncert. (ppm): 0.13

$\lambda\!\!\!^{-}_e = \hbar/m_e c = r_e\alpha^{-1}$ **electron Compton wavelength**
Value: 3.861 593 23(35)$\times 10^{-13}$ m
Uncert. (ppm): 0.089

$a_\infty = 4\pi\epsilon_0\hbar^2/m_e e^2 = r_e\alpha^{-2}$ **Bohr radius ($m_{\text{nucleus}} = \infty$)**
Value: 0.529 177 249(24)$\times 10^{-10}$ m
Uncert. (ppm): 0.045

hc/e **wavelength of 1 eV/c particle**
Value: 1.239 842 44(37)$\times 10^{-6}$ m
Uncert. (ppm): 0.30

$hcR_\infty = m_e e^4/2(4\pi\epsilon_0)^2\hbar^2 = m_e c^2\alpha^2/2$ **Rydberg energy**
Value: 13.605 698 1(40) eV
Uncert. (ppm): 0.30

$\sigma_T = 8\pi r_e^2/3$ **Thomson cross section**
Value: 0.665 246 16(18) barn
Uncert. (ppm): 0.27

$\mu_B = e\hbar/2m_e$ **Bohr magneton**
Value: $5.788\,382\,63(52)\times10^{-11}$ MeV T^{-1}
Uncert. (ppm): 0.089

$\mu_N = e\hbar/2m_p$ **nuclear magneton**
Value: $3.152\,451\,66(28)\times10^{-14}$ MeV T^{-1}
Uncert. (ppm): 0.089

$\omega^e_{\rm cycl}/B = e/m_e$ **electron cyclotron freq./field**
Value: $1.758\,819\,62(53)\times10^{11}$ rad s^{-1} T^{-1}
Uncert. (ppm): 0.30

$\omega^p_{\rm cycl}/B = e/m_p$ **proton cyclotron freq./field**
Value: $9.578\,830\,9(29)\times10^{7}$ rad s^{-1} T^{-1}
Uncert. (ppm): 0.30

G_N **gravitational constant**‡
Value: $6.672\,59(85)\times10^{-11}$ m^3 kg^{-1} s^{-2}
Uncert. (ppm): 128

Value: $= 6.707\,11(86)\times10^{-39}\ \hbar c\ (\mathrm{GeV}/c^2)^{-2}$
Uncert. (ppm): 128

‡ Absolute lab measurements of G_N were performed only on scales of $10^{-1\pm1}$ m.

g **standard grav. accel., sea level**
Value: 9.806 65 m s^{-2}
Uncert. (ppm): exact

N_A **Avogadro constant**
Value: $6.022\,136\,7(36)\times10^{23}$ mol^{-1}
Uncert. (ppm): 0.59

k **Boltzmann constant**
Value: $1.380\,658(12)\times10^{-23}$ J K^{-1}
Uncert. (ppm): 8.5

Value: $= 8.617\,385(73)\times10^{-5}$ eV K^{-1}
Uncert. (ppm): 8.4

$N_A k(273.15\ \mathrm{K})/(101\,325\ \mathrm{Pa})$ **molar volume, ideal gas at STP**
Value: $22.414\,10(19)\times10^{-3}$ m^3 mol^{-1}
Uncert. (ppm): 8.4

$b = \lambda_{\max} T$ **Wien displacement law constant**
Value: $2.897\,756(24)\times10^{-3}$ m K
Uncert. (ppm): 8.4

$\sigma = \pi^2 k^4/60\hbar^3 c^2$ **Stefan-Boltzmann constant**
Value: $5.670\,51(19)\times10^{-8}$ W m^{-2} K^{-4}
Uncert. (ppm): 34

$G_F/(\hbar c)^3$ **Fermi coupling constant****
Value: $1.166\,39(1)\times10^{-5}$ GeV^{-2}
Uncert. (ppm): 9

** See discussion in Sec. 10 "Electroweak model and constraints on new physics."

$\sin^2\widehat{\theta}(M_Z)$ ($\overline{\text{MS}}$) **weak mixing angle**
Value: 0.23124(24)
Uncert. (ppm): 1000

m_W $\boldsymbol{W^\pm}$ **boson mass**
Value: 80.41(10) GeV/c^2
Uncert. (ppm): 1200

m_Z $\boldsymbol{Z^0}$ **boson mass**
Value: 91.187(7) GeV/c^2
Uncert. (ppm): 77

$\alpha_s(m_Z)$ **strong coupling constant**
Value: 0.119(2)
Uncert. (ppm): 17000

$\pi = 3.141\ 592\ 653\ 589\ 793\ 238$
$e = 2.718\ 281\ 828\ 459\ 045\ 235$
$\gamma = 0.577\ 215\ 664\ 901\ 532\ 861$

$$1 \text{ in} \equiv 0.0254 \text{ m}$$
$$1 \text{ Å} \equiv 10^{-10} \text{ m}$$
$$1 \text{ barn} \equiv 10^{-28} \text{ m}^2$$

$$1 \text{ G} \equiv 10^{-4} \text{ T}$$
$$1 \text{ dyne} \equiv 10^{-5} \text{ N}$$
$$1 \text{ erg} \equiv 10^{-7} \text{ J}$$

$$1 \text{ eV} = 1.602\ \ 177\ \ 33(49) \times 10^{-19} \text{ J}$$
$$1 \text{ eV}/c^2 = 1.782\ \ 662\ \ 70(54) \times 10^{-36} \text{ kg}$$
$$2.997\ \ 924\ \ 58 \times 10^9 \text{ esu} = 1 \text{ C}$$

$$kT \text{ at } 300 \text{ K} = [38.681\ \ 49(33)]^{-1} \text{ eV}$$
$$0\ \ ^\circ\text{C} \equiv 273.15 \text{ K}$$
$$1 \text{ atmosphere} \equiv 760 \text{ torr} \equiv 101\ \ 325 \text{ Pa}$$