

$\psi(2S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

### $\psi(2S)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>3686.00 ± 0.09 OUR AVERAGE</b>				
3686.02 ± 0.09 ± 0.27		ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3686.00 ± 0.10	413	ZHOLENTZ 80	OLYA	$e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3684 ± 2		GRIBUSHIN 96	FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$
3683 ± 5	77	ANTONIAZZI 94	E705	300 $\pi^\pm, p\text{Li} \rightarrow J/\psi \pi^+ \pi^- X$

### $m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>589.07 ± 0.13 OUR AVERAGE</b>			
589.7 ± 1.2	LEMOIGNE 82	GOLI	190 $\pi^- \text{Be} \rightarrow 2\mu$
589.07 ± 0.13	<sup>1</sup> ZHOLENTZ 80	OLYA	$e^+e^-$
588.7 ± 0.8	LUTH 75	MRK1	

<sup>1</sup> Redundant with data in mass above.

### $\psi(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>277 ± 31 OUR AVERAGE</b>	Error includes scale factor of 1.1.		
306 ± 36 ± 16	ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
243 ± 43	<sup>2</sup> PDG	92	RVUE

<sup>2</sup> Uses  $\Gamma(ee)$  from ALEXANDER 89 and  $B(ee) = (88 \pm 13) \times 10^{-4}$  from FELDMAN 77.

### $\psi(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ hadrons	(98.10 ± 0.30) %	
$\Gamma_2$ virtual $\gamma \rightarrow$ hadrons	( 2.9 ± 0.4 ) %	
$\Gamma_3$ $e^+e^-$	( 8.5 ± 0.7 ) × 10 <sup>-3</sup>	
$\Gamma_4$ $\mu^+\mu^-$	( 7.7 ± 1.7 ) × 10 <sup>-3</sup>	

#### Decays into $J/\psi(1S)$ and anything

$\Gamma_5$ $J/\psi(1S)$ anything	(54.2 ± 3.0) %	
$\Gamma_6$ $J/\psi(1S)$ neutrals	(22.8 ± 1.7) %	
$\Gamma_7$ $J/\psi(1S)\pi^+\pi^-$	(30.2 ± 1.9) %	
$\Gamma_8$ $J/\psi(1S)\pi^0\pi^0$	(17.9 ± 1.8) %	
$\Gamma_9$ $J/\psi(1S)\eta$	( 2.7 ± 0.4 ) %	S=1.7
$\Gamma_{10}$ $J/\psi(1S)\pi^0$	( 9.7 ± 2.1 ) × 10 <sup>-4</sup>	
$\Gamma_{11}$ $J/\psi(1S)\mu^+\mu^-$	(10.0 ± 3.3) × 10 <sup>-3</sup>	

**Hadronic decays**

$\Gamma_{12}$	$3(\pi^+\pi^-\pi^0)$	$(3.5 \pm 1.6) \times 10^{-3}$	
$\Gamma_{13}$	$2(\pi^+\pi^-\pi^0)$	$(3.0 \pm 0.8) \times 10^{-3}$	
$\Gamma_{14}$	$\pi^+\pi^-K^+K^-$	$(1.6 \pm 0.4) \times 10^{-3}$	
$\Gamma_{15}$	$\pi^+\pi^-p\bar{p}$	$(8.0 \pm 2.0) \times 10^{-4}$	
$\Gamma_{16}$	$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$	
$\Gamma_{17}$	$2(\pi^+\pi^-)$	$(4.5 \pm 1.0) \times 10^{-4}$	
$\Gamma_{18}$	$\rho^0\pi^+\pi^-$	$(4.2 \pm 1.5) \times 10^{-4}$	
$\Gamma_{19}$	$\bar{p}p$	$(1.9 \pm 0.5) \times 10^{-4}$	
$\Gamma_{20}$	$3(\pi^+\pi^-)$	$(1.5 \pm 1.0) \times 10^{-4}$	
$\Gamma_{21}$	$\bar{p}p\pi^0$	$(1.4 \pm 0.5) \times 10^{-4}$	
$\Gamma_{22}$	$K^+K^-$	$(1.0 \pm 0.7) \times 10^{-4}$	
$\Gamma_{23}$	$\pi^+\pi^-\pi^0$	$(9 \pm 5) \times 10^{-5}$	
$\Gamma_{24}$	$\rho\pi$	$< 8.3 \times 10^{-5}$	CL=90%
$\Gamma_{25}$	$\pi^+\pi^-$	$(8 \pm 5) \times 10^{-5}$	
$\Gamma_{26}$	$\Lambda\bar{\Lambda}$	$< 4 \times 10^{-4}$	CL=90%
$\Gamma_{27}$	$\Xi^-\Xi^+$	$< 2 \times 10^{-4}$	CL=90%
$\Gamma_{28}$	$K^+K^-\pi^0$	$< 2.96 \times 10^{-5}$	CL=90%
$\Gamma_{29}$	$K^+\bar{K}^*(892)^- + \text{c.c.}$	$< 5.4 \times 10^{-5}$	CL=90%

**Radiative decays**

$\Gamma_{30}$	$\gamma\chi_{c0}(1P)$	$(9.3 \pm 0.9) \%$	
$\Gamma_{31}$	$\gamma\chi_{c1}(1P)$	$(8.7 \pm 0.8) \%$	
$\Gamma_{32}$	$\gamma\chi_{c2}(1P)$	$(7.8 \pm 0.8) \%$	
$\Gamma_{33}$	$\gamma\eta_c(1S)$	$(2.8 \pm 0.6) \times 10^{-3}$	
$\Gamma_{34}$	$\gamma\eta_c(2S)$		
$\Gamma_{35}$	$\gamma\pi^0$		
$\Gamma_{36}$	$\gamma\eta'(958)$	$< 1.1 \times 10^{-3}$	CL=90%
$\Gamma_{37}$	$\gamma\eta$		
$\Gamma_{38}$	$\gamma\gamma$	$< 1.6 \times 10^{-4}$	CL=90%
$\Gamma_{39}$	$\gamma\eta(1440) \rightarrow \gamma K\bar{K}\pi$	$< 1.2 \times 10^{-4}$	CL=90%

**Mode needed for fitting purposes**

$\Gamma_{40}$	1. – other fit modes	$(22.4 \pm 3.3) \%$	
---------------	----------------------	---------------------	--

---

## CONSTRAINED FIT INFORMATION

An overall fit to 9 branching ratios uses 17 measurements and one constraint to determine 8 parameters. The overall fit has a  $\chi^2 = 8.9$  for 10 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_8$	25						
$x_9$	2	-8					
$x_{11}$	19	5	0				
$x_{30}$	0	0	0	0			
$x_{31}$	2	-5	-1	0	0		
$x_{32}$	1	-2	0	0	0	0	
$x_{40}$	-75	-66	-10	-24	-26	-22	-23
	$x_7$	$x_8$	$x_9$	$x_{11}$	$x_{30}$	$x_{31}$	$x_{32}$

## $\psi(2S)$ PARTIAL WIDTHS

### $\Gamma(\text{hadrons})$

 $\Gamma_1$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$224 \pm 56$	LUTH	75	MRK1 $e^+ e^-$

### $\Gamma(e^+ e^-)$

 $\Gamma_3$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b><math>2.14 \pm 0.21</math></b>	ALEXANDER	89	RVUE See $\Upsilon$ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$2.0 \pm 0.3$	BRANDELIK	79C	DASP $e^+ e^-$
$2.1 \pm 0.3$	<sup>3</sup> LUTH	75	MRK1 $e^+ e^-$

<sup>3</sup>From a simultaneous fit to  $e^+ e^-$ ,  $\mu^+ \mu^-$ , and hadronic channels assuming  $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$ .

### $\Gamma(\gamma\gamma)$

 $\Gamma_{38}$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;43</b>	90	BRANDELIK	79C	DASP $e^+ e^-$

### $\psi(2S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into  $e^+ e^-$  and with the total width is obtained from the integrated cross section into channel<sub>l</sub> in the  $e^+ e^-$  annihilation. We list only data that have not been used to determine the partial width  $\Gamma(l)$  or the branching ratio  $\Gamma(l)/\text{total}$ .

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_1 \Gamma_3/\Gamma$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.2 ± 0.4	ABRAMS	75	MRK1	$e^+ e^-$

### $\psi(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.981 ± 0.003	<sup>4</sup> LUTH	75	MRK1	$e^+ e^-$

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$				$\Gamma_2/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.029 ± 0.004	<sup>5</sup> LUTH	75	MRK1	$e^+ e^-$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_3/\Gamma$
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>85 ± 7 OUR AVERAGE</b>				
83 ± 5 ± 7	<sup>6</sup> ARMSTRONG	97	E760	$\bar{p}p \rightarrow \psi(2S)X$
88 ± 13	<sup>7</sup> FELDMAN	77	RVUE	$e^+ e^-$

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$				$\Gamma_4/\Gamma$
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
77 ± 17	<sup>8</sup> HILGER	75	SPEC	$e^+ e^-$

$\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$				$\Gamma_4/\Gamma_3$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.89 ± 0.16	BOYARSKI	75C	MRK1	$e^+ e^-$

<sup>4</sup> Includes cascade decay into  $J/\psi(1S)$ .

<sup>5</sup> Included in  $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ .

<sup>6</sup> Using  $B(J/\psi \rightarrow e^+ e^-) = 0.0599 \pm 0.0025$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \text{ anything}) = 0.04$ .

<sup>7</sup> From an overall fit assuming equal partial widths for  $e^+ e^-$  and  $\mu^+ \mu^-$ . For a measurement of the ratio see the entry  $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$  below. Includes LUTH 75, HILGER 75, BURMESTER 77.

<sup>8</sup> Restated by us using  $B(\psi(2S) \rightarrow J/\psi(1S) \text{ anything}) = 0.55$ .

DECAYS INTO  $J/\psi(1S)$  AND ANYTHING

$$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma = (\Gamma_7 + \Gamma_8 + \Gamma_9 + 0.273\Gamma_{31} + 0.135\Gamma_{32})/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.542 ± 0.030 OUR FIT</b>			
<b>0.55 ± 0.07 OUR AVERAGE</b>			
0.51 ± 0.12	BRANDELIK	79c DASP	$e^+e^- \rightarrow \mu^+\mu^-X$
0.57 ± 0.08	ABRAMS	75B MRK1	$e^+e^- \rightarrow \mu^+\mu^-X$

$$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma = (0.9761\Gamma_8 + 0.715\Gamma_9 + 0.273\Gamma_{31} + 0.135\Gamma_{32})/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.228 ± 0.017 OUR FIT</b>			

$$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\text{anything}) \quad \Gamma_6/\Gamma_5 = (0.9761\Gamma_8 + 0.715\Gamma_9 + 0.273\Gamma_{31} + 0.135\Gamma_{32})/(\Gamma_7 + \Gamma_8 + \Gamma_9 + 0.273\Gamma_{31} + 0.135\Gamma_{32})$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.421 ± 0.021 OUR FIT</b>			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.44 ± 0.03	<sup>9</sup> ABRAMS	75B MRK1	$e^+e^- \rightarrow J/\psi X$
-------------	---------------------	----------	-------------------------------

$$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_6/\Gamma_7 = (0.9761\Gamma_8 + 0.715\Gamma_9 + 0.273\Gamma_{31} + 0.135\Gamma_{32})/\Gamma_7$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.76 ± 0.07 OUR FIT</b>			
<b>0.73 ± 0.09</b>	<sup>9</sup> TANENBAUM	76 MRK1	$e^+e^-$

$$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.302 ± 0.019 OUR FIT</b>				
<b>0.296 ± 0.023 OUR AVERAGE</b>				
0.283 ± 0.021 ± 0.020	363	<sup>10</sup> ARMSTRONG	97 E760	$\bar{p}p \rightarrow \psi(2S)X$
0.32 ± 0.04		ABRAMS	75B MRK1	$e^+e^- \rightarrow J/\psi\pi^+\pi^-$

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.179 ± 0.018 OUR FIT</b>				
<b>0.184 ± 0.019 ± 0.013</b>	157	<sup>10</sup> ARMSTRONG	97 E760	$\bar{p}p \rightarrow \psi(2S)X$

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_8/\Gamma_7$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.59 ± 0.06 OUR FIT</b>			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.53 ± 0.06	<sup>11</sup> TANENBAUM	76 MRK1	$e^+e^-$
0.64 ± 0.15	<sup>12</sup> HILGER	75 SPEC	$e^+e^-$

$$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\mu^+\mu^-) \quad \Gamma_7/\Gamma_{11}$$


VALUE	DOCUMENT ID	TECN	COMMENT
<b>30 ± 10 OUR FIT</b>			
<b>30.2 ± 7.1 ± 6.8</b>	<sup>13</sup> GRIBUSHIN	96 FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$

$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$

$\Gamma_9/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.027 ± 0.004 OUR FIT</b>				Error includes scale factor of 1.7.
<b>0.027 ± 0.004 OUR AVERAGE</b>				Error includes scale factor of 1.6. See the ideogram below.
0.025 ± 0.006	166	HIMEL	80 MRK2	$e^+e^-$
0.0218 ± 0.0014 ± 0.0035	386	OREGLIA	80 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
0.036 ± 0.005	164	BARTEL	78B CNTR	$e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.032 ± 0.010 ± 0.002	36	14 ARMSTRONG	97 E760	$\bar{p}p \rightarrow \psi(2S)X$
0.035 ± 0.009	17	14 BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
0.043 ± 0.008	44	14 TANENBAUM	76 MRK1	$e^+e^-$

WEIGHTED AVERAGE  
0.027±0.004 (Error scaled by 1.6)



$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{total}$

$\Gamma_{10}/\Gamma$

<u>VALUE (units 10<sup>-4</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.7 ± 2.1 OUR AVERAGE</b>				
15 ± 6	7	HIMEL	80 MRK2	$e^+e^-$
9 ± 2 ± 1	23	OREGLIA	80 CBAL	$\psi(2S) \rightarrow J/\psi 2\gamma$

<sup>9</sup> The ABRAMS 75B measurement of  $\Gamma_6/\Gamma_5$  and the TANENBAUM 76 result for  $\Gamma_6/\Gamma_7$  are not independent. The TANENBAUM 76 result is used in the fit because it includes more accurate corrections for angular distributions.

<sup>10</sup> Using  $B(J/\psi \rightarrow e^+e^-) = 0.0599 \pm 0.0025$ .

<sup>11</sup> Not independent of the TANENBAUM 76 result for  $\Gamma_6/\Gamma_7$ .

<sup>12</sup> Ignoring the  $J/\psi(1S)\eta$  and  $J/\psi(1S)\gamma\gamma$  decays.

<sup>13</sup> Using  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$ .

<sup>14</sup> Low statistics data removed from average.

## HADRONIC DECAYS

$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$		$\Gamma_{12}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>35±16</b>	6	FRANKLIN 83	MRK2	$e^+e^- \rightarrow \text{hadrons}$
$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$		$\Gamma_{13}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>30±8</b>	42	FRANKLIN 83	MRK2	$e^+e^-$
$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$		$\Gamma_{14}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>16±4</b>		<sup>15</sup> TANENBAUM 78	MRK1	$e^+e^-$
$\Gamma(\pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}$		$\Gamma_{15}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8 ±2</b>		<sup>15</sup> TANENBAUM 78	MRK1	$e^+e^-$
$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma_{\text{total}}$		$\Gamma_{16}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>6.7±2.5</b>		TANENBAUM 78	MRK1	$e^+e^-$
$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$		$\Gamma_{17}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.5±1.0</b>		TANENBAUM 78	MRK1	$e^+e^-$
$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$		$\Gamma_{18}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.2±1.5</b>		TANENBAUM 78	MRK1	$e^+e^-$
$\Gamma(\bar{\rho}\rho)/\Gamma_{\text{total}}$		$\Gamma_{19}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.9±0.5 OUR AVERAGE</b>				
1.4±0.8	4	BRANDELIK 79C	DASP	$e^+e^-$
2.3±0.7		FELDMAN 77	MRK1	$e^+e^-$
$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$		$\Gamma_{20}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.5±1.0</b>		<sup>15</sup> TANENBAUM 78	MRK1	$e^+e^-$
$\Gamma(\bar{\rho}\rho\pi^0)/\Gamma_{\text{total}}$		$\Gamma_{21}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.4±0.5</b>	9	FRANKLIN 83	MRK2	$e^+e^-$

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<b>1.0±0.7</b>		BRANDELIK	79c DASP	$e^+ e^-$
----------------	--	-----------	----------	-----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.5	90	FELDMAN	77 MRK1	$e^+ e^-$
------	----	---------	---------	-----------

 $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<b>0.8±0.5</b>		BRANDELIK	79c DASP	$e^+ e^-$
----------------	--	-----------	----------	-----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.5	90	FELDMAN	77 MRK1	$e^+ e^-$
------	----	---------	---------	-----------

 $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

<b>0.85±0.46</b>	4	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow$ hadrons
------------------	---	----------	---------	-------------------------------

 $\Gamma(\Lambda \bar{\Lambda})/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<b>&lt;4</b>	90	FELDMAN	77 MRK1	$e^+ e^-$
--------------	----	---------	---------	-----------

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<b>&lt;2</b>	90	FELDMAN	77 MRK1	$e^+ e^-$
--------------	----	---------	---------	-----------

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

<b>&lt; 0.83</b>	90	1	FRANKLIN	83 MRK2	$e^+ e^-$
------------------	----	---	----------	---------	-----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10	90		BARTEL	76 CNTR	$e^+ e^-$
-----	----	--	--------	---------	-----------

<10	90	16	ABRAMS	75 MRK1	$e^+ e^-$
-----	----	----	--------	---------	-----------

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$ 

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

<b>&lt;2.96</b>	90	1	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow$ hadrons
-----------------	----	---	----------	---------	-------------------------------

 $\Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$ 

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<b>&lt;5.4</b>	90	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow$ hadrons
----------------	----	----------	---------	-------------------------------

<sup>15</sup> Assuming entirely strong decay.

<sup>16</sup> Final state  $\rho^0 \pi^0$ .



————— **RADIATIVE DECAYS** —————

**$\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$**   **$\Gamma_{30}/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.3±0.9 OUR FIT</b>			
<b>9.3±0.8 OUR AVERAGE</b>			
9.9±0.5±0.8	17 GAISER	86 CBAL	$e^+e^- \rightarrow \gamma X$
7.2±2.3	17 BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$
7.5±2.6	17 WHITAKER	76 MRK1	$e^+e^-$

**$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$**   **$\Gamma_{31}/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.7±0.8 OUR FIT</b>			
<b>8.7±0.8 OUR AVERAGE</b>			
9.0±0.5±0.7	18 GAISER	86 CBAL	$e^+e^- \rightarrow \gamma X$
7.1±1.9	19 BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$

**$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$**   **$\Gamma_{32}/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>7.8±0.8 OUR FIT</b>			
<b>7.8±0.8 OUR AVERAGE</b>			
8.0±0.5±0.7	20 GAISER	86 CBAL	$e^+e^- \rightarrow \gamma X$
7.0±2.0	19 BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$

**$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$**   **$\Gamma_{33}/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.28±0.06</b>	GAISER	86 CBAL	$e^+e^- \rightarrow \gamma X$

**$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$**   **$\Gamma_{34}/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.2 to 1.3	95	EDWARDS	82c CBAL	$e^+e^- \rightarrow \gamma X$

**$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{35}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 54	95	21 LIBERMAN	75 SPEC	$e^+e^-$
<100	90	WIJK	75 DASP	$e^+e^-$

**$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$**   **$\Gamma_{36}/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.11</b>	90	22 BARTEL	76 CNTR	$e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.6	90	23 BRAUNSCH...	77 DASP	$e^+e^-$

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$					$\Gamma_{37}/\Gamma$
VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.02	90	YAMADA	77	DASP	$e^+e^- \rightarrow 3\gamma$

$\Gamma(\gamma\eta(1440) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$					$\Gamma_{39}/\Gamma$
VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.12</b>	90	<sup>24</sup> SCHARRE	80	MRK1	$e^+e^-$
<sup>17</sup> Angular distribution $(1+\cos^2\theta)$ assumed. <sup>18</sup> Angular distribution $(1-0.189 \cos^2\theta)$ assumed. <sup>19</sup> Valid for isotropic distribution of the photon. <sup>20</sup> Angular distribution $(1-0.052 \cos^2\theta)$ assumed. <sup>21</sup> Restated by us using $B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.0077$ . <sup>22</sup> The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$ . <sup>23</sup> Restated by us using total decay width 228 keV. <sup>24</sup> Includes unknown branching fraction $\eta(1440) \rightarrow K \bar{K} \pi$ .					

### $\psi(2S)$ REFERENCES

ARMSTRONG	97	PR D55 1153	+Bettoni, Bharadwaj+	(E760 Collab.)
GRIBUSHIN	96	PR D53 4723	+Abramov, Antipov+	(E672 Collab., E706 Collab.)
ANTONIAZZI	94	PR D50 4258	+Arenton+	(E705 Collab.)
ARMSTRONG	93B	PR D47 772	+Bettoni, Bharadwaj+	(FNAL E760 Collab.)
PDG	92	PR D45, 1 June, Part II	Hikasa, Barnett, Stone+	(KEK, LBL, BOST+)
ALEXANDER	89	NP B320 45	+Bonvicini, Drell, Frey, Luth	(LBL, MICH, SLAC)
GAISER	86	PR D34 711	+Bloom, Bulos, Godfrey+	(Crystal Ball Collab.)
FRANKLIN	83	PRL 51 963	+Franklin, Feldman, Abrams, Alam+	(LBL, SLAC)
EDWARDS	82C	PRL 48 70	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
LEMOIGNE	82	PL 113B 509	+Barate, Astbury+	(SACL, LOIC, SHMP, IND)
HIMEL	80	PRL 44 920	+Abrams, Alam, Blocker+	(LBL, SLAC)
OREGLIA	80	PRL 45 959	+Partridge+	(SLAC, CIT, HARV, PRIN, STAN)
SCHARRE	80	PL 97B 329	+Trilling, Abrams, Alam, Blocker+	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	+Kurdadze, Lechuk, Mishnev+	(NOVO)
Also	81	SJNP 34 814	Zholentz, Kurdadze, Lechuk+	(NOVO)
Translated from YAF 34 1471.				
BRANDELIK	79B	NP B160 426	+Cords+	(DASP Collab.)
BRANDELIK	79C	ZPHY C1 233	+Cords+	(DASP Collab.)
BARTEL	78B	PL 79B 492	+Dittmann, Duinker, Olsson, O'Neill+	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	+Alam, Boyarski+	(SLAC, LBL)
BIDDICK	77	PRL 38 1324	+Burnett+	(UCSD, UMD, PAVI, PRIN, SLAC, STAN)
BRAUNSCH...	77	PL 67B 249	Braunschweig+	(DASP Collab.)
BURMESTER	77	PL 66B 395	+Criegee+	(DESY, HAMB, SIEG, WUPP)
FELDMAN	77	PRPL 33C 285	+Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69		(DASP Collab.)
BARTEL	76	PL 64B 483	+Duinker, Olsson, Steffen, Heintze+	(DESY, HEIDP)
TANENBAUM	76	PRL 36 402	+Abrams, Boyarski, Bulos+	(SLAC, LBL) IG
WHITAKER	76	PRL 37 1596	+Tanenbaum, Abrams, Alam+	(SLAC, LBL)
ABRAMS	75	Stanford Symp. 25		(LBL)
ABRAMS	75B	PRL 34 1181	+Briggs, Chinowsky, Friedberg+	(LBL, SLAC)
BOYARSKI	75C	Palermo Conf. 54	+Breidenbach, Bulos, Abrams, Briggs+	(SLAC, LBL)
HILGER	75	PRL 35 625	+Beron, Ford, Hofstadter, Howell+	(STAN, PENN)
LIBERMAN	75	Stanford Symp. 55		(STAN)
LUTH	75	PRL 35 1124	+Boyarski, Lynch, Breidenbach+	(SLAC, LBL) JPC
WIIK	75	Stanford Symp. 69		(DESY)

————— **OTHER RELATED PAPERS** —————

HOU	97	PR D55 6952	Wei-Shu Hou	
BARATE	83	PL 121B 449	+Bareyre, Bonamy+	(SACL, LOIC, SHMP, IND)
AUBERT	75B	PRL 33 1624	+Becker, Biggs, Burger, Glenn+	(MIT, BNL)
BRAUNSCH...	75B	PL 57B 407	Braunschweig, Konigs+	(DASP Collab.)
CAMERINI	75	PRL 35 483	+Learned, Prepost, Ash, Anderson+	(WISC, SLAC)
FELDMAN	75B	PRL 35 821	+Jean-Marie, Sadoulet, Vannucci+	(LBL, SLAC)
GRECO	75	PL 56B 367	+Pancheri-Srivastava, Srivastava	(FRAS)
JACKSON	75	NIM 128 13	+Scharre	(LBL)
SIMPSON	75	PRL 35 699	+Beron, Ford, Hilger, Hofstadter+	(STAN, PENN)
ABRAMS	74	PRL 33 1453	+Briggs, Augustin, Boyarski+	(LBL, SLAC)

---