

$\chi_{b1}(2P)$

$$J^G(J^{PC}) = 0^+(1^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

 $\chi_{b1}(2P)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.2552 ± 0.0005 OUR AVERAGE			
10.2547 ± 0.0004 ± 0.0010	¹ HEINTZ 92	CSB2	$e^+e^- \rightarrow \gamma X, \ell^+\ell^- \gamma\gamma$
10.2553 ± 0.0005	² MORRISON 91	CLE2	$e^+e^- \rightarrow \gamma X$

¹ From the average photon energy for inclusive and exclusive events and assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. Supersedes HEINTZ 91 and NARAIN 91.

² From γ energy below assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. The error on the $\Upsilon(3S)$ mass is not included in the individual measurements. It is included in the final evaluation.

 $m\chi_{b1}(2P) - m\chi_{b0}(2P)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
23.5 ± 0.7 ± 0.7	³ HEINTZ 92	CSB2	$e^+e^- \rightarrow \gamma X, \ell^+\ell^- \gamma\gamma$

³ From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

 γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
99.90 ± 0.26 OUR AVERAGE				
99 ± 1	169	CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^- \gamma\gamma$
100.1 ± 0.4	11147	⁴ HEINTZ 92	CSB2	$e^+e^- \rightarrow \gamma X$
100.2 ± 0.5	223	⁵ HEINTZ 92	CSB2	$e^+e^- \rightarrow \ell^+\ell^- \gamma\gamma$
99.5 ± 0.1 ± 0.5	25759	MORRISON 91	CLE2	$e^+e^- \rightarrow \gamma X$

⁴ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

⁵ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

 $\chi_{b1}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
$\Gamma_1 \quad \gamma \Upsilon(2S)$	(21 ± 4) %	1.5
$\Gamma_2 \quad \gamma \Upsilon(1S)$	(8.5 ± 1.3) %	1.3

$\chi_{b1}(2P)$ BRANCHING RATIOS $\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.21 ± 0.04 OUR AVERAGE Error includes scale factor of 1.5.0.356 ± 0.042 ± 0.092 ⁶ CRAWFORD 92B CLE2 $e^+e^- \rightarrow l^+l^-\gamma\gamma$ 0.199 ± 0.020 ± 0.022 ⁷ HEINTZ 92 CSB2 $e^+e^- \rightarrow l^+l^-\gamma\gamma$

⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma\gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (10.23 \pm 1.20 \pm 1.26) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

⁷ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.085 ± 0.013 OUR AVERAGE Error includes scale factor of 1.3.0.120 ± 0.021 ± 0.021 ⁸ CRAWFORD 92B CLE2 $e^+e^- \rightarrow l^+l^-\gamma\gamma$ 0.080 ± 0.009 ± 0.007 ⁹ HEINTZ 92 CSB2 $e^+e^- \rightarrow l^+l^-\gamma\gamma$

⁸ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma\gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (6.47 \pm 1.12 \pm 0.82) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

 $\chi_{b1}(2P)$ REFERENCES

CRAWFORD	92B	PL B294 139	+Fulton	(CLEO Collab.)
HEINTZ	92	PR D46 1928	+Lee, Franzini+	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	+Kaarsberg+	(CUSB Collab.)
MORRISON	91	PRL 67 1696	+Schmidt+	(CLEO Collab.)
NARAIN	91	PRL 66 3113	+Loveloock+	(CUSB Collab.)

OTHER RELATED PAPERS

EIGEN	82	PRL 49 1616	+Bohringer, Herb+	(CUSB Collab.)
HAN	82	PRL 49 1612	+Horstkotte, Imlay+	(CUSB Collab.)