

N(1710) P_{11}

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

The various partial-wave analyses do not agree very well.

N(1710) BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1680 to 1740 (\approx 1710) OUR ESTIMATE			
1717 \pm 28	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
1700 \pm 50	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1723 \pm 9	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1720 \pm 10	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
1766 \pm 34	¹ BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
1706	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
1692	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
1730	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
1690	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$
1650 to 1680	BAKER	78	DPWA $\pi^- p \rightarrow \Lambda K^0$
1721	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
1625 \pm 10	² BAKER	77	IPWA $\pi^- p \rightarrow \Lambda K^0$
1650	² BAKER	77	DPWA $\pi^- p \rightarrow \Lambda K^0$
1720	³ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
1670	KNASEL	75	DPWA $\pi^- p \rightarrow \Lambda K^0$
1710	⁴ LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

N(1710) BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
50 to 250 (\approx 100) OUR ESTIMATE			
480 \pm 230	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
93 \pm 30	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
90 \pm 30	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
120 \pm 15	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
105 \pm 10	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
185 \pm 61	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
540	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$
200	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
550	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
97	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$
90 to 150	BAKER	78	DPWA $\pi^- p \rightarrow \Lambda K^0$

167		BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$
160 ± 6		² BAKER	77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
95		² BAKER	77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
120		³ LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
174		KNASEL	75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
75		⁴ LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1710) POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1670 to 1770 (≈ 1720) OUR ESTIMATE			
1770	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1690	⁵ HOEHLER	93	SPED $\pi N \rightarrow \pi N$
1698	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
1690 ± 20	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1636	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
1708 or 1712	⁶ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
1720 or 1711	³ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

− 2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
80 to 380 (≈ 230) OUR ESTIMATE			
378	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
200	⁵ HOEHLER	93	SPED $\pi N \rightarrow \pi N$
88	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
80 ± 20	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
544	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
17 or 22	⁶ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
123 or 115	³ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

N(1710) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
37	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
15	HOEHLER	93	SPED $\pi N \rightarrow \pi N$
9	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
8 ± 2	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
149	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
− 167	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
− 167	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
175 ± 35	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
149	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

N(1710) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10–20 %
Γ_2 $N\eta$	
Γ_3 ΛK	5–25 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	40–90 %
Γ_6 $\Delta\pi$	15–40 %
Γ_7 $\Delta(1232)\pi$, P -wave	
Γ_8 $N\rho$	5–25 %
Γ_9 $N\rho$, $S=1/2$, P -wave	
Γ_{10} $N\rho$, $S=3/2$, P -wave	
Γ_{11} $N(\pi\pi)_{S\text{-wave}}^{I=0}$	10–40 %
Γ_{12} $p\gamma$	0.002–0.05%
Γ_{13} $p\gamma$, helicity=1/2	0.002–0.05%
Γ_{14} $n\gamma$	0.0–0.02%
Γ_{15} $n\gamma$, helicity=1/2	0.0–0.02%

N(1710) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.10 to 0.20 OUR ESTIMATE			
0.09±0.04	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
0.20±0.04	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
0.12±0.04	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.08±0.14	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$

$\Gamma(N\eta)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.16±0.10	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow N\eta$ $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.22	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$
+0.383	FELTESSE	75	DPWA Soln A; see BAKER 79

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow \Lambda K$	$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.12 to +0.18 OUR ESTIMATE			
+0.16	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$
+0.14	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.12	7 BAKER	78	DPWA See SAXON 80
-0.05 ± 0.03	2 BAKER	77	IPWA $\pi^- p \rightarrow \Lambda K^0$
-0.10	2 BAKER	77	DPWA $\pi^- p \rightarrow \Lambda K^0$
0.10	KNASEL	75	DPWA $\pi^- p \rightarrow \Lambda K^0$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow \Sigma K$	$(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.034	LIVANOS	80	DPWA $\pi p \rightarrow \Sigma K$
0.075 to 0.203	8 DEANS	75	DPWA $\pi N \rightarrow \Sigma K$

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow \Delta(1232)\pi$, <i>P-wave</i>	$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
±0.16 to ±0.22 OUR ESTIMATE			
-0.21 ± 0.04	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.17	3 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
+0.20	4 LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow N\rho$, <i>S=1/2, P-wave</i>	$(\Gamma_1 \Gamma_9)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
±0.09 to ±0.19 OUR ESTIMATE			
+0.05 ± 0.06	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
+0.19	3 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
-0.20	4 LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow N\rho$, <i>S=3/2, P-wave</i>	$(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.31	3 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow N(\pi\pi)_{S=0}^{I=0}$, <i>S-wave</i>	$(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
±0.14 to ±0.22 OUR ESTIMATE			
+0.04 ± 0.05	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.26	3 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
-0.28	4 LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

N(1710) PHOTON DECAY AMPLITUDES

N(1710) → pγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.009±0.022 OUR ESTIMATE			
0.007±0.015	ARNDT	96	IPWA γN → πN
0.006±0.018	CRAWFORD	83	IPWA γN → πN
0.028±0.009	AWAJI	81	DPWA γN → πN
-0.009±0.006	ARAI	80	DPWA γN → πN (fit 1)
-0.012±0.005	ARAI	80	DPWA γN → πN (fit 2)
0.015±0.025	CRAWFORD	80	DPWA γN → πN
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-0.037±0.002	LI	93	IPWA γN → πN
+0.001±0.039	BARBOUR	78	DPWA γN → πN
+0.053±0.019	FELLER	76	DPWA γN → πN

N(1710) → nγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.002±0.014 OUR ESTIMATE			
-0.002±0.015	ARNDT	96	IPWA γN → πN
0.000±0.018	AWAJI	81	DPWA γN → πN
-0.001±0.003	FUJII	81	DPWA γN → πN
0.005±0.013	ARAI	80	DPWA γN → πN (fit 1)
0.011±0.021	ARAI	80	DPWA γN → πN (fit 2)
-0.017±0.020	CRAWFORD	80	DPWA γN → πN
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.052±0.003	LI	93	IPWA γN → πN
-0.028±0.045	BARBOUR	78	DPWA γN → πN

N(1710) γp → ΛK⁺ AMPLITUDES

(Γ_iΓ_f)^{1/2}/Γ_{total} in pγ → N(1710) → ΛK⁺ (M₁₋ amplitude)

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●		
-10.6 ±0.4	WORKMAN	90 DPWA
- 7.21	TANABE	89 DPWA

pγ → N(1710) → ΛK⁺ phase angle θ (M₁₋ amplitude)

VALUE (degrees)	DOCUMENT ID	TECN
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●		
215 ±3	WORKMAN	90 DPWA
176.3	TANABE	89 DPWA

N(1710) FOOTNOTES

- ¹ BATINIC 95 finds a second state with a 6 MeV mass difference.
- ² The two BAKER 77 entries are from an IPWA using the Barrelet-zero method and from a conventional energy-dependent analysis.
- ³ LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ⁴ From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ⁵ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ⁶ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- ⁷ The overall phase of BAKER 78 couplings has been changed to agree with previous conventions.
- ⁸ The range given for DEANS 75 is from the four best solutions.

N(1710) REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

ARNDT	96	PR C53 430	+Strakovsky, Workman	(VPI)
ARNDT	95	PR C52 2120	+Strakovsky, Workman, Pavan	(VPI, BRCO)
BATINIC	95	PR C51 2310	+Slaus, Svarc, Nefkens	(BOSK, UCLA)
Also	98	PR C57 1004 (erratum)	M. Batinic+	
HOEHLER	93	πN Newsletter 9 1		(KARL)
LI	93	PR C47 2759	+Arndt, Roper, Workman	(VPI)
MANLEY	92	PR D45 4002	+Saleski	(KENT) IJP
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
ARNDT	91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
CUTKOSKY	90	PR D42 235	+Wang	(CMU)
WORKMAN	90	PR C42 781		(VPI)
TANABE	89	PR C39 741	+Kohno, Bennhold	(MANZ)
Also	89	NC 102A 193	Kohno, Tanabe, Bennhold	(MANZ)
BELL	83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+	(RL) IJP
CRAWFORD	83	NP B211 1	+Morton	(GLAS)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
FUJII	81	NP B187 53	+Hayashii, Iwata, Kajikawa+	(NAGO, OSAK)
ARAI	80	Toronto Conf. 93		(INUS)
Also	82	NP B194 251	Arai, Fujii	(INUS)
CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	+Baton, Coutures, Kochowski, Neveu	(SACL) IJP
SAXON	80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
BAKER	79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
BAKER	78	NP B141 29	+Blissett, Bloodworth, Broome+	(RL, CAVE) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons	(GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
BAKER	77	NP B126 365	+Blissett, Bloodworth, Broome, Hart+	(RHEL) IJP
LONGACRE	77	NP B122 493	+Dolbeau	(SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet	(SACL) IJP
FELLER	76	NP B104 219	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
DEANS	75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
FELTESSE	75	NP B93 242	+Ayed, Bareyre, Borgeaud, David+	(SACL) IJP
KNASEL	75	PR D11 1	+Lindquist, Nelson+	(CHIC, WUSL, OSU, ANL) IJP
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP
