

Δ(1600) P₃₃

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

The various analyses are not in good agreement.

Δ(1600) BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1550 to 1700 (≈ 1600) OUR ESTIMATE			
1706 ± 10	MANLEY	92	IPWA π N → π N & N π π
1600 ± 50	CUTKOSKY	80	IPWA π N → π N
1522 ± 13	HOEHLER	79	IPWA π N → π N
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1672 ± 15	ARNDT	96	IPWA γ N → π N
1706	LI	93	IPWA γ N → π N
1690	BARNHAM	80	IPWA π N → N π π
1560	¹ LONGACRE	77	IPWA π N → N π π
1640	² LONGACRE	75	IPWA π N → N π π

Δ(1600) BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
250 to 450 (≈ 350) OUR ESTIMATE			
430 ± 73	MANLEY	92	IPWA π N → π N & N π π
300 ± 100	CUTKOSKY	80	IPWA π N → π N
220 ± 40	HOEHLER	79	IPWA π N → π N
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
315 ± 20	ARNDT	96	IPWA γ N → π N
215	LI	93	IPWA γ N → π N
250	BARNHAM	80	IPWA π N → N π π
180	¹ LONGACRE	77	IPWA π N → N π π
300	² LONGACRE	75	IPWA π N → N π π

Δ(1600) POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1500 to 1700 (≈ 1600) OUR ESTIMATE			
1675	ARNDT	95	DPWA π N → N π
1550	³ HOEHLER	93	SPED π N → π N
1550 ± 40	CUTKOSKY	80	IPWA π N → π N
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1612	ARNDT	91	DPWA π N → π N Soln SM90
1609 or 1610	⁴ LONGACRE	78	IPWA π N → N π π
1541 or 1542	¹ LONGACRE	77	IPWA π N → N π π

– 2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
200 to 400 (≈ 300) OUR ESTIMATE			
386	ARNDT	95 DPWA	$\pi N \rightarrow N\pi$
200±60	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
230	ARNDT	91 DPWA	$\pi N \rightarrow \pi N$ Soln SM90
323 or 325	⁴ LONGACRE	78 IPWA	$\pi N \rightarrow N\pi\pi$
178 or 178	¹ LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$

Δ(1600) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
52	ARNDT	95 DPWA	$\pi N \rightarrow N\pi$
17±4	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
16	ARNDT	91 DPWA	$\pi N \rightarrow \pi N$ Soln SM90
PHASE θ			
VALUE (°)	DOCUMENT ID	TECN	COMMENT
+ 14	ARNDT	95 DPWA	$\pi N \rightarrow N\pi$
– 150±30	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
– 73	ARNDT	91 DPWA	$\pi N \rightarrow \pi N$ Soln SM90

Δ(1600) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10–25 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	75–90 %
Γ_4 $\Delta\pi$	40–70 %
Γ_5 $\Delta(1232)\pi$, <i>P</i> -wave	
Γ_6 $\Delta(1232)\pi$, <i>F</i> -wave	
Γ_7 $N\rho$	<25 %
Γ_8 $N\rho$, <i>S</i> =1/2, <i>P</i> -wave	
Γ_9 $N\rho$, <i>S</i> =3/2, <i>P</i> -wave	
Γ_{10} $N\rho$, <i>S</i> =3/2, <i>F</i> -wave	
Γ_{11} $N(1440)\pi$	10–35 %
Γ_{12} $N(1440)\pi$, <i>P</i> -wave	
Γ_{13} $N\gamma$	0.001–0.02 %
Γ_{14} $N\gamma$, helicity=1/2	0.0–0.02 %
Γ_{15} $N\gamma$, helicity=3/2	0.001–0.005 %

$\Delta(1600)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ Γ_1/Γ
VALUE DOCUMENT ID TECN COMMENT

0.10 to 0.25 OUR ESTIMATE

0.12±0.02	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$
0.18±0.04	CUTKOSKY	80	IPWA	$\pi N \rightarrow \pi N$
0.21±0.06	HOEHLER	79	IPWA	$\pi N \rightarrow \pi N$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow \Sigma K$ $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

-0.36 to -0.28 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.006 to 0.042	⁵ DEANS	75	DPWA	$\pi N \rightarrow \Sigma K$
----------------	--------------------	----	------	------------------------------

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620)$ S_{31} coupling to $\Delta(1232)\pi$.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow \Delta(1232)\pi$, *P-wave* $(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.27 to +0.33 OUR ESTIMATE

+0.29±0.02	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$
+0.24±0.05	BARNHAM	80	IPWA	$\pi N \rightarrow N\pi\pi$
+0.34	^{1,6} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.30	² LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow \Delta(1232)\pi$, *F-wave* $(\Gamma_1\Gamma_6)^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

-0.15 to -0.03 OUR ESTIMATE

-0.07	^{1,6} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
-------	-------------------------	----	------	-----------------------------

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow N\rho$, *S=1/2, P-wave* $(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.10	^{1,6} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
-------	-------------------------	----	------	-----------------------------

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow N\rho$, *S=3/2, P-wave* $(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.10	^{1,6} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
-------	-------------------------	----	------	-----------------------------

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow N(1440)\pi$, *P-wave* $(\Gamma_1\Gamma_{12})^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.15 to +0.23 OUR ESTIMATE

+0.16±0.02	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$
+0.23±0.04	BARNHAM	80	IPWA	$\pi N \rightarrow N\pi\pi$

$\Delta(1600)$ PHOTON DECAY AMPLITUDES

$\Delta(1600) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
−0.023±0.020 OUR ESTIMATE			
−0.018±0.015	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
−0.039±0.030	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
−0.046±0.013	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.005±0.020	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
−0.026±0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
−0.200	⁷ WADA	84	DPWA Compton scattering
0.000±0.030	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
0.0 ±0.020	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

$\Delta(1600) \rightarrow N\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
−0.009±0.021 OUR ESTIMATE			
−0.025±0.015	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
−0.013±0.014	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.025±0.031	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
−0.009±0.020	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
−0.016±0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
0.023	WADA	84	DPWA Compton scattering
0.000±0.045	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
0.0 ±0.015	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

$\Delta(1600)$ FOOTNOTES

¹ LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

² From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

³ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

⁴ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

⁵ The range given is from the four best solutions. DEANS 75 disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.

⁶ LONGACRE 77 considers this coupling to be well determined.

⁷ WADA 84 is inconsistent with other analyses — see the Note on N and Δ Resonances.

Δ(1600) REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

ARNDT	96	PR C53 430	+Strakovsky, Workman	(VPI)
ARNDT	95	PR C52 2120	+Strakovsky, Workman, Pavan	(VPI, BRCO)
HOEHLER	93	π <i>N</i> Newsletter 9 1		(KARL)
LI	93	PR C47 2759	+Arndt, Roper, Workman	(VPI)
MANLEY	92	PR D45 4002	+Saleski	(KENT) IJP
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
ARNDT	91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
WADA	84	NP B247 313	+Egawa, Imanishi, Ishii, Kato, Ukai+	(INUS)
CRAWFORD	83	NP B211 1	+Morton	(GLAS)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
BARNHAM	80	NP B168 243	+Glickman, Mier-Jedrzejowicz+	(LOIC)
CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons	(GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
LONGACRE	77	NP B122 493	+Dolbeau	(SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet	(SACL) IJP
WINNIK	77	NP B128 66	+Toaff, Revel, Goldberg, Berny	(HAIF) I
FELLER	76	NP B104 219	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
DEANS	75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP