

$\Lambda_c(2625)^+$

$I(J^P) = 0(\frac{3}{2}^-)$ Status: ***

Seen in $\Lambda_c^+ \pi^+ \pi^-$ but not in $\Lambda_c^+ \pi^0$ so this is indeed an excited Λ_c^+ rather than a Σ_c^+ . The spin-parity has not been measured but is expected to be $3/2^-$: this is presumably the charm counterpart of the strange $\Lambda(1520)$.


$\Lambda_c(2625)^+$ MASS

The mass is obtained from the $\Lambda_c(2625)^+ - \Lambda_c^+$ mass-difference measurements below.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-----------------------|---------|-------------------------------------|
| 2626.6 ± 0.8 OUR FIT | | | | Error includes scale factor of 1.2. |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2626.6 ± 0.5 ± 1.5 | 42 | ¹ ALBRECHT | 93F ARG | See ALBRECHT 97 |
| ¹ ALBRECHT 93F claims a signal of 42.4 ± 8.8 events. | | | | |

$\Lambda_c(2625)^+ - \Lambda_c^+$ MASS DIFFERENCE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-----------------------|---------|---|
| 341.7 ± 0.6 OUR FIT | | | | Error includes scale factor of 1.6. |
| 341.7 ± 0.6 OUR AVERAGE | | | | Error includes scale factor of 1.6. See the ideogram below. |
| 342.1 ± 0.5 ± 0.5 | 51 | ALBRECHT | 97 ARG | $e^+ e^- \approx 10$ GeV |
| 342.2 ± 0.2 ± 0.5 | 245 | ² EDWARDS | 95 CLE2 | $e^+ e^- \approx 10.5$ GeV |
| 340.4 ± 0.6 ± 0.3 | 40 | ³ FRABETTI | 94 E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |
| ² EDWARDS 95 claims a signal of 244.6 ± 19.0 events in $\Lambda_c^+ \pi^+ \pi^-$. | | | | |
| ³ FRABETTI 94 claims a signal of 39.7 ± 8.7 events. | | | | |



$$m_{\Lambda_c(2625)^+} - m_{\Lambda_c^+}$$

$\Lambda_c(2625)^+$ WIDTH

| VALUE (MeV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|---------|------------------------------------|
| <1.9 | 90 | 245 | EDWARDS | 95 CLE2 | $e^+ e^- \approx 10.5 \text{ GeV}$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <3.2 | 90 | | ALBRECHT | 93F ARG | $e^+ e^- \approx \Upsilon(4S)$ |

$\Lambda_c(2625)^+$ DECAY MODES

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma(2455) \pi$ are the only strong decays allowed to an excited Λ_c^+ having this mass.

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|---|--------------------------------|------------------|
| Γ_1 $\Lambda_c^+ \pi^+ \pi^-$ | [a] $\approx 67\%$ | |
| Γ_2 $\Sigma_c(2455)^{++} \pi^-$ | <5 | 90% |
| Γ_3 $\Sigma_c(2455)^0 \pi^+$ | <5 | 90% |
| Γ_4 $\Lambda_c^+ \pi^+ \pi^-$ 3-body | large | |
| Γ_5 $\Lambda_c^+ \pi^0$ | not seen | |
| Γ_6 $\Lambda_c^+ \gamma$ | not seen | |

[a] Assuming isospin conservation, so that the other third is $\Lambda_c^+ \pi^0 \pi^0$.

$\Lambda_c(2625)^+$ BRANCHING RATIOS

$$\Gamma(\Sigma_c(2455)^{++}\pi^-)/\Gamma(\Lambda_c^+\pi^+\pi^-) \quad \Gamma_2/\Gamma_1$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|---------|---------------------------|
| <0.08 | 90 | EDWARDS | 95 CLE2 | $e^+e^- \approx 10.5$ GeV |

$$\Gamma(\Sigma_c(2455)^0\pi^+)/\Gamma(\Lambda_c^+\pi^+\pi^-) \quad \Gamma_3/\Gamma_1$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|---------|---------------------------|
| <0.07 | 90 | EDWARDS | 95 CLE2 | $e^+e^- \approx 10.5$ GeV |

$$[\Gamma(\Sigma_c(2455)^{++}\pi^-) + \Gamma(\Sigma_c(2455)^0\pi^+)]/\Gamma(\Lambda_c^+\pi^+\pi^-) \quad (\Gamma_2+\Gamma_3)/\Gamma_1$$

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|-----|------|-------------|------|---------|
|-------|-----|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------|----|----------|---------|---|
| <0.36 | 90 | FRABETTI | 94 E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |
| 0.46 ± 0.14 | 21 | ALBRECHT | 93F ARG | $e^+e^- \approx \Upsilon(4S)$ |

$$\Gamma(\Lambda_c^+\pi^+\pi^- \text{ 3-body})/\Gamma(\Lambda_c^+\pi^+\pi^-) \quad \Gamma_4/\Gamma_1$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------|----|----------|---------|-------------------------------|
| 0.54 ± 0.14 | 16 | ALBRECHT | 93F ARG | $e^+e^- \approx \Upsilon(4S)$ |
|-----------------|----|----------|---------|-------------------------------|

$$\Gamma(\Lambda_c^+\pi^0)/\Gamma(\Lambda_c^+\pi^+\pi^-) \quad \Gamma_5/\Gamma_1$$

$\Lambda_c^+\pi^0$ decay is forbidden by isospin conservation if this state is in fact a Λ_c .

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|-------|----|---------|---------|---------------------------|
| <0.91 | 90 | EDWARDS | 95 CLE2 | $e^+e^- \approx 10.5$ GeV |
|-------|----|---------|---------|---------------------------|

$$\Gamma(\Lambda_c^+\gamma)/\Gamma(\Lambda_c^+\pi^+\pi^-) \quad \Gamma_6/\Gamma_1$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|-------|----|---------|---------|---------------------------|
| <0.52 | 90 | EDWARDS | 95 CLE2 | $e^+e^- \approx 10.5$ GeV |
|-------|----|---------|---------|---------------------------|

$\Lambda_c(2625)^+$ REFERENCES

| | | | | |
|----------|-----|-------------|-----------------------------|---------------------|
| ALBRECHT | 97 | PL B402 207 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| EDWARDS | 95 | PRL 74 3331 | K.W. Edwards <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 94 | PRL 72 961 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| ALBRECHT | 93F | PL B317 227 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |