BOTTOM BARYONS ($B = -1$)

$\Lambda_b^0 = udb$, $\Xi_b^0 = usb$, $\Xi_b^- = dsb$

Λ_b^0

$I(J^P) = 0(\frac{1}{2}^+)$

$I(J^P)$ not yet measured; $0(\frac{1}{2}^+)$ is the quark model prediction.

Mass $m = 5624 \pm 9$ MeV (S = 1.8)

Mean life $\tau = (1.229 \pm 0.080) \times 10^{-12}$ s

$c\tau = 368$ μm

These branching fractions are actually an average over weakly decaying b-baryons weighted by their production rates in Z decay (or high-energy $p\overline{p}$), branching ratios, and detection efficiencies. They scale with the LEP b-baryon production fraction B($b \rightarrow$ b-baryon) and are evaluated for our value B($b \rightarrow$ b-baryon) = (11.6 ± 2.0)%.

The branching fractions B(b-baryon $\rightarrow$ $\Lambda\ell^-\overline{\nu}_\ell$anything) and B($\Lambda_b^0 \rightarrow$ $\Lambda_c^+\ell^-\overline{\nu}_\ell$anything) are not pure measurements because the underlying measured products of these with B($b \rightarrow$ b-baryon) were used to determine B($b \rightarrow$ b-baryon), as described in the note "Production and Decay of b-Flavored Hadrons."

Λ_b^0 DECAY MODES		Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$J/\psi(1S)\Lambda$		$(4.7\pm2.8) \times 10^{-4}$		1744
$\Lambda_c^+\pi^-$		seen		2345
$\Lambda_c^+ a_1(1260)^-$		seen		2156
$\Lambda_c^+\ell^-\overline{\nu}_\ell$anything	[r]	(7.9±1.9) %		–
$p\pi^-$		$< 5.0 \times 10^{-5}$	90%	2732
pK^-		$< 5.0 \times 10^{-5}$	90%	2711

b-baryon ADMIXTURE (Λ_b, Ξ_b, Σ_b, Ω_b)

Mean life $\tau = (1.208 \pm 0.051) \times 10^{-12}$ s

These branching fractions are actually an average over weakly decaying b-baryons weighted by their production rates in Z decay (or high-energy $p\overline{p}$), branching ratios, and detection efficiencies. They scale with the LEP b-baryon production fraction B($b \rightarrow$ b-baryon) and are evaluated for our value B($b \rightarrow$ b-baryon) = (11.6 ± 2.0)%.

The branching fractions B(b-baryon $\rightarrow$ $\Lambda\ell^-\overline{\nu}_\ell$anything) and B($\Lambda_b^0 \rightarrow \Lambda_c^+\ell^-\overline{\nu}_\ell$anything) are not pure measurements because the underlying measured products of these with B($b \rightarrow$ b-baryon) were used to determine B($b \rightarrow$ b-baryon), as described in the note "Production and Decay of b-Flavored Hadrons."

***b*-baryon ADMIXTURE (Λ_b,Ξ_b,Σ_b,Ω_b)**	Fraction (Γ_i/Γ)	p (MeV/c)
$p\mu^-\overline{\nu}$anything	($4.2^{+\ 1.8}_{-\ 1.5}$) %	–
$p\ell\overline{\nu}_\ell$anything	(4.1± 1.0) %	–
$p$anything	(51 ±17) %	–
$\Lambda\ell^-\overline{\nu}_\ell$anything	(2.7± 0.8) %	–
$\Lambda/\overline{\Lambda}$anything	(28 ± 7) %	–
$\Xi^-\ell^-\overline{\nu}_\ell$anything	(4.8± 1.3) $\times 10^{-3}$	–