

$K_4^*(2045)$

$$I(J^P) = \frac{1}{2}(4^+)$$

 $K_4^*(2045)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
2045 ± 9 OUR AVERAGE		Error includes scale factor of 1.1.			
2062 ± 14 ± 13		¹ ASTON	86 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
2039 ± 10	400	^{2,3} CLELAND	82 SPEC	±	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
2070 ⁺¹⁰⁰ ₋₄₀		⁴ ASTON	81C LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2079 ± 7	431	TORRES	86 MPSF		400 $pA \rightarrow 4KX$
2088 ± 20	650	BAUBILLIER	82 HBC	-	8.25 $K^- p \rightarrow$ $K_S^0 \pi^- p$
2115 ± 46	488	CARMONY	77 HBC	0	9 $K^+ d \rightarrow K^+ \pi^+ s X$
¹ From a fit to all moments.					
² From a fit to 8 moments.					
³ Number of events evaluated by us.					
⁴ From energy-independent partial-wave analysis.					

 $K_4^*(2045)$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
198 ± 30 OUR AVERAGE					
221 ± 48 ± 27		⁵ ASTON	86 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
189 ± 35	400	^{6,7} CLELAND	82 SPEC	±	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
61 ± 58	431	TORRES	86 MPSF		400 $pA \rightarrow 4KX$
170 ⁺¹⁰⁰ ₋₅₀	650	BAUBILLIER	82 HBC	-	8.25 $K^- p \rightarrow$ $K_S^0 \pi^- p$
240 ⁺⁵⁰⁰ ₋₁₀₀		⁸ ASTON	81C LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
300 ± 200		CARMONY	77 HBC	0	9 $K^+ d \rightarrow K^+ \pi^+ s X$
⁵ From a fit to all moments.					
⁶ From a fit to 8 moments.					
⁷ Number of events evaluated by us.					
⁸ From energy-independent partial-wave analysis.					

$K_4^*(2045)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	$(9.9 \pm 1.2) \%$
Γ_2 $K^*(892)\pi\pi$	$(9 \pm 5) \%$
Γ_3 $K^*(892)\pi\pi\pi$	$(7 \pm 5) \%$
Γ_4 $\rho K\pi$	$(5.7 \pm 3.2) \%$
Γ_5 $\omega K\pi$	$(5.0 \pm 3.0) \%$
Γ_6 $\phi K\pi$	$(2.8 \pm 1.4) \%$
Γ_7 $\phi K^*(892)$	$(1.4 \pm 0.7) \%$

$K_4^*(2045)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.099 ± 0.012	ASTON 88 LASS 0 11 $K^- p \rightarrow K^- \pi^+ n$
$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$	Γ_2/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.89 ± 0.53	BAUBILLIER 82 HBC - 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$
$\Gamma(K^*(892)\pi\pi\pi)/\Gamma(K\pi)$	Γ_3/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.75 ± 0.49	BAUBILLIER 82 HBC - 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$
$\Gamma(\rho K\pi)/\Gamma(K\pi)$	Γ_4/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.58 ± 0.32	BAUBILLIER 82 HBC - 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$
$\Gamma(\omega K\pi)/\Gamma(K\pi)$	Γ_5/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.50 ± 0.30	BAUBILLIER 82 HBC - 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$
$\Gamma(\phi K\pi)/\Gamma_{\text{total}}$	Γ_6/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.028 ± 0.014	⁹ TORRES 86 MPSF 400 $pA \rightarrow 4KX$
$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$	Γ_7/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.014 ± 0.007	⁹ TORRES 86 MPSF 400 $pA \rightarrow 4KX$

⁹ Error determination is model dependent.

K_4^* (2045) REFERENCES

ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
TORRES	86	PR 34 707	S. Torres <i>et al.</i>	(VPI, ARIZ, FNAL, FSU+)
BAUBILLIER	82	PL 118B 447	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
CARMONY	77	PR D16 1251	D.D. Carmony <i>et al.</i>	(PURD, UCD, IUPU)

OTHER RELATED PAPERS

ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BROMBERG	80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)
CARMONY	71	PRL 27 1160	D.D. Carmony <i>et al.</i>	(PURD, UCD, IUPU)
