$\psi(2S)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\psi(2S)$ MASS

VALUE (MeV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
3685.96±0.09 OUR AVERAGE					
3685.95±0.10	413	[1] ARTAMONOV	00	OLYA	$e^+e^- \to$ hadrons
3686.02±0.09±0.27		ARMSTRONG	93B	E760	$\overline{p}p \to e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3684 ±2		GRIBUSHIN	96	FMPS	515 π^- Be $\to 2\mu X$
3683 ±5	77	ANTONIAZZI	94	E705	300 $\pi^\pm$, p Li $\to$ $J/\psi\pi^+\pi^-$ X
3686.00±0.10	413	[2] ZHOLENTZ	80	OLYA	e^+e^-

[1] Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).
[2] Superseded by ARTAMONOV 00.

$m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
589.07±0.13 OUR AVERAGE				
589.7 ±1.2	LEMOIGNE	82	GOLI	190 π^- Be $\to 2\mu$
589.07±0.13	[3] ZHOLENTZ	80	OLYA	e^+e^-
588.7 ±0.8	LUTH	75	MRK1	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
588 ±1	[4] BAI	98E	BES	e^+e^-

[3] Redundant with data in mass above.
[4] Systematic errors not evaluated.

$\psi(2S)$ WIDTH

VALUE (keV)	*DOCUMENT ID*		*TECN*	*COMMENT*
277±31 OUR AVERAGE Error includes scale factor of 1.1.				
306±36±16	ARMSTRONG	93B	E760	$\overline{p}p \to e^+e^-$
243±43	[5] PDG	92	RVUE	

[5] Uses $\Gamma(ee)$ from ALEXANDER 89 and $B(ee) = (88 \pm 13) \times 10^{-4}$ from FELDMAN 77.

$\psi(2S)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1	hadrons	(98.10±0.30) %	
Γ_2	virtual $\gamma \to$ hadrons	(2.9 ±0.4) %	
Γ_3	e^+e^-	(8.8 ±1.3) $\times 10^{-3}$	
Γ_4	$\mu^+\mu^-$	(1.03±0.35) %	

Decays into $J/\psi(1S)$ and anything

Γ_5	$J/\psi(1S)$ anything	(55 ±5) %	
Γ_6	$J/\psi(1S)$ neutrals	(23.1 ±2.3) %	
Γ_7	$J/\psi(1S)\pi^+\pi^-$	(31.0 ±2.8) %	
Γ_8	$J/\psi(1S)\pi^0\pi^0$	(18.2 ±2.3) %	
Γ_9	$J/\psi(1S)\eta$	(2.7 ±0.4) %	S=1.6
Γ_{10}	$J/\psi(1S)\pi^0$	(9.7 ±2.1) $\times 10^{-4}$	

Hadronic decays

Γ_{11}	$3(\pi^+\pi^-)\pi^0$	(3.5 ±1.6) $\times 10^{-3}$	
Γ_{12}	$2(\pi^+\pi^-)\pi^0$	(3.0 ±0.8) $\times 10^{-3}$	
Γ_{13}	$\omega f_2(1270)$	$< 1.7 \times 10^{-4}$	CL=90%
Γ_{14}	$\rho a_2(1320)$	$< 2.3 \times 10^{-4}$	CL=90%
Γ_{15}	$\pi^+\pi^- K^+ K^-$	(1.6 ±0.4) $\times 10^{-3}$	
Γ_{16}	$K^*(892)\overline{K}_2^*(1430)^0$	$< 1.2 \times 10^{-4}$	CL=90%
Γ_{17}	$K_1(1270)^\pm K^\mp$	(1.00±0.28) $\times 10^{-3}$	
Γ_{18}	$\pi^+\pi^- p\overline{p}$	(8.0 ±2.0) $\times 10^{-4}$	
Γ_{19}	$K^+\overline{K}^*(892)^0\pi^-$ + c.c.	(6.7 ±2.5) $\times 10^{-4}$	
Γ_{20}	$b_1^\pm\pi^\mp$	(5.2 ±1.3) $\times 10^{-4}$	
Γ_{21}	$2(\pi^+\pi^-)$	(4.5 ±1.0) $\times 10^{-4}$	
Γ_{22}	$\rho^0\pi^+\pi^-$	(4.2 ±1.5) $\times 10^{-4}$	
Γ_{23}	$\overline{p}p$	(1.9 ±0.5) $\times 10^{-4}$	
Γ_{24}	$3(\pi^+\pi^-)$	(1.5 ±1.0) $\times 10^{-4}$	
Γ_{25}	$\overline{p}p\pi^0$	(1.4 ±0.5) $\times 10^{-4}$	
Γ_{26}	K^+K^-	(1.0 ±0.7) $\times 10^{-4}$	
Γ_{27}	$\pi^+\pi^-\pi^0$	(8 ±5) $\times 10^{-5}$	
Γ_{28}	$\rho\pi$	$< 8.3 \times 10^{-5}$	CL=90%
Γ_{29}	$\pi^+\pi^-$	(8 ±5) $\times 10^{-5}$	
Γ_{30}	$\Lambda\overline{\Lambda}$	$< 4 \times 10^{-4}$	CL=90%
Γ_{31}	$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%
Γ_{32}	$\Xi^-\overline{\Xi}^+$	$< 2 \times 10^{-4}$	CL=90%
Γ_{33}	$K^+K^-\pi^0$	$< 2.96 \times 10^{-5}$	CL=90%
Γ_{34}	$K^+\overline{K}^*(892)^-$ + c.c.	$< 5.4 \times 10^{-5}$	CL=90%
Γ_{35}	$\phi f_2'(1525)$	$< 4.5 \times 10^{-5}$	CL=90%

Radiative decays

Γ_{36}	$\gamma\chi_{c0}(1P)$	(9.3 ±0.9) %	
Γ_{37}	$\gamma\chi_{c1}(1P)$	(8.7 ±0.8) %	
Γ_{38}	$\gamma\chi_{c2}(1P)$	(7.8 ±0.8) %	
Γ_{39}	$\gamma\eta_c(1S)$	(2.8 ±0.6) $\times 10^{-3}$	
Γ_{40}	$\gamma\eta_c(2S)$		

Γ_{41}	$\gamma\pi^0$			
Γ_{42}	$\gamma\eta'(958)$	$(\ 1.5\ \pm 0.4\ )\times 10^{-4}$		
Γ_{43}	$\gamma\eta$	$<\ 9$	$\times 10^{-5}$	CL=90%
Γ_{44}	$\gamma\gamma$	$<\ 1.6$	$\times 10^{-4}$	CL=90%
Γ_{45}	$\gamma\eta(1440)\rightarrow\ \gamma K\overline{K}\pi$	$<\ 1.2$	$\times 10^{-4}$	CL=90%

Mode needed for fitting purposes

Γ_{46}	1. − other fit modes	(21 ±5) %	

CONSTRAINED FIT INFORMATION

An overall fit to 10 branching ratios uses 17 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 9.0$ for 10 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\left\langle \delta x_i \delta x_j \right\rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\rm total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_7	27						
x_8	17	63					
x_9	2	9	3				
x_{36}	0	0	0	0			
x_{37}	0	−1	−5	0	0		
x_{38}	0	0	−2	0	0	0	
x_{46}	−30	−89	−83	−15	−17	−13	−15
	x_4	x_7	x_8	x_9	x_{36}	x_{37}	x_{38}

$\psi(2S)$ PARTIAL WIDTHS

Γ(hadrons) Γ_1

VALUE (keV)	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •				
224±56	LUTH	75	MRK1	e^+e^-

$\Gamma(e^+e^-)$ Γ_3

VALUE (keV)	*DOCUMENT ID*		*TECN*	*COMMENT*
2.12±0.18 OUR AVERAGE				
2.07±0.32	[6] BAI	98E	BES	e^+e^-
2.14±0.21	ALEXANDER	89	RVUE	See Υ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.0 ±0.3	BRANDELIK	79C	DASP	e^+e^-
2.1 ±0.3	[7] LUTH	75	MRK1	e^+e^-

[6] Value includes radiative corrections computed by ALEXANDER 89.

[7] From a simultaneous fit to e^+e^-, $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

$\Gamma(\gamma\gamma)$ Γ_{44}

VALUE (eV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<43	90	BRANDELIK	79C	DASP	e^+e^-

$\psi(2S)$ Γ(i)$\Gamma(e^+e^-)/\Gamma$(total)

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel$_I$ in the e^+e^- annihilation. We list only data that have not been used to determine the partial width Γ(I) or the branching ratio Γ(I)/total.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_3/\Gamma$

VALUE (keV)	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.2±0.4	ABRAMS	75	MRK1	e^+e^-

$\psi(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.981 ±0.003	[8] LUTH	75	MRK1	e^+e^-

$\Gamma(\text{virtual}\,\gamma \to \text{hadrons})/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.029±0.004	[9] LUTH	75	MRK1	e^+e^-

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10^{-4})	*DOCUMENT ID*		*TECN*	*COMMENT*
88±13	[10] FELDMAN	77	RVUE	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
83± 5±7	[11] ARMSTRONG	97	E760	$\overline{p}p \to \psi$(2S)X

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-4})	*DOCUMENT ID*		*TECN*	*COMMENT*
77±17	[12] HILGER	75	SPEC	e^+e^-

$\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_4/Γ_3

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.89±0.16	BOYARSKI	75C	MRK1	e^+e^-

[8] Includes cascade decay into $J/\psi(1S)$.
[9] Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.
[10] From an overall fit assuming equal partial widths for e^+e^- and $\mu^+\mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.
[11] Using B($J/\psi \to e^+e^-$) = 0.0599 ± 0.0025 and B($\psi(2S) \to J/\psi(1S)$anything) = 0.57 ± 0.04. Not an independent measurement, see GU 99.
[12] Restated by us using B($\psi(2S) \to J/\psi(1S)$anything) = 0.55.

DECAYS INTO $J/\psi(1S)$ AND ANYTHING

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma = (\Gamma_7+\Gamma_8+\Gamma_9+0.273\Gamma_{37}+0.135\Gamma_{38})/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
0.55±0.05 OUR FIT				
0.55±0.07 OUR AVERAGE				
0.51±0.12	BRANDELIK	79C	DASP	$e^+e^- \to \mu^+\mu^-$X
0.57±0.08	ABRAMS	75B	MRK1	$e^+e^- \to \mu^+\mu^-$X

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma = (0.9761\Gamma_8+0.715\Gamma_9+0.273\Gamma_{37}+0.135\Gamma_{38})/\Gamma$

VALUE	DOCUMENT ID
0.231±0.023 OUR FIT	

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\text{anything})$ $\Gamma_6/\Gamma_5 = (0.9761\Gamma_8+0.715\Gamma_9+0.273\Gamma_{37}+0.135\Gamma_{38})/(\Gamma_7+\Gamma_8+\Gamma_9+0.273\Gamma_{37}+0.135\Gamma_{38})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.418±0.019 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.44 ±0.03	[13] ABRAMS	75B	MRK1	$e^+e^- \to J/\psi$X

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$ $\Gamma_6/\Gamma_7 = (0.9761\Gamma_8+0.715\Gamma_9+0.273\Gamma_{37}+0.135\Gamma_{38})/\Gamma_7$

VALUE	DOCUMENT ID		TECN	COMMENT
0.75±0.06 OUR FIT				
0.73±0.09	[13] TANENBAUM	76	MRK1	e^+e^-

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.310±0.028 OUR FIT					
0.32 ±0.04		ABRAMS	75B	MRK1	$e^+e^- \to J/\psi\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.283±0.021±0.020	363	[14] ARMSTRONG	97	E760	$\overline{p}p \to \psi$(2S)X

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.182±0.023 OUR FIT					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.184±0.019±0.013	157	[14] ARMSTRONG	97	E760	$\overline{p}p \to \psi$(2S)X

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_8/Γ_7

VALUE	DOCUMENT ID		TECN	COMMENT
0.59 ±0.06 OUR FIT				
0.609±0.079	[15] GU	99	RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.53 ±0.06	[16] TANENBAUM	76	MRK1	e^+e^-
0.64 ±0.15	[17] HILGER	75	SPEC	e^+e^-

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(\mu^+\mu^-)$ Γ_7/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
30 ±10 OUR FIT			
30.2± 7.1±6.8	[18] GRIBUSHIN 96	FMPS	515 π^- Be $\rightarrow 2\mu X$

$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$ Γ_9/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.027 ±0.004 OUR FIT Error includes scale factor of 1.6.				
0.027 ±0.004 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.				
0.025 ±0.006	166	HIMEL 80	MRK2	e^+e^-
0.0218±0.0014±0.0035	386	OREGLIA 80	CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
0.036 ±0.005	164	BARTEL 78B	CNTR	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.032 ±0.010 ±0.002	36	[19] ARMSTRONG 97	E760	$\overline{p}p \rightarrow \psi(2S)X$
0.035 ±0.009	17	[19] BRANDELIK 79B	DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
0.043 ±0.008	44	[19] TANENBAUM 76	MRK1	e^+e^-



$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$

$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$ $\Gamma_9/\Gamma_5 = \Gamma_9/(\Gamma_7+\Gamma_8+\Gamma_9+0.273\Gamma_{37}+0.135\Gamma_{38})$

VALUE	DOCUMENT ID	TECN
0.049±0.008 OUR FIT Error includes scale factor of 1.3.		
0.062±0.016	[15] GU 99	RVUE

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{total}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
9.7±2.1 OUR AVERAGE					
15 ±6	7	HIMEL	80	MRK2	e^+e^-
9 ±2 ±1	23	OREGLIA	80	CBAL	$\psi(2S) \to J/\psi 2\gamma$

[13] The ABRAMS 75B measurement of Γ_6/Γ_5 and the TANENBAUM 76 result for Γ_6/Γ_7 are not independent. The TANENBAUM 76 result is used in the fit because it includes more accurate corrections for angular distributions.
[14] Using B($J/\psi \to e^+e^-$) = 0.0599 ± 0.0025 and B($\psi(2S) \to J/\psi(1S)$anything)= 0.57 ± 0.04.
[15] Using data from ARMSTRONG 97.
[16] Not independent of the TANENBAUM 76 result for Γ_6/Γ_7.
[17] Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.
[18] Using B($J/\psi(1S) \to \mu^+\mu^-$) = 0.0597 ± 0.0025.
[19] Low statistics data removed from average.

HADRONIC DECAYS

$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{total}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
35±16	6	FRANKLIN	83	MRK2	$e^+e^- \to$ hadrons

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{total}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
30±8	42	FRANKLIN	83	MRK2	e^+e^-

$\Gamma(\pi^+\pi^- K^+K^-)/\Gamma_{total}$ Γ_{15}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
16±4	[20] TANENBAUM	78	MRK1	e^+e^-

$\Gamma(K_1(1270)^{\pm}K^{\mp})/\Gamma_{total}$ Γ_{17}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
10.0±1.8±2.1	[21] BAI	99C	BES	e^+e^-

$\Gamma(\pi^+\pi^- p\overline{p})/\Gamma_{total}$ Γ_{18}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
8 ±2	[20] TANENBAUM	78	MRK1	e^+e^-

$\Gamma(K^+\overline{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma_{total}$ Γ_{19}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
6.7±2.5	TANENBAUM	78	MRK1	e^+e^-

$\Gamma(b_1^{\pm}\pi^{\mp})/\Gamma_{total}$ Γ_{20}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
5.2±0.8±1.0	[22] BAI	99C	BES	e^+e^-

$\Gamma(2(\pi^+\pi^-))/\Gamma_{total}$ Γ_{21}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
4.5±1.0	TANENBAUM	78	MRK1	e^+e^-

$\Gamma(\omega f_2(1270))/\Gamma_{total}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<1.7	90	BAI	98J	BES	e^+e^-

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{total}$ Γ_{22}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
4.2±1.5	TANENBAUM	78	MRK1	e^+e^-

$\Gamma(\rho a_2(1320))/\Gamma_{total}$ Γ_{14}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2.3	90	BAI	98J	BES	e^+e^-

$\Gamma(\bar{p}p)/\Gamma_{total}$ Γ_{23}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
1.9±0.5 OUR AVERAGE					
1.4±0.8	4	BRANDELIK	79C	DASP	e^+e^-
2.3±0.7		FELDMAN	77	MRK1	e^+e^-

$\Gamma(3(\pi^+\pi^-))/\Gamma_{total}$ Γ_{24}/Γ

VALUE (units 10^{-4})		DOCUMENT ID		TECN	COMMENT
1.5±1.0	[20]	TANENBAUM	78	MRK1	e^+e^-

$\Gamma(\bar{p}p\pi^0)/\Gamma_{total}$ Γ_{25}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
1.4±0.5	9	FRANKLIN	83	MRK2	e^+e^-

$\Gamma(K^+K^-)/\Gamma_{total}$ Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
1.0±0.7		BRANDELIK	79C	DASP	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.5	90	FELDMAN	77	MRK1	e^+e^-

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$ Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
0.8±0.5		BRANDELIK	79C	DASP	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.5	90	FELDMAN	77	MRK1	e^+e^-

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{27}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
0.85±0.46	4	FRANKLIN	83	MRK2	$e^+e^- \rightarrow$ hadrons

$\Gamma(\Lambda\overline{\Lambda})/\Gamma_{total}$ Γ_{30}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<4	90	FELDMAN	77	MRK1	e^+e^-

$\Gamma(K_1(1400)^{\pm}K^{\mp})/\Gamma_{total}$ Γ_{31}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<3.1	90	[23] BAI	99C	BES	e^+e^-

$\Gamma(\Xi^-\overline{\Xi}^+)/\Gamma_{total}$ Γ_{32}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2	90	FELDMAN	77	MRK1	e^+e^-

$\Gamma(\rho\pi)/\Gamma_{total}$ Γ_{28}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
< 0.83	90	1	FRANKLIN	83	MRK2	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<10	90		BARTEL	76	CNTR	e^+e^-
<10	90		[24] ABRAMS	75	MRK1	e^+e^-

$\Gamma(K^+K^-\pi^0)/\Gamma_{total}$ Γ_{33}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
<2.96	90	1	FRANKLIN	83	MRK2	$e^+e^- \rightarrow$ hadrons

$\Gamma(K^+\overline{K}^*(892)^- + \text{c.c.})/\Gamma_{total}$ Γ_{34}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<5.4	90	FRANKLIN	83	MRK2	$e^+e^- \rightarrow$ hadrons

$\Gamma(K^*(892)\overline{K}_2^*(1430)^0)/\Gamma_{total}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<1.2	90	BAI	98J	BES	e^+e^-

$\Gamma(\phi f_2'(1525))/\Gamma_{total}$ Γ_{35}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<0.45	90	BAI	98J	BES	$e^+e^- \rightarrow 2(K^+K^-)$

[20] Assuming entirely strong decay.
[21] Assuming B($K_1(1270) \rightarrow K\rho$)=0.42 ± 0.06
[22] Assuming B($b_1 \rightarrow \omega\pi$)=1.
[23] Assuming B($K_1(1400) \rightarrow K^*\pi$)=0.94 ± 0.06
[24] Final state $\rho^0\pi^0$.

RADIATIVE DECAYS

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{total}$ Γ_{36}/Γ

VALUE (units 10^{-2})	DOCUMENT ID		TECN	COMMENT
9.3±0.9 OUR FIT				
9.3±0.8 OUR AVERAGE				
9.9±0.5±0.8	[25] GAISER	86	CBAL	$e^+e^- \rightarrow \gamma X$
7.2±2.3	[25] BIDDICK	77	CNTR	$e^+e^- \rightarrow \gamma X$
7.5±2.6	[25] WHITAKER	76	MRK1	e^+e^-

$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{total}$ Γ_{37}/Γ

VALUE (units 10^{-2})	*DOCUMENT ID*		*TECN*	*COMMENT*
8.7±0.8 OUR FIT				
8.7±0.8 OUR AVERAGE				
9.0±0.5±0.7	[26] GAISER	86	CBAL	$e^+e^- \to \gamma X$
7.1±1.9	[27] BIDDICK	77	CNTR	$e^+e^- \to \gamma X$

$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{total}$ Γ_{38}/Γ

VALUE (units 10^{-2})	*DOCUMENT ID*		*TECN*	*COMMENT*
7.8±0.8 OUR FIT				
7.8±0.8 OUR AVERAGE				
8.0±0.5±0.7	[28] GAISER	86	CBAL	$e^+e^- \to \gamma X$
7.0±2.0	[27] BIDDICK	77	CNTR	$e^+e^- \to \gamma X$

$\Gamma(\gamma\eta_c(1S))/\Gamma_{total}$ Γ_{39}/Γ

VALUE (units 10^{-2})	*DOCUMENT ID*		*TECN*	*COMMENT*
0.28±0.06	GAISER	86	CBAL	$e^+e^- \to \gamma X$

$\Gamma(\gamma\eta_c(2S))/\Gamma_{total}$ Γ_{40}/Γ

VALUE (units 10^{-2})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.2 to 1.3	95	EDWARDS	82C	CBAL	$e^+e^- \to \gamma X$

$\Gamma(\gamma\pi^0)/\Gamma_{total}$ Γ_{41}/Γ

VALUE (units 10^{-4})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 54	95	[29] LIBERMAN	75	SPEC	e^+e^-
<100	90	WIIK	75	DASP	e^+e^-

$\Gamma(\gamma\eta'(958))/\Gamma_{total}$ Γ_{42}/Γ

VALUE (units 10^{-4})	*CL%*	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
1.54±0.31±0.20		~ 43	BAI	98F	BES	$\psi(2S) \to \pi^+\pi^- 2\gamma$, $\pi^+\pi^- 3\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<60	90		[30] BRAUNSCH...	77	DASP	e^+e^-
<11	90		[31] BARTEL	76	CNTR	e^+e^-

$\Gamma(\gamma\eta)/\Gamma_{total}$ Γ_{43}/Γ

VALUE (units 10^{-4})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<0.9	90	BAI	98F	BES	$\psi(2S) \to \pi^+\pi^- 3\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2	90	YAMADA	77	DASP	$e^+e^- \to 3\gamma$

$\Gamma(\gamma\eta(1440) \to \gamma K\overline{K}\pi)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-3})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<0.12	90	[32] SCHARRE	80	MRK1	$e^+ e^-$

[25] Angular distribution $(1+\cos^2\theta)$ assumed.
[26] Angular distribution $(1-0.189\ \cos^2\theta)$ assumed.
[27] Valid for isotropic distribution of the photon.
[28] Angular distribution $(1-0.052\ \cos^2\theta)$ assumed.
[29] Restated by us using $B(\psi(2S) \to \mu^+\mu^-) = 0.0077$.
[30] Restated by us using total decay width 228 keV.
[31] The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.
[32] Includes unknown branching fraction $\eta(1440) \to K\overline{K}\pi$.

$\psi(2S)$ REFERENCES

ARTAMONOV	00	PL B474 427	A.S. Artamonov *et al.*	
BAI	99C	PRL 83 1918	J.Z. Bai *et al.*	(BES Collab.)
GU	99	PL B449 361	Y.F. Gu, X.H. Li	
BAI	98E	PR D57 3854	J.Z. Bai *et al.*	(BES Collab.)
BAI	98F	PR D58 097101	J.Z. Bai *et al.*	(BES Collab.)
BAI	98J	PRL 81 5080	J.Z. Bai *et al.*	(BES Collab.)
ARMSTRONG	97	PR D55 1153	T.A. Armstrong *et al.*	(E760 Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin *et al.*	
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi *et al.*	(E705 Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong *et al.*	(FNAL E760 Collab.)
PDG	92	PR D45, 1 June, Part II	K. Hikasa *et al.*	(KEK, LBL, BOST+)
ALEXANDER	89	NP B320 45	J.P. Alexander *et al.*	(LBL, MICH, SLAC)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
GAISER	86	PR D34 711	J. Gaiser *et al.*	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
FRANKLIN	83	PRL 51 963	M.E.B. Franklin *et al.*	(LBL, SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards *et al.*	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne *et al.*	(SACL, LOIC, SHMP+)
HIMEL	80	PRL 44 920	T. Himel *et al.*	(LBL, SLAC)
OREGLIA	80	PRL 45 959	M.J. Oreglia *et al.*	(SLAC, CIT, HARV+)
SCHARRE	80	PL 97B 329	D.L. Scharre *et al.*	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents *et al.*	(NOVO)
Also	81	SJNP 34 814	A.A. Zholents *et al.*	(NOVO)
		Translated from YAF 34 1471.		
BRANDELIK	79B	NP B160 426	R. Brandelik *et al.*	(DASP Collab.)
BRANDELIK	79C	ZPHY C1 233	R. Brandelik *et al.*	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel *et al.*	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum *et al.*	(SLAC, LBL)
BIDDICK	77	PRL 38 1324	C.J. Biddick *et al.*	(UCSD, UMD, PAVI+)
BRAUNSCH...	77	PL 67B 249	W. Braunschweig *et al.*	(DASP Collab.)
BURMESTER	77	PL 66B 395	J. Burmester *et al.*	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
BARTEL	76	PL 64B 483	W. Bartel *et al.*	(DESY, HEIDP)
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum *et al.*	(SLAC, LBL) IG
WHITAKER	76	PRL 37 1596	J.S. Whitaker *et al.*	(SLAC, LBL)
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)
ABRAMS	75B	PRL 34 1181	G.S. Abrams *et al.*	(LBL, SLAC)
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski *et al.*	(SLAC, LBL)
HILGER	75	PRL 35 625	E. Hilger *et al.*	(STAN, PENN)
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)
LUTH	75	PRL 35 1124	V. Luth *et al.*	(SLAC, LBL) JPC
WIIK	75	Stanford Symp. 69	B.H. Wiik	(DESY)

OTHER RELATED PAPERS

CHEN	98	PRL 80 5060	Y.Q. Chen, E. Braaten	
SUZUKI	98	PR D57 5717	M. Suzuki	
HOU	97	PR D55 6952	W.-S. Hou	
BARATE	83	PL 121B 449	R. Barate *et al.*	(SACL, LOIC, SHMP, IND)
AUBERT	75B	PRL 33 1624	J.J. Aubert *et al.*	(MIT, BNL)
BRAUNSCH...	75B	PL 57B 407	W. Braunschweig *et al.*	(DASP Collab.)
CAMERINI	75	PRL 35 483	U. Camerini *et al.*	(WISC, SLAC)
FELDMAN	75B	PRL 35 821	G.J. Feldman *et al.*	(LBL, SLAC)
GRECO	75	PL 56B 367	M. Greco, G. Pancheri-Srivastava, Y. Srivastava	
JACKSON	75	NIM 128 13	J.D. Jackson, D.L. Scharre	(LBL)
SIMPSON	75	PRL 35 699	J.W. Simpson *et al.*	(STAN, PENN)
ABRAMS	74	PRL 33 1453	G.S. Abrams *et al.*	(LBL, SLAC)