$\omega(1420)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\omega(1420)$ MASS

VALUE (MeV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
1419± 31	315	[1] ANTONELLI	92	DM2	1.34–2.4 $e^+e^- \to \rho\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1170± 10		[2] ACHASOV	99E	RVUE	0.75–1.80 $e^+e^- \to \pi^+\pi^-\pi^0$
1400^{+100}_{-200}		[3] ACHASOV	98H	RVUE	$e^+e^- \to \pi^+\pi^-\pi^0$
~ 1400		[4] ACHASOV	98H	RVUE	$e^+e^- \to \omega\pi^+\pi^-$
~ 1460		[5] ACHASOV	98H	RVUE	$e^+e^- \to K^+K^-$
1440± 70		[6] CLEGG	94	RVUE	

[1] From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed (+,−,+) phases.
[2] Using the data of DOLINSKY 91, ANTONELLI 92, AKHMETSHIN 98, and ACHASOV 99E. From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed (+,−,+) phases.
[3] Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.
[4] Using the data from ANTONELLI 92.
[5] Using the data from IVANOV 81 and BISELLO 88B.
[6] Using data published by ANTONELLI 92.

$\omega(1420)$ WIDTH

VALUE (MeV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
174±59	315	[7] ANTONELLI	92	DM2	1.34–2.4 $e^+e^- \to \rho\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
187±15		[8] ACHASOV	99E	RVUE	0.75–1.80 $e^+e^- \to \pi^+\pi^-\pi^0$
240±70		[9] CLEGG	94	RVUE	

[7] From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed (+,−,+) phases.
[8] Using the data of DOLINSKY 91, ANTONELLI 92, AKHMETSHIN 98, and ACHASOV 99E. From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed (+,−,+) phases.
[9] Using data published by ANTONELLI 92.

$\omega(1420)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)
Γ_1	$\rho\pi$	dominant
Γ_2	e^+e^-	

$\omega(1420)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma(total)$

$\Gamma(\rho\pi) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_1\Gamma_2/\Gamma$

VALUE (eV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
81±31	315	[10] ANTONELLI	92	DM2	1.34–2.4 $e^+e^- \to \rho\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

137± 3±15	[11] ACHASOV	99E	RVUE	0.75–1.80 $e^+e^- \to \pi^+\pi^-\pi^0$

[10] From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed (+,−,+) phases.

[11] Using the data of DOLINSKY 91, ANTONELLI 92, AKHMETSHIN 98, and ACHASOV 99E. From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed (+,−,+) phases.

$\omega(1420)$ REFERENCES

ACHASOV	99E	PL B462 365	M.N. Achasov *et al.*	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
AKHMETSHIN	98	PL B434 426	R.R. Akhmetshin *et al.*	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli *et al.*	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky *et al.*	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello *et al.*	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov *et al.*	(NOVO)
		Translated from ZETFP 46 132.		
IVANOV	81	PL 107B 297	P.M. Ivanov *et al.*	(NOVO)

OTHER RELATED PAPERS

ABELE	99D	PL B468 178	A. Abele *et al.*	(Crystal Barrel Collab.)
BELOZEROVA	98	PPN 29 63	T.S. Belozerova, V.K. Henner	
		Translated from FECAY 29 148.		
ACHASOV	97F	PAN 60 2029	N.N. Achasov, A.A. Kozhevnikov	(NOVM)
		Translated from YAF 60 2212.		
ATKINSON	87	ZPHY C34 157	M. Atkinson *et al.*	(BONN, CERN, GLAS+)
ATKINSON	84	NP B231 15	M. Atkinson *et al.*	(BONN, CERN, GLAS+)
ATKINSON	83B	PL 127B 132	M. Atkinson *et al.*	(BONN, CERN, GLAS+)