

$$I(J^P) = \frac{1}{2}(0^-)$$

D^\pm MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , and $D_s^{*\pm}$ mass and mass difference measurements.

| <u>VALUE (MeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|------------------------|-------------|------------------------------|
| 1869.3 ± 0.5 OUR FIT | Error includes scale factor of 1.1. | | | |
| 1869.4 ± 0.5 OUR AVERAGE | | | | |
| 1870.0 ± 0.5 ± 1.0 | 317 | BARLAG | 90C ACCM | π^- Cu 230 GeV |
| 1863 ± 4 | | DERRICK | 84 HRS | $e^+ e^-$ 29 GeV |
| 1869.4 ± 0.6 | | ¹ TRILLING | 81 RVUE | $e^+ e^-$ 3.77 GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 1875 ± 10 | 9 | ADAMOVICH | 87 EMUL | Photoproduction |
| 1860 ± 16 | 6 | ADAMOVICH | 84 EMUL | Photoproduction |
| 1868.4 ± 0.5 | | ¹ SCHINDLER | 81 MRK2 | $e^+ e^-$ 3.77 GeV |
| 1874 ± 5 | | GOLDHABER | 77 MRK1 | D^0 , D^+ recoil spectra |
| 1868.3 ± 0.9 | | ¹ PERUZZI | 77 MRK1 | $e^+ e^-$ 3.77 GeV |
| 1874 ± 11 | | PICCOLO | 77 MRK1 | $e^+ e^-$ 4.03, 4.41 GeV |
| 1876 ± 15 | 50 | PERUZZI | 76 MRK1 | $K^\mp \pi^\pm \pi^\pm$ |

¹ PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision $J/\psi(1S)$ and $\psi(2S)$ measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted.

D^\pm MEAN LIFE

Measurements with an error $> 0.1 \times 10^{-12}$ s are omitted from the average, and those with an error $> 0.2 \times 10^{-12}$ s have been omitted from the Listings.

| <u>VALUE (10^{-12} s)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--|---------------------|-------------|--|
| 1.051 ± 0.013 OUR NEW AVERAGE | [[$(1.057 \pm 0.015) \times 10^{-12}$ s OUR 1998 AVERAGE] | | | |
| 1.0336 ± 0.0221 ^{+0.0099} / _{-0.0127} | 3777 | BONVICINI | 99 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |
| 1.048 ± 0.015 ± 0.011 | 9k | FRABETTI | 94D E687 | $D^+ \rightarrow K^- \pi^+ \pi^+$ |
| 1.075 ± 0.040 ± 0.018 | 2455 | FRABETTI | 91 E687 | γ Be, $D^+ \rightarrow K^- \pi^+ \pi^+$ |
| 1.03 ± 0.08 ± 0.06 | 200 | ALVAREZ | 90 NA14 | γ , $D^+ \rightarrow K^- \pi^+ \pi^+$ |
| 1.05 ^{+0.077} / _{-0.072} | 317 | ² BARLAG | 90C ACCM | π^- Cu 230 GeV |
| 1.05 ± 0.08 ± 0.07 | 363 | ALBRECHT | 88I ARG | $e^+ e^-$ 10 GeV |
| 1.090 ± 0.030 ± 0.025 | 2992 | RAAB | 88 E691 | Photoproduction |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|------|--|------------|--------------------|----------|---------------------------|------------------|
| 1.12 | $\begin{matrix} +0.14 \\ -0.11 \end{matrix}$ | 149 | AGUILAR-... | 87D HYBR | $\pi^- p$ and pp | |
| 1.09 | $\begin{matrix} +0.19 \\ -0.15 \end{matrix}$ | 59 | BARLAG | 87B ACCM | K^- and π^- 200 GeV | |
| 1.14 | ± 0.16 | ± 0.07 | 247 | CSORNA | 87 CLEO | $e^+ e^-$ 10 GeV |
| 1.09 | ± 0.14 | 74 | ³ PALKA | 87B SILI | π Be 200 GeV | |
| 0.86 | $\begin{matrix} \pm 0.13 \\ +0.07 \\ -0.03 \end{matrix}$ | 48 | ABE | 86 HYBR | γp 20 GeV | |

² BARLAG 90C estimates the systematic error to be negligible.

³ PALKA 87B observes this in $D^+ \rightarrow \bar{K}^*(892)e\nu$.

D^+ DECAY MODES

D^- modes are charge conjugates of the modes below.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|---|--|-----------------------------------|
| Inclusive modes | | |
| Γ_1 e^+ anything | $(17.2 \pm 1.9) \%$ | |
| Γ_2 K^- anything | $(24.2 \pm 2.8) \%$ | S=1.4 |
| Γ_3 \bar{K}^0 anything + K^0 anything | $(59 \pm 7) \%$ | |
| Γ_4 K^+ anything | $(5.8 \pm 1.4) \%$ | |
| Γ_5 η anything | [a] < 13 % | CL=90% |
| Γ_6 μ^+ anything | | |
| Leptonic and semileptonic modes | | |
| Γ_7 $\mu^+ \nu_\mu$ | $(8 \begin{matrix} +17 \\ -5 \end{matrix}) \times 10^{-4}$ | |
| Γ_8 $\bar{K}^0 \ell^+ \nu_\ell$ | [b] $(6.8 \pm 0.8) \%$ | |
| Γ_9 $\bar{K}^0 e^+ \nu_e$ | $(6.7 \pm 0.9) \%$ | |
| Γ_{10} $\bar{K}^0 \mu^+ \nu_\mu$ | $(7.0 \begin{matrix} +3.0 \\ -2.0 \end{matrix}) \%$ | |
| Γ_{11} $K^- \pi^+ e^+ \nu_e$ | $(4.1 \begin{matrix} +0.9 \\ -0.7 \end{matrix}) \%$ | |
| Γ_{12} $\bar{K}^*(892)^0 e^+ \nu_e$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$ | $(3.2 \pm 0.33) \%$ | |
| Γ_{13} $K^- \pi^+ e^+ \nu_e$ nonresonant | < 7 $\times 10^{-3}$ | CL=90% |
| Γ_{14} $K^- \pi^+ \mu^+ \nu_\mu$ | $(3.2 \pm 0.4) \%$ | S=1.1 |
| In the fit as $\frac{2}{3}\Gamma_{26} + \Gamma_{16}$, where $\frac{2}{3}\Gamma_{26} = \Gamma_{15}$. | | |
| Γ_{15} $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$ | $(2.9 \pm 0.4) \%$ | |

| | | | |
|---------------|---|--------------------------------------|-------------------------|
| Γ_{16} | $K^- \pi^+ \mu^+ \nu_\mu$ nonresonant | $(2.7 \pm 1.1) \times 10^{-3}$ | |
| Γ_{17} | $\bar{K}^0 \pi^+ \pi^- e^+ \nu_e$ | | |
| Γ_{18} | $K^- \pi^+ \pi^0 e^+ \nu_e$ | | |
| Γ_{19} | $(\bar{K}^*(892)\pi)^0 e^+ \nu_e$ | < 1.2 | % CL=90% |
| Γ_{20} | $(\bar{K}\pi\pi)^0 e^+ \nu_e$ non- $\bar{K}^*(892)$ | < 9 | $\times 10^{-3}$ CL=90% |
| Γ_{21} | $K^- \pi^+ \pi^0 \mu^+ \nu_\mu$ | < 1.4 | $\times 10^{-3}$ CL=90% |
| Γ_{22} | $\pi^0 \ell^+ \nu_\ell$ | [c] $(3.1 \pm 1.5) \times 10^{-3}$ | |
| Γ_{23} | $\pi^+ \pi^- e^+ \nu_e$ | | |

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

| | | | |
|---------------|-------------------------------------|----------------------------------|-------------------------|
| Γ_{24} | $\bar{K}^*(892)^0 \ell^+ \nu_\ell$ | [b] $(4.7 \pm 0.4) \%$ | |
| Γ_{25} | $\bar{K}^*(892)^0 e^+ \nu_e$ | $(4.8 \pm 0.5) \%$ | |
| Γ_{26} | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ | $(4.4 \pm 0.6) \%$ | S=1.1 |
| Γ_{27} | $\bar{K}_1(1270)^0 \mu^+ \nu_\mu$ | < 3.5 | % CL=95% |
| Γ_{28} | $\bar{K}^*(1410)^0 \mu^+ \nu_\mu$ | < 2.7 | % CL=95% |
| Γ_{29} | $\bar{K}_2^*(1430)^0 \mu^+ \nu_\mu$ | < 8 | $\times 10^{-3}$ CL=95% |
| Γ_{30} | $\rho^0 e^+ \nu_e$ | $(2.2 \pm 0.8) \times 10^{-3}$ | |
| Γ_{31} | $\rho^0 \mu^+ \nu_\mu$ | $(2.7 \pm 0.7) \times 10^{-3}$ | |
| Γ_{32} | $\phi e^+ \nu_e$ | < 2.09 | % CL=90% |
| Γ_{33} | $\phi \mu^+ \nu_\mu$ | < 3.72 | % CL=90% |
| Γ_{34} | $\eta \ell^+ \nu_\ell$ | < 5 | $\times 10^{-3}$ CL=90% |
| Γ_{35} | $\eta'(958) \mu^+ \nu_\mu$ | < 9 | $\times 10^{-3}$ CL=90% |

Hadronic modes with a \bar{K} or $\bar{K}K\bar{K}$

| | | | |
|---------------|--|----------------------------------|-------|
| Γ_{36} | $\bar{K}^0 \pi^+$ | $(2.89 \pm 0.26) \%$ | S=1.1 |
| Γ_{37} | $K^- \pi^+ \pi^+$ | [d] $(9.0 \pm 0.6) \%$ | |
| Γ_{38} | $\bar{K}^*(892)^0 \pi^+$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$ | $(1.27 \pm 0.13) \%$ | |
| Γ_{39} | $\bar{K}_0^*(1430)^0 \pi^+$ $\times B(\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+)$ | $(2.3 \pm 0.3) \%$ | |
| Γ_{40} | $\bar{K}^*(1680)^0 \pi^+$ $\times B(\bar{K}^*(1680)^0 \rightarrow K^- \pi^+)$ | $(3.7 \pm 0.8) \times 10^{-3}$ | |
| Γ_{41} | $K^- \pi^+ \pi^+$ nonresonant | $(8.5 \pm 0.8) \%$ | |
| Γ_{42} | $\bar{K}^0 \pi^+ \pi^0$ | [d] $(9.7 \pm 3.0) \%$ | S=1.1 |
| Γ_{43} | $\bar{K}^0 \rho^+$ | $(6.6 \pm 2.5) \%$ | |
| Γ_{44} | $\bar{K}^*(892)^0 \pi^+$ $\times B(\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0)$ | $(6.3 \pm 0.4) \times 10^{-3}$ | |
| Γ_{45} | $\bar{K}^0 \pi^+ \pi^0$ nonresonant | $(1.3 \pm 1.1) \%$ | |
| Γ_{46} | $K^- \pi^+ \pi^+ \pi^0$ | [d] $(6.4 \pm 1.1) \%$ | |
| Γ_{47} | $\bar{K}^*(892)^0 \rho^+$ total $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$ | $(1.4 \pm 0.9) \%$ | |
| Γ_{48} | $\bar{K}_1(1400)^0 \pi^+$ $\times B(\bar{K}_1(1400)^0 \rightarrow K^- \pi^+ \pi^0)$ | $(2.2 \pm 0.6) \%$ | |

| | | | |
|-----------------|--|--|--------|
| Γ ₄₉ | $K^- \rho^+ \pi^+$ total | (3.1 ± 1.1) % | |
| Γ ₅₀ | $K^- \rho^+ \pi^+$ 3-body | (1.1 ± 0.4) % | |
| Γ ₅₁ | $\bar{K}^*(892)^0 \pi^+ \pi^0$ total | (4.5 ± 0.9) % | |
| | × B($\bar{K}^{*0} \rightarrow K^- \pi^+$) | | |
| Γ ₅₂ | $\bar{K}^*(892)^0 \pi^+ \pi^0$ 3-body | (2.8 ± 0.9) % | |
| | × B($\bar{K}^{*0} \rightarrow K^- \pi^+$) | | |
| Γ ₅₃ | $K^*(892)^- \pi^+ \pi^+$ 3-body | (7 ± 3) × 10 ⁻³ | |
| | × B($K^{*-} \rightarrow K^- \pi^0$) | | |
| Γ ₅₄ | $K^- \pi^+ \pi^+ \pi^0$ nonresonant | [e] (1.2 ± 0.6) % | |
| Γ ₅₅ | $\bar{K}^0 \pi^+ \pi^+ \pi^-$ | [d] (7.0 ± 0.9) % | |
| Γ ₅₆ | $\bar{K}^0 a_1(1260)^+$ | (4.0 ± 0.9) % | |
| | × B($a_1(1260)^+ \rightarrow \pi^+ \pi^+ \pi^-$) | | |
| Γ ₅₇ | $\bar{K}_1(1400)^0 \pi^+$ | (2.2 ± 0.6) % | |
| | × B($\bar{K}_1(1400)^0 \rightarrow \bar{K}^0 \pi^+ \pi^-$) | | |
| Γ ₅₈ | $K^*(892)^- \pi^+ \pi^+$ 3-body | (1.4 ± 0.6) % | |
| | × B($K^{*-} \rightarrow \bar{K}^0 \pi^-$) | | |
| Γ ₅₉ | $\bar{K}^0 \rho^0 \pi^+$ total | (4.2 ± 0.9) % | |
| Γ ₆₀ | $\bar{K}^0 \rho^0 \pi^+$ 3-body | (5 ± 5) × 10 ⁻³ | |
| Γ ₆₁ | $\bar{K}^0 \pi^+ \pi^+ \pi^-$ nonresonant | (8 ± 4) × 10 ⁻³ | |
| Γ ₆₂ | $K^- \pi^+ \pi^+ \pi^+ \pi^-$ | [d] (7.2 ± 1.0) × 10 ⁻³ | |
| Γ ₆₃ | $\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$ | (5.4 ± 2.3) × 10 ⁻³ | |
| | × B($\bar{K}^{*0} \rightarrow K^- \pi^+$) | | |
| Γ ₆₄ | $\bar{K}^*(892)^0 \rho^0 \pi^+$ | (1.9 ^{+ 1.1} _{- 1.0}) × 10 ⁻³ | |
| | × B($\bar{K}^{*0} \rightarrow K^- \pi^+$) | | |
| Γ ₆₅ | $\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$ no- ρ | (2.9 ± 1.1) × 10 ⁻³ | |
| | × B($\bar{K}^{*0} \rightarrow K^- \pi^+$) | | |
| Γ ₆₆ | $K^- \rho^0 \pi^+ \pi^+$ | (3.1 ± 0.9) × 10 ⁻³ | |
| Γ ₆₇ | $K^- \pi^+ \pi^+ \pi^+ \pi^-$ nonresonant | < 2.3 × 10 ⁻³ | CL=90% |
| Γ ₆₈ | $K^- \pi^+ \pi^+ \pi^0 \pi^0$ | (2.2 ^{+ 5.0} _{- 0.9}) % | |
| Γ ₆₉ | $\bar{K}^0 \pi^+ \pi^+ \pi^- \pi^0$ | (5.4 ^{+ 3.0} _{- 1.4}) % | |
| Γ ₇₀ | $\bar{K}^0 \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ | (8 ± 7) × 10 ⁻⁴ | |
| Γ ₇₁ | $K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^0$ | (2.0 ± 1.8) × 10 ⁻³ | |
| Γ ₇₂ | $\bar{K}^0 \bar{K}^0 K^+$ | (1.8 ± 0.8) % | |

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

| | | | |
|-----------------|----------------------------------|------------------------|--------|
| Γ ₇₃ | $\bar{K}^0 \rho^+$ | (6.6 ± 2.5) % | |
| Γ ₇₄ | $\bar{K}^0 a_1(1260)^+$ | (8.0 ± 1.7) % | |
| Γ ₇₅ | $\bar{K}^0 a_2(1320)^+$ | < 3 × 10 ⁻³ | CL=90% |
| Γ ₇₆ | $\bar{K}^*(892)^0 \pi^+$ | (1.90 ± 0.19) % | |
| Γ ₇₇ | $\bar{K}^*(892)^0 \rho^+$ total | [e] (2.1 ± 1.3) % | |
| Γ ₇₈ | $\bar{K}^*(892)^0 \rho^+$ S-wave | [e] (1.6 ± 1.6) % | |
| Γ ₇₉ | $\bar{K}^*(892)^0 \rho^+$ P-wave | < 1 × 10 ⁻³ | CL=90% |

| | | | |
|---------------|--|--|--------|
| Γ_{80} | $\bar{K}^*(892)^0 \rho^+ D\text{-wave}$ | $(10 \pm 7) \times 10^{-3}$ | |
| Γ_{81} | $\bar{K}^*(892)^0 \rho^+ D\text{-wave longitudinal}$ | $< 7 \times 10^{-3}$ | CL=90% |
| Γ_{82} | $\bar{K}_1(1270)^0 \pi^+$ | $< 7 \times 10^{-3}$ | CL=90% |
| Γ_{83} | $\bar{K}_1(1400)^0 \pi^+$ | $(4.9 \pm 1.2) \%$ | |
| Γ_{84} | $\bar{K}^*(1410)^0 \pi^+$ | $< 7 \times 10^{-3}$ | CL=90% |
| Γ_{85} | $\bar{K}_0^*(1430)^0 \pi^+$ | $(3.7 \pm 0.4) \%$ | |
| Γ_{86} | $\bar{K}^*(1680)^0 \pi^+$ | $(1.43 \pm 0.30) \%$ | |
| Γ_{87} | $\bar{K}^*(892)^0 \pi^+ \pi^0 \text{ total}$ | $(6.7 \pm 1.4) \%$ | |
| Γ_{88} | $\bar{K}^*(892)^0 \pi^+ \pi^0 \text{ 3-body}$ | [e] $(4.2 \pm 1.4) \%$ | |
| Γ_{89} | $K^*(892)^- \pi^+ \pi^+ \text{ total}$ | | |
| Γ_{90} | $K^*(892)^- \pi^+ \pi^+ \text{ 3-body}$ | $(2.0 \pm 0.9) \%$ | |
| Γ_{91} | $K^- \rho^+ \pi^+ \text{ total}$ | $(3.1 \pm 1.1) \%$ | |
| Γ_{92} | $K^- \rho^+ \pi^+ \text{ 3-body}$ | $(1.1 \pm 0.4) \%$ | |
| Γ_{93} | $\bar{K}^0 \rho^0 \pi^+ \text{ total}$ | $(4.2 \pm 0.9) \%$ | CL=90% |
| Γ_{94} | $\bar{K}^0 \rho^0 \pi^+ \text{ 3-body}$ | $(5 \pm 5) \times 10^{-3}$ | |
| Γ_{95} | $\bar{K}^0 f_0(980) \pi^+$ | $< 5 \times 10^{-3}$ | CL=90% |
| Γ_{96} | $\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$ | $(8.1 \pm 3.4) \times 10^{-3}$ | S=1.7 |
| Γ_{97} | $\bar{K}^*(892)^0 \rho^0 \pi^+$ | $(2.9 \pm_{-1.5}^{+1.7}) \times 10^{-3}$ | S=1.8 |
| Γ_{98} | $\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^- \text{ no-}\rho$ | $(4.3 \pm 1.7) \times 10^{-3}$ | |
| Γ_{99} | $K^- \rho^0 \pi^+ \pi^+$ | $(3.1 \pm 0.9) \times 10^{-3}$ | |

Pionic modes

| | | | |
|----------------|--|--|--------|
| Γ_{100} | $\pi^+ \pi^0$ | $(2.5 \pm 0.7) \times 10^{-3}$ | |
| Γ_{101} | $\pi^+ \pi^+ \pi^-$ | $(3.6 \pm 0.4) \times 10^{-3}$ | |
| Γ_{102} | $\rho^0 \pi^+$ | $(1.05 \pm 0.31) \times 10^{-3}$ | |
| Γ_{103} | $\pi^+ \pi^+ \pi^- \text{ nonresonant}$ | $(2.2 \pm 0.4) \times 10^{-3}$ | |
| Γ_{104} | $\pi^+ \pi^+ \pi^- \pi^0$ | $(1.9 \pm_{-1.2}^{+1.5}) \%$ | |
| Γ_{105} | $\eta \pi^+ \times \text{B}(\eta \rightarrow \pi^+ \pi^- \pi^0)$ | $(6.9 \pm 1.4) \times 10^{-4}$ | |
| Γ_{106} | $\omega \pi^+ \times \text{B}(\omega \rightarrow \pi^+ \pi^- \pi^0)$ | $< 6 \times 10^{-3}$ | CL=90% |
| Γ_{107} | $\pi^+ \pi^+ \pi^+ \pi^- \pi^-$ | $(2.1 \pm 0.4) \times 10^{-3}$ | |
| Γ_{108} | $\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$ | $(2.9 \pm_{-2.0}^{+2.9}) \times 10^{-3}$ | |

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

| | | | |
|----------------|---------------------|----------------------------------|--------|
| Γ_{109} | $\eta \pi^+$ | $(3.0 \pm 0.6) \times 10^{-3}$ | |
| Γ_{110} | $\rho^0 \pi^+$ | $(1.05 \pm 0.31) \times 10^{-3}$ | |
| Γ_{111} | $\omega \pi^+$ | $< 7 \times 10^{-3}$ | CL=90% |
| Γ_{112} | $\eta \rho^+$ | $< 7 \times 10^{-3}$ | CL=90% |
| Γ_{113} | $\eta'(958) \pi^+$ | $(5.0 \pm 1.0) \times 10^{-3}$ | |
| Γ_{114} | $\eta'(958) \rho^+$ | $< 5 \times 10^{-3}$ | CL=90% |

Hadronic modes with a $K\bar{K}$ pair

| | | | | |
|----------------|---|-----|--------------------------------|--------|
| Γ_{115} | $K^+\bar{K}^0$ | | $(7.4 \pm 1.0) \times 10^{-3}$ | |
| Γ_{116} | $K^+K^-\pi^+$ | [d] | $(8.7 \pm 0.7) \times 10^{-3}$ | |
| Γ_{117} | $\phi\pi^+ \times B(\phi \rightarrow K^+K^-)$ | | $(3.0 \pm 0.3) \times 10^{-3}$ | |
| Γ_{118} | $K^+\bar{K}^*(892)^0$ $\times B(\bar{K}^{*0} \rightarrow K^-\pi^+)$ | | $(2.8 \pm 0.4) \times 10^{-3}$ | |
| Γ_{119} | $K^+K^-\pi^+$ nonresonant | | $(4.5 \pm 0.9) \times 10^{-3}$ | |
| Γ_{120} | $K^0\bar{K}^0\pi^+$ | | — | |
| Γ_{121} | $K^*(892)^+\bar{K}^0$ $\times B(K^{*+} \rightarrow K^0\pi^+)$ | | $(2.1 \pm 1.0) \%$ | |
| Γ_{122} | $K^+K^-\pi^+\pi^0$ | | — | |
| Γ_{123} | $\phi\pi^+\pi^0 \times B(\phi \rightarrow K^+K^-)$ | | $(1.1 \pm 0.5) \%$ | |
| Γ_{124} | $\phi\rho^+ \times B(\phi \rightarrow K^+K^-)$ | | $< 7 \times 10^{-3}$ | CL=90% |
| Γ_{125} | $K^+K^-\pi^+\pi^0$ non- ϕ | | $(1.5 \pm_{-0.6}^{+0.7}) \%$ | |
| Γ_{126} | $K^+\bar{K}^0\pi^+\pi^-$ | | $< 2 \%$ | CL=90% |
| Γ_{127} | $K^0K^-\pi^+\pi^+$ | | $(1.0 \pm 0.6) \%$ | |
| Γ_{128} | $K^*(892)^+\bar{K}^*(892)^0$ $\times B^2(K^{*+} \rightarrow K^0\pi^+)$ | | $(1.2 \pm 0.5) \%$ | |
| Γ_{129} | $K^0K^-\pi^+\pi^+$ non- $K^{*+}\bar{K}^{*0}$ | | $< 7.9 \times 10^{-3}$ | CL=90% |
| Γ_{130} | $K^+K^-\pi^+\pi^+\pi^-$ | | — | |
| Γ_{131} | $\phi\pi^+\pi^+\pi^-$ $\times B(\phi \rightarrow K^+K^-)$ | | $< 1 \times 10^{-3}$ | CL=90% |
| Γ_{132} | $K^+K^-\pi^+\pi^+\pi^-$ nonresonant | | $< 3 \%$ | CL=90% |

Fractions of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

| | | | | |
|----------------|------------------------------|--|--------------------------------|--------|
| Γ_{133} | $\phi\pi^+$ | | $(6.1 \pm 0.6) \times 10^{-3}$ | |
| Γ_{134} | $\phi\pi^+\pi^0$ | | $(2.3 \pm 1.0) \%$ | |
| Γ_{135} | $\phi\rho^+$ | | $< 1.4 \%$ | CL=90% |
| Γ_{136} | $\phi\pi^+\pi^+\pi^-$ | | $< 2 \times 10^{-3}$ | CL=90% |
| Γ_{137} | $K^+\bar{K}^*(892)^0$ | | $(4.2 \pm 0.5) \times 10^{-3}$ | |
| Γ_{138} | $K^*(892)^+\bar{K}^0$ | | $(3.2 \pm 1.5) \%$ | |
| Γ_{139} | $K^*(892)^+\bar{K}^*(892)^0$ | | $(2.6 \pm 1.1) \%$ | |

Doubly Cabibbo suppressed (DC) modes, $\Delta C = 1$ weak neutral current (C1) modes, or Lepton Family number (LF) or Lepton number (L) violating modes

| | | | | |
|----------------|-----------------------------|----|--------------------------------|--------|
| Γ_{140} | $K^+\pi^+\pi^-$ | DC | $(6.8 \pm 1.5) \times 10^{-4}$ | |
| Γ_{141} | $K^+\rho^0$ | DC | $(2.5 \pm 1.2) \times 10^{-4}$ | |
| Γ_{142} | $K^*(892)^0\pi^+$ | DC | $(3.6 \pm 1.6) \times 10^{-4}$ | |
| Γ_{143} | $K^+\pi^+\pi^-$ nonresonant | DC | $(2.4 \pm 1.2) \times 10^{-4}$ | |
| Γ_{144} | $K^+K^+K^-$ | DC | $< 1.4 \times 10^{-4}$ | CL=90% |
| Γ_{145} | ϕK^+ | DC | $< 1.3 \times 10^{-4}$ | CL=90% |
| Γ_{146} | $\pi^+e^+e^-$ | C1 | $< 5.2 \times 10^{-5}$ | CL=90% |

| | | | | | |
|----------------|--------------------------|-----------|----------------------|------------------|--------|
| Γ_{147} | $\pi^+ \mu^+ \mu^-$ | <i>CI</i> | < 1.5 | $\times 10^{-5}$ | CL=90% |
| Γ_{148} | $\rho^+ \mu^+ \mu^-$ | <i>CI</i> | < 5.6 | $\times 10^{-4}$ | CL=90% |
| Γ_{149} | $K^+ e^+ e^-$ | | [<i>f</i>] < 2.0 | $\times 10^{-4}$ | CL=90% |
| Γ_{150} | $K^+ \mu^+ \mu^-$ | | [<i>f</i>] < 4.4 | $\times 10^{-5}$ | CL=90% |
| Γ_{151} | $\pi^+ e^\pm \mu^\mp$ | <i>LF</i> | [<i>g</i>] < 3.4 | $\times 10^{-5}$ | CL=90% |
| Γ_{152} | $\pi^+ e^+ \mu^-$ | | | | |
| Γ_{153} | $\pi^+ e^- \mu^+$ | | | | |
| Γ_{154} | $K^+ e^\pm \mu^\mp$ | <i>LF</i> | [<i>g</i>] < 6.8 | $\times 10^{-5}$ | CL=90% |
| Γ_{155} | $K^+ e^+ \mu^-$ | | | | |
| Γ_{156} | $K^+ e^- \mu^+$ | | | | |
| Γ_{157} | $\pi^- e^+ e^+$ | <i>L</i> | < 9.6 | $\times 10^{-5}$ | CL=90% |
| Γ_{158} | $\pi^- \mu^+ \mu^+$ | <i>L</i> | < 1.7 | $\times 10^{-5}$ | CL=90% |
| Γ_{159} | $\pi^- e^+ \mu^+$ | <i>L</i> | < 5.0 | $\times 10^{-5}$ | CL=90% |
| Γ_{160} | $\rho^- \mu^+ \mu^+$ | <i>L</i> | < 5.6 | $\times 10^{-4}$ | CL=90% |
| Γ_{161} | $K^- e^+ e^+$ | <i>L</i> | < 1.2 | $\times 10^{-4}$ | CL=90% |
| Γ_{162} | $K^- \mu^+ \mu^+$ | <i>L</i> | < 1.2 | $\times 10^{-4}$ | CL=90% |
| Γ_{163} | $K^- e^+ \mu^+$ | <i>L</i> | < 1.3 | $\times 10^{-4}$ | CL=90% |
| Γ_{164} | $K^*(892)^- \mu^+ \mu^+$ | <i>L</i> | < 8.5 | $\times 10^{-4}$ | CL=90% |

Γ_{165} A dummy mode used by the fit. (33 ± 5) %

[a] This is a weighted average of D^\pm (44%) and D^0 (56%) branching fractions. See " D^+ and $D^0 \rightarrow (\eta \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$ " under " D^+ Branching Ratios" in these Particle Listings.

[b] This value averages the e^+ and μ^+ branching fractions, after making a small phase-space adjustment to the μ^+ fraction to be able to use it as an e^+ fraction; hence our ℓ^+ here is really an e^+ .

[c] An ℓ indicates an e or a μ mode, not a sum over these modes.

[d] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.

[e] The two experiments measuring this fraction are in serious disagreement. See the Particle Listings.

[f] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

[g] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 32 branching ratios uses 54 measurements and one constraint to determine 20 parameters. The overall fit has a $\chi^2 = 20.8$ for 35 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | | | | | | | | |
|-----------|----------|----------|----------|----------|----------|----------|-----------|-----------|-----------|----------|--|
| x_{11} | 5 | | | | | | | | | | |
| x_{16} | 4 | 2 | | | | | | | | | |
| x_{25} | 18 | 29 | 8 | | | | | | | | |
| x_{26} | 14 | 7 | 31 | 25 | | | | | | | |
| x_{36} | 38 | 9 | 8 | 31 | 25 | | | | | | |
| x_{37} | 32 | 16 | 14 | 56 | 45 | 55 | | | | | |
| x_{42} | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | | | |
| x_{46} | 7 | 4 | 3 | 13 | 10 | 12 | 23 | 0 | | | |
| x_{55} | 9 | 5 | 4 | 17 | 14 | 16 | 30 | 0 | 18 | | |
| x_{62} | 15 | 8 | 7 | 28 | 22 | 27 | 49 | 0 | 11 | 15 | |
| x_{76} | 21 | 11 | 9 | 37 | 29 | 36 | 65 | 0 | 15 | 20 | |
| x_{83} | 5 | 3 | 2 | 9 | 7 | 8 | 16 | 0 | 31 | 37 | |
| x_{90} | 3 | 1 | 1 | 5 | 4 | 5 | 9 | 0 | 29 | 13 | |
| x_{96} | 5 | 2 | 2 | 9 | 7 | 8 | 15 | 0 | 3 | 5 | |
| x_{97} | 3 | 2 | 1 | 6 | 5 | 6 | 11 | 0 | 2 | 3 | |
| x_{101} | 19 | 10 | 9 | 35 | 28 | 33 | 61 | 0 | 14 | 18 | |
| x_{103} | 11 | 5 | 5 | 19 | 15 | 18 | 34 | 0 | 8 | 10 | |
| x_{115} | 22 | 7 | 6 | 23 | 18 | 53 | 41 | 0 | 9 | 12 | |
| x_{165} | -35 | -26 | -12 | -41 | -34 | -38 | -55 | -58 | -46 | -45 | |
| | x_9 | x_{11} | x_{16} | x_{25} | x_{26} | x_{36} | x_{37} | x_{42} | x_{46} | x_{55} | |
| x_{76} | 32 | | | | | | | | | | |
| x_{83} | 8 | 10 | | | | | | | | | |
| x_{90} | 4 | 6 | 12 | | | | | | | | |
| x_{96} | 29 | 10 | 2 | 1 | | | | | | | |
| x_{97} | 8 | 7 | 2 | 1 | 15 | | | | | | |
| x_{101} | 30 | 40 | 10 | 5 | 9 | 7 | | | | | |
| x_{103} | 16 | 22 | 5 | 3 | 5 | 4 | 43 | | | | |
| x_{115} | 20 | 26 | 6 | 4 | 6 | 4 | 25 | 14 | | | |
| x_{165} | -30 | -38 | -46 | -32 | -16 | -10 | -35 | -19 | -27 | | |
| | x_{62} | x_{76} | x_{83} | x_{90} | x_{96} | x_{97} | x_{101} | x_{103} | x_{115} | | |

D^+ BRANCHING RATIOS

See the "Note on D Mesons" above. Some now-obsolete measurements have been omitted from these Listings.

c-quark decays **$\Gamma(c \rightarrow e^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$**

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-----------------------|-------------|----------------------------|
| $0.103 \pm 0.009^{+0.009}_{-0.008}$ | 378 | ⁴ ABBIENDI | 99K OPAL | $Z^0 \rightarrow c\bar{c}$ |

⁴ ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed $D^*(2020)^+ \rightarrow D^0\pi^+$ decays in $Z^0 \rightarrow c\bar{c}$.

 $\Gamma(c \rightarrow \mu^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$

This is the average branching ratio for charm $\rightarrow \mu^+ X$. The mixture of charmed particles is unknown. We don't put this result in the Summary Table.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--|-------------------|----------------|
| 0.085 ± 0.007 OUR NEW AVERAGE | | [0.081 ^{+0.010} _{-0.009} | OUR 1998 AVERAGE] | |

| | | | | |
|-------------------------------------|-----|-----------------------|----------|----------------------------|
| $0.090 \pm 0.007^{+0.007}_{-0.006}$ | 476 | ⁵ ABBIENDI | 99K OPAL | $Z^0 \rightarrow c\bar{c}$ |
| $0.086 \pm 0.017^{+0.008}_{-0.007}$ | 69 | ⁶ ALBRECHT | 92F ARG | $e^+e^- \approx 10$ GeV |
| $0.078 \pm 0.009 \pm 0.012$ | | ONG | 88 MRK2 | $e^+e^- 29$ GeV |
| $0.078 \pm 0.015 \pm 0.02$ | | BARTEL | 87 JADE | $e^+e^- 34.6$ GeV |
| $0.082 \pm 0.012^{+0.02}_{-0.01}$ | | ALTHOFF | 84G TASS | $e^+e^- 34.5$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|--|--------|----------|---------------|
| $0.089 \pm 0.018 \pm 0.025$ | | BARTEL | 85J JADE | See BARTEL 87 |
|-----------------------------|--|--------|----------|---------------|

⁵ ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed $D^*(2020)^+ \rightarrow D^0\pi^+$ decays in $Z^0 \rightarrow c\bar{c}$.

⁶ ALBRECHT 92F uses the excess of right-sign over wrong-sign leptons in a sample of events tagged by fully reconstructed $D^*(2010)^+ \rightarrow D^0\pi^+$ decays.

 $\Gamma(c \rightarrow \ell^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$

This is an average (not a sum) of e^+ and μ^+ measurements.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-----------------------|-------------|----------------------------|
| $0.095 \pm 0.006^{+0.007}_{-0.006}$ | 854 | ⁷ ABBIENDI | 99K OPAL | $Z^0 \rightarrow c\bar{c}$ |

⁷ ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed $D^*(2020)^+ \rightarrow D^0\pi^+$ decays in $Z^0 \rightarrow c\bar{c}$.

Inclusive modes **$\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$** **$\Gamma_1/\Gamma$**

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------|
| 0.172 ± 0.019 OUR AVERAGE | | | | |
| $0.20^{+0.09}_{-0.07}$ | | AGUILAR-... | 87E HYBR | $\pi p, pp 360, 400$ GeV |
| $0.170 \pm 0.019 \pm 0.007$ | 158 | BALTRUSAIT.. | 85B MRK3 | $e^+e^- 3.77$ GeV |
| 0.168 ± 0.064 | 23 | SCHINDLER | 81 MRK2 | $e^+e^- 3.771$ GeV |
| $0.220^{+0.044}_{-0.022}$ | | BACINO | 80 DLCO | $e^+e^- 3.77$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

D^+ and $D^0 \rightarrow (e^+ \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$

If measured at the $\psi(3770)$, this quantity is a weighted average of D^+ (44%) and D^0 (56%) branching fractions. Only experiments at $E_{\text{cm}} = 3.77$ GeV are included in the average here. We don't put this result in the Meson Summary Table.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-------------------------------------|-------------|-------------------------|
| 0.110±0.011 OUR AVERAGE | | Error includes scale factor of 1.1. | | |
| 0.117±0.011 | 295 | BALTRUSAIT..85B | MRK3 | e^+e^- 3.77 GeV |
| 0.10 ±0.032 | | ⁸ SCHINDLER | 81 MRK2 | e^+e^- 3.771 GeV |
| 0.072±0.028 | | FELLER | 78 MRK1 | e^+e^- 3.772 GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.096±0.004±0.011 | 2207 | ⁹ ALBRECHT | 96C ARG | $e^+e^- \approx$ 10 GeV |
| 0.134±0.015±0.010 | | ¹⁰ ABE | 93E VNS | e^+e^- 58 GeV |
| 0.098±0.009 ^{+0.006} _{-0.005} | 240 | ¹¹ ALBRECHT | 92F ARG | $e^+e^- \approx$ 10 GeV |
| 0.096±0.007±0.015 | | ¹² ONG | 88 MRK2 | e^+e^- 29 GeV |
| 0.116 ^{+0.011} _{-0.009} | | ¹² PAL | 86 DLCO | e^+e^- 29 GeV |
| 0.091±0.009±0.013 | | ¹² AIHARA | 85 TPC | e^+e^- 29 GeV |
| 0.092±0.022±0.040 | | ¹² ALTHOFF | 84J TASS | e^+e^- 34.6 GeV |
| 0.091±0.013 | | ¹² KOOP | 84 DLCO | See PAL 86 |
| 0.08 ±0.015 | | ¹³ BACINO | 79 DLCO | e^+e^- 3.772 GeV |

⁸ Isolates D^+ and $D^0 \rightarrow e^+X$ and weights for relative production (44%–56%).

⁹ ALBRECHT 96C uses e^- in the hemisphere opposite to $D^{*+} \rightarrow D^0\pi^+$ events.

¹⁰ ABE 93E also measures forward-backward asymmetries and fragmentation functions for c and b quarks.

¹¹ ALBRECHT 92F uses the excess of right-sign over wrong-sign leptons in a sample of events tagged by fully reconstructed $D^*(2010)^+ \rightarrow D^0\pi^+$ decays.

¹² Average BR for charm $\rightarrow e^+X$. Unlike at $E_{\text{cm}} = 3.77$ GeV, the admixture of charmed mesons is unknown.

¹³ Not independent of BACINO 80 measurements of $\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$ for the D^+ and D^0 separately.

$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$


Γ_2/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---|-------------|--------------------------|
| 0.242±0.028 OUR AVERAGE | | Error includes scale factor of 1.4. See the ideogram below. | | |
| 0.278 ^{+0.036} _{-0.031} | | ¹⁴ BARLAG | 92C ACCM | π^- Cu 230 GeV |
| 0.271±0.023±0.024 | | COFFMAN | 91 MRK3 | e^+e^- 3.77 GeV |
| 0.17 ±0.07 | | AGUILAR-... | 87E HYBR | $\pi p, pp$ 360, 400 GeV |
| 0.19 ±0.05 | 26 | SCHINDLER | 81 MRK2 | e^+e^- 3.771 GeV |
| 0.10 ±0.07 | 3 | VUILLEMIN | 78 MRK1 | e^+e^- 3.772 GeV |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

| | | | | |
|--|--|-------------|----------|-------------------------|
| 0.16 ^{+0.08} _{-0.07} | | AGUILAR-... | 86B HYBR | See AGUILAR-BENITEZ 87E |
|--|--|-------------|----------|-------------------------|

¹⁴ BARLAG 92C computes the branching fraction using topological normalization.



$$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$$

$$\frac{[\Gamma(K^0 \text{ anything}) + \Gamma(K^+ \text{ anything})]}{\Gamma_{\text{total}}} \quad \Gamma_3/\Gamma$$

| VALUE | EVTs | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|------|-------------------------|
| 0.59 ± 0.07 | | | | OUR AVERAGE |
| 0.612 ± 0.065 ± 0.043 | | COFFMAN | 91 | MRK3 e^+e^- 3.77 GeV |
| 0.52 ± 0.18 | 15 | SCHINDLER | 81 | MRK2 e^+e^- 3.771 GeV |
| 0.39 ± 0.29 | 3 | VUILLEMIN | 78 | MRK1 e^+e^- 3.772 GeV |

$$\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

| VALUE | EVTs | DOCUMENT ID | TECN | COMMENT |
|-------------------------|------|-------------|------|-------------------------------|
| 0.058 ± 0.014 | | | | OUR AVERAGE |
| 0.055 ± 0.013 ± 0.009 | | COFFMAN | 91 | MRK3 e^+e^- 3.77 GeV |
| 0.08 $^{+0.06}_{-0.05}$ | | AGUILAR-... | 87E | HYBR $\pi p, pp$ 360, 400 GeV |
| 0.06 ± 0.04 | 12 | SCHINDLER | 81 | MRK2 e^+e^- 3.771 GeV |
| 0.06 ± 0.06 | 2 | VUILLEMIN | 78 | MRK1 e^+e^- 3.772 GeV |

D^+ and $D^0 \rightarrow (\eta \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$

If measured at the $\psi(3770)$, this quantity is a weighted average of D^+ (44%) and D^0 (56%) branching fractions. Only the experiment at $E_{\text{cm}} = 3.77$ GeV is used.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------|-------------|------|------------------------|
| <0.13 | PARTRIDGE | 81 | CBAL e^+e^- 3.77 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.02 ¹⁵BRANDELIK 79 DASP e^+e^- 4.03 GeV

¹⁵The BRANDELIK 79 result is based on the absence of an η signal at $E_{\text{cm}} = 4.03$ GeV. PARTRIDGE 81 observes a substantially higher η cross section at 4.03 GeV.

————— Leptonic and semileptonic modes —————

$\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_7/Γ

See the "Note on Pseudoscalar-Meson Decay Constants" in the Listings for the π^\pm .

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------|----------------------------------|----------------------------------|-------------|-------------------|--|
| 0.0008 | +0.0016 -0.0005 | +0.0005 -0.0002 | 1 | ¹⁶ BAI | 98B BES $e^+ e^- \rightarrow D^{*+} D^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------|----|---|----------------------|----------|---------------------|
| < 0.00072 | 90 | | ADLER | 88B MRK3 | $e^+ e^-$ 3.77 GeV |
| < 0.02 | 90 | 0 | ¹⁷ AUBERT | 83 SPEC | μ^+ Fe, 250 GeV |

¹⁶ BAI 98B obtains $f_D = (300^{+180+80}_{-150-40})$ MeV from this measurement.

¹⁷ AUBERT 83 obtains an upper limit 0.014 assuming the final state contains equal amounts of (D^+, D^-) , (D^+, \bar{D}^0) , (D^-, D^0) , and (D^0, \bar{D}^0) . We quote the limit they get under more general assumptions.

$\Gamma(\bar{K}^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_8/Γ

We average our $\bar{K}^0 e^+ \nu_e$ and $\bar{K}^0 \mu^+ \nu_\mu$ branching fractions, after multiplying the latter by a phase-space factor of 1.03 to be able to use it with the $\bar{K}^0 e^+ \nu_e$ fraction.

Hence our ℓ^+ here is really an e^+ .

| VALUE | DOCUMENT ID | COMMENT |
|---|-------------|--|
| 0.068 ± 0.008 OUR AVERAGE | | |
| 0.067 ± 0.009 | PDG 00 | Our $\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma_{\text{total}}$ |
| 0.072 ^{+0.031} _{-0.020} | PDG 00 | 1.03 × our $\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ |

$\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_9/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|--|-------------|--------|---------------------------------|
| 0.067 ± 0.009 OUR FIT | | | | |
| 0.06 | +0.022 -0.013 ± 0.007 | 13 | BAI 91 | MRK3 $e^+ e^- \approx 3.77$ GeV |

$\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma(\bar{K}^0 \pi^+)$ Γ_9/Γ_{36}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|--------------------|----------|--------------------------------|
| 2.32 ± 0.31 OUR FIT | | | | |
| 2.60 ± 0.35 ± 0.26 | 186 | ¹⁸ BEAN | 93C CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

¹⁸ BEAN 93C uses $\bar{K}^0 \mu^+ \nu_\mu$ as well as $\bar{K}^0 e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

$\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma(K^- \pi^+ \pi^+)$ Γ_9/Γ_{37}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------|----------|------------------------|
| 0.74 ± 0.10 OUR FIT | | | |
| 0.66 ± 0.09 ± 0.14 | ANJOS | 91C E691 | γ Be 80–240 GeV |

$\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{10}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|--|-------------|--------|---------------------------------|
| 0.07 | +0.028 -0.016 ± 0.012 | 14 | BAI 91 | MRK3 $e^+ e^- \approx 3.77$ GeV |

$\Gamma(\bar{K}^0 \mu^+ \nu_\mu) / \Gamma(\mu^+ \text{ anything})$

Γ_{10} / Γ_6

| VALUE | EVTS | DOCUMENT ID | COMMENT |
|-------|------|-------------|---------|
|-------|------|-------------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------|----|--------------------|---------------------|
| 0.76 ± 0.06 | 84 | ¹⁹ AOKI | 88 π^- emulsion |
|-------------|----|--------------------|---------------------|

¹⁹ From topological branching ratios in emulsion with an identified muon.

$\Gamma(K^- \pi^+ e^+ \nu_e) / \Gamma_{\text{total}}$

Γ_{11} / Γ

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|-----|------|-------------|------|---------|
|-------|-----|------|-------------|------|---------|

0.041^{+0.009}_{-0.007} OUR FIT

| | | | | | |
|--|----|-------------------|---------|------------------------|-----|
| 0.035^{+0.012}_{-0.007} ± 0.004 | 14 | ²⁰ BAI | 91 MRK3 | $e^+ e^- \approx 3.77$ | GeV |
|--|----|-------------------|---------|------------------------|-----|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------|----|---------------------------|----------|-------------|--------------|
| < 0.057 | 90 | ²¹ AGUILAR-... | 87F HYBR | $\pi p, pp$ | 360, 400 GeV |
|---------|----|---------------------------|----------|-------------|--------------|

²⁰ BAI 91 finds that a fraction $0.79^{+0.15+0.09}_{-0.17-0.03}$ of combined D^+ and D^0 decays to $\bar{K} \pi e^+ \nu_e$ (24 events) are $\bar{K}^*(892) e^+ \nu_e$.

²¹ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(\bar{K}^*(892)^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$

Γ_{24} / Γ

We average our $\bar{K}^{*0} e^+ \nu_e$ and $\bar{K}^{*0} \mu^+ \nu_\mu$ branching fractions, after multiplying the latter by a phase-space factor of 1.05 to be able to use it with the $\bar{K}^{*0} e^+ \nu_e$ fraction. Hence our ℓ^+ here is really an e^+ .

| VALUE | DOCUMENT ID | COMMENT |
|-------|-------------|---------|
|-------|-------------|---------|

0.047 ± 0.004 OUR AVERAGE

| | | |
|---------------|-----|---|
| 0.048 ± 0.005 | PDG | 00 Our $\Gamma(\bar{K}^{*0} e^+ \nu_e) / \Gamma_{\text{total}}$ |
| 0.046 ± 0.006 | PDG | 00 $1.05 \times$ our $\Gamma(\bar{K}^{*0} \mu^+ \nu_\mu) / \Gamma_{\text{total}}$ |

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e) / \Gamma(K^- \pi^+ e^+ \nu_e)$

$\Gamma_{25} / \Gamma_{11}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

1.16^{+0.21}_{-0.24} OUR FIT

| | | | | |
|------------------|----|-----------|---------|-----------------|
| 1.0 ± 0.3 | 35 | ADAMOVICH | 91 OMEG | π^- 340 GeV |
|------------------|----|-----------|---------|-----------------|

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e) / \Gamma(K^- \pi^+ \pi^+)$

$\Gamma_{25} / \Gamma_{37}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.53 ± 0.05 OUR FIT

0.54 ± 0.05 OUR AVERAGE

| | | | | |
|--------------------|-----|--------------------|----------|--------------------------------|
| 0.67 ± 0.09 ± 0.07 | 710 | ²² BEAN | 93C CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.62 ± 0.15 ± 0.09 | 35 | ADAMOVICH | 91 OMEG | π^- 340 GeV |
| 0.55 ± 0.08 ± 0.10 | 880 | ALBRECHT | 91 ARG | $e^+ e^- \approx 10.4$ GeV |
| 0.49 ± 0.04 ± 0.05 | | ANJOS | 89B E691 | Photoproduction |

²² BEAN 93C uses $\bar{K}^{*0} \mu^+ \nu_\mu$ as well as $\bar{K}^{*0} e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

$\Gamma(K^- \pi^+ e^+ \nu_e \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{13}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--------|-----|-------------|----------|-----------------|
| <0.007 | 90 | 23 ANJOS | 89B E691 | Photoproduction |

²³ ANJOS 89B assumes a $\Gamma(D^+ \rightarrow K^- \pi^+ \pi^+)/\Gamma_{\text{total}} = 9.1 \pm 1.3 \pm 0.4\%$.

$\Gamma(K^- \pi^+ \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ $\Gamma_{14}/\Gamma = (\Gamma_{16} + \frac{2}{3}\Gamma_{26})/\Gamma$

| VALUE | DOCUMENT ID |
|------------------------------|-------------------------------------|
| 0.032 ± 0.004 OUR FIT | Error includes scale factor of 1.1. |

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{26}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|-------------------------------------|
| 0.044 ± 0.006 OUR FIT | | | | Error includes scale factor of 1.1. |

0.0325 ± 0.0071 ± 0.0075 224 ²⁴ KODAMA 92C E653 π^- emulsion 600 GeV

²⁴ KODAMA 92C measures $\Gamma(D^+ \rightarrow \bar{K}^{*0} \mu^+ \nu_\mu)/\Gamma(D^0 \rightarrow K^- \mu^+ \nu_\mu) = 0.43 \pm 0.09 \pm 0.09$ and then uses $\Gamma(D^0 \rightarrow K^- \mu^+ \nu_\mu) = (7.0 \pm 0.7) \times 10^{10} \text{ s}^{-1}$ to get the quoted branching fraction. See also the footnote to KODAMA 92C in the next data block.

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{26}/Γ_{37}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|------|---------|
| 0.49 ± 0.06 OUR FIT | | | | |
| 0.53 ± 0.06 OUR AVERAGE | | | | |

0.56 ± 0.04 ± 0.06 875 FRABETTI 93E E687 γ Be $\bar{E}_\gamma \approx 200$ GeV

0.46 ± 0.07 ± 0.08 224 ²⁵ KODAMA 92C E653 π^- emulsion 600 GeV

²⁵ KODAMA 92C uses the same $\bar{K}^{*0} \mu^+ \nu_\mu$ events normalizing instead with $D^0 \rightarrow K^- \mu^+ \nu_\mu$ events, as reported in the preceding data block.

$\Gamma(K^- \pi^+ \mu^+ \nu_\mu \text{ nonresonant})/\Gamma(K^- \pi^+ \mu^+ \nu_\mu)$ $\Gamma_{16}/\Gamma_{14} = \Gamma_{16}/(\Gamma_{16} + \frac{2}{3}\Gamma_{26})$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|------|---------|
| 0.083 ± 0.029 OUR FIT | | | |

0.083 ± 0.029 FRABETTI 93E E687 < 0.12 (90% CL)

$\Gamma(\bar{K}^0 \pi^+ \pi^- e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{17}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---|
| • • • | | | | We do not use the following data for averages, fits, limits, etc. • • • |

0.022 $^{+0.047}_{-0.006}$ ± 0.004 1 ²⁶ AGUILAR-... 87F HYBR $\pi p, pp$ 360, 400 GeV

²⁶ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(K^- \pi^+ \pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{18}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---|
| • • • | | | | We do not use the following data for averages, fits, limits, etc. • • • |

0.044 $^{+0.052}_{-0.013}$ ± 0.007 2 ²⁷ AGUILAR-... 87F HYBR $\pi p, pp$ 360, 400 GeV

²⁷ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma((\bar{K}^*(892)\pi)^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{19}/Γ

Unseen decay modes of the $\bar{K}^*(892)$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--------|-----|-------------|---------|-----------------|
| <0.012 | 90 | ANJOS | 92 E691 | Photoproduction |

$\Gamma((\bar{K}\pi\pi)^0 e^+ \nu_e \text{ non-}\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{20}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--------|-----|-------------|---------|-----------------|
| <0.009 | 90 | ANJOS | 92 E691 | Photoproduction |

$\Gamma(K^- \pi^+ \pi^0 \mu^+ \nu_\mu)/\Gamma(K^- \pi^+ \mu^+ \nu_\mu)$ $\Gamma_{21}/\Gamma_{14} = \Gamma_{21}/(\Gamma_{16} + \frac{2}{3}\Gamma_{26})$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--------|-----|-------------|----------|---|
| <0.042 | 90 | FRABETTI | 93E E687 | $\gamma \text{ Be } \bar{E}_\gamma \approx 200 \text{ GeV}$ |

$\Gamma(\bar{K}_1(1270)^0 \mu^+ \nu_\mu)/\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$ Γ_{27}/Γ_{26}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|---------|--------------------|
| <0.78 | 95 | ABE | 99P CDF | $\bar{p}p$ 1.8 TeV |

$\Gamma(\bar{K}^*(1410)^0 \mu^+ \nu_\mu)/\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$ Γ_{28}/Γ_{26}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|---------|--------------------|
| <0.60 | 95 | ABE | 99P CDF | $\bar{p}p$ 1.8 TeV |

$\Gamma(\bar{K}_2^*(1430)^0 \mu^+ \nu_\mu)/\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$ Γ_{29}/Γ_{26}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|---------|--------------------|
| <0.19 | 95 | ABE | 99P CDF | $\bar{p}p$ 1.8 TeV |

$\Gamma(\pi^0 \ell^+ \nu_\ell)/\Gamma(\bar{K}^0 \ell^+ \nu_\ell)$ Γ_{22}/Γ_8

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------|-----------------------|---------|------------------------------|
| $0.046 \pm 0.014 \pm 0.017$ | 100 | ²⁸ BARTELT | 97 CLE2 | $e^+ e^- \approx \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|----|--------------------|---------|----------------|
| $0.085 \pm 0.027 \pm 0.014$ | 53 | ²⁹ ALAM | 93 CLE2 | See BARTELT 97 |
|-----------------------------|----|--------------------|---------|----------------|

²⁸ BARTELT 97 thus directly measures the product of ratios squared of CKM matrix elements and form factors at $q^2=0$: $|V_{cd}/V_{cs}|^2 \cdot |f_+^\pi(0)/f_+^K(0)|^2 = 0.046 \pm 0.014 \pm 0.017$.

²⁹ ALAM 93 thus directly measures the product of ratios squared of CKM matrix elements and form factors at $q^2=0$: $|V_{cd}/V_{cs}|^2 \cdot |f_+^\pi(0)/f_+^K(0)|^2 = 0.085 \pm 0.027 \pm 0.014$.

$\Gamma(\pi^+ \pi^- e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{23}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--------|-----|---------------------------|----------|---------------------------|
| <0.057 | 90 | ³⁰ AGUILAR-... | 87F HYBR | $\pi p, p p$ 360, 400 GeV |

³⁰ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(\rho^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{30}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---------|-----|-------------|---------|------------------------------------|
| <0.0037 | 90 | BAI | 91 MRK3 | $e^+ e^- \approx 3.77 \text{ GeV}$ |

$\Gamma(\rho^0 e^+ \nu_e) / \Gamma(\bar{K}^*(892)^0 e^+ \nu_e)$ $\Gamma_{30} / \Gamma_{25}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|----------------------|---------|--------------------------|
| 0.045 ± 0.014 ± 0.009 | 49 | ³¹ AITALA | 97 E791 | π^- nucleus, 500 GeV |

³¹ AITALA 97 explicitly subtracts $D^+ \rightarrow \eta' e^+ \nu_e$ and other backgrounds to get this result.

$\Gamma(\rho^0 \mu^+ \nu_\mu) / \Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$ $\Gamma_{31} / \Gamma_{26}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|------------------------|---------|---|
| 0.061 ± 0.014 OUR AVERAGE | | | | |
| 0.051 ± 0.015 ± 0.009 | 54 | ³² AITALA | 97 E791 | π^- nucleus, 500 GeV |
| 0.079 ± 0.019 ± 0.013 | 39 | ³³ FRABETTI | 97 E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---|---|----------------------|----------|--------------------------|
| 0.044 ^{+0.031} _{-0.025} ± 0.014 | 4 | ³⁴ KODAMA | 93C E653 | π^- emulsion 600 GeV |
|---|---|----------------------|----------|--------------------------|

³² AITALA 97 explicitly subtracts $D^+ \rightarrow \eta' \mu^+ \nu_\mu$ and other backgrounds to get this result.

³³ Because the reconstruction efficiency for photons is low, this FRABETTI 97 result also includes any $D^+ \rightarrow \eta' \mu^+ \nu_\mu \rightarrow \gamma \rho^0 \mu^+ \nu_\mu$ events in the numerator.

³⁴ This KODAMA 93C result is based on a final signal of $4.0^{+2.8}_{-2.3} \pm 1.3$ events; the estimates of backgrounds that affect this number are somewhat model dependent.

$\Gamma(\phi e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{32} / Γ

Decay modes of the ϕ not included in the search are corrected for.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------|-----|-------------|---------|----------------------------|
| <0.0209 | 90 | BAI | 91 MRK3 | $e^+ e^- \approx 3.77$ GeV |

$\Gamma(\phi \mu^+ \nu_\mu) / \Gamma_{\text{total}}$ Γ_{33} / Γ

Decay modes of the ϕ not included in the search are corrected for.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------|-----|-------------|---------|----------------------------|
| <0.0372 | 90 | BAI | 91 MRK3 | $e^+ e^- \approx 3.77$ GeV |

$\Gamma(\eta \ell^+ \nu_\ell) / \Gamma(\pi^0 \ell^+ \nu_\ell)$ $\Gamma_{34} / \Gamma_{22}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------|-----|-------------|---------|--------------------------------|
| <1.5 | 90 | BARTELT | 97 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\eta'(958) \mu^+ \nu_\mu) / \Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$ $\Gamma_{35} / \Gamma_{26}$

Decay modes of the $\eta'(958)$ not included in the search are corrected for.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|-------------|----------|--------------------------|
| <0.20 | 90 | KODAMA | 93B E653 | π^- emulsion 600 GeV |

————— Hadronic modes with a \bar{K} or $\bar{K}K\bar{K}$ —————

$\Gamma(\bar{K}^0 \pi^+) / \Gamma_{\text{total}}$ Γ_{36} / Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|-------------|------|-------------------------------------|
| 0.0289 ± 0.0026 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.032 ± 0.004 OUR AVERAGE | | | | |

| | | | | |
|-----------------------|-----|-------|----------|--------------------|
| 0.032 ± 0.005 ± 0.002 | 161 | ADLER | 88C MRK3 | $e^+ e^-$ 3.77 GeV |
|-----------------------|-----|-------|----------|--------------------|

| | | | | |
|---------------|----|-------------------------|---------|---------------------|
| 0.033 ± 0.009 | 36 | ³⁵ SCHINDLER | 81 MRK2 | $e^+ e^-$ 3.771 GeV |
|---------------|----|-------------------------|---------|---------------------|

| | | | | |
|---------------|----|-----------------------|---------|--------------------|
| 0.033 ± 0.013 | 17 | ³⁶ PERUZZI | 77 MRK1 | $e^+ e^-$ 3.77 GeV |
|---------------|----|-----------------------|---------|--------------------|

³⁵ SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.14 ± 0.03 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

³⁶ PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.14 ± 0.05 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

$\Gamma(\bar{K}^0\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{36}/Γ_{37}

It is generally assumed for modes such as $D^+ \rightarrow \bar{K}^0\pi^+$ that

$$\Gamma(D^+ \rightarrow \bar{K}^0\pi^+) = 2\Gamma(D^+ \rightarrow K_S^0\pi^+);$$

it is the latter Γ that is actually measured. BIGI 95 points out that interference between Cabibbo-allowed and doubly Cabibbo-suppressed amplitudes, where both occur, could invalidate this assumption by a few percent.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|------|----------------------|----------|-------------------------------------|
| 0.321±0.025 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.32 ±0.04 OUR AVERAGE | | | | Error includes scale factor of 1.4. |
| 0.348±0.024±0.022 | 473 | ³⁷ BISHAI | 97 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| 0.274±0.030±0.031 | 264 | ANJOS | 90C E691 | Photoproduction |

³⁷ See BISHAI 97 for an isospin analysis of $D^+ \rightarrow \bar{K}\pi$ amplitudes.

$\Gamma(K^-\pi^+\pi^+)/\Gamma_{\text{total}}$ Γ_{37}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------------------|----------|-------------------------------|
| 0.090±0.006 OUR FIT | | | | |
| 0.091±0.007 OUR AVERAGE | | | | |
| 0.093±0.006±0.008 | 1502 | ³⁸ BALEST | 94 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| 0.091±0.013±0.004 | 1164 | ADLER | 88C MRK3 | e^+e^- 3.77 GeV |
| 0.091±0.019 | 239 | ³⁹ SCHINDLER | 81 MRK2 | e^+e^- 3.771 GeV |
| 0.086±0.020 | 85 | ⁴⁰ PERUZZI | 77 MRK1 | e^+e^- 3.77 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--|---|---------------------------|----------|--------------------------|
| 0.064 ^{+0.015} _{-0.014} | | ⁴¹ BARLAG | 92C ACCM | π^- Cu 230 GeV |
| 0.063 ^{+0.028} _{-0.014} ±0.011 | 8 | ⁴¹ AGUILAR-... | 87F HYBR | $\pi p, pp$ 360, 400 GeV |

³⁸ BALEST 94 measures the ratio of $D^+ \rightarrow K^-\pi^+\pi^+$ and $D^0 \rightarrow K^-\pi^+$ branching fractions to be $2.35 \pm 0.16 \pm 0.16$ and uses their absolute measurement of the $D^0 \rightarrow K^-\pi^+$ fraction (AKERIB 93).

³⁹ SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.38 ± 0.05 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

⁴⁰ PERUZZI 77 (MARK-1) measures $\sigma(e^+e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.36 ± 0.06 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

⁴¹ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(\bar{K}^*(892)^0\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{76}/Γ_{37}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-----|-------------|----------|---|
| 0.212±0.016 OUR FIT | | | | |
| 0.210±0.015 OUR AVERAGE | | | | |
| 0.206±0.009±0.014 | | FRABETTI | 94G E687 | $\gamma\text{Be}, \bar{E}_\gamma \approx 220$ GeV |
| 0.255±0.014±0.050 | | ANJOS | 93 E691 | γBe 90–260 GeV |
| 0.21 ±0.06 ±0.06 | | ALVAREZ | 91B NA14 | Photoproduction |
| 0.20 ±0.02 ±0.11 | | ADLER | 87 MRK3 | e^+e^- 3.77 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|-----------|---------|--------------------|
| <0.053 | 90 | SCHINDLER | 81 MRK2 | e^+e^- 3.771 GeV |
|--------|----|-----------|---------|--------------------|

$\Gamma(\bar{K}_0^*(1430)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{85} / \Gamma_{37}$

Unseen decay modes of the $\bar{K}_0^*(1430)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------|----------|---|
| 0.41 ± 0.04 OUR AVERAGE | | | |
| 0.458 ± 0.035 ± 0.094 | FRABETTI | 94G E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| 0.400 ± 0.031 ± 0.027 | ANJOS | 93 E691 | γ Be 90–260 GeV |

$\Gamma(\bar{K}^*(1680)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{86} / \Gamma_{37}$

Unseen decay modes of the $\bar{K}^*(1680)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-------------|----------|---|
| 0.160 ± 0.032 OUR AVERAGE | | | Error includes scale factor of 1.1. |
| 0.182 ± 0.023 ± 0.028 | FRABETTI | 94G E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| 0.113 ± 0.015 ± 0.050 | ANJOS | 93 E691 | γ Be 90–260 GeV |

$\Gamma(K^- \pi^+ \pi^+ \text{ nonresonant}) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{41} / \Gamma_{37}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------|----------|---|
| 0.95 ± 0.07 OUR AVERAGE | | | |
| 0.998 ± 0.037 ± 0.072 | FRABETTI | 94G E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| 0.838 ± 0.088 ± 0.275 | ANJOS | 93 E691 | γ Be 90–260 GeV |
| 0.79 ± 0.07 ± 0.15 | ADLER | 87 MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}^0 \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{42} / Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|-------------------------|----------|-------------------------------------|
| 0.097 ± 0.030 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.107 ± 0.029 OUR AVERAGE | | | | |
| 0.102 ± 0.025 ± 0.016 | 159 | ADLER | 88C MRK3 | $e^+ e^-$ 3.77 GeV |
| 0.19 ± 0.12 | 10 | ⁴² SCHINDLER | 81 MRK2 | $e^+ e^-$ 3.771 GeV |

⁴²SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.78 ± 0.48 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

$\Gamma(\bar{K}^0 \rho^+) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{43} / \Gamma_{42}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|---------|--------------------|
| 0.68 ± 0.08 ± 0.12 | ADLER | 87 MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}^*(892)^0 \pi^+) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{76} / \Gamma_{42}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------|---------|--------------------|
| 0.20 ± 0.06 OUR FIT | | | |
| 0.57 ± 0.18 ± 0.18 | ADLER | 87 MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}^0 \pi^+ \pi^0 \text{ nonresonant}) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{45} / \Gamma_{42}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|---------|--------------------|
| 0.13 ± 0.07 ± 0.08 | ADLER | 87 MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(K^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{46}/Γ

| <u>VALUE</u> | <u>EVTs</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------|
| 0.064±0.011 OUR FIT | | | | |
| 0.058±0.012±0.012 | 142 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.034 ^{+0.056} _{-0.070} | | 43 BARLAG | 92C ACCM | π^- Cu 230 GeV |
| 0.022 ^{+0.047} _{-0.006} ±0.004 | 1 | 43 AGUILAR-... | 87F HYBR | $\pi p, pp$ 360, 400 GeV |
| 0.063 ^{+0.014} _{-0.013} ±0.012 | 175 | BALTRUSAIT..86E | MRK3 | See COFFMAN 92B |

⁴³ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(K^- \pi^+ \pi^+ \pi^0)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{46}/Γ_{37}

| <u>VALUE</u> | <u>EVTs</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|------------------------|
| 0.71±0.12 OUR FIT | | | | |
| 0.76±0.11±0.12 | 91 | ANJOS | 92C E691 | γ Be 90–260 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.69±0.10±0.16 | | ANJOS | 89E E691 | See ANJOS 92C |
| 0.57 ^{+0.65} _{-0.17} | 1 | AGUILAR-... | 83B HYBR | $\pi^- p$, 360 GeV |

$\Gamma(\bar{K}^*(892)^0 \rho^+ \text{ total})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{77}/Γ_{46}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|--------------------|-------------|------------------------|
| 0.33±0.165±0.12 | 44 ANJOS | 92C E691 | γ Be 90–260 GeV |

⁴⁴ See, however, the next entry, where the two experiments disagree completely.

$\Gamma(\bar{K}^*(892)^0 \rho^+ S\text{-wave})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{78}/Γ_{46}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included. The two experiments here disagree completely.

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------|-------------------------------------|-------------|------------------------|
| 0.26 ±0.25 OUR AVERAGE | Error includes scale factor of 3.1. | | |
| 0.15 ±0.075±0.045 | ANJOS | 92C E691 | γ Be 90–260 GeV |
| 0.833±0.116±0.165 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}^*(892)^0 \rho^+ P\text{-wave})/\Gamma_{\text{total}}$ Γ_{79}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|------------------------|
| <0.001 | 90 | ANJOS | 92C E691 | γ Be 90–260 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.005 | 90 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}^*(892)^0 \rho^+ D\text{-wave})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{80}/Γ_{46}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|--------------------|-------------|------------------------|
| 0.15±0.09±0.045 | ANJOS | 92C E691 | γ Be 90–260 GeV |

$\Gamma(\bar{K}^*(892)^0 \rho^+ D\text{-wave longitudinal})/\Gamma_{\text{total}}$ Γ_{81}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--------|-----|-------------|----------|--------------------|
| <0.007 | 90 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}_1(1400)^0 \pi^+)/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{83}/Γ_{46}

Unseen decay modes of the $\bar{K}_1(1400)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|----------|--------------------|
| 0.77 ± 0.20 OUR FIT | | | |
| 0.907 ± 0.218 ± 0.180 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(K^- \rho^+ \pi^+ \text{total})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{91}/Γ_{46}

This includes $\bar{K}^*(892)^0 \rho^+$, etc. The next entry gives the specifically 3-body fraction.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|----------|------------------------|
| 0.48 ± 0.13 ± 0.09 | ANJOS | 92C E691 | γ Be 90–260 GeV |

$\Gamma(K^- \rho^+ \pi^+ \text{3-body})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{92}/Γ_{46}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------|----------|------------------------|
| 0.17 ± 0.06 OUR AVERAGE | | | |
| 0.18 ± 0.08 ± 0.04 | ANJOS | 92C E691 | γ Be 90–260 GeV |
| 0.159 ± 0.065 ± 0.060 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{total})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{87}/Γ_{46}

This includes $\bar{K}^*(892)^0 \rho^+$, etc. The next two entries give the specifically 3-body fraction. Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|----------|------------------------|
| 1.05 ± 0.11 ± 0.08 | ANJOS | 92C E691 | γ Be 90–260 GeV |

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{3-body})/\Gamma_{\text{total}}$ Γ_{88}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--------|-----|-----------------------|----------|--------------------|
| <0.008 | 90 | ⁴⁵ COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁴⁵See, however, the next entry: ANJOS 92C sees a large signal in this channel.

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{3-body})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{88}/Γ_{46}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|----------|------------------------|
| 0.66 ± 0.09 ± 0.17 | ANJOS | 92C E691 | γ Be 90–260 GeV |

$\Gamma(K^*(892)^- \pi^+ \pi^+ \text{3-body})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{90}/Γ_{46}

Unseen decay modes of the $K^*(892)^-$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------|----------|-------------------------------------|
| 0.32 ± 0.14 OUR FIT | | | Error includes scale factor of 1.1. |
| 0.24 ± 0.12 ± 0.09 | ANJOS | 92C E691 | γ Be 90–260 GeV |

$\Gamma(K^- \pi^+ \pi^+ \pi^0 \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{54}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|---------------------|----------|------------------------|
| <0.002 | 90 | ⁴⁶ ANJOS | 92C E691 | γ Be 90–260 GeV |
|--------|----|---------------------|----------|------------------------|

⁴⁶ Whereas ANJOS 92C finds no signal here, COFFMAN 92B finds a fairly large one; see the next entry.

$\Gamma(K^- \pi^+ \pi^+ \pi^0 \text{ nonresonant})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{54}/Γ_{46}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|------------------------------|---------|----------|--------------------|
| 0.184 ± 0.070 ± 0.050 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |
|------------------------------|---------|----------|--------------------|

$\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{55}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.070 ± 0.009 OUR FIT

0.071 ± 0.016 OUR AVERAGE

| | | | | |
|-----------------------|-----|-------|----------|--------------------|
| 0.066 ± 0.015 ± 0.005 | 168 | ADLER | 88C MRK3 | $e^+ e^-$ 3.77 GeV |
|-----------------------|-----|-------|----------|--------------------|

| | | | | |
|-------------|----|-------------------------|---------|---------------------|
| 0.12 ± 0.05 | 21 | ⁴⁷ SCHINDLER | 81 MRK2 | $e^+ e^-$ 3.771 GeV |
|-------------|----|-------------------------|---------|---------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---|--|----------------------|----------|--------------------|
| 0.042 ^{+0.019} _{-0.017} | | ⁴⁸ BARLAG | 92C ACCM | π^- Cu 230 GeV |
|---|--|----------------------|----------|--------------------|

| | | | | |
|---|----|---------------------------|----------|---------------------------|
| 0.243 ^{+0.064} _{-0.041} ± 0.041 | 11 | ⁴⁸ AGUILAR-... | 87F HYBR | $\pi p, p p$ 360, 400 GeV |
|---|----|---------------------------|----------|---------------------------|

⁴⁷ SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.51 ± 0.08 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

⁴⁸ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{55}/Γ_{37}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.78 ± 0.10 OUR FIT

| | | | | |
|---------------------------|-----|-------|----------|------------------------|
| 0.77 ± 0.07 ± 0.11 | 229 | ANJOS | 92C E691 | γ Be 90–260 GeV |
|---------------------------|-----|-------|----------|------------------------|

$\Gamma(\bar{K}^0 a_1(1260)^+)/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{74}/Γ_{55}

Unseen decay modes of the $a_1(1260)^+$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

1.15 ± 0.19 OUR AVERAGE Error includes scale factor of 1.1.

| | | | |
|--------------------|-------|----------|------------------------|
| 1.66 ± 0.28 ± 0.40 | ANJOS | 92C E691 | γ Be 90–260 GeV |
|--------------------|-------|----------|------------------------|

| | | | |
|-----------------------|---------|----------|--------------------|
| 1.078 ± 0.114 ± 0.140 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |
|-----------------------|---------|----------|--------------------|

$\Gamma(\bar{K}^0 a_2(1320)^+)/\Gamma_{\text{total}}$ Γ_{75}/Γ

Unseen decay modes of the $a_2(1320)^+$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|------------------|----|-------|----------|------------------------|
| <0.003 | 90 | ANJOS | 92C E691 | γ Be 90–260 GeV |
|------------------|----|-------|----------|------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|---------|----------|--------------------|
| <0.008 | 90 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |
|--------|----|---------|----------|--------------------|

$\Gamma(\bar{K}_1(1270)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{82}/Γ

Unseen decay modes of the $\bar{K}_1(1270)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|------------------------|
| <0.007 | 90 | ANJOS | 92C E691 | γ Be 90–260 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.011 | 90 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}_1(1400)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{83}/Γ

Unseen decay modes of the $\bar{K}_1(1400)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|---------------------|----------|------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.009 | 90 | ⁴⁹ ANJOS | 92C E691 | γ Be 90–260 GeV |
| ⁴⁹ ANJOS 92C sees no evidence for $\bar{K}_1(1400)^0 \pi^+$ in either the $\bar{K}^0 \pi^+ \pi^+ \pi^-$ or $K^- \pi^+ \pi^+ \pi^0$ channels, whereas COFFMAN 92B finds the $\bar{K}_1(1400)^0 \pi^+$ branching fraction to be large; see the next entry. | | | | |

$\Gamma(\bar{K}_1(1400)^0 \pi^+)/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{83}/Γ_{55}

Unseen decay modes of the $\bar{K}_1(1400)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|----------|--------------------|
| 0.70 ± 0.17 OUR FIT | | | |
| 0.623 ± 0.106 ± 0.180 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\bar{K}^*(1410)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{84}/Γ

Unseen decay modes of the $\bar{K}^*(1410)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-------------|----------|--------------------|
| <0.007 | 90 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(K^*(892)^- \pi^+ \pi^+ \text{total})/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{89}/Γ_{55}

Unseen decay modes of the $K^*(892)^-$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|----------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.41 ± 0.14 | 14 | ALEEV | 94 BIS2 | nN 20–70 GeV |

$\Gamma(K^*(892)^- \pi^+ \pi^+ \text{3-body})/\Gamma_{\text{total}}$ Γ_{90}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|--------------------|
| 0.020 ± 0.009 OUR FIT | | | | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.013 | 90 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(K^*(892)^- \pi^+ \pi^+ \text{3-body})/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{90}/Γ_{55}

Unseen decay modes of the $K^*(892)^-$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------|----------|-------------------------------------|
| 0.29 ± 0.13 OUR FIT | | | Error includes scale factor of 1.1. |
| 0.50 ± 0.09 ± 0.21 | ANJOS | 92C E691 | γ Be 90–260 GeV |

$\Gamma(\overline{K}^0 \rho^0 \pi^+ \text{total}) / \Gamma(\overline{K}^0 \pi^+ \pi^+ \pi^-)$ $\Gamma_{93} / \Gamma_{55}$

This includes $\overline{K}^0 a_1(1260)^+$. The next two entries give the specifically 3-body reaction.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|-------------|----------|------------------------|
| 0.60 ± 0.10 ± 0.17 | 90 | ANJOS | 92C E691 | γ Be 90–260 GeV |

$\Gamma(\overline{K}^0 \rho^0 \pi^+ \text{3-body}) / \Gamma_{\text{total}}$ Γ_{94} / Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|---------|----------|--------------------|
| <0.004 | 90 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |
|--------|----|---------|----------|--------------------|

$\Gamma(\overline{K}^0 \rho^0 \pi^+ \text{3-body}) / \Gamma(\overline{K}^0 \pi^+ \pi^+ \pi^-)$ $\Gamma_{94} / \Gamma_{55}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|---------------------------|-------|----------|------------------------|
| 0.07 ± 0.04 ± 0.06 | ANJOS | 92C E691 | γ Be 90–260 GeV |
|---------------------------|-------|----------|------------------------|

$\Gamma(\overline{K}^0 f_0(980) \pi^+) / \Gamma_{\text{total}}$ Γ_{95} / Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|------------------|----|-------|----------|------------------------|
| <0.005 | 90 | ANJOS | 92C E691 | γ Be 90–260 GeV |
|------------------|----|-------|----------|------------------------|

$\Gamma(\overline{K}^0 \pi^+ \pi^+ \pi^- \text{nonresonant}) / \Gamma(\overline{K}^0 \pi^+ \pi^+ \pi^-)$ $\Gamma_{61} / \Gamma_{55}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.12 ± 0.06 OUR AVERAGE

| | | | |
|--------------------|-------|----------|------------------------|
| 0.10 ± 0.04 ± 0.06 | ANJOS | 92C E691 | γ Be 90–260 GeV |
|--------------------|-------|----------|------------------------|

| | | | |
|----------------------|---------|----------|--------------------|
| 0.17 ± 0.056 ± 0.100 | COFFMAN | 92B MRK3 | $e^+ e^-$ 3.77 GeV |
|----------------------|---------|----------|--------------------|

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{62} / Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--|----------------------|----------|--------------------|
| 0.0037 ^{+0.0012} _{-0.0010} | ⁵⁰ BARLAG | 92C ACCM | π^- Cu 230 GeV |
|--|----------------------|----------|--------------------|

⁵⁰BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{62} / \Gamma_{37}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.080 ± 0.009 OUR FIT

0.083 ± 0.009 OUR AVERAGE

| | | | | |
|-----------------------|-----|----------|----------|--|
| 0.077 ± 0.008 ± 0.010 | 239 | FRABETTI | 97C E687 | γ Be, $\overline{E}_\gamma \approx 200$ GeV |
|-----------------------|-----|----------|----------|--|

| | | | | |
|--------------------|-----|-------|----------|-----------------|
| 0.09 ± 0.01 ± 0.01 | 113 | ANJOS | 90D E691 | Photoproduction |
|--------------------|-----|-------|----------|-----------------|

$\Gamma(\overline{K}^*(892)^0 \pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{96} / \Gamma_{62}$

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

1.1 ± 0.4 OUR FIT Error includes scale factor of 1.8.

| | | | |
|---------------------------|-------|----------|-----------------|
| 1.25 ± 0.12 ± 0.23 | ANJOS | 90D E691 | Photoproduction |
|---------------------------|-------|----------|-----------------|

$\Gamma(\overline{K}^*(892)^0 \rho^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{97} / \Gamma_{37}$

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.032^{+0.019}_{-0.017} OUR FIT Error includes scale factor of 1.8.

| | | | |
|------------------------------|----------|----------|--|
| 0.023 ± 0.010 ± 0.006 | FRABETTI | 97C E687 | γ Be, $\overline{E}_\gamma \approx 200$ GeV |
|------------------------------|----------|----------|--|

$\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+) / \Gamma(\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-)$ $\Gamma_{97} / \Gamma_{96}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

$0.36^{+0.24}_{-0.20}$ OUR FIT Error includes scale factor of 1.8.

$0.75 \pm 0.17 \pm 0.19$ ANJOS 90D E691 Photoproduction

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^- \text{no-}\rho) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{98} / \Gamma_{37}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

$0.048 \pm 0.015 \pm 0.011$ FRABETTI 97C E687 γ Be, $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(K^- \rho^0 \pi^+ \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{66} / \Gamma_{37}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

$0.034 \pm 0.009 \pm 0.005$ FRABETTI 97C E687 γ Be, $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^- \text{nonresonant}) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{67} / \Gamma_{37}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

< 0.026 90 FRABETTI 97C E687 γ Be, $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(K^- \pi^+ \pi^+ \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{68} / Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

$0.022^{+0.047}_{-0.008} \pm 0.004$ 1 ⁵¹ AGUILAR-... 87F HYBR πp , $p p$ 360, 400 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.015 ⁵¹ BARLAG 92C ACCM π^- Cu 230 GeV

⁵¹ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{69} / Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

$0.054^{+0.030}_{-0.014}$ OUR AVERAGE

$0.099^{+0.036}_{-0.070}$ ⁵² BARLAG 92C ACCM π^- Cu 230 GeV

$0.044^{+0.052}_{-0.013} \pm 0.007$ 2 ⁵² AGUILAR-... 87F HYBR πp , $p p$ 360, 400 GeV

⁵² AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^+ \pi^- \pi^-) / \Gamma_{\text{total}}$ Γ_{70} / Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.0008 ± 0.0007 ⁵³ BARLAG 92C ACCM π^- Cu 230 GeV

⁵³ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{71} / Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.0020 ± 0.0018 ⁵⁴ BARLAG 92C ACCM π^- Cu 230 GeV

⁵⁴ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\overline{K}^0 \overline{K}^0 K^+)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{72}/Γ_{37}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------|-------------|-------------------------------------|-------------|----------------------------|
| 0.20±0.09 OUR AVERAGE | | Error includes scale factor of 2.4. | | |
| 0.14±0.04±0.02 | 39 | ALBRECHT | 94I ARG | $e^+ e^- \approx 10$ GeV |
| 0.34±0.07 | 70 | AMMAR | 91 CLEO | $e^+ e^- \approx 10.5$ GeV |

————— Pionic modes —————

$\Gamma(\pi^+ \pi^0)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{100}/Γ_{37}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|------------------------------|
| 0.028±0.006±0.005 | 34 | SELEN | 93 CLE2 | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\pi^+ \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{101}/Γ_{37}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|-------------|--------------------|-------------|-------------------------------|
| 0.0406±0.0034 OUR FIT | | | | |
| 0.0403±0.0035 OUR AVERAGE | | | | |
| 0.043 ±0.003 ±0.003 | 236 | FRABETTI | 97D E687 | γ Be ≈ 200 GeV |
| 0.032 ±0.011 ±0.003 | 20 | ADAMOVICH | 93 WA82 | π^- 340 GeV |
| 0.035 ±0.007 ±0.003 | | ANJOS | 89 E691 | Photoproduction |
| 0.042 ±0.016 ±0.010 | 57 | BALTRUSAIT..85E | MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(\rho^0 \pi^+)/\Gamma(\pi^+ \pi^+ \pi^-)$ $\Gamma_{102}/\Gamma_{101}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|------------------------|-------------|-------------------------------|
| 0.289±0.055±0.058 | ⁵⁵ FRABETTI | 97D E687 | γ Be ≈ 200 GeV |

⁵⁵ FRABETTI 97D also includes $f_2(1270)\pi^+$ and $f_0(980)\pi^+$ modes in the fit, but the resulting decay fractions are not statistically significant.

$\Gamma(\rho^0 \pi^+)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{102}/Γ_{37}

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|-------|---------|-----------------|
| <0.015 | 90 | ANJOS | 89 E691 | Photoproduction |
|--------|----|-------|---------|-----------------|

$\Gamma(\pi^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma(\pi^+ \pi^+ \pi^-)$ $\Gamma_{103}/\Gamma_{101}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

0.62 ±0.11 OUR FIT

| | | | |
|--------------------------|------------------------|----------|-------------------------------|
| 0.589±0.105±0.081 | ⁵⁶ FRABETTI | 97D E687 | γ Be ≈ 200 GeV |
|--------------------------|------------------------|----------|-------------------------------|

⁵⁶ FRABETTI 97D also includes $f_2(1270)\pi^+$ and $f_0(980)\pi^+$ modes in the fit, but the resulting decay fractions are not statistically significant.

$\Gamma(\pi^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma(K^- \pi^+ \pi^+)$ Γ_{103}/Γ_{37}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

0.025±0.005 OUR FIT

| | | | |
|--------------------------|-------|---------|-----------------|
| 0.027±0.007±0.002 | ANJOS | 89 E691 | Photoproduction |
|--------------------------|-------|---------|-----------------|

$\Gamma(\pi^+ \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{104}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

| | | | |
|--|----------------------|----------|--------------------|
| 0.019^{+0.015}_{-0.012} | ⁵⁷ BARLAG | 92C ACCM | π^- Cu 230 GeV |
|--|----------------------|----------|--------------------|

⁵⁷ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+\pi^+)$ Γ_{104}/Γ_{37}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|-------|----------|-----------------|
| <0.4 | 90 | ANJOS | 89E E691 | Photoproduction |
|------|----|-------|----------|-----------------|

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+)$ $\Gamma_{109}/\Gamma_{133}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

| | | | | |
|------------------|-----|--------|---------|-----------------------------|
| 0.49±0.08 | 275 | JESSOP | 98 CLE2 | $e^+e^- \approx \gamma(4S)$ |
|------------------|-----|--------|---------|-----------------------------|

$\Gamma(\eta\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{109}/Γ_{37}

Unseen decay modes of the η are included.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|-----|------|-------------|------|---------|
|-------|-----|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------------|----|----|--------|----------|-----------------|
| $0.083 \pm 0.023 \pm 0.014$ | | 99 | DAOUDI | 92 CLE2 | See JESSOP 98 |
| <0.12 | 90 | | ANJOS | 89E E691 | Photoproduction |

$\Gamma(\omega\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{111}/Γ_{37}

Unseen decay modes of the ω are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|-----------------|----|-------|----------|-----------------|
| <0.08 | 90 | ANJOS | 89E E691 | Photoproduction |
|-----------------|----|-------|----------|-----------------|

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{107}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|------------------------------|----|--------|-----------------------------|
| $0.0010^{+0.0008}_{-0.0007}$ | 58 | BARLAG | 92C ACCM π^- Cu 230 GeV |
|------------------------------|----|--------|-----------------------------|

⁵⁸ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma(K^-\pi^+\pi^+)$ Γ_{107}/Γ_{37}

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|-----|------|-------------|------|---------|
|-------|-----|------|-------------|------|---------|

| | | | | | |
|--------------------------|--|----|----------|----------|--|
| 0.023±0.004±0.002 | | 58 | FRABETTI | 97C E687 | $\gamma\text{Be}, \bar{E}_\gamma \approx 200$ GeV |
|--------------------------|--|----|----------|----------|--|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------|----|--|-------|---------|-----------------|
| <0.019 | 90 | | ANJOS | 89 E691 | Photoproduction |
|--------|----|--|-------|---------|-----------------|

$\Gamma(\eta\rho^+)/\Gamma(\phi\pi^+)$ $\Gamma_{112}/\Gamma_{133}$

Unseen decay modes of the η are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|-----------------|----|--------|---------|-----------------------------|
| <1.11 | 90 | JESSOP | 98 CLE2 | $e^+e^- \approx \gamma(4S)$ |
|-----------------|----|--------|---------|-----------------------------|

$\Gamma(\eta\rho^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{112}/Γ_{37}

Unseen decay modes of the η are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|--------|---------|---------------|
| <0.13 | 90 | DAOUDI | 92 CLE2 | See JESSOP 98 |
|-------|----|--------|---------|---------------|

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{108}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|------------------------------|----|--------|-----------------------------|
| $0.0029^{+0.0029}_{-0.0020}$ | 59 | BARLAG | 92C ACCM π^- Cu 230 GeV |
|------------------------------|----|--------|-----------------------------|

⁵⁹ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$ $\Gamma_{113}/\Gamma_{133}$

Unseen decay modes of the $\eta'(958)$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------|------|-------------|---------|-----------------------------|
| 0.82±0.14 | 126 | JESSOP | 98 CLE2 | $e^+e^- \approx \gamma(4S)$ |

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{113}/Γ_{37}

Unseen decay modes of the $\eta'(958)$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|---------|----------|---|
| <0.1 | 90 | DAOUDI | 92 CLE2 | See JESSOP 98 |
| <0.1 | 90 | ALVAREZ | 91 NA14 | Photoproduction |
| <0.13 | 90 | ANJOS | 91B E691 | $\gamma\text{Be}, \bar{E}_\gamma \approx 145 \text{ GeV}$ |

$\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$ $\Gamma_{114}/\Gamma_{133}$

Unseen decay modes of the $\eta'(958)$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|-------------|---------|-----------------------------|
| <0.86 | 90 | JESSOP | 98 CLE2 | $e^+e^- \approx \gamma(4S)$ |

$\Gamma(\eta'(958)\rho^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{114}/Γ_{37}

Unseen decay modes of the $\eta'(958)$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|--------|---------|---------------|
| <0.17 | 90 | DAOUDI | 92 CLE2 | See JESSOP 98 |
|-------|----|--------|---------|---------------|

————— **Hadronic modes with a $K\bar{K}$ pair** —————

$\Gamma(K^+\bar{K}^0)/\Gamma(\bar{K}^0\pi^+)$ Γ_{115}/Γ_{36}

It is generally assumed for modes such as $D^+ \rightarrow \bar{K}^0\pi^+$ that

$$\Gamma(D^+ \rightarrow \bar{K}^0\pi^+) = 2\Gamma(D^+ \rightarrow K_S^0\pi^+);$$

it is the latter Γ that is actually measured. BIGI 95 points out that interference between Cabibbo-allowed and doubly Cabibbo-suppressed amplitudes, where both occur, could invalidate this assumption by a few percent.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.255±0.029 OUR FIT

0.263±0.035 OUR AVERAGE

| | | | | |
|-------------------|-----|-----------------|----------|--|
| 0.25 ±0.04 ±0.02 | 129 | FRABETTI | 95 E687 | $\gamma\text{Be} \bar{E}_\gamma \approx 200 \text{ GeV}$ |
| 0.271±0.065±0.039 | 69 | ANJOS | 90C E691 | γBe |
| 0.317±0.086±0.048 | 31 | BALTRUSAIT..85E | MRK3 | $e^+e^- 3.77 \text{ GeV}$ |
| 0.25 ±0.15 | 6 | SCHINDLER | 81 MRK2 | $e^+e^- 3.771 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------|----|----------------------|---------|-----------------------------|
| 0.222±0.041±0.029 | 70 | ⁶⁰ BISHAI | 97 CLE2 | $e^+e^- \approx \gamma(4S)$ |
|-------------------|----|----------------------|---------|-----------------------------|

⁶⁰This BISHAI 97 result is redundant with results elsewhere in the Listings.

$\Gamma(K^+\bar{K}^0)/\Gamma(K^-\pi^+\pi^+)$ Γ_{115}/Γ_{37}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.082±0.010 OUR FIT

| | | | | |
|--------------------------|----|----------------------|---------|-----------------------------|
| 0.077±0.014±0.007 | 70 | ⁶¹ BISHAI | 97 CLE2 | $e^+e^- \approx \gamma(4S)$ |
|--------------------------|----|----------------------|---------|-----------------------------|

⁶¹See BISHAI 97 for an isospin analysis of $D^+ \rightarrow K\bar{K}$ amplitudes.

$\Gamma(K^+ K^- \pi^+)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{116}/Γ_{37}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|----------|-----------------------------|
| 0.097 ± 0.006 OUR NEW AVERAGE [0.098 ± 0.006 OUR 1998 AVERAGE] | | | |
| 0.093 ± 0.010 ^{+0.008} / _{-0.006} | JUN | 00 SELX | Σ^- nucleus, 600 GeV |
| 0.0976 ± 0.0042 ± 0.0046 | FRABETTI | 95B E687 | Dalitz plot analysis |

$\Gamma(\phi \pi^+)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{133}/Γ_{37}

Unseen decay modes of the ϕ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|-----------------|----------|----------------------------|
| 0.068 ± 0.005 OUR AVERAGE | | | | |
| 0.058 ± 0.006 ± 0.006 | | FRABETTI | 95B E687 | Dalitz plot analysis |
| 0.062 ± 0.017 ± 0.006 | 19 | ADAMOVICH | 93 WA82 | π^- 340 GeV |
| 0.077 ± 0.011 ± 0.005 | 128 | DAOUDI | 92 CLE2 | $e^+ e^- \approx 10.5$ GeV |
| 0.098 ± 0.032 ± 0.014 | 12 | ALVAREZ | 90C NA14 | Photoproduction |
| 0.071 ± 0.008 ± 0.007 | 84 | ANJOS | 88 E691 | Photoproduction |
| 0.084 ± 0.021 ± 0.011 | 21 | BALTRUSAIT..85E | MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(K^+ \bar{K}^*(892)^0)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{137}/Γ_{37}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|------------------------|----------|----------------------|
| 0.047 ± 0.005 OUR AVERAGE Error includes scale factor of 1.2. | | | | |
| 0.044 ± 0.003 ± 0.004 | | ⁶² FRABETTI | 95B E687 | Dalitz plot analysis |
| 0.058 ± 0.009 ± 0.006 | 73 | ANJOS | 88 E691 | Photoproduction |
| 0.048 ± 0.021 ± 0.011 | 14 | BALTRUSAIT..85E | MRK3 | $e^+ e^-$ 3.77 GeV |

⁶²See FRABETTI 95B for evidence also of $\bar{K}_0^*(1430)^0 K^+$ in the $D^+ \rightarrow K^+ K^- \pi^+$ Dalitz plot.

$\Gamma(K^+ K^- \pi^+ \text{ nonresonant})/\Gamma(K^- \pi^+ \pi^+)$ Γ_{119}/Γ_{37}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|-----------------|---------|--------------------|
| 0.050 ± 0.009 OUR AVERAGE | | | | |
| 0.049 ± 0.008 ± 0.006 | 95 | ANJOS | 88 E691 | Photoproduction |
| 0.059 ± 0.026 ± 0.009 | 37 | BALTRUSAIT..85E | MRK3 | $e^+ e^-$ 3.77 GeV |

$\Gamma(K^*(892)^+ \bar{K}^0)/\Gamma(\bar{K}^0 \pi^+)$ Γ_{138}/Γ_{36}

Unseen decay modes of the $K^*(892)^+$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------|------|-------------|---------|--|
| 1.1 ± 0.3 ± 0.4 | 67 | FRABETTI | 95 E687 | γ Be $\bar{E}_\gamma \approx 200$ GeV |

$\Gamma(\phi \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{134}/Γ

Unseen decay modes of the ϕ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------|----------------------|----------|--------------------|
| 0.023 ± 0.010 | ⁶³ BARLAG | 92C ACCM | π^- Cu 230 GeV |

⁶³BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\phi \pi^+ \pi^0)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{134}/Γ_{37}

Unseen decay modes of the ϕ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|-----------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.58 | 90 | ALVAREZ | 90C NA14 | Photoproduction |
| <0.28 | 90 | ANJOS | 89E E691 | Photoproduction |

$\Gamma(\phi\rho^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{135}/Γ_{37}

Unseen decay modes of the ϕ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|---------|---------------------------|
| <0.16 | 90 | DAOUDI | 92 CLE2 | $e^+e^- \approx 10.5$ GeV |

$\Gamma(K^+K^-\pi^+\pi^0 \text{ non-}\phi)/\Gamma_{\text{total}}$ Γ_{125}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|----------------------|----------|--------------------|
| $0.015^{+0.007}_{-0.006}$ | ⁶⁴ BARLAG | 92C ACCM | π^- Cu 230 GeV |

⁶⁴ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^+K^-\pi^+\pi^0 \text{ non-}\phi)/\Gamma(K^-\pi^+\pi^+)$ Γ_{125}/Γ_{37}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|-------|----------|-----------------|
| <0.25 | 90 | ANJOS | 89E E691 | Photoproduction |
|-------|----|-------|----------|-----------------|

$\Gamma(K^+\bar{K}^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{126}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|---------|--------------------------|
| <0.02 | 90 | ALBRECHT | 92B ARG | $e^+e^- \simeq 10.4$ GeV |

$\Gamma(K^0K^-\pi^+\pi^+)/\Gamma_{\text{total}}$ Γ_{127}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------|---------|--------------------------|
| $0.01 \pm 0.005 \pm 0.003$ | ALBRECHT | 92B ARG | $e^+e^- \simeq 10.4$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------|----------------------|----------|--------------------|
| <0.003 | ⁶⁵ BARLAG | 92C ACCM | π^- Cu 230 GeV |
|--------|----------------------|----------|--------------------|

⁶⁵ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^*(892)^+\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{139}/Γ

Unseen decay modes of the $K^*(892)$'s are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|---------|--------------------------|
| $0.026 \pm 0.008 \pm 0.007$ | ALBRECHT | 92B ARG | $e^+e^- \simeq 10.4$ GeV |

$\Gamma(K^0K^-\pi^+\pi^+ \text{ non-}K^*\bar{K}^0)/\Gamma_{\text{total}}$ Γ_{129}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---------|-----|-------------|---------|--------------------------|
| <0.0079 | 90 | ALBRECHT | 92B ARG | $e^+e^- \simeq 10.4$ GeV |

$\Gamma(\phi\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{136}/Γ

Unseen decay modes of the ϕ are included.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------|-----|------|-------------|---------|-----------------|
| <0.002 | 90 | 0 | ANJOS | 88 E691 | Photoproduction |

$\Gamma(\phi\pi^+\pi^+\pi^-)/\Gamma(K^-\pi^+\pi^+)$ Γ_{136}/Γ_{37}

Unseen decay modes of the ϕ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|---------|----------|-----------------|
| <0.031 | 90 | ALVAREZ | 90C NA14 | Photoproduction |
|--------|----|---------|----------|-----------------|

$\Gamma(\phi\pi^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ $\Gamma_{136}/\Gamma_{133}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|---------|-------------|
| <0.6 | 90 | FRABETTI | 92 E687 | γ Be |

$\Gamma(K^+K^-\pi^+\pi^+\pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{132}/Γ

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|-----|------|-------------|---------|-----------------|
| <0.03 | 90 | 12 | ANJOS | 88 E691 | Photoproduction |

————— Rare or forbidden modes —————

$\Gamma(K^+\pi^+\pi^-)/\Gamma(K^-\pi^+\pi^+)$ Γ_{140}/Γ_{37}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|-------------|----------|---|
| 0.0075±0.0016 OUR AVERAGE | | | | |
| 0.0077±0.0017±0.0008 | 59 | AITALA | 97C E791 | π^- nucleus, 500 GeV |
| 0.0072±0.0023±0.0017 | 21 | FRABETTI | 95E E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |

$\Gamma(K^+\rho^0)/\Gamma(K^+\pi^+\pi^-)$ $\Gamma_{141}/\Gamma_{140}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------|--------------------------|
| 0.37±0.14±0.07 | AITALA | 97C E791 | π^- nucleus, 500 GeV |

$\Gamma(K^+\rho^0)/\Gamma(K^-\pi^+\pi^+)$ Γ_{141}/Γ_{37}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---------|-----|-------------|----------|---|
| <0.0067 | 90 | FRABETTI | 95E E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |

$\Gamma(K^*(892)^0\pi^+)/\Gamma(K^+\pi^+\pi^-)$ $\Gamma_{142}/\Gamma_{140}$

Unseen decay modes of the $K^*(892)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------|--------------------------|
| 0.53±0.21±0.02 | AITALA | 97C E791 | π^- nucleus, 500 GeV |

$\Gamma(K^*(892)^0\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{142}/Γ_{37}

Unseen decay modes of the $K^*(892)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---------|-----|-------------|----------|---|
| <0.0021 | 90 | FRABETTI | 95E E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |

$\Gamma(K^+\pi^+\pi^- \text{ nonresonant})/\Gamma(K^+\pi^+\pi^-)$ $\Gamma_{143}/\Gamma_{140}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------|--------------------------|
| 0.36±0.14±0.07 | AITALA | 97C E791 | π^- nucleus, 500 GeV |

$\Gamma(K^+K^+K^-)/\Gamma(K^-\pi^+\pi^+)$ Γ_{144}/Γ_{37}

A doubly Cabibbo-suppressed decay with no simple spectator process possible.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------|-----|------|------------------------|----------|---|
| <0.0016 | 90 | | ⁶⁶ FRABETTI | 95F E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.057 ±0.020±0.007 13 ADAMOVICH 93 WA82 π^- 340 GeV

⁶⁶ Using the $\phi\pi^+$ mode to normalize, FRABETTI 95F gets $\Gamma(K^+K^+K^-)/\Gamma(\phi\pi^+) < 0.025$.

$\Gamma(\phi K^+)/\Gamma(\phi\pi^+)$ $\Gamma_{145}/\Gamma_{133}$

A doubly Cabibbo-suppressed decay with no simple spectator process possible.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|------|-------------|----------|---|
| <0.021 | 90 | | FRABETTI | 95F E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------------------------------|--|---|---------------------|----------|---|
| $0.058^{+0.032}_{-0.026} \pm 0.007$ | | 4 | ⁶⁷ ANJOS | 92D E691 | γ Be, $\bar{E}_\gamma = 145$ GeV |
|-------------------------------------|--|---|---------------------|----------|---|

⁶⁷ The evidence of ANJOS 92D is a small excess of events ($4.5^{+2.4}_{-2.0}$).

$\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{146}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|------|--|
| <5.2 × 10⁻⁵ (CL = 90%) | | | | | [<6.6 × 10 ⁻⁵ (CL = 90%) OUR 1998 BEST LIMIT] |

| | | | | | |
|----------------------------------|----|--|--------|----------|-------------------|
| <5.2 × 10⁻⁵ | 90 | | AITALA | 99G E791 | $\pi^- N$ 500 GeV |
|----------------------------------|----|--|--------|----------|-------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------------------|----|----|----------|----------|---|
| <1.1 × 10 ⁻⁴ | 90 | | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <6.6 × 10 ⁻⁵ | 90 | | AITALA | 96 E791 | $\pi^- N$ 500 GeV |
| <2.5 × 10 ⁻³ | 90 | | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |
| <2.6 × 10 ⁻³ | 90 | 39 | HAAS | 88 CLEO | $e^+ e^-$ 10 GeV |

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{147}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|------|--|
| <1.5 × 10⁻⁵ (CL = 90%) | | | | | [<1.8 × 10 ⁻⁵ (CL = 90%) OUR 1998 BEST LIMIT] |

| | | | | | |
|----------------------------------|----|--|--------|----------|-------------------|
| <1.5 × 10⁻⁵ | 90 | | AITALA | 99G E791 | $\pi^- N$ 500 GeV |
|----------------------------------|----|--|--------|----------|-------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------------------|----|----|----------|----------|---|
| <8.9 × 10 ⁻⁵ | 90 | | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <1.8 × 10 ⁻⁵ | 90 | | AITALA | 96 E791 | $\pi^- N$ 500 GeV |
| <2.2 × 10 ⁻⁴ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |
| <5.9 × 10 ⁻³ | 90 | | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |
| <2.9 × 10 ⁻³ | 90 | 36 | HAAS | 88 CLEO | $e^+ e^-$ 10 GeV |

$\Gamma(\rho^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{148}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|------|-------------|---------|--------------------------|
| <5.6 × 10⁻⁴ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{149}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-------------|----------|---|
| <2.0 × 10⁻⁴ | 90 | AITALA | 99G E791 | $\pi^- N$ 500 GeV |
| <2.0 × 10⁻⁴ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------------|----|------|----------|------------------|
| <4.8 × 10 ⁻³ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |
|-------------------------|----|------|----------|------------------|

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{150}/Γ

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|----------|--|
| <4.4 × 10⁻⁵ (CL = 90%) | | | | | [<9.7 × 10 ⁻⁵ (CL = 90%) OUR 1998 BEST LIMIT] |
| <4.4 × 10⁻⁵ | 90 | | AITALA | 99G E791 | $\pi^- N$ 500 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <9.7 × 10 ⁻⁵ | 90 | | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <3.2 × 10 ⁻⁴ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |
| <9.2 × 10 ⁻³ | 90 | | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(\pi^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{151}/Γ

A test of lepton-family-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-------------|----------|-------------------|
| <3.4 × 10⁻⁵ | 90 | AITALA | 99G E791 | $\pi^- N$ 500 GeV |

$\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{152}/Γ

A test of lepton-family-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <1.1 × 10 ⁻⁴ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <3.3 × 10 ⁻³ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{153}/Γ

A test of lepton-family-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <1.3 × 10 ⁻⁴ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <3.3 × 10 ⁻³ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(K^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{154}/Γ

A test of lepton-family-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-------------|----------|-------------------|
| <6.8 × 10⁻⁵ | 90 | AITALA | 99G E791 | $\pi^- N$ 500 GeV |

$\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{155}/Γ

A test of lepton-family-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <1.3 × 10 ⁻⁴ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <3.4 × 10 ⁻³ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{156}/Γ

A test of lepton-family-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <1.2 × 10 ⁻⁴ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <3.4 × 10 ⁻³ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{157}/Γ

A test of lepton-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------------------|----------------------|---|
| <9.6 × 10⁻⁵ (CL = 90%) | | [<1.1 × 10 ⁻⁴ (CL = 90%) | OUR 1998 BEST LIMIT] | |
| <9.6 × 10⁻⁵ | 90 | AITALA | 99G E791 | $\pi^- N$ 500 GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| <1.1 × 10 ⁻⁴ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <4.8 × 10 ⁻³ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{158}/Γ

A test of lepton-number conservation.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------------------------------|----------------------|---|
| <1.7 × 10⁻⁵ (CL = 90%) | | | [<8.7 × 10 ⁻⁵ (CL = 90%) | OUR 1998 BEST LIMIT] | |
| <1.7 × 10⁻⁵ | 90 | | AITALA | 99G E791 | $\pi^- N$ 500 GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <8.7 × 10 ⁻⁵ | 90 | | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <2.2 × 10 ⁻⁴ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |
| <6.8 × 10 ⁻³ | 90 | | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{159}/Γ

A test of lepton-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------------------|----------------------|---|
| <5.0 × 10⁻⁵ (CL = 90%) | | [<1.1 × 10 ⁻⁴ (CL = 90%) | OUR 1998 BEST LIMIT] | |
| <5.0 × 10⁻⁵ | 90 | AITALA | 99G E791 | $\pi^- N$ 500 GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| <1.1 × 10 ⁻⁴ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| <3.7 × 10 ⁻³ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(\rho^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{160}/Γ

A test of lepton-number conservation.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|------|-------------|---------|--------------------------|
| <5.6 × 10⁻⁴ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |

$\Gamma(K^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{161}/Γ

A test of lepton-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|---|
| <1.2 × 10⁻⁴ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| <9.1 × 10 ⁻³ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{162}/Γ

A test of lepton-number conservation.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|----------|---|
| <1.2 × 10⁻⁴ | 90 | | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <3.2 × 10 ⁻⁴ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |
| <4.3 × 10 ⁻³ | 90 | | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{163}/Γ

A test of lepton-number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|---|
| $<1.3 \times 10^{-4}$ | 90 | FRABETTI | 97B E687 | γ Be, $\bar{E}_\gamma \approx 220$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $<4.0 \times 10^{-3}$ | 90 | WEIR | 90B MRK2 | $e^+ e^-$ 29 GeV |

$\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{164}/Γ

A test of lepton-number conservation.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|------|-------------|---------|--------------------------|
| $<8.5 \times 10^{-4}$ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |

D^\pm CP-VIOLATING DECAY-RATE ASYMMETRIES

$A_{CP}(K^+ K^- \pi^\pm)$ in $D^\pm \rightarrow K^+ K^- \pi^\pm$

This is the difference between D^+ and D^- partial widths for these modes divided by the sum of the widths.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|------------------------|----------|-------------------------------------|
| -0.017 ± 0.027 OUR AVERAGE | | | |
| -0.014 ± 0.029 | ⁶⁸ AITALA | 97B E791 | $-0.062 < A_{CP} < +0.034$ (90% CL) |
| -0.031 ± 0.068 | ⁶⁸ FRABETTI | 94I E687 | $-0.14 < A_{CP} < +0.081$ (90% CL) |
| ⁶⁸ FRABETTI 94I and AITALA 97B measure $N(D^+ \rightarrow K^- K^+ \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$, the ratio of numbers of events observed, and similarly for the D^- . | | | |

$A_{CP}(K^\pm K^{*0})$ in $D^+ \rightarrow K^+ \bar{K}^{*0}$, $D^- \rightarrow K^- K^{*0}$

This is the difference between D^+ and D^- partial widths for these modes divided by the sum of the widths.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|----------|-------------------------------------|
| -0.02 ± 0.05 OUR AVERAGE | | | |
| -0.010 ± 0.050 | ⁶⁹ AITALA | 97B E791 | $-0.092 < A_{CP} < +0.072$ (90% CL) |
| -0.12 ± 0.13 | ⁶⁹ FRABETTI | 94I E687 | $-0.33 < A_{CP} < +0.094$ (90% CL) |
| ⁶⁹ FRABETTI 94I and AITALA 97B measure $N(D^+ \rightarrow K^+ \bar{K}^*(892)^0)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$, the ratio of numbers of events observed, and similarly for the D^- . | | | |

$A_{CP}(\phi \pi^\pm)$ in $D^\pm \rightarrow \phi \pi^\pm$

This is the difference between D^+ and D^- partial widths for these modes divided by the sum of the widths.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|----------|-------------------------------------|
| -0.014 ± 0.033 OUR AVERAGE | | | |
| -0.028 ± 0.036 | ⁷⁰ AITALA | 97B E791 | $-0.087 < A_{CP} < +0.031$ (90% CL) |
| $+0.066 \pm 0.086$ | ⁷⁰ FRABETTI | 94I E687 | $-0.075 < A_{CP} < +0.21$ (90% CL) |
| ⁷⁰ FRABETTI 94I and AITALA 97B measure $N(D^+ \rightarrow \phi \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$, the ratio of numbers of events observed, and similarly for the D^- . | | | |

$A_{CP}(\pi^+ \pi^- \pi^\pm)$ in $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm$

This is the difference between D^+ and D^- partial widths for these modes divided by the sum of the widths.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|----------|-------------------------------------|
| -0.017 ± 0.042 | ⁷¹ AITALA | 97B E791 | $-0.086 < A_{CP} < +0.052$ (90% CL) |
| ⁷¹ AITALA 97B measure $N(D^+ \rightarrow \pi^+ \pi^- \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$, the ratio of numbers of events observed, and similarly for the D^- . | | | |

D^\pm PRODUCTION CROSS SECTION AT $\psi(3770)$

A compilation of the cross sections for the direct production of D^\pm mesons at or near the $\psi(3770)$ peak in e^+e^- production.

| <u>VALUE (nanobarns)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|--------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 4.2 ± 0.6 ± 0.3 | 72 ADLER | 88C MRK3 | e^+e^- 3.768 GeV |
| 5.5 ± 1.0 | 73 PARTRIDGE | 84 CBAL | e^+e^- 3.771 GeV |
| 6.00 ± 0.72 ± 1.02 | 74 SCHINDLER | 80 MRK2 | e^+e^- 3.771 GeV |
| 9.1 ± 2.0 | 75 PERUZZI | 77 MRK1 | e^+e^- 3.774 GeV |

⁷²This measurement compares events with one detected D to those with two detected D mesons, to determine the the absolute cross section. ADLER 88C measure the ratio of cross sections (neutral to charged) to be $1.36 \pm 0.23 \pm 0.14$. This measurement does not include the decays of the $\psi(3770)$ not associated with charmed particle production.

⁷³This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. PARTRIDGE 84 measures 6.4 ± 1.15 nb for the cross section. We take the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and we assume that the $\psi(3770)$ is an isosinglet to evaluate the cross sections. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

⁷⁴This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. SCHINDLER 80 assume the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and that the $\psi(3770)$ is an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

⁷⁵This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. The phase space division of neutral and charged D mesons in $\psi(3770)$ decay is taken to be 1.33, and $\psi(3770)$ is assumed to be an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction. We exclude this measurement from the average because of uncertainties in the contamination from τ lepton pairs. Also see RAPIDIS 77.

$D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$ FORM FACTORS

$r_\nu \equiv V(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------------------|-------------|----------------------------------|
| 1.82 ± 0.09 OUR NEW AVERAGE | | [1.85 ± 0.12 OUR 1998 AVERAGE] | | |
| 1.45 ± 0.23 ± 0.07 | 763 | ADAMOVICH 99 | BEAT | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| 1.90 ± 0.11 ± 0.09 | 3000 | ⁷⁶ AITALA 98B | E791 | $\bar{K}^*(892)^0 e^+ \nu_e$ |
| 1.84 ± 0.11 ± 0.09 | 3034 | AITALA 98F | E791 | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| 1.74 ± 0.27 ± 0.28 | 874 | FRABETTI 93E | E687 | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| 2.00 ^{+0.34} _{-0.32} ± 0.16 | 305 | KODAMA 92 | E653 | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| 2.0 ± 0.6 ± 0.3 | 183 | ANJOS 90E | E691 | $\bar{K}^*(892)^0 e^+ \nu_e$ |

⁷⁶This is slightly different from the AITALA 98B value: see ref. [5] in AITALA 98F.

$r_2 \equiv A_2(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------------------|-------------|------------------------------------|
| 0.78±0.07 OUR NEW AVERAGE | | [0.72 ± 0.09 OUR 1998 AVERAGE] | | |
| 1.00±0.15±0.03 | 763 | ADAMOVICH 99 | BEAT | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 0.71±0.08±0.09 | 3000 | AITALA 98B | E791 | $\bar{K}^*(892)^0_{e^+ \nu_e}$ |
| 0.75±0.08±0.09 | 3034 | AITALA 98F | E791 | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 0.78±0.18±0.10 | 874 | FRABETTI 93E | E687 | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 0.82 ^{+0.22} _{-0.23} ±0.11 | 305 | KODAMA 92 | E653 | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 0.0 ±0.5 ±0.2 | 183 | ANJOS 90E | E691 | $\bar{K}^*(892)^0_{e^+ \nu_e}$ |

$r_3 \equiv A_3(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$


| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|------------------------------------|
| 0.04±0.33±0.29 | 3034 | AITALA 98F | E791 | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |

Γ_L/Γ_T in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------------------|-------------|------------------------------------|
| 1.14±0.08 OUR NEW AVERAGE | | [1.23 ± 0.13 OUR 1998 AVERAGE] | | |
| 1.09±0.10±0.02 | 763 | ADAMOVICH 99 | BEAT | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 1.20±0.13±0.13 | 874 | FRABETTI 93E | E687 | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 1.18±0.18±0.08 | 305 | KODAMA 92 | E653 | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 1.8 ^{+0.6} _{-0.4} ±0.3 | 183 | ANJOS 90E | E691 | $\bar{K}^*(892)^0_{e^+ \nu_e}$ |

Γ_+/Γ_- in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--|-------------|------------------------------------|
| 0.21±0.04 OUR NEW AVERAGE | | Error includes scale factor of 1.3. See the ideogram below. [0.16 ± 0.04 OUR 1998 AVERAGE] | | |
| 0.28±0.05±0.02 | 763 | ADAMOVICH 99 | BEAT | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 0.16±0.05±0.02 | 305 | KODAMA 92 | E653 | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 0.15 ^{+0.07} _{-0.05} ±0.03 | 183 | ANJOS 90E | E691 | $\bar{K}^*(892)^0_{e^+ \nu_e}$ |



$$\Gamma_+/\Gamma_- \text{ in } D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$$

D^\pm REFERENCES

| | | | | |
|-----------|-----|---------------|-----------------------------|---------------------------|
| JUN | 00 | PRL 84 1857 | S.Y. Jun <i>et al.</i> | (FNAL SELEX Collab.) |
| PDG | 00 | EPJ C15 1 | | |
| ABBIENDI | 99K | EPJ C8 573 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABE | 99P | PR D60 092005 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ADAMOVICH | 99 | EPJ C6 35 | M. Adamovich <i>et al.</i> | (CERN BEATRICE Collab.) |
| AITALA | 99G | PL B462 401 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| BONVICINI | 99 | PRL 82 4586 | G. Bonvicini <i>et al.</i> | (CLEO Collab.) |
| AITALA | 98B | PRL 80 1393 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| AITALA | 98F | PL B440 435 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| BAI | 98B | PL B429 188 | J.Z. Bai <i>et al.</i> | (BEPC BES Collab.) |
| JESSOP | 98 | PR D58 052002 | C.P. Jessop <i>et al.</i> | (CLEO Collab.) |
| AITALA | 97 | PL B397 325 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| AITALA | 97B | PL B403 377 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| AITALA | 97C | PL B404 187 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| BARTELT | 97 | PL B405 373 | J. Bartelt <i>et al.</i> | (CLEO Collab.) |
| BISHAI | 97 | PRL 78 3261 | M. Bishai <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 97 | PL B391 235 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 97B | PL B398 239 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 97C | PL B401 131 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 97D | PL B407 79 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| AITALA | 96 | PRL 76 364 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| ALBRECHT | 96C | PL B374 249 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BIGI | 95 | PL B349 363 | I.I. Bigi, H. Yamamoto | (NDAM, HARV) |
| FRABETTI | 95 | PL B346 199 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 95B | PL B351 591 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 95E | PL B359 403 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 95F | PL B363 259 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| KODAMA | 95 | PL B345 85 | K. Kodama <i>et al.</i> | (FNAL E653 Collab.) |
| ALBRECHT | 94I | ZPHY C64 375 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEEV | 94 | PAN 57 1370 | A.N. Aleev <i>et al.</i> | (Serpukhov BIS-2 Collab.) |

Translated from YF 57 1443.

| | | | | |
|-------------|-----|--------------|----------------------------------|-----------------------|
| BALEST | 94 | PRL 72 2328 | R. Balest <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 94D | PL B323 459 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 94G | PL B331 217 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 94I | PR D50 R2953 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| ABE | 93E | PL B313 288 | K. Abe <i>et al.</i> | (VENUS Collab.) |
| ADAMOVICH | 93 | PL B305 177 | M.I. Adamovich <i>et al.</i> | (CERN WA82 Collab.) |
| AKERIB | 93 | PRL 71 3070 | D.S. Akerib <i>et al.</i> | (CLEO Collab.) |
| ALAM | 93 | PRL 71 1311 | M.S. Alam <i>et al.</i> | (CLEO Collab.) |
| ANJOS | 93 | PR D48 56 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| BEAN | 93C | PL B317 647 | A. Bean <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 93E | PL B307 262 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| KODAMA | 93B | PL B313 260 | K. Kodama <i>et al.</i> | (FNAL E653 Collab.) |
| KODAMA | 93C | PL B316 455 | K. Kodama <i>et al.</i> | (FNAL E653 Collab.) |
| SELEN | 93 | PRL 71 1973 | M.A. Selen <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 92B | ZPHY C53 361 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 92F | PL B278 202 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ANJOS | 92 | PR D45 R2177 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ANJOS | 92C | PR D46 1941 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ANJOS | 92D | PRL 69 2892 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| BARLAG | 92C | ZPHY C55 383 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| Also | 90D | ZPHY C48 29 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| COFFMAN | 92B | PR D45 2196 | D.M. Coffman <i>et al.</i> | (Mark III Collab.) |
| DAOUDI | 92 | PR D45 3965 | M. Daoudi <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 92 | PL B281 167 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| KODAMA | 92 | PL B274 246 | K. Kodama <i>et al.</i> | (FNAL E653 Collab.) |
| KODAMA | 92C | PL B286 187 | K. Kodama <i>et al.</i> | (FNAL E653 Collab.) |
| ADAMOVICH | 91 | PL B268 142 | M.I. Adamovich <i>et al.</i> | (WA82 Collab.) |
| ALBRECHT | 91 | PL B255 634 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALVAREZ | 91 | PL B255 639 | M.P. Alvarez <i>et al.</i> | (CERN NA14/2 Collab.) |
| ALVAREZ | 91B | ZPHY C50 11 | M.P. Alvarez <i>et al.</i> | (CERN NA14/2 Collab.) |
| AMMAR | 91 | PR D44 3383 | R. Ammar <i>et al.</i> | (CLEO Collab.) |
| ANJOS | 91B | PR D43 R2063 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ANJOS | 91C | PRL 67 1507 | J.C. Anjos <i>et al.</i> | (FNAL-TPS Collab.) |
| BAI | 91 | PRL 66 1011 | Z. Bai <i>et al.</i> | (Mark III Collab.) |
| COFFMAN | 91 | PL B263 135 | D.M. Coffman <i>et al.</i> | (Mark III Collab.) |
| FRABETTI | 91 | PL B263 584 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| ALVAREZ | 90 | ZPHY C47 539 | M.P. Alvarez <i>et al.</i> | (CERN NA14/2 Collab.) |
| ALVAREZ | 90C | PL B246 261 | M.P. Alvarez <i>et al.</i> | (CERN NA14/2 Collab.) |
| ANJOS | 90C | PR D41 2705 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ANJOS | 90D | PR D42 2414 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ANJOS | 90E | PRL 65 2630 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| BARLAG | 90C | ZPHY C46 563 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| WEIR | 90B | PR D41 1384 | A.J. Weir <i>et al.</i> | (Mark II Collab.) |
| ANJOS | 89 | PRL 62 125 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ANJOS | 89B | PRL 62 722 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ANJOS | 89E | PL B223 267 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ADLER | 88B | PRL 60 1375 | J. Adler <i>et al.</i> | (Mark III Collab.) |
| ADLER | 88C | PRL 60 89 | J. Adler <i>et al.</i> | (Mark III Collab.) |
| ALBRECHT | 88I | PL B210 267 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ANJOS | 88 | PRL 60 897 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| AOKI | 88 | PL B209 113 | S. Aoki <i>et al.</i> | (WA75 Collab.) |
| HAAS | 88 | PRL 60 1614 | P. Haas <i>et al.</i> | (CLEO Collab.) |
| ONG | 88 | PRL 60 2587 | R.A. Ong <i>et al.</i> | (Mark II Collab.) |
| RAAB | 88 | PR D37 2391 | J.R. Raab <i>et al.</i> | (FNAL E691 Collab.) |
| ADAMOVICH | 87 | EPL 4 887 | M.I. Adamovich <i>et al.</i> | |
| ADLER | 87 | PL B196 107 | J. Adler <i>et al.</i> | (Mark III Collab.) |
| AGUILAR-... | 87D | PL B193 140 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | 88B | ZPHY C40 321 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| AGUILAR-... | 87E | ZPHY C36 551 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | 88B | ZPHY C40 321 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| AGUILAR-... | 87F | ZPHY C36 559 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | 88 | ZPHY C38 520 | erratum | |
| BARLAG | 87B | ZPHY C37 17 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| BARTEL | 87 | ZPHY C33 339 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| CSORNA | 87 | PL B191 318 | S.E. Csorna <i>et al.</i> | (CLEO Collab.) |
| PALKA | 87B | ZPHY C35 151 | H. Palka <i>et al.</i> | (ACCMOR Collab.) |
| ABE | 86 | PR D33 1 | K. Abe <i>et al.</i> | |
| AGUILAR-... | 86B | ZPHY C31 491 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |

| | | | | |
|---------------|-----|---------------------|----------------------------------|------------------------|
| BALTRUSAIT... | 86E | PRL 56 2140 | R.M. Baltrusaitis <i>et al.</i> | (Mark III Collab.) |
| PAL | 86 | PR D33 2708 | T. Pal <i>et al.</i> | (DELCO Collab.) |
| AIHARA | 85 | ZPHY C27 39 | H. Aihara <i>et al.</i> | (TPC Collab.) |
| BALTRUSAIT... | 85B | PRL 54 1976 | R.M. Baltrusaitis <i>et al.</i> | (Mark III Collab.) |
| BALTRUSAIT... | 85E | PRL 55 150 | R.M. Baltrusaitis <i>et al.</i> | (Mark III Collab.) |
| BARTEL | 85J | PL 163B 277 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| ADAMOVIICH | 84 | PL 140B 119 | M.I. Adamovich <i>et al.</i> | (CERN WA58 Collab.) |
| ALTHOFF | 84G | ZPHY C22 219 | M. Althoff <i>et al.</i> | (TASSO Collab.) |
| ALTHOFF | 84J | PL 146B 443 | M. Althoff <i>et al.</i> | (TASSO Collab.) |
| DERRICK | 84 | PRL 53 1971 | M. Derrick <i>et al.</i> | (HRS Collab.) |
| KOOP | 84 | PRL 52 970 | D.E. Koop <i>et al.</i> | (DELCO Collab.) |
| PARTRIDGE | 84 | Thesis CALT-68-1150 | R.A. Partridge | (Crystal Ball Collab.) |
| AGUILAR-... | 83B | PL 123B 98 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| AUBERT | 83 | NP B213 31 | J.J. Aubert <i>et al.</i> | (EMC Collab.) |
| PARTRIDGE | 81 | PRL 47 760 | R. Partridge <i>et al.</i> | (Crystal Ball Collab.) |
| SCHINDLER | 81 | PR D24 78 | R.H. Schindler <i>et al.</i> | (Mark II Collab.) |
| TRILLING | 81 | PRPL 75 57 | G.H. Trilling | (LBL, UCB) J |
| BACINO | 80 | PRL 45 329 | W.J. Bacino <i>et al.</i> | (DELCO Collab.) |
| SCHINDLER | 80 | PR D21 2716 | R.H. Schindler <i>et al.</i> | (Mark II Collab.) |
| ZHOLENTZ | 80 | PL 96B 214 | A.A. Zholents <i>et al.</i> | (NOVO) |
| Also | 81 | SJNP 34 814 | A.A. Zholents <i>et al.</i> | (NOVO) |
| BACINO | 79 | PRL 43 1073 | W.J. Bacino <i>et al.</i> | (DELCO Collab.) |
| BRANDELIK | 79 | PL 80B 412 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| FELLER | 78 | PRL 40 274 | J.M. Feller <i>et al.</i> | (Mark I Collab.) |
| VUILLEMIN | 78 | PRL 41 1149 | V. Vuillemin <i>et al.</i> | (Mark I Collab.) |
| GOLDHABER | 77 | PL 69B 503 | G. Goldhaber <i>et al.</i> | (Mark I Collab.) |
| PERUZZI | 77 | PRL 39 1301 | I. Peruzzi <i>et al.</i> | (Mark I Collab.) |
| PICCOLO | 77 | PL 70B 260 | M. Piccolo <i>et al.</i> | (Mark I Collab.) |
| RAPIDIS | 77 | PRL 39 526 | P.A. Rapidis <i>et al.</i> | (Mark I Collab.) |
| PERUZZI | 76 | PRL 37 569 | I. Peruzzi <i>et al.</i> | (Mark I Collab.) |

OTHER RELATED PAPERS

| | | | | |
|---------|----|-------------|----------------------------|--------------|
| RICHMAN | 95 | RMP 67 893 | J.D. Richman, P.R. Burchat | (UCSB, STAN) |
| ROSNER | 95 | CNPP 21 369 | J. Rosner | (CHIC) |