

Average Hadron Multiplicities in Hadronic e^+e^- Annihilation Events

Table 38.1: Average hadron multiplicities per hadronic e^+e^- annihilation event at $\sqrt{s} = 10$, 29–35, 91, and 130–200 GeV. The rates given include decay products from resonances with $c\tau < 10$ cm, and include the corresponding anti-particle state. Correlations of the systematic uncertainties were considered for the calculation of the averages. (Updated July 2001 by O. Biebel.)

Particle	$\sqrt{s} = 10$ GeV	$\sqrt{s} = 29\text{--}35$ GeV	$\sqrt{s} = 91$ GeV	$\sqrt{s} = 130\text{--}200$ GeV
Pseudoscalar mesons:				
π^+	6.6 ± 0.2	10.3 ± 0.4	16.99 ± 0.27	21.24 ± 0.39
π^0	3.2 ± 0.3	5.83 ± 0.28	9.42 ± 0.32	
K^+	0.90 ± 0.04	1.48 ± 0.09	2.242 ± 0.063	2.81 ± 0.19
K^0	0.91 ± 0.05	1.48 ± 0.07	2.049 ± 0.026	2.10 ± 0.12
η	0.20 ± 0.04	0.61 ± 0.07	0.946 ± 0.075	
$\eta'(958)$	0.03 ± 0.01	0.26 ± 0.10	0.152 ± 0.020	
D^+	0.16 ± 0.03	0.17 ± 0.03	0.175 ± 0.016	
D^0	0.37 ± 0.06	0.45 ± 0.07	0.454 ± 0.030	
D_s^+	0.13 ± 0.02	$0.45 \pm 0.20^{(a)}$	0.131 ± 0.021	
B^+, B_d^0	—	—	$0.165 \pm 0.026^{(b)}$	
B_s^0	—	—	$0.057 \pm 0.013^{(b)}$	
Scalar mesons:				
$f_0(980)$	0.024 ± 0.006	$0.05 \pm 0.02^{(c)}$	0.146 ± 0.012	
$a_0(980)^\pm$	—	—	$0.27 \pm 0.11^{(d)}$	
Vector mesons:				
$\rho(770)^0$	0.35 ± 0.04	0.81 ± 0.08	1.231 ± 0.098	
$\rho(770)^\pm$	—	—	$2.40 \pm 0.43^{(d)}$	
$\omega(782)$	0.30 ± 0.08	—	1.08 ± 0.12	
$K^*(892)^+$	0.27 ± 0.03	0.64 ± 0.05	0.715 ± 0.059	
$K^*(892)^0$	0.29 ± 0.03	0.56 ± 0.06	0.738 ± 0.024	
$\phi(1020)$	0.044 ± 0.003	0.085 ± 0.011	0.0963 ± 0.0032	
$D^*(2010)^+$	0.22 ± 0.04	0.43 ± 0.07	$0.1973 \pm 0.0057^{(e)}$	
$D^*(2007)^0$	0.23 ± 0.06	0.27 ± 0.11	—	
$D_s^*(2112)^+$	—	—	$0.101 \pm 0.048^{(f)}$	
$B^*(g)$	—	—	0.288 ± 0.026	
$J/\psi(1S)$	—	—	$0.0052 \pm 0.0004^{(h)}$	
$\psi(2S)$	—	—	$0.0023 \pm 0.0004^{(h)}$	
$\Upsilon(1S)$	—	—	$0.00014 \pm 0.00007^{(h)}$	
Pseudovector mesons:				
$\chi_{c1}(3510)$	—	—	$0.0041 \pm 0.0011^{(h)}$	
Tensor mesons:				
$f_2(1270)$	0.09 ± 0.02	0.14 ± 0.04	0.166 ± 0.020	
$f_2'(1525)$	—	—	0.012 ± 0.006	
$K_2^*(1430)^+$	—	0.09 ± 0.03	—	
$K_2^*(1430)^0$	—	0.12 ± 0.06	$0.084 \pm 0.022^{(h)}$	
$B^{**}(i)$	—	—	0.118 ± 0.024	
Baryons:				
p	0.253 ± 0.016	0.640 ± 0.050	1.048 ± 0.045	1.41 ± 0.18
Λ	0.080 ± 0.007	0.205 ± 0.010	0.3915 ± 0.0065	0.39 ± 0.03
Σ^0	0.023 ± 0.008	—	0.076 ± 0.011	
Σ^-	—	—	0.081 ± 0.010	
Σ^+	—	—	0.107 ± 0.011	
Σ^\pm	—	—	0.174 ± 0.009	
Ξ^-	0.0059 ± 0.0007	0.0176 ± 0.0027	0.0258 ± 0.0010	
$\Delta(1232)^{++}$	0.040 ± 0.010	—	0.085 ± 0.014	
$\Sigma(1385)^-$	0.006 ± 0.002	0.017 ± 0.004	0.0240 ± 0.0017	
$\Sigma(1385)^+$	0.005 ± 0.001	0.017 ± 0.004	0.0239 ± 0.0015	
$\Sigma(1385)^\pm$	0.0106 ± 0.0020	0.033 ± 0.008	0.0462 ± 0.0028	
$\Xi(1530)^0$	0.0015 ± 0.0006	—	0.0055 ± 0.0005	
Ω^-	0.0007 ± 0.0004	0.014 ± 0.007	0.0016 ± 0.0003	
Λ_c^+	$0.100 \pm 0.030^{(j)}$	0.110 ± 0.050	0.078 ± 0.017	
Λ_b^0	—	—	0.031 ± 0.016	
$\Sigma_c^{++}, \Sigma_c^0$	0.014 ± 0.007	—	—	
$\Lambda(1520)$	0.008 ± 0.002	—	0.0222 ± 0.0027	

Notes for Table 38.1:

- (a) $B(D_s \rightarrow \eta\pi, \eta'\pi)$ was used (RPP1994).
- (b) The Standard Model $B(Z \rightarrow b\bar{b}) = 0.217$ was used.
- (c) $x_p = p/p_{\text{beam}} > 0.1$ only.
- (d) Both charge states.
- (e) $B(D^*(2010)^+ \rightarrow D^0\pi^+) \times B(D^0 \rightarrow K^-\pi^+)$ has been used (RPP2000).
- (f) $B(D_s^* \rightarrow D_s^+\gamma)$, $B(D_s^+ \rightarrow \phi\pi^+)$, $B(\phi \rightarrow K^+K^-)$ have been used (RPP1998).
- (g) Any charge state (*i.e.*, B_d^* , B_u^* , or B_s^*).
- (h) $B(Z \rightarrow \text{hadrons}) = 0.699$ was used (RPP1994).
- (i) Any charge state (*i.e.*, B_d^{**} , B_u^{**} , or B_s^{**}).
- (j) The value was derived from the cross section of $A_c^+ \rightarrow p\pi K$, assuming the branching fraction to be $(3.2 \pm 0.7)\%$ (RPP1992).

References for Table 38.1:

- RPP1992**: Phys. Rev. **D45** (1992) and references therein.
- RPP1994**: Phys. Rev. **D50**, 1173 (1994) and references therein.
- RPP1996**: Phys. Rev. **D54**, 1 (1996) and references therein.
- RPP1998**: Eur. Phys. J. **C3**, 1 (1998) and references therein.
- RPP2000**: Eur. Phys. J. **C15**, 1 (2000) and references therein.
- R. Marshall, Rep. Prog. Phys. **52**, 1329 (1989)
- A. De Angelis, J. Phys. **G19**, 1233 (1993) and references therein.
- ALEPH**: D. Buskulic *et al.*: Phys. Lett. **B295**, 396 (1992); Z. Phys. **C64**, 361 (1994); **C69**, 15 (1996); **C69**, 379 (1996); **C73**, 409 (1997); and
R. Barate *et al.*: Z. Phys. **C74**, 451 (1997); Phys. Reports **294**, 1 (1998); Eur. Phys. J. **C5**, 205 (1998); **C16**, 597 (2000); **C16**, 613 (2000).
- ARGUS**: H. Albrecht *et al.*: Phys. Lett. **230B**, 169 (1989); Z. Phys. **C44**, 547 (1989); **C46**, 15 (1990); **C54**, 1 (1992); **C58**, 199 (1993); **C61**, 1 (1994); Phys. Rep. **276**, 223 (1996).
- CELLO**: H.J. Behrend *et al.*: Z. Phys. **C46**, 397 (1990); **C47**, 1 (1990).
- CLEO**: D. Bortoletto *et al.*, Phys. Rev. **D37**, 1719 (1988).
- Crystal Ball**: Ch. Bieler *et al.*, Z. Phys. **C49**, 225 (1991).
- DELPHI**: P. Abreu *et al.*: Z. Phys. **C57**, 181 (1993); **C59**, 533 (1993); **C61**, 407 (1994); **C65**, 587 (1995); **C67**, 543 (1995); **C68**, 353 (1995); **C73**, 61 (1996); Nucl. Phys. **B444**, 3 (1995); Phys. Lett. **B341**, 109 (1994); **B345**, 598 (1995); **B361**, 207 (1995); **B372**, 172 (1996); **B379**, 309 (1996); **B416**, 233 (1998); **B449**, 364 (1999); **B475**, 429 (2000); Eur. Phys. J. **C6**, 19 (1999); **C5**, 585 (1998); **C18**, 203 (2000).
- HRS**: S. Abachi *et al.*, Phys. Rev. Lett. **57**, 1990 (1986); and
M. Derrick *et al.*, Phys. Rev. **D35**, 2639 (1987).
- L3**: M. Acciarri *et al.*: Phys. Lett. **B328**, 223 (1994); **B345**, 589 (1995); **B371**, 126 (1996); **B371**, 137 (1996); **B393**, 465 (1997); **B404**, 390 (1997); **B407**, 351 (1997); **B407**, 389 (1997), erratum *ibid.* **B427**, 409 (1998); **B453**, 94 (1999); **B479**, 79 (2000).
- MARK II**: H. Schellman *et al.*, Phys. Rev. **D31**, 3013 (1985); and
G. Wormser *et al.*, Phys. Rev. Lett. **61**, 1057 (1988).
- JADE**: W. Bartel *et al.*, Z. Phys. **C20**, 187 (1983); and
D.D. Pietzl *et al.*, Z. Phys. **C46**, 1 (1990).
- OPAL**: R. Akers *et al.*: Z. Phys. **C63**, 181 (1994); **C66**, 555 (1995); **C67**, 389 (1995); **C68**, 1 (1995); and
G. Alexander *et al.*: Phys. Lett. **B358**, 162 (1995); Z. Phys. **C70**, 197 (1996); **C72**, 1 (1996); **C72**, 191 (1996); **C73**, 569 (1997); **C73**, 587 (1997); Phys. Lett. **B370**, 185 (1996); and
K. Ackerstaff *et al.*: Z. Phys. **C75**, 192 (1997); Phys. Lett. **B412**, 210 (1997); Eur. Phys. J. **C1**, 439 (1998); **C4**, 19 (1998); **C5**, 1 (1998); **C5**, 411 (1998); and
G. Abbiendi *et al.*: Eur. Phys. J. **C16**, 185 (2000); **C16**, 185 (2000).
- PLUTO**: Ch. Berger *et al.*, Phys. Lett. **104B**, 79 (1981).
- SLD**: K. Abe, Phys. Rev. **D59**, 052001 (1999).
- TASSO**: H. Aihara *et al.*, Z. Phys. **C27**, 27 (1985).
- TPC**: H. Aihara *et al.*, Phys. Rev. Lett. **53**, 2378 (1984).