

$f_2(2150)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called T_0 .

$f_2(2150)$ MASS

$f_2(2150)$ MASS, COMBINED MODES (MeV)

VALUE (MeV) DOCUMENT ID
2156 ± 11 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

$\eta\eta$ MODE

VALUE (MeV) DOCUMENT ID TECN COMMENT
 The data in this block is included in the average printed for a previous datablock.

2157 ± 12 OUR AVERAGE

2151 ± 16	BARBERIS	00E	450	$p\bar{p} \rightarrow p_f \eta\eta p_S$
2175 ± 20	PROKOSHKIN	95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p\bar{p} \rightarrow p\bar{p} 2\eta$
2130 ± 35	SINGOVSKI	94	GAM4	450 $p\bar{p} \rightarrow p\bar{p} 2\eta$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 1 2140 ± 30 ABELE 99B CBAR
- 2 seen ANISOVICH 99B SPEC 1.35–1.94 $\bar{p}p \rightarrow \eta\eta\pi^0$
- 3 2105 ± 10 ANISOVICH 99K RVUE 0.6–1.94 $\bar{p}p \rightarrow \eta\eta, \eta\eta'$
- 4 2104 ± 20 ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$

¹ Spin not determined.
² $J^{PC} = 0^{++}$
³ Using preliminary CBAR data. PWA gives $J^{PC} = 0^{++}$.
⁴ No J^{PC} determination.

$\eta\pi\pi$ MODE

VALUE (MeV) DOCUMENT ID TECN CHG COMMENT
 The data in this block is included in the average printed for a previous datablock.

2135 ± 20 ± 45	ADOMEIT	96	CBAR	0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
-----------------------	---------	----	------	---	---

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2226	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2090	⁵ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2120	⁶ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2170	⁷ MARTIN	80B	RVUE	
~ 2150	⁷ MARTIN	80C	RVUE	
~ 2150	⁸ DULUDE	78B	OSPK	1–2 $\bar{p}p \rightarrow \pi^0\pi^0$

⁵ OAKDEN 94 makes an amplitude analysis of LEAR data on $\bar{p}p \rightarrow \pi\pi$ using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.

⁶ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$.
⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.
⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2139^{+8}_{-9}	⁹ EVANGELISTA 97	SPEC		0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
~ 2190	¹⁰ CUTTS	78B	CNTR	0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{10,11} COUPLAND	77	CNTR	0
2193 ± 2	^{10,12} ALSPECTOR	73	CNTR	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$ $\bar{p}p$ S channel

- ⁹ Isospin 0 and 1 not separated.
- ¹⁰ Isospins 0 and 1 not separated.
- ¹¹ From a fit to the total elastic cross section.
- ¹² Referred to as T or T region by ALSPECTOR 73.


$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2130\pm35	BARBERIS	99	OMEG 450 $p\bar{p} \rightarrow p_S p_f K^+ K^-$

$f_2(2150)$ WIDTH

$f_2(2150)$ WIDTH, COMBINED MODES (MeV)

167 \pm 30 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.



$f_2(2150)$ WIDTH, COMBINED MODES (MeV)

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

250 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 226	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 70	17 OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	18 MARTIN	80B	RVUE
~ 250	18 MARTIN	80C	RVUE
~ 250	19 DULUDE	78B	OSPK 1–2 $\bar{p}p \rightarrow \pi^0\pi^0$

¹⁷ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

¹⁸ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

¹⁹ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p, \bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

56^{+31}_{-16}	20 EVANGELISTA 97	SPEC		0.6–2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
135 ± 75	21,22 COUPLAND 77	CNTR 0		0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98 ± 8	22 ALSPECTOR 73	CNTR		$\bar{p}p$ S channel

²⁰ Isospin 0 and 2 not separated.

²¹ From a fit to the total elastic cross section.

²² Isospins 0 and 1 not separated.

$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
270 ± 50	BARBERIS	99 OMEG	450 $p\bar{p} \rightarrow p_S p_f K^+ K^-$

$f_2(2150)$ DECAY MODES

Mode
$\Gamma_1 \quad \pi\pi$
$\Gamma_2 \quad \eta\eta$
$\Gamma_3 \quad K\bar{K}$
$\Gamma_4 \quad f_2(1270)\eta$
$\Gamma_5 \quad a_2(1320)\pi$

$f_2(2150)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma(\eta\eta)$	Γ_3/Γ_2			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
1.28 ± 0.23		BARBERIS	00E	450 $p\bar{p} \rightarrow p_f \eta \eta p_S$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	95	23 PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p\bar{p} \rightarrow p p 2\eta$
------	----	-------------------	------	---

²³ Using data from ARMSTRONG 89D.

