

$\rho(1700)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

A REVIEW GOES HERE – Check our WWW List of Reviews

$\rho(1700)$ MASS

$\eta\rho^0$ AND $\pi^+\pi^-$ MODES


VALUE (MeV)

DOCUMENT ID

1700±20 OUR ESTIMATE

1723±11 OUR AVERAGE

Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.3. See the ideogram below.



$\rho(1700)$ mass, $\eta\rho^0$ and $\pi^+\pi^-$ modes (MeV)

$\eta\rho^0$ MODE

VALUE (MeV)

DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.

1740±20	ANTONELLI	88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1701±15	² FUKUI	88	SPEC	$8.95\pi^-p \rightarrow \eta\pi^+\pi^-n$

$\pi\pi$ MODE

VALUE (MeV)

DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.

1780 $\begin{matrix} +37 \\ -29 \end{matrix}$	³ ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
1719 ±15	³ BERTIN	97C	OBLX	$0.0\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
1730 ±30	CLEGG	94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1768 ± 21	BISELLO	89	DM2	$e^+ e^- \rightarrow \pi^+ \pi^-$
1745.7 ± 91.9	DUBNICKA	89	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
1546 ± 26	GESHKEN...	89	RVUE	
1650	⁴ ERKAL	85	RVUE	20–70 $\gamma p \rightarrow \gamma \pi$
1550 ± 70	ABE	84B	HYBR	20 $\gamma p \rightarrow \pi^+ \pi^- p$
1590 ± 20	⁵ ASTON	80	OMEG	20–70 $\gamma p \rightarrow p 2\pi$
1600 ± 10	⁶ ATIYA	79B	SPEC	50 $\gamma C \rightarrow C 2\pi$
1598 $\begin{smallmatrix} +24 \\ -22 \end{smallmatrix}$	BECKER	79	ASPK	17 $\pi^- p$ polarized
1659 ± 25	⁴ LANG	79	RVUE	
1575	⁴ MARTIN	78C	RVUE	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1610 ± 30	⁴ FROGGATT	77	RVUE	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1590 ± 20	⁷ HYAMS	73	ASPK	17 $\pi^- p \rightarrow \pi^+ \pi^- n$

$\pi\omega$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	----------------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1550 to 1620	⁸ ACHASOV	00I	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
1580 to 1710	⁹ ACHASOV	00I	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
1710 ± 90	ACHASOV	97	RVUE	$e^+ e^- \rightarrow \omega \pi^0$

$K\bar{K}$ MODE

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	------------	----------------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1740.8 ± 22.2	27k	¹ ABELE	99D	CBAR	± 0.0 $\bar{p} p \rightarrow K^+ K^- \pi^0$
1582 ± 36	1600	CLELAND	82B	SPEC	± 50 $\pi p \rightarrow K_S^0 K^\pm p$

¹ K-matrix pole. Isospin not determined, could be $\omega(1650)$ or $\phi(1680)$.

$2(\pi^+ \pi^-)$ MODE

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	----------------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1851 $\begin{smallmatrix} +27 \\ -24 \end{smallmatrix}$		ACHASOV	97	RVUE	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1570 ± 20		¹⁰ CORDIER	82	DM1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1520 ± 30		⁵ ASTON	81E	OMEG	20–70 $\gamma p \rightarrow p 4\pi$
1654 ± 25		¹¹ DIBIANCA	81	DBC	$\pi^+ d \rightarrow p p 2(\pi^+ \pi^-)$
1666 ± 39		¹⁰ BACCI	80	FRAG	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1780	34	KILLIAN	80	SPEC	11 $e^- p \rightarrow 2(\pi^+ \pi^-)$
1500		¹² ATIYA	79B	SPEC	50 $\gamma C \rightarrow C 4\pi^\pm$
1570 ± 60	65	¹³ ALEXANDER	75	HBC	7.5 $\gamma p \rightarrow p 4\pi$
1550 ± 60		⁵ CONVERSI	74	OSPK	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1550 ± 50	160	SCHACHT	74	STRC	5.5–9 $\gamma p \rightarrow p 4\pi$
1450 ± 100	340	SCHACHT	74	STRC	9–18 $\gamma p \rightarrow p 4\pi$
1430 ± 50	400	BINGHAM	72B	HBC	9.3 $\gamma p \rightarrow p 4\pi$

$\pi^+\pi^-\pi^0\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1660 ± 30	ATKINSON	85B	OMEG $20-70 \gamma p$

$3(\pi^+\pi^-)$ AND $2(\pi^+\pi^-\pi^0)$ MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1783 ± 15	CLEGG	90	RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

² Assuming $\rho^+ f_0(1370)$ decay mode interferes with $a_1(1260)^+ \pi$ background. From a two Breit-Wigner fit.

³ T-matrix pole.

⁴ From phase shift analysis of HYAMS 73 data.

⁵ Simple relativistic Breit-Wigner fit with constant width.

⁶ An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

⁷ Included in BECKER 79 analysis.

⁸ Taking into account both $\rho(1450)$ and $\rho(1700)$ contributions. Using the data of ACHASOV 00i on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^-\nu_\tau$. $\rho(1450)$ mass and width fixed at 1400 MeV and 500 MeV respectively.


⁹ Taking into account the $\rho(1700)$ contribution only. Using the data of ACHASOV 00i on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^-\nu_\tau$.

¹⁰ Simple relativistic Breit-Wigner fit with model dependent width.

¹¹ One peak fit result.

¹² Parameters roughly estimated, not from a fit.

¹³ Skew mass distribution compensated by Ross-Stodolsky factor.

$\rho(1700)$ WIDTH **$\eta\rho^0$ AND $\pi^+\pi^-$ MODES**VALUE (MeV)DOCUMENT ID**240±60 OUR ESTIMATE****240±40 OUR AVERAGE** Includes data from the 2 datablocks that follow this one. Error includes scale factor of 2.0. See the ideogram below. $\rho(1700)$ width, $\eta\rho^0$ and $\pi^+\pi^-$ modes (MeV) **$\eta\rho^0$ MODE**VALUE (MeV)DOCUMENT IDTECNCOMMENT

The data in this block is included in the average printed for a previous datablock.

150±30

16 ANTONELLI 88 DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$

282±44

15 FUKUI 88 SPEC 8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$ **$\pi\pi$ MODE**VALUE (MeV)DOCUMENT IDTECNCOMMENT

The data in this block is included in the average printed for a previous datablock.

275 ± 45

16 ABELE 97 CBAR $\bar{p}n \rightarrow \pi^-\pi^0\pi^0$

310 ± 40

16 BERTIN 97C OBLX 0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$

400 ± 100

CLEGG 94 RVUE $e^+e^- \rightarrow \pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

224 ± 22		BISELLO	89	DM2	$e^+ e^- \rightarrow \pi^+ \pi^-$
242.5 ± 163.0		DUBNICKA	89	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
620 ± 60		GESHKEN...	89	RVUE	
<315		17 ERKAL	85	RVUE	20–70 $\gamma p \rightarrow \gamma \pi$
280 + 30 – 80		ABE	84B	HYBR	20 $\gamma p \rightarrow \pi^+ \pi^- p$
230 ± 80		18 ASTON	80	OMEG	20–70 $\gamma p \rightarrow p 2\pi$
283 ± 14		19 ATIYA	79B	SPEC	50 $\gamma C \rightarrow C 2\pi$
175 + 98 – 53		BECKER	79	ASPK	17 $\pi^- p$ polarized
232 ± 34		17 LANG	79	RVUE	
340		17 MARTIN	78C	RVUE	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
300 ± 100		17 FROGGATT	77	RVUE	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
180 ± 50		20 HYAMS	73	ASPK	17 $\pi^- p \rightarrow \pi^+ \pi^- n$

$K\bar{K}$ MODE

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

187.2 ± 26.7	27k	14 ABELE	99D	CBAR	± 0.0 $\bar{p} p \rightarrow K^+ K^- \pi^0$
265 ± 120	1600	CLELAND	82B	SPEC	± 50 $\pi p \rightarrow K_S^0 K^\pm p$

¹⁴ K-matrix pole. Isospin not determined, could be $\omega(1650)$ or $\phi(1680)$.

$2(\pi^+ \pi^-)$ MODE

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 40		21 CORDIER	82	DM1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
400 ± 50		18 ASTON	81E	OMEG	20–70 $\gamma p \rightarrow p 4\pi$
400 ± 146		22 DIBIANCA	81	DBC	$\pi^+ d \rightarrow p p 2(\pi^+ \pi^-)$
700 ± 160		21 BACCI	80	FRAG	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
100	34	KILLIAN	80	SPEC	11 $e^- p \rightarrow 2(\pi^+ \pi^-)$
600		23 ATIYA	79B	SPEC	50 $\gamma C \rightarrow C 4\pi^\pm$
340 ± 160	65	24 ALEXANDER	75	HBC	7.5 $\gamma p \rightarrow p 4\pi$
360 ± 100		18 CONVERSI	74	OSPK	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
400 ± 120	160	25 SCHACHT	74	STRC	5.5–9 $\gamma p \rightarrow p 4\pi$
850 ± 200	340	25 SCHACHT	74	STRC	9–18 $\gamma p \rightarrow p 4\pi$
650 ± 100	400	BINGHAM	72B	HBC	9.3 $\gamma p \rightarrow p 4\pi$

$\pi^+ \pi^- \pi^0 \pi^0$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

300 ± 50	ATKINSON	85B	OMEG 20–70 γp
----------	----------	-----	-----------------------

$\omega \pi^0$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

350 to 580	26 ACHASOV	00I	SND $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
490 to 1040	27 ACHASOV	00I	SND $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

3($\pi^+\pi^-$) AND 2($\pi^+\pi^-\pi^0$) MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

285 ± 20	CLEGG	90	RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$
----------	-------	----	---

¹⁵ Assuming $\rho^+ f_0(1370)$ decay mode interferes with $a_1(1260)^+\pi$ background. From a two Breit-Wigner fit.

¹⁶ T-matrix pole.

¹⁷ From phase shift analysis of HYAMS 73 data.

¹⁸ Simple relativistic Breit-Wigner fit with constant width.

¹⁹ An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

²⁰ Included in BECKER 79 analysis.

²¹ Simple relativistic Breit-Wigner fit with model-dependent width.

²² One peak fit result.

²³ Parameters roughly estimated, not from a fit.

²⁴ Skew mass distribution compensated by Ross-Stodolsky factor.

²⁵ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

²⁶ Taking into account both $\rho(1450)$ and $\rho(1700)$ contributions. Using the data of ACHASOV 00I on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^-\nu_\tau$. $\rho(1450)$ mass and width fixed at 1400 MeV and 500 MeV respectively.

²⁷ Taking into account the $\rho(1700)$ contribution only. Using the data of ACHASOV 00I on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^-\nu_\tau$.

$\rho(1700)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi\pi$	dominant
Γ_2 $\rho^0\pi^+\pi^-$	large
Γ_3 $\rho^0\pi^0\pi^0$	
Γ_4 $\rho^\pm\pi^\mp\pi^0$	large
Γ_5 $2(\pi^+\pi^-)$	large
Γ_6 $\pi^+\pi^-$	seen
Γ_7 $\pi^-\pi^0$	seen
Γ_8 $K\bar{K}^*(892) + \text{c.c.}$	seen
Γ_9 $\eta\rho$	seen
Γ_{10} $a_2(1320)\pi$	not seen
Γ_{11} $K\bar{K}$	seen
Γ_{12} e^+e^-	seen
Γ_{13} $\pi^0\omega$	seen

$\rho(1700) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the cross-section into channel_l in e^+e^- annihilation.

$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_5\Gamma_{12}/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.83±0.42	BACCI	80 FRAG	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.6 ±0.2	DELCOURT	81B DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$

$\Gamma(\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_6\Gamma_{12}/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.13	²⁸ DIEKMAN	88 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
0.029 ^{+0.016} _{-0.012}	KURDADZE	83 OLYA	0.64–1.4 $e^+e^- \rightarrow \pi^+\pi^-$

²⁸ Using total width = 220 MeV.

$\Gamma(K\bar{K}^*(892)+\text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_8\Gamma_{12}/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.305±0.071	²⁹ BIZOT	80 DM1	e^+e^-

$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_9\Gamma_{12}/\Gamma$

<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7 ±3	ANTONELLI	88 DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$

$\Gamma(K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{11}\Gamma_{12}/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.035±0.029	²⁹ BIZOT	80 DM1	e^+e^-

$\Gamma(\rho\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_1\Gamma_{12}/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.510±0.090	²⁹ BIZOT	80 DM1	e^+e^-

²⁹ Model dependent.

$\rho(1700)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.287 ^{+0.043} _{-0.042}	BECKER	79	ASPK	17 $\pi^- p$ polarized
0.15 to 0.30	³⁰ MARTIN	78C	RVUE	17 $\pi^- p \rightarrow \pi^+\pi^- n$
<0.20	³¹ COSTA...	77B	RVUE	$e^+e^- \rightarrow 2\pi, 4\pi$
0.30 \pm 0.05	³⁰ FROGGATT	77	RVUE	17 $\pi^- p \rightarrow \pi^+\pi^- n$
<0.15	³² EISENBERG	73	HBC	5 $\pi^+ p \rightarrow \Delta^{++} 2\pi$
0.25 \pm 0.05	³³ HYAMS	73	ASPK	17 $\pi^- p \rightarrow \pi^+\pi^- n$

³⁰ From phase shift analysis of HYAMS 73 data.

³¹ Estimate using unitarity, time reversal invariance, Breit-Wigner.

³² Estimated using one-pion-exchange model.

³³ Included in BECKER 79 analysis.

$\Gamma(\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$ Γ_6/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13 \pm 0.05	ASTON	80	OMEG	20-70 $\gamma p \rightarrow p 2\pi$
<0.14	³⁴ DAVIER	73	STRC	6-18 $\gamma p \rightarrow p 4\pi$
<0.2	³⁵ BINGHAM	72B	HBC	9.3 $\gamma p \rightarrow p 2\pi$

³⁴ Upper limit is estimate.

³⁵ 2σ upper limit.

$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(2(\pi^+\pi^-))$ Γ_8/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.15 \pm 0.03	³⁶ DELCOURT	81B	DM1	$e^+e^- \rightarrow \bar{K} K \pi$
-----------------	------------------------	-----	-----	------------------------------------

³⁶ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

$\Gamma(\eta\rho)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.04 DONNACHIE 87B RVUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen		AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
<0.02	58	ATKINSON	86B	OMEG 20-70 γp

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AMELIN	00	VES	37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
----------	--------	----	-----	---

$\Gamma(\eta\rho)/\Gamma(2(\pi^+\pi^-))$ Γ_9/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.123 \pm 0.027	DELCOURT	82	DM1	$e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
\sim 0.1	ASTON	80	OMEG	20-70 γp

$\Gamma(\pi^+ \pi^- \text{ neutrals})/\Gamma(2(\pi^+ \pi^-))$ $(\Gamma_3+\Gamma_4+0.714\Gamma_9)/\Gamma_5$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.4 ³⁷ BALLAM 74 HBC 9.3 γp

³⁷ Upper limit. Background not subtracted.

$\Gamma(\pi^0 \omega)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ACHASOV 97 RVUE $e^+ e^- \rightarrow \omega \pi^0$

$\Gamma(K \bar{K})/\Gamma(2(\pi^+ \pi^-))$ Γ_{11}/Γ_5

VALUE CL% DOCUMENT ID TECN CHG COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.015 ± 0.010 ³⁸ DELCOURT 81B DM1 $e^+ e^- \rightarrow \bar{K} K$

<0.04 95 BINGHAM 72B HBC 0 9.3 γp

³⁸ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

$\Gamma(K \bar{K})/\Gamma(K \bar{K}^*(892) + \text{c.c.})$ Γ_{11}/Γ_8

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.052 ± 0.026 BUON 82 DM1 $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-))$ Γ_2/Γ_5

VALUE EVTS DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1.0 DELCOURT 81B DM1 $e^+ e^- \rightarrow 2(\pi^+ \pi^-)$

0.7 ± 0.1 500 SCHACHT 74 STRC 5.5–18 $\gamma p \rightarrow p 4\pi$

0.80 ³⁹ BINGHAM 72B HBC 9.3 $\gamma p \rightarrow p 4\pi$

³⁹ The $\pi\pi$ system is in *S*-wave.

$\Gamma(\rho^0 \pi^0 \pi^0)/\Gamma(\rho^\pm \pi^\mp \pi^0)$ Γ_3/Γ_4

VALUE DOCUMENT ID TECN CHG COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10 ATKINSON 85B OMEG 20–70 γp

<0.15 ATKINSON 82 OMEG 0 20–70 $\gamma p \rightarrow p 4\pi$

$\rho(1700)$ REFERENCES

ACHASOV	00I	PL B486 29	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AMELIN	00	NP B668 83	D. Amelin <i>et al.</i>	(VES Collab.)
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ACHASOV	97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)
DUBNICKA	89	JPG 15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)
GESHKEN...	89	ZPHY 45 351	B.V. Geshkenbein	(ITEP)
ANTONELLI	88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)
DIEKMAN	88	PRPL 159 101	B. Diekmann	(BONN)
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
DONNACHIE	87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)
ATKINSON	86B	ZPHY C30 531	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	85B	ZPHY C26 499	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ERKAL	85	ZPHY C29 485	C. Erkal, M.G. Olsson	(WISC)
ABE	84B	PRL 53 751	K. Abe <i>et al.</i>	
KURDADZE	83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)
		Translated from ZETFP 37 613.		
ATKINSON	82	PL 108B 55	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
CORDIER	82	PL 109B 129	A. Cordier <i>et al.</i>	(LALO)
DELCOURT	82	PL 113B 93	B. Delcourt <i>et al.</i>	(LALO)
ASTON	81E	NP B189 15	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
DELCOURT	81B	Bonn Conf. 205	B. Delcourt	(ORSAY)
Also	82	PL 109B 129	A. Cordier <i>et al.</i>	(LALO)
DIBIANCA	81	PR D23 595	F.A. di Bianca <i>et al.</i>	(CASE, CMU)
ASTON	80	PL 92B 215	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
BACCI	80	PL 95B 139	C. Bacci <i>et al.</i>	(ROMA, FRAS)
BIZOT	80	Madison Conf. 546	J.C. Bizot <i>et al.</i>	(LALO, MONP)
KILLIAN	80	PR D21 3005	T.J. Killian <i>et al.</i>	(CORN)
ATIYA	79B	PRL 43 1691	M.S. Atiya <i>et al.</i>	(COLU, ILL, FNAL)
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)
MARTIN	78C	ANP 114 1	A.D. Martin, M.R. Pennington	(CERN)
COSTA...	77B	PL 71B 345	B. Costa de Beauregard, B. Pire, T.N. Truong	(EPOL)
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
ALEXANDER	75	PL 57B 487	G. Alexander <i>et al.</i>	(TELA)
BALLAM	74	NP B76 375	J. Ballam <i>et al.</i>	(SLAC, LBL, MPIM)
CONVERSI	74	PL 52B 493	M. Conversi <i>et al.</i>	(ROMA, FRAS)
SCHACHT	74	NP B81 205	P. Schacht <i>et al.</i>	(MPIM)
DAVIER	73	NP B58 31	M. Davier <i>et al.</i>	(SLAC)
EISENBERG	73	PL 43B 149	Y. Eisenberg <i>et al.</i>	(REHO)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
BINGHAM	72B	PL 41B 635	H.H. Bingham <i>et al.</i>	(LBL, UCB, SLAC) IGJP

OTHER RELATED PAPERS

ACHASOV	00J	PR D62 117503	N.N. Achasov, A.A. Kozhevnikov	
ANDERSON	00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ABELE	99C	PL B450 275	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
DONNACHIE	99	PR D60 114011	A. Donnachie, Yu.S. Kalashnikova	
KULZINGER	99	EPJ C7 73	G. Kulzinger <i>et al.</i>	
BELOZEROVA	98	PPN 29 63	T.S. Belozerova, V.K. Henner	
		Translated from FECAY 29 148.		
BARNES	97	PR D55 4157	T. Barnes <i>et al.</i>	(ORNL, RAL, MCHS)
CLOSE	97C	PR D56 1584	F.E. Close <i>et al.</i>	(RAL, MCHS)
URHEIM	97	NPBPS 55C 359	J. Urheim	(CLEO Collab.)
ACHASOV	96B	PAN 59 1262	N.N. Achasov, G.N. Shestakov	(NOVM)
		Translated from YAF 59 1319.		
AMSLER	93B	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
LANDSBERG	92	SJNP 55 1051	L.G. Landsberg	(SERP)
		Translated from YAF 55 1896.		

ASTON	91B	NPBPS 21 105	D. Aston <i>et al.</i>	(LASS Collab.)
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)
ACHASOV	88C	PL B209 373	N.N. Achasov, A.A. Kozhevnikov	(NOVM)
BRAU	88	PR D37 2379	J.E. Brau <i>et al.</i>	JP
CLEGG	88	ZPHY C40 313	A.B. Clegg, A. Donnachie	(MCHS, LANC)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ERKAL	86	ZPHY C31 615	C. Erkal, M.G. Olsson	(WISC)
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)
ATKINSON	84C	NP B243 1	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+) JP
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	83C	NP B229 269	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
AUGUSTIN	83	LAL 83-21	J.E. Augustin <i>et al.</i>	(LALO, PADO, FRAS)
SHAMBROOM	82	PR D26 1	W.D. Shambroom <i>et al.</i>	(HARV, EFI, ILL+)
ASTON	80C	PL 92B 211	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
BARBER	80C	ZPHY C4 169	D.P. Barber <i>et al.</i>	(DARE, LANC, SHEF)
KILLIAN	80	PR D21 3005	T.J. Killian <i>et al.</i>	(CORN)
COSME	76	PL 63B 352	G. Cosme <i>et al.</i>	(ORSAY)
FRENKIEL	72	NP B47 61	P. Frenkiel <i>et al.</i>	(CDEF, CERN)
ALVENSLEB...	71	PRL 26 273	H. Alvensleben <i>et al.</i>	(DESY, MIT) G
BRAUN	71	NP B30 213	H.M. Braun <i>et al.</i>	(STRB) G
BULOS	71	PRL 26 149	F. Bulos <i>et al.</i>	(SLAC, UMD, IBM, LBL) G
LAYSSAC	71	NC 6A 134	J. Layssac, F.M. Renard	(MONP)
