

$D_{sJ}(2573)^\pm$

$$I(J^P) = 0(?^?)$$

J^P is natural, width and decay modes consistent with 2^+ .

$D_{sJ}(2573)^\pm$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
2573.5 ± 1.7 OUR AVERAGE					
2574.5 ± 3.3 ± 1.6		ALBRECHT	96	ARG	$e^+e^- \rightarrow D^0 K^+ X$
2573.2 ^{+1.7} _{-1.6} ± 0.9	217	KUBOTA	94	CLE2 +	$e^+e^- \sim 10.5$ GeV

$D_{sJ}(2573)^\pm$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
15⁺⁵₋₄ OUR AVERAGE					
10.4 ± 8.3 ± 3.0		ALBRECHT	96	ARG	$e^+e^- \rightarrow D^0 K^+ X$
16 ⁺⁵ ₋₄ ± 3	217	KUBOTA	94	CLE2 +	$e^+e^- \sim 10.5$ GeV

$D_{sJ}(2573)^+$ DECAY MODES

$D_{sJ}(2573)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 K^+$	seen
Γ_2 $D^*(2007)^0 K^+$	not seen

$D_{sJ}(2573)^+$ BRANCHING RATIOS

$\Gamma(D^0 K^+)/\Gamma_{\text{total}}$	Γ_1/Γ				
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
seen	217	KUBOTA	94	CLE2 ±	$e^+e^- \sim 10.5$ GeV

$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^0 K^+)$	Γ_2/Γ_1				
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<0.33	90	KUBOTA	94	CLE2 +	$e^+e^- \sim 10.5$ GeV

$D_{sJ}(2573)^\pm$ REFERENCES

ALBRECHT	96	ZPHY C69 405	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KUBOTA	94	PRL 72 1972	Y. Kubota <i>et al.</i>	(CLEO Collab.)

OTHER RELATED PAPERS

SEMENOV	99	SPU 42 847	S.V. Semenov
Translated from UFN 42 937.			