

$f_0(2060)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

$f_0(2060)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 2050	¹ OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2060	² OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$

¹ From solution A of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

² From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

$f_0(2060)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 120	³ OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 50	⁴ OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$

³ From solution A of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

⁴ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

$f_0(2060)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi^+\pi^-$	seen

$f_0(2060)$ REFERENCES

KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
OAKDEN	94	NPA 574 731	M.N. Oakden, M.R. Pennington	(DURH)

OTHER RELATED PAPERS

SEMENOV	99	SPU 42 847	S.V. Semenov	
		Translated from UFN 42 937.		