

# $a_2(1660)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

OMITTED FROM SUMMARY TABLE

### $a_2(1660)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1660 ± 40</b>	ABELE	99B CBAR	1.94 $\bar{p}p \rightarrow \pi^0 \eta \eta$

### $a_2(1660)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>280 ± 70</b>	ABELE	99B CBAR	1.94 $\bar{p}p \rightarrow \pi^0 \eta \eta$

### $a_2(1660)$ DECAY MODES

<u>Mode</u>	<u>Fraction (<math>\Gamma_i/\Gamma</math>)</u>
$\Gamma_1 \quad \eta \pi$	seen

### $a_2(1660)$ REFERENCES

ABELE      99B EPJ C8 67      A. Abele *et al.*      (Crystal Barrel Collab.)

### OTHER RELATED PAPERS

BARBERIS      00H PL B488 225      D. Barberis *et al.*      (WA 102 Collab.)