D^0

$I(J^P) = \frac{1}{2}(0^-)$

D^0 MASS

The fit includes $D^\pm$, D^0, $D_s^\pm$, $D^{*\pm}$, D^{*0}, and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
1864.5± 0.5 OUR FIT Error includes scale factor of 1.1.					
1864.1± 1.0 OUR AVERAGE					
1864.6± 0.3± 1.0	641	BARLAG	90C	ACCM	π^- Cu 230 GeV
1852 ± 7	16	ADAMOVICH	87	EMUL	Photoproduction
1861 ± 4		DERRICK	84	HRS	e^+e^- 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1856 ±36	22	ADAMOVICH	84B	EMUL	Photoproduction
1847 ± 7	1	FIORINO	81	EMUL	$\gamma N \rightarrow \overline{D}^0 +$
1863.8± 0.5		[1] SCHINDLER	81	MRK2	e^+e^- 3.77 GeV
1864.7± 0.6		[1] TRILLING	81	RVUE	e^+e^- 3.77 GeV
1863.0± 2.5	238	ASTON	80E	OMEG	$\gamma p \rightarrow \overline{D}^0$
1860 ± 2	143	[2] AVERY	80	SPEC	$\gamma N \rightarrow D^{*+}$
1869 ± 4	35	[2] AVERY	80	SPEC	$\gamma N \rightarrow D^{*+}$
1854 ± 6	94	[2] ATIYA	79	SPEC	$\gamma N \rightarrow D^0\overline{D}^0$
1850 ±15	64	BALTAY	78C	HBC	$\nu N \rightarrow K^0\pi\pi$
1863 ± 3		GOLDHABER	77	MRK1	D^0, D^+ recoil spectra
1863.3± 0.9		[1] PERUZZI	77	MRK1	e^+e^- 3.77 GeV
1868 ±11		PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV
1865 ±15	234	GOLDHABER	76	MRK1	$K\pi$ and $K3\pi$

[1] PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision $J/\psi(1S)$ and $\psi(2S)$ measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted. TRILLING 81 enters the fit in the $D^\pm$ mass, and PERUZZI 77 and SCHINDLER 81 enter in the $m_{D^\pm} - m_{D^0}$, below.

[2] Error does not include possible systematic mass scale shift, estimated to be less than 5 MeV.

$m_{D^\pm} - m_{D^0}$

The fit includes $D^\pm$, D^0, $D_s^\pm$, $D^{*\pm}$, D^{*0}, and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
4.79±0.10 OUR FIT Error includes scale factor of 1.1.				
4.74±0.28 OUR AVERAGE				
4.7 ±0.3	[3] SCHINDLER	81	MRK2	e^+e^- 3.77 GeV
5.0 ±0.8	[3] PERUZZI	77	MRK1	e^+e^- 3.77 GeV

[3] See the footnote on TRILLING 81 in the D^0 and $D^\pm$ sections on the mass.

D^0 MEAN LIFE

Measurements with an error $> 0.05 \times 10^{-12}$ s are omitted from the average, and those with an error $> 0.1 \times 10^{-12}$ s or that have been superseded by later results have been removed from the Listings.

VALUE (10^{-12} s)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.4126±0.0028 OUR AVERAGE					
0.413 ±0.003 ±0.004	35k	AITALA	99E	E791	$K^-\pi^+$
$0.4085 \pm 0.0041 \,^{+0.0035}_{-0.0034}$	25k	BONVICINI	99	CLE2	$e^+e^- \approx \Upsilon(4S)$
0.413 ±0.004 ±0.003	16k	FRABETTI	94D	E687	$K^-\pi^+$, $K^-\pi^+\pi^+\pi^-$
0.424 ±0.011 ±0.007	5118	FRABETTI	91	E687	$K^-\pi^+$, $K^-\pi^+\pi^+\pi^-$
0.417 ±0.018 ±0.015	890	ALVAREZ	90	NA14	$K^-\pi^+$, $K^-\pi^+\pi^+\pi^-$
$0.388 \,^{+0.023}_{-0.021}$	641	[4] BARLAG	90C	ACCM	π^- Cu 230 GeV
0.48 ±0.04 ±0.03	776	ALBRECHT	88I	ARG	e^+e^- 10 GeV
0.422 ±0.008 ±0.010	4212	RAAB	88	E691	Photoproduction
0.42 ±0.05	90	BARLAG	87B	ACCM	K^- and π^- 200 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.34 \,^{+0.06}_{-0.05}$ ±0.03	58	AMENDOLIA	88	SPEC	Photoproduction
$0.46 \,^{+0.06}_{-0.05}$	145	AGUILAR-...	87D	HYBR	$\pi^- p$ and pp
0.50 ±0.07 ±0.04	317	CSORNA	87	CLEO	e^+e^- 10 GeV
0.61 ±0.09 ±0.03	50	ABE	86	HYBR	γp 20 GeV
$0.47 \,^{+0.09}_{-0.08}$ ±0.05	74	GLADNEY	86	MRK2	e^+e^- 29 GeV
$0.43 \,^{+0.07}_{-0.05} \,^{+0.01}_{-0.02}$	58	USHIDA	86B	EMUL	ν wideband
$0.37 \,^{+0.10}_{-0.07}$	26	BAILEY	85	SILI	π^- Be 200 GeV

[4] BARLAG 90C estimate systematic error to be negligible.

THERE WILL BE A NOTE ON Δm AND $\Delta\Gamma$ IN THE 2002 EDITION.

$\left| m_{D_1^0} - m_{D_2^0} \right|$

The D_1^0 and D_2^0 are the mass eigenstates of the D^0 meson. To calculate the following limits, we use $\Delta m = [2r/(1-r)]^{1/2}\,\hbar/4.126 \times 10^{-13}$ s, where r is the experimental D^0-$\overline{D}^0$ mixing ratio.

VALUE ($10^{10}\ \hbar$ s^{-1})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
< 7	95	[5] GODANG	00	CLE2	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

<32	90	[6,7]	AITALA	98	E791	π^- nucleus, 500 GeV
<24	90	[8]	AITALA	96C	E791	π^- nucleus, 500 GeV
<21	90	[7,9]	ANJOS	88C	E691	Photoproduction

[5] This GODANG 00 limit is inferred from the D^0-$\overline{D}^0$ mixing ratio $\Gamma(K^+\pi^-$ (via $\overline{D}^0))/\Gamma(K^-\pi^+)$ given near the end of this D^0 Listings. Decay-time information is used to distinguish DCS decays from D^0-$\overline{D}^0$ mixing. The limit allows interference between the DCS and mixing ratios, and also allows *CP* violation. The strong phase between $D^0 \to K^+\pi^-$ and $\overline{D}^0 \to K^+\pi^-$ is assumed to be small.

[6] AITALA 98 allows interference between the doubly Cabibbo-suppressed and mixing amplitudes, and also allows *CP* violation in this term.

[7] This limit is inferred from the D^0-$\overline{D}^0$ mixing ratio $\Gamma(K^+\pi^-$ or $K^+\pi^-\pi^+\pi^-$ (via $\overline{D}^0))/\Gamma(K^-\pi^+$ or $K^-\pi^+\pi^+\pi^-)$ near the end of the D^0 Listings. Decay-time information is used to distinguish doubly Cabibbo-suppressed decays from D^0-$\overline{D}^0$ mixing.

[8] This limit is inferred from the D^0-$\overline{D}^0$ mixing ratio $\Gamma(K^+\ell^-\overline{\nu}_\ell$(via $\overline{D}^0$) $)/\Gamma(K^-\ell^+\nu_\ell)$ given near the end of the D^0 Listings.

[9] ANJOS 88C assumes no interference between doubly Cabibbo-suppressed and mixing amplitudes. When interference is allowed, the limit degrades by about a factor of two.

THERE WILL BE A NOTE ON Δm AND $\Delta\Gamma$ IN THE 2002 EDITION.

$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma_{D^0} = 2y$

The D_1^0 and D_2^0 are the mass eigenstates of the D^0 meson. AITALA 99E and LINK 00 use a difference in directly measured decay rates to obtain their limits. The other experiments infer the limits here from limits on mixing, using $\Delta\Gamma/\Gamma = [8r/(1+r)]^{1/2}$, where r is the experimental D^0-$\overline{D}^0$ mixing ratio. See the footnotes to the entries below.

VALUE	*CL%*	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.0684±0.0278±0.0148		10k	[10] LINK	00	FOCS	γ nucleus

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.116 < \Delta\Gamma/\Gamma < 0.020$	95	[11] GODANG	00	CLE2	e^+e^-
$-0.08 < \Delta\Gamma/\Gamma < 0.12$	90	[12] AITALA	99E	E791	$K^-\pi^+$, K^+K^-
$\lvert\Delta\Gamma\rvert/\Gamma < 0.26$	90	[13,14] AITALA	98	E791	π^- nucleus, 500 GeV
$\lvert\Delta\Gamma\rvert/\Gamma < 0.20$	90	[15] AITALA	96C	E791	π^- nucleus, 500 GeV
$\lvert\Delta\Gamma\rvert/\Gamma < 0.17$	90	[14,16] ANJOS	88C	E691	Photoproduction

[10] LINK 00 measures the lifetime difference between $D^0 \to K^-K^+$ (*CP* even) decays and $D^0 \to K^-\pi^+$ (*CP* mixed) decays, or $y_{CP} = [\Gamma(CP+)-\Gamma(CP-)]/[\Gamma(CP+)+\Gamma(CP-)]$. We list $2y_{CP}=\Delta\Gamma/\Gamma$.

[11] This GODANG 00 limit is inferred from the D^0-$\overline{D}^0$ mixing ratio $\Gamma(K^+\pi^-$ (via $\overline{D}^0))/\Gamma(K^-\pi^+)$ given near the end of this D^0 Listings. Decay-time information is used to distinguish DCS decays from D^0-$\overline{D}^0$ mixing. The limit allows interference between the DCS and mixing ratios, and also allows *CP* violation. The phase between $D^0 \to K^+\pi^-$ and $\overline{D}^0 \to K^+\pi^-$ is assumed to be small. This is a measurement of y' and is not the same as the y_{CP} of LINK 00.

[12] AITALA 99E measures $\Delta\Gamma = 2[\Gamma(D^0 \to K^+K^-) - \Gamma(D^0 \to K^-\pi^+)] = +0.04 \pm 0.14 \pm 0.05\ \text{ps}^{-1}$ and thus gets 90%-confidence-level limits $-0.20 < \Delta\Gamma < +0.28\ \text{ps}^{-1}$.

[13] AITALA 98 allows interference between the doubly Cabibbo-suppressed and mixing amplitudes, and also allows *CP* violation in this term.

[14] This limit is inferred from the D^0-$\overline{D}^0$ mixing ratio $\Gamma(K^+\pi^-$ or $K^+\pi^-\pi^+\pi^-$ (via $\overline{D}^0$))/$\Gamma(K^-\pi^+$ or $K^-\pi^+\pi^+\pi^-$) near the end of the D^0 Listings. Decay-time information is used to distinguish doubly Cabibbo-suppressed decays from D^0-$\overline{D}^0$ mixing.

[15] This limit is inferred from the D^0-$\overline{D}^0$ mixing ratio $\Gamma(K^+\ell^-\overline{\nu}_\ell$(via $\overline{D}^0$))/$\Gamma(K^-\ell^+\nu_\ell)$ given near the end of the D^0 Listings.

[16] ANJOS 88C assumes no interference between doubly Cabibbo-suppressed and mixing amplitudes. When interference is allowed, the limit degrades by about a factor of two.

D^0 DECAY MODES

$\overline{D}^0$ modes are charge conjugates of the modes below.

	Mode		Fraction (Γ_i/Γ)	Scale factor/ Confidence level
	Inclusive modes			
Γ_1	e^+ anything		(6.75 ± 0.29) %	
Γ_2	μ^+ anything		(6.6 ± 0.8) %	
Γ_3	K^- anything		(53 ± 4) %	S=1.3
Γ_4	$\overline{K}^0$ anything + K^0 anything		(42 ± 5) %	
Γ_5	K^+ anything		(3.4 $^{+0.6}_{-0.4}$) %	
Γ_6	η anything	[a]	< 13 %	CL=90%
Γ_7	ϕ anything		(1.7 ± 0.8) %	
	Semileptonic modes			
Γ_8	$K^-\ell^+\nu_\ell$	[b]	(3.47 ± 0.17) %	S=1.3
Γ_9	$K^-e^+\nu_e$		(3.64 ± 0.18) %	
Γ_{10}	$K^-\mu^+\nu_\mu$		(3.22 ± 0.17) %	
Γ_{11}	$K^-\pi^0e^+\nu_e$		(1.6 $^{+1.3}_{-0.5}$) %	
Γ_{12}	$\overline{K}^0\pi^-e^+\nu_e$		(2.8 $^{+1.7}_{-0.9}$) %	
Γ_{13}	$\overline{K}^*(892)^-e^+\nu_e$ × B($K^{*-}\to\overline{K}^0\pi^-$)		(1.35 ± 0.22) %	
Γ_{14}	$K^*(892)^-\ell^+\nu_\ell$			
Γ_{15}	$\overline{K}^*(892)^0\pi^-e^+\nu_e$			
Γ_{16}	$K^-\pi^+\pi^-\mu^+\nu_\mu$		< 1.2 × 10^{-3}	CL=90%
Γ_{17}	$(\overline{K}^*(892)\pi)^-\mu^+\nu_\mu$		< 1.4 × 10^{-3}	CL=90%
Γ_{18}	$\pi^-e^+\nu_e$		(3.7 ± 0.6) × 10^{-3}	
	A fraction of the following resonance mode has already appeared above as a submode of a charged-particle mode.			
Γ_{19}	$K^*(892)^-e^+\nu_e$		(2.02 ± 0.33) %	

Hadronic modes with a $\overline{K}$ or $\overline{K}K\overline{K}$

Γ_{20}	$K^-\pi^+$		(3.83±0.09) %	
Γ_{21}	$\overline{K}^0\pi^0$		(2.11±0.21) %	S=1.1
Γ_{22}	$\overline{K}^0\pi^+\pi^-$	[c]	(5.4 ±0.4) %	S=1.2
Γ_{23}	$\overline{K}^0\rho^0$		(1.21±0.17) %	
Γ_{24}	$\overline{K}^0 f_0(980)$		(3.0 ±0.8) $\times 10^{-3}$	
	$\times$ B($f_0 \to \pi^+\pi^-$)			
Γ_{25}	$\overline{K}^0 f_2(1270)$		(2.4 ±0.9) $\times 10^{-3}$	
	$\times$ B($f_2 \to \pi^+\pi^-$)			
Γ_{26}	$\overline{K}^0 f_0(1370)$		(4.3 ±1.3) $\times 10^{-3}$	
	$\times$ B($f_0 \to \pi^+\pi^-$)			
Γ_{27}	$K^*(892)^-\pi^+$		(3.4 ±0.3) %	
	$\times$ B($K^{*-} \to \overline{K}^0\pi^-$)			
Γ_{28}	$K_0^*(1430)^-\pi^+$		(6.4 ±1.6) $\times 10^{-3}$	
	$\times$ B($K_0^*(1430)^- \to \overline{K}^0\pi^-$)			
Γ_{29}	$\overline{K}^0\pi^+\pi^-$ nonresonant		(1.47±0.24) %	
Γ_{30}	$K^-\pi^+\pi^0$	[c]	(13.9 ±0.9) %	S=1.3
Γ_{31}	$K^-\rho^+$		(10.8 ±1.0) %	
Γ_{32}	$K^*(892)^-\pi^+$		(1.7 ±0.2) %	
	$\times$ B($K^{*-} \to K^-\pi^0$)			
Γ_{33}	$\overline{K}^*(892)^0\pi^0$		(2.1 ±0.3) %	
	$\times$ B($\overline{K}^{*0} \to K^-\pi^+$)			
Γ_{34}	$K^-\pi^+\pi^0$ nonresonant		(6.9 ±2.5) $\times 10^{-3}$	
Γ_{35}	$\overline{K}^0\pi^0\pi^0$		—	
Γ_{36}	$\overline{K}^*(892)^0\pi^0$		(1.1 ±0.2) %	
	$\times$ B($\overline{K}^{*0} \to \overline{K}^0\pi^0$)			
Γ_{37}	$\overline{K}^0\pi^0\pi^0$ nonresonant		(7.8 ±2.0) $\times 10^{-3}$	
Γ_{38}	$K^-\pi^+\pi^+\pi^-$	[c]	(7.49±0.31) %	
Γ_{39}	$K^-\pi^+\rho^0$ total		(6.3 ±0.4) %	
Γ_{40}	$K^-\pi^+\rho^0$ 3-body		(4.7 ±2.1) $\times 10^{-3}$	
Γ_{41}	$\overline{K}^*(892)^0\rho^0$		(9.8 ±2.2) $\times 10^{-3}$	
	$\times$ B($\overline{K}^{*0} \to K^-\pi^+$)			
Γ_{42}	$K^- a_1(1260)^+$		(3.6 ±0.6) %	
	$\times$ B($a_1(1260)^+ \to \pi^+\pi^+\pi^-$)			
Γ_{43}	$\overline{K}^*(892)^0\pi^+\pi^-$ total		(1.5 ±0.4) %	
	$\times$ B($\overline{K}^{*0} \to K^-\pi^+$)			
Γ_{44}	$\overline{K}^*(892)^0\pi^+\pi^-$ 3-body		(9.5 ±2.1) $\times 10^{-3}$	
	$\times$ B($\overline{K}^{*0} \to K^-\pi^+$)			
Γ_{45}	$K_1(1270)^-\pi^+$	[d]	(3.6 ±1.0) $\times 10^{-3}$	
	$\times$ B($K_1(1270)^- \to K^-\pi^+\pi^-$)			
Γ_{46}	$K^-\pi^+\pi^+\pi^-$ nonresonant		(1.74±0.25) %	
Γ_{47}	$\overline{K}^0\pi^+\pi^-\pi^0$	[c]	(10.0 ±1.2) %	

Γ_{48}	$\overline{K}^0\eta \times$ B($\eta \to \pi^+\pi^-\pi^0$)	(1.6 ±0.3) $\times 10^{-3}$	
Γ_{49}	$\overline{K}^0\omega \times$ B($\omega \to \pi^+\pi^-\pi^0$)	(1.9 ±0.4) %	
Γ_{50}	$K^*(892)^-\rho^+$	(4.1 ±1.6) %	
	$\times$ B($K^{*-} \to \overline{K}^0\pi^-$)		
Γ_{51}	$\overline{K}^*(892)^0\rho^0$	(4.9 ±1.1) $\times 10^{-3}$	
	$\times$ B($\overline{K}^{*0} \to \overline{K}^0\pi^0$)		
Γ_{52}	$K_1(1270)^-\pi^+$ [d]	(5.1 ±1.4) $\times 10^{-3}$	
	$\times$ B($K_1(1270)^- \to \overline{K}^0\pi^-\pi^0$)		
Γ_{53}	$\overline{K}^*(892)^0\pi^+\pi^-$ 3-body	(4.8 ±1.1) $\times 10^{-3}$	
	$\times$ B($\overline{K}^{*0} \to \overline{K}^0\pi^0$)		
Γ_{54}	$\overline{K}^0\pi^+\pi^-\pi^0$ nonresonant	(2.1 ±2.1) %	
Γ_{55}	$K^-\pi^+\pi^0\pi^0$	—	
Γ_{56}	$K^-\pi^+\pi^+\pi^-\pi^0$	(4.0 ±0.4) %	
Γ_{57}	$\overline{K}^*(892)^0\pi^+\pi^-\pi^0$	(1.2 ±0.6) %	
	$\times$ B($\overline{K}^{*0} \to K^-\pi^+$)		
Γ_{58}	$\overline{K}^*(892)^0\eta$	(2.9 ±0.8) $\times 10^{-3}$	
	$\times$ B($\overline{K}^{*0} \to K^-\pi^+$)		
	$\times$ B($\eta \to \pi^+\pi^-\pi^0$)		
Γ_{59}	$K^-\pi^+\omega \times$ B($\omega \to \pi^+\pi^-\pi^0$)	(2.7 ±0.5) %	
Γ_{60}	$\overline{K}^*(892)^0\omega$	(7 ±3) $\times 10^{-3}$	
	$\times$ B($\overline{K}^{*0} \to K^-\pi^+$)		
	$\times$ B($\omega \to \pi^+\pi^-\pi^0$)		
Γ_{61}	$\overline{K}^0\pi^+\pi^+\pi^-\pi^-$	(5.8 ±1.6) $\times 10^{-3}$	
Γ_{62}	$\overline{K}^0\pi^+\pi^-\pi^0\pi^0(\pi^0)$	—	
Γ_{63}	$\overline{K}^0K^+K^-$	(9.4 ±1.0) $\times 10^{-3}$	
	In the fit as $\frac{1}{2}\Gamma_{75} + \Gamma_{65}$, where $\frac{1}{2}\Gamma_{75} = \Gamma_{64}$.		
Γ_{64}	$\overline{K}^0\phi \times$ B($\phi \to K^+K^-$)	(4.3 ±0.5) $\times 10^{-3}$	
Γ_{65}	$\overline{K}^0K^+K^-$ non-ϕ	(5.1 ±0.8) $\times 10^{-3}$	
Γ_{66}	$K^0_SK^0_SK^0_S$	(8.3 ±1.5) $\times 10^{-4}$	
Γ_{67}	$K^+K^-K^-\pi^+$	(2.1 ±0.5) $\times 10^{-4}$	
Γ_{68}	$K^+K^-\overline{K}^0\pi^0$	(7.2 $^{+4.8}_{-3.5}$) $\times 10^{-3}$	

Fractions of many of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. (Modes for which there are only upper limits and $\overline{K}^*(892)\rho$ submodes only appear below.)

Γ_{69}	$\overline{K}^0\eta$	(7.0 ±1.0) $\times 10^{-3}$	
Γ_{70}	$\overline{K}^0\rho^0$	(1.21±0.17) %	
Γ_{71}	$K^-\rho^+$	(10.8 ±0.9) %	S=1.2
Γ_{72}	$\overline{K}^0\omega$	(2.1 ±0.4) %	
Γ_{73}	$\overline{K}^0\eta'(958)$	(1.71±0.26) %	
Γ_{74}	$\overline{K}^0f_0(980)$	(5.7 ±1.6) $\times 10^{-3}$	
Γ_{75}	$\overline{K}^0\phi$	(8.6 ±1.0) $\times 10^{-3}$	
Γ_{76}	$K^-a_1(1260)^+$	(7.3 ±1.1) %	

Γ_{77}	$\overline{K}^0 a_1(1260)^0$		$<$ 1.9	%	CL=90%
Γ_{78}	$\overline{K}^0 f_2(1270)$		(4.1 $\pm$1.5) $\times 10^{-3}$		
Γ_{79}	$K^- a_2(1320)^+$		$<$ 2	$\times 10^{-3}$	CL=90%
Γ_{80}	$\overline{K}^0 f_0(1370)$		(6.9 $\pm$2.1) $\times 10^{-3}$		
Γ_{81}	$K^*(892)^- \pi^+$		(5.0 $\pm$0.4) %		S=1.2
Γ_{82}	$\overline{K}^*(892)^0 \pi^0$		(3.1 $\pm$0.4) %		
Γ_{83}	$\overline{K}^*(892)^0 \pi^+ \pi^-$ total		(2.2 $\pm$0.5) %		
Γ_{84}	$\overline{K}^*(892)^0 \pi^+ \pi^-$ 3-body		(1.42$\pm$0.32) %		
Γ_{85}	$K^- \pi^+ \rho^0$ total		(6.3 $\pm$0.4) %		
Γ_{86}	$K^- \pi^+ \rho^0$ 3-body		(4.7 $\pm$2.1) $\times 10^{-3}$		
Γ_{87}	$\overline{K}^*(892)^0 \rho^0$		(1.46$\pm$0.32) %		
Γ_{88}	$\overline{K}^*(892)^0 \rho^0$ transverse		(1.5 $\pm$0.5) %		
Γ_{89}	$\overline{K}^*(892)^0 \rho^0$ S-wave		(2.8 $\pm$0.6) %		
Γ_{90}	$\overline{K}^*(892)^0 \rho^0$ S-wave long.		$<$ 3	$\times 10^{-3}$	CL=90%
Γ_{91}	$\overline{K}^*(892)^0 \rho^0$ P-wave		$<$ 3	$\times 10^{-3}$	CL=90%
Γ_{92}	$\overline{K}^*(892)^0 \rho^0$ D-wave		(1.9 $\pm$0.6) %		
Γ_{93}	$K^*(892)^- \rho^+$		(6.1 $\pm$2.4) %		
Γ_{94}	$K^*(892)^- \rho^+$ longitudinal		(2.9 $\pm$1.2) %		
Γ_{95}	$K^*(892)^- \rho^+$ transverse		(3.2 $\pm$1.8) %		
Γ_{96}	$K^*(892)^- \rho^+$ P-wave		$<$ 1.5	%	CL=90%
Γ_{97}	$K^- \pi^+ f_0(980)$		$<$ 1.1	%	CL=90%
Γ_{98}	$\overline{K}^*(892)^0 f_0(980)$		$<$ 7	$\times 10^{-3}$	CL=90%
Γ_{99}	$K_1(1270)^- \pi^+$	[d]	(1.06$\pm$0.29) %		
Γ_{100}	$K_1(1400)^- \pi^+$		$<$ 1.2	%	CL=90%
Γ_{101}	$\overline{K}_1(1400)^0 \pi^0$		$<$ 3.7	%	CL=90%
Γ_{102}	$K^*(1410)^- \pi^+$		$<$ 1.2	%	CL=90%
Γ_{103}	$K_0^*(1430)^- \pi^+$		(1.04$\pm$0.26) %		
Γ_{104}	$K_2^*(1430)^- \pi^+$		$<$ 8	$\times 10^{-3}$	CL=90%
Γ_{105}	$\overline{K}_2^*(1430)^0 \pi^0$		$<$ 4	$\times 10^{-3}$	CL=90%
Γ_{106}	$\overline{K}^*(892)^0 \pi^+ \pi^- \pi^0$		(1.8 $\pm$0.9) %		
Γ_{107}	$\overline{K}^*(892)^0 \eta$		(1.9 $\pm$0.5) %		
Γ_{108}	$K^- \pi^+ \omega$		(3.0 $\pm$0.6) %		
Γ_{109}	$\overline{K}^*(892)^0 \omega$		(1.1 $\pm$0.4) %		
Γ_{110}	$K^- \pi^+ \eta'(958)$		(7.0 $\pm$1.8) $\times 10^{-3}$		
Γ_{111}	$\overline{K}^*(892)^0 \eta'(958)$		$<$ 1.0	$\times 10^{-3}$	CL=90%

Pionic modes

Γ_{112}	$\pi^+ \pi^-$	(1.52$\pm$0.09) $\times 10^{-3}$		
Γ_{113}	$\pi^0 \pi^0$	(8.4 $\pm$2.2) $\times 10^{-4}$		
Γ_{114}	$\pi^+ \pi^- \pi^0$	(1.1 $\pm$0.4) %		
Γ_{115}	$\pi^+ \pi^+ \pi^- \pi^-$	(7.3 $\pm$0.5) $\times 10^{-3}$		
Γ_{116}	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	(1.9 $\pm$0.4) %		
Γ_{117}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-$	(4.0 $\pm$3.0) $\times 10^{-4}$		

Hadronic modes with a $K\overline{K}$ pair

Γ_{118}	K^+K^-		$(\ 4.25\pm0.16)\times10^{-3}$		
Γ_{119}	$K^0\overline{K}^0$		$(\ 6.5\ \pm1.8\ )\times10^{-4}$		S=1.2
Γ_{120}	$K^0K^-\pi^+$		$(\ 6.4\ \pm1.0\ )\times10^{-3}$		S=1.1
Γ_{121}	$\overline{K}^*(892)^0K^0$		$<\ 1.1$	$\times10^{-3}$	CL=90%
	$\times$ B($\overline{K}^{*0}\to\ K^-\pi^+$)				
Γ_{122}	$K^*(892)^+K^-$		$(\ 2.3\ \pm0.5\ )\times10^{-3}$		
	$\times$ B($K^{*+}\to\ K^0\pi^+$)				
Γ_{123}	$K^0K^-\pi^+$ nonresonant		$(\ 2.3\ \pm2.3\ )\times10^{-3}$		
Γ_{124}	$\overline{K}^0K^+\pi^-$		$(\ 5.0\ \pm1.0\ )\times10^{-3}$		
Γ_{125}	$K^*(892)^0\overline{K}^0$		$<\ 5$	$\times10^{-4}$	CL=90%
	$\times$ B($K^{*0}\to\ K^+\pi^-$)				
Γ_{126}	$K^*(892)^-K^+$		$(\ 1.2\ \pm0.7\ )\times10^{-3}$		
	$\times$ B($K^{*-}\to\ \overline{K}^0\pi^-$)				
Γ_{127}	$\overline{K}^0K^+\pi^-$ nonresonant		$(\ 3.8\ ^{+2.3}_{-1.9}\ )\times10^{-3}$		
Γ_{128}	$K^+K^-\pi^0$		$(\ 1.3\ \pm0.4\ )\times10^{-3}$		
Γ_{129}	$K^0_SK^0_S\pi^0$		$<\ 5.9$	$\times10^{-4}$	
Γ_{130}	$K^+K^-\pi^+\pi^-$	[e]	$(\ 2.50\pm0.23)\times10^{-3}$		
Γ_{131}	$\phi\pi^+\pi^-\times$ B($\phi\to\ K^+K^-$)		$(\ 5.3\ \pm1.4\ )\times10^{-4}$		
Γ_{132}	$\phi\rho^0\times$ B($\phi\to\ K^+K^-$)		$(\ 3.0\ \pm1.6\ )\times10^{-4}$		
Γ_{133}	$K^+K^-\rho^0$ 3-body		$(\ 9.0\ \pm2.3\ )\times10^{-4}$		
Γ_{134}	$K^*(892)^0K^-\pi^+$ +c.c.	[f]	$<\ 5$	$\times10^{-4}$	
	$\times$ B($K^{*0}\to\ K^+\pi^-$)				
Γ_{135}	$K^*(892)^0\overline{K}^*(892)^0$		$(\ 6\ \ \pm2\ \ )\times10^{-4}$		
	$\times$ B^2($K^{*0}\to\ K^+\pi^-$)				
Γ_{136}	$K^+K^-\pi^+\pi^-$ non-ϕ		—		
Γ_{137}	$K^+K^-\pi^+\pi^-$ nonresonant		$<\ 8$	$\times10^{-4}$	CL=90%
Γ_{138}	$K^0\overline{K}^0\pi^+\pi^-$		$(\ 6.8\ \pm2.7\ )\times10^{-3}$		
Γ_{139}	$K^+K^-\pi^+\pi^-\pi^0$		$(\ 3.1\ \pm2.0\ )\times10^{-3}$		

Fractions of most of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

Γ_{140}	$\overline{K}^*(892)^0K^0$		$<\ 1.6$	$\times10^{-3}$	CL=90%
Γ_{141}	$K^*(892)^+K^-$		$(\ 3.5\ \pm0.8\ )\times10^{-3}$		
Γ_{142}	$K^*(892)^0\overline{K}^0$		$<\ 8$	$\times10^{-4}$	CL=90%
Γ_{143}	$K^*(892)^-K^+$		$(\ 1.8\ \pm1.0\ )\times10^{-3}$		
Γ_{144}	$\phi\pi^0$		$<\ 1.4$	$\times10^{-3}$	CL=90%
Γ_{145}	$\phi\eta$		$<\ 2.8$	$\times10^{-3}$	CL=90%
Γ_{146}	$\phi\omega$		$<\ 2.1$	$\times10^{-3}$	CL=90%

Γ_{147}	$\phi\pi^+\pi^-$			$(\ 1.07\pm0.28)\times10^{-3}$		
Γ_{148}	$\phi\rho^0$			$(\ 6\ \ \pm3\ \ )\times10^{-4}$		
Γ_{149}	$\phi\pi^+\pi^-$ 3-body			$(\ 7\ \ \pm5\ \ )\times10^{-4}$		
Γ_{150}	$K^*(892)^0K^-\pi^+$ + c.c.		[f]	$<\ 7$	$\times10^{-4}$	CL=90%
Γ_{151}	$K^*(892)^0K^-\pi^+$					
Γ_{152}	$\overline{K}^*(892)^0K^+\pi^-$					
Γ_{153}	$K^*(892)^0\overline{K}^*(892)^0$			$(\ 1.4\ \pm0.5\ )\times10^{-3}$		

Radiative modes

Γ_{154}	$\rho^0\gamma$			$<\ 2.4$	$\times10^{-4}$	CL=90%
Γ_{155}	$\omega\gamma$			$<\ 2.4$	$\times10^{-4}$	CL=90%
Γ_{156}	$\phi\gamma$			$<\ 1.9$	$\times10^{-4}$	CL=90%
Γ_{157}	$\overline{K}^*(892)^0\gamma$			$<\ 7.6$	$\times10^{-4}$	CL=90%

Doubly Cabibbo suppressed (*DC*) modes, $\Delta C=2$ forbidden via mixing (*C2M*) modes, $\Delta C=1$ weak neutral current (*C1*) modes, or Lepton Family number (*LF*) violating modes

Γ_{158}	$K^+\ell^-\overline{\nu}_\ell$(via $\overline{D}^0$)	*C2M*		$<\ 1.7$	$\times10^{-4}$	CL=90%
Γ_{159}	$K^+\pi^-$	*DC*		$(\ 1.46\pm0.30)\times10^{-4}$		
Γ_{160}	$K^+\pi^-$(via $\overline{D}^0$)	*C2M*		$<\ 1.6$	$\times10^{-5}$	CL=95%
Γ_{161}	$K^+\pi^-\pi^+\pi^-$	*DC*		$(\ 1.9\ \pm2.6\ )\times10^{-4}$		
Γ_{162}	$K^+\pi^-\pi^+\pi^-$(via $\overline{D}^0$)	*C2M*		$<\ 4$	$\times10^{-4}$	CL=90%
Γ_{163}	$K^+\pi^-$ or $K^+\pi^-\pi^+\pi^-$(via $\overline{D}^0$)			$<\ 1.0$	$\times10^{-3}$	CL=90%
Γ_{164}	μ^- anything (via $\overline{D}^0$)	*C2M*		$<\ 4$	$\times10^{-4}$	CL=90%
Γ_{165}	e^+e^-	*C1*		$<\ 6.2$	$\times10^{-6}$	CL=90%
Γ_{166}	$\mu^+\mu^-$	*C1*		$<\ 4.1$	$\times10^{-6}$	CL=90%
Γ_{167}	$\pi^0e^+e^-$	*C1*		$<\ 4.5$	$\times10^{-5}$	CL=90%
Γ_{168}	$\pi^0\mu^+\mu^-$	*C1*		$<\ 1.8$	$\times10^{-4}$	CL=90%
Γ_{169}	ηe^+e^-	*C1*		$<\ 1.1$	$\times10^{-4}$	CL=90%
Γ_{170}	$\eta\mu^+\mu^-$	*C1*		$<\ 5.3$	$\times10^{-4}$	CL=90%
Γ_{171}	$\rho^0e^+e^-$	*C1*		$<\ 1.0$	$\times10^{-4}$	CL=90%
Γ_{172}	$\rho^0\mu^+\mu^-$	*C1*		$<\ 2.3$	$\times10^{-4}$	CL=90%
Γ_{173}	ωe^+e^-	*C1*		$<\ 1.8$	$\times10^{-4}$	CL=90%
Γ_{174}	$\omega\mu^+\mu^-$	*C1*		$<\ 8.3$	$\times10^{-4}$	CL=90%
Γ_{175}	ϕe^+e^-	*C1*		$<\ 5.2$	$\times10^{-5}$	CL=90%
Γ_{176}	$\phi\mu^+\mu^-$	*C1*		$<\ 4.1$	$\times10^{-4}$	CL=90%
Γ_{177}	$\overline{K}^0e^+e^-$		[g]	$<\ 1.1$	$\times10^{-4}$	CL=90%
Γ_{178}	$\overline{K}^0\mu^+\mu^-$		[g]	$<\ 2.6$	$\times10^{-4}$	CL=90%
Γ_{179}	$\overline{K}^*(892)^0e^+e^-$		[g]	$<\ 1.4$	$\times10^{-4}$	CL=90%
Γ_{180}	$\overline{K}^*(892)^0\mu^+\mu^-$		[g]	$<\ 1.18$	$\times10^{-3}$	CL=90%

Γ_{181}	$\pi^+\pi^-\pi^0\mu^+\mu^-$	*C1*	$<$	8.1	$\times 10^{-4}$	CL=90%
Γ_{182}	$\mu^\pm e^\mp$	*LF*	[*h*] $<$	8.1	$\times 10^{-6}$	CL=90%
Γ_{183}	$\pi^0 e^\pm \mu^\mp$	*LF*	[*h*] $<$	8.6	$\times 10^{-5}$	CL=90%
Γ_{184}	$\eta e^\pm \mu^\mp$	*LF*	[*h*] $<$	1.0	$\times 10^{-4}$	CL=90%
Γ_{185}	$\rho^0 e^\pm \mu^\mp$	*LF*	[*h*] $<$	4.9	$\times 10^{-5}$	CL=90%
Γ_{186}	$\omega e^\pm \mu^\mp$	*LF*	[*h*] $<$	1.2	$\times 10^{-4}$	CL=90%
Γ_{187}	$\phi e^\pm \mu^\mp$	*LF*	[*h*] $<$	3.4	$\times 10^{-5}$	CL=90%
Γ_{188}	$\overline{K}^0 e^\pm \mu^\mp$	*LF*	[*h*] $<$	1.0	$\times 10^{-4}$	CL=90%
Γ_{189}	$\overline{K}^*(892)^0 e^\pm \mu^\mp$	*LF*	[*h*] $<$	1.0	$\times 10^{-4}$	CL=90%

Γ_{190} A dummy mode used by the fit. (17.2 $\pm$3.4) % S=1.1

[*a*] This is a weighted average of $D^\pm$ (44%) and D^0 (56%) branching fractions. See "D^+ and $D^0 \to$ (η anything) / (total D^+ and D^0)" under "D^+ Branching Ratios" in these Particle Listings.

[*b*] This value averages the e^+ and μ^+ branching fractions, after making a small phase-space adjustment to the μ^+ fraction to be able to use it as an e^+ fraction; hence our ℓ^+ here is really an e^+.

[*c*] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.

[*d*] The two experiments measuring this fraction are in serious disagreement. See the Particle Listings.

[*e*] The experiments on the division of this charge mode amongst its submodes disagree, and the submode branching fractions here add up to considerably more than the charged-mode fraction.

[*f*] However, these upper limits are in serious disagreement with values obtained in another experiment.

[*g*] This mode is not a useful test for a ΔC=1 weak neutral current because both quarks must change flavor in this decay.

[*h*] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 51 branching ratios uses 122 measurements and one constraint to determine 28 parameters. The overall fit has a $\chi^2 = 64.5$ for 95 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\left\langle \delta x_i \delta x_j \right\rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\rm total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_9	6									
x_{10}	32	19								
x_{18}	1	24	5							
x_{19}	1	8	3	2						
x_{20}	13	46	42	11	6					
x_{21}	1	5	3	1	24	8				
x_{22}	1	6	4	2	36	10	66			
x_{30}	3	11	9	3	7	23	16	18		
x_{38}	5	18	17	4	3	40	4	5	9	
x_{47}	1	3	2	1	18	6	33	51	9	4
x_{56}	3	9	8	2	1	19	2	2	4	28
x_{65}	1	3	2	1	16	5	30	46	8	2
x_{69}	1	3	2	1	17	5	58	47	11	2
x_{72}	1	2	2	1	13	4	24	37	6	2
x_{75}	1	4	3	1	21	6	39	60	10	3
x_{81}	1	6	4	1	30	9	56	84	18	4
x_{82}	1	5	4	1	7	10	24	18	43	4
x_{84}	1	3	3	1	0	7	1	1	2	18
x_{88}	1	2	2	0	2	4	3	5	2	9
x_{99}	0	2	1	0	7	3	13	20	4	3
x_{107}	1	3	3	1	2	6	4	4	23	3
x_{118}	8	28	25	7	4	60	5	6	14	24
x_{119}	0	2	1	0	9	3	17	25	4	1
x_{120}	1	4	3	1	14	6	26	39	7	3
x_{124}	1	3	2	1	11	6	20	30	6	2
x_{141}	0	2	1	0	11	3	20	30	5	1
x_{190}	−28	−21	−23	−7	−34	−32	−53	−70	−50	−26
	x_2	x_9	x_{10}	x_{18}	x_{19}	x_{20}	x_{21}	x_{22}	x_{30}	x_{38}

x_{56}	1									
x_{65}	23	1								
x_{69}	24	1	21							
x_{72}	43	1	17	17						
x_{75}	30	1	7	28	22					
x_{81}	43	2	38	40	31	50				
x_{82}	9	2	8	14	7	11	17			
x_{84}	1	5	0	0	0	0	1	1		
x_{88}	9	3	2	2	4	3	4	1	2	
x_{99}	40	1	9	9	17	12	17	4	1	4
x_{107}	2	1	2	2	2	2	4	10	0	0
x_{118}	3	12	3	3	2	4	6	6	4	2
x_{119}	13	1	11	12	9	15	21	5	0	1
x_{120}	20	1	18	18	14	23	33	7	0	2
x_{124}	15	1	13	14	11	18	25	6	0	2
x_{141}	15	1	14	14	11	18	25	6	0	1
x_{190}	−68	−20	−33	−38	−45	−43	−64	−39	−14	−23
	x_{47}	x_{56}	x_{65}	x_{69}	x_{72}	x_{75}	x_{81}	x_{82}	x_{84}	x_{88}

x_{107}	1						
x_{118}	2	4					
x_{119}	5	1	2				
x_{120}	8	2	4	10			
x_{124}	6	1	3	7	12		
x_{141}	6	1	2	8	12	9	
x_{190}	−34	−25	−20	−18	−30	−24	−23
	x_{99}	x_{107}	x_{118}	x_{119}	x_{120}	x_{124}	x_{141}

D^0 BRANCHING RATIOS

See the "Note on D Mesons" in the $D^{\pm}$ Listings.

Some older now obsolete results have been omitted from these Listings.

Inclusive modes

$\Gamma(e^+ \text{anything})/\Gamma_{\text{total}}$ **Γ_1/Γ**

VALUE	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.0675±0.0029 OUR AVERAGE					
0.069 ±0.003 ±0.005	1670	ALBRECHT	96C	ARG	$e^+e^- \approx$ 10 GeV
0.0664±0.0018±0.0029	4609	[17] KUBOTA	96B	CLE2	$e^+e^- \approx \Upsilon(4S)$
0.075 ±0.011 ±0.004	137	BALTRUSAIT..85B		MRK3	e^+e^- 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.15 ±0.05			AGUILAR-...	87E	HYBR	πp, pp 360, 400 GeV
0.055 ±0.037		12	SCHINDLER	81	MRK2	$e^+ e^-$ 3.771 GeV

[17] KUBOTA 96B uses $D^{*+} \rightarrow D^0 \pi^+$ (and charge conjugate) events in which the D^0 subsequently decays to $X e^+ \nu_e$.

$\Gamma(\mu^+ \text{anything})/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.066±0.008 OUR FIT					
0.060±0.007±0.012	310	ALBRECHT	96C	ARG	$e^+ e^-$ ≈ 10 GeV

$\Gamma(K^- \text{anything})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.53 ±0.04 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
$0.546^{+0.039}_{-0.038}$		[18] BARLAG	92C	ACCM	π^- Cu 230 GeV
0.609±0.032±0.052		COFFMAN	91	MRK3	$e^+ e^-$ 3.77 GeV
0.42 ±0.08		AGUILAR-...	87E	HYBR	πp, pp 360, 400 GeV
0.55 ±0.11	121	SCHINDLER	81	MRK2	$e^+ e^-$ 3.771 GeV
0.35 ±0.10	19	VUILLEMIN	78	MRK1	$e^+ e^-$ 3.772 GeV

[18] BARLAG 92C computes the branching fraction using topological normalization.



$\Gamma(K^- \text{anything})/\Gamma_{\text{total}}$

$[\Gamma(\overline{K^0}\text{anything}) + \Gamma(K^0\text{anything})]/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.42 ±0.05 OUR AVERAGE					
0.455±0.050±0.032		COFFMAN	91	MRK3	e^+e^- 3.77 GeV
0.29 ±0.11	13	SCHINDLER	81	MRK2	e^+e^- 3.771 GeV
0.57 ±0.26	6	VUILLEMIN	78	MRK1	e^+e^- 3.772 GeV

$\Gamma(K^+\text{anything})/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$0.034^{+0.006}_{-0.004}$ OUR AVERAGE					
$0.034^{+0.007}_{-0.005}$		[19] BARLAG	92C	ACCM	π^- Cu 230 GeV
0.028±0.009±0.004		COFFMAN	91	MRK3	e^+e^- 3.77 GeV
$0.03\ ^{+0.05}_{-0.02}$		AGUILAR-...	87E	HYBR	πp, pp 360, 400 GeV
0.08 ±0.03	25	SCHINDLER	81	MRK2	e^+e^- 3.771 GeV

[19] BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\phi\text{ anything})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$0.0171^{+0.0076}_{-0.0071}\pm0.0017$	9	[20] BAI	00C	BES	$e^+e^- \to D\overline{D}^*$, $D^*\overline{D}^*$

[20] BAI 00C finds the average (ϕ anything) branching fraction for the 4.03-GeV mix of D^+ and D^0 mesons to be (1.34 ± 0.52 ± 0.12)%.

Semileptonic modes

$\Gamma(K^-\ell^+\nu_\ell)/\Gamma_{\text{total}}$ Γ_8/Γ

We average our $K^-e^+\nu_e$ and $K^-\mu^+\nu_\mu$ branching fractions, after multiplying the latter by a phase-space factor of 1.03 to be able to use it with the $K^-e^+\nu_e$ fraction. Hence our ℓ^+ here is really an e^+.

VALUE	DOCUMENT ID		COMMENT
0.0348±0.0017 OUR AVERAGE	Error includes scale factor of 1.3.		
0.0364±0.0018	PDG	00	Our $\Gamma(K^-e^+\nu_e)/\Gamma_{\text{total}}$
0.0331±0.0018	PDG	00	1.03 × our $\Gamma(K^-\mu^+\nu_\mu)/\Gamma_{\text{total}}$

$\Gamma(K^-e^+\nu_e)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0364±0.0018 OUR FIT					
0.034 ±0.005 ±0.004	55	ADLER	89	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-e^+\nu_e)/\Gamma(K^-\pi^+)$ Γ_9/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.95 ±0.04 OUR FIT					
0.95 ±0.04 OUR AVERAGE					
0.978±0.027±0.044	2510	[21] BEAN	93C	CLE2	$e^+e^- \approx \Upsilon(4S)$
0.90 ±0.06 ±0.06	584	[22] CRAWFORD	91B	CLEO	$e^+e^- \approx$ 10.5 GeV
0.91 ±0.07 ±0.11	250	[23] ANJOS	89F	E691	Photoproduction

[21] BEAN 93C uses $K^- \mu^+ \nu_\mu$ as well as $K^- e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events. A pole mass of $2.00 \pm 0.12 \pm 0.18$ GeV/c^2 is obtained from the q^2 dependence of the decay rate.

[22] CRAWFORD 91B uses $K^- e^+ \nu_e$ and $K^- \mu^+ \nu_\mu$ candidates to measure a pole mass of $2.1 \,^{+0.4}_{-0.2} \,^{+0.3}_{-0.2}$ GeV/c^2 from the q^2 dependence of the decay rate.

[23] ANJOS 89F measures a pole mass of $2.1 \,^{+0.4}_{-0.2} \pm 0.2$ GeV/c^2 from the q^2 dependence of the decay rate.

$\Gamma(K^- \mu^+ \nu_\mu)/\Gamma(K^- \pi^+)$ Γ_{10}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.84 ±0.04 OUR FIT					
0.84 ±0.04 OUR AVERAGE					
0.852±0.034±0.028	1897	[24] FRABETTI	95G	E687	γ Be $\overline{E}_\gamma$= 220 GeV
0.82 ±0.13 ±0.13	338	[25] FRABETTI	93I	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.79 ±0.08 ±0.09	231	[26] CRAWFORD	91B	CLEO	$e^+ e^-$ ≈ 10.5 GeV

[24] FRABETTI 95G extracts the ratio of form factors $f_-(0)/f_+(0) = -1.3 \,^{+3.6}_{-3.4} \pm 0.6$, and measures a pole mass of $1.87 \,^{+0.11}_{-0.08} \,^{+0.07}_{-0.06}$ GeV/c^2 from the q^2 dependence of the decay rate.

[25] FRABETTI 93I measures a pole mass of $2.1 \,^{+0.7}_{-0.3} \,^{+0.7}_{-0.3}$ GeV/c^2 from the q^2 dependence of the decay rate.

[26] CRAWFORD 91B measures a pole mass of $2.00 \pm 0.12 \pm 0.18$ GeV/c^2 from the q^2 dependence of the decay rate.

$\Gamma(K^- \mu^+ \nu_\mu)/\Gamma(\mu^+ \text{anything})$ Γ_{10}/Γ_{2}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.49 ±0.06 OUR FIT					
0.472±0.051±0.040	232	KODAMA	94	E653	π^- emulsion 600 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.32 ±0.05 ±0.05	124	KODAMA	91	EMUL	pA 800 GeV

$\Gamma(K^- \pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$0.016 \,^{+0.013}_{-0.005} \pm 0.002$	4	[27] BAI	91	MRK3	$e^+ e^-$ ≈ 3.77 GeV

[27] BAI 91 finds that a fraction $0.79 \,^{+0.15}_{-0.17} \,^{+0.09}_{-0.03}$ of combined D^+ and D^0 decays to $\overline{K} \pi e^+ \nu_e$ (24 events) are $\overline{K}^*(892) e^+ \nu_e$. BAI 91 uses 56 $K^- e^+ \nu_e$ events to measure a pole mass of $1.8 \pm 0.3 \pm 0.2$ GeV/c^2 from the q^2 dependence of the decay rate.

$\Gamma(\overline{K}^0 \pi^- e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$0.028 \,^{+0.017}_{-0.008} \pm 0.003$	6	[28] BAI	91	MRK3	$e^+ e^-$ ≈ 3.77 GeV

[28] BAI 91 finds that a fraction $0.79 \,^{+0.15}_{-0.17} \,^{+0.09}_{-0.03}$ of combined D^+ and D^0 decays to $\overline{K} \pi e^+ \nu_e$ (24 events) are $\overline{K}^*(892) e^+ \nu_e$.

$\Gamma(K^*(892)^- e^+ \nu_e)/\Gamma(K^- e^+ \nu_e)$ Γ_{19}/Γ_9

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.55±0.09 OUR FIT			
0.51±0.18±0.06	CRAWFORD 91B	CLEO	$e^+e^-\approx$ 10.5 GeV

$\Gamma(K^*(892)^- e^+ \nu_e)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ Γ_{19}/Γ_{22}

Unseen decay modes of the $\overline{K}^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.37±0.06 OUR FIT				
0.38±0.06±0.03	152	[29] BEAN 93C	CLE2	$e^+e^-\approx \Upsilon(4S)$

[29] BEAN 93C uses $K^{*-}\mu^+\nu_\mu$ as well as $K^{*-}e^+\nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

$\Gamma(K^*(892)^- \ell^+ \nu_\ell)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ Γ_{14}/Γ_{22}

This an average of the $K^*(892)^- e^+\nu_e$ and $K^*(892)^-\mu^+\nu_\mu$ ratios. Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.24±0.07±0.06	137	[30] ALEXANDER 90B	CLEO	e^+e^- 10.5–11 GeV

[30] ALEXANDER 90B cannot exclude extra π^0's in the final state. See nearby data blocks for more detailed results.

$\Gamma(\overline{K}^*(892)^0 \pi^- e^+ \nu_e)/\Gamma(K^*(892)^- e^+ \nu_e)$ Γ_{15}/Γ_{19}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.64	90	[31] CRAWFORD 91B	CLEO	$e^+e^-\approx$ 10.5 GeV

[31] The limit on $(\overline{K}^*(892)\pi)^-\ \mu^+\nu_\mu$ below is much stronger.

$\Gamma(K^-\pi^+\pi^-\mu^+\nu_\mu)/\Gamma(K^-\mu^+\nu_\mu)$ Γ_{16}/Γ_{10}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.037	90	KODAMA 93B	E653	π^- emulsion 600 GeV

$\Gamma((\overline{K}^*(892)\pi)^-\mu^+\nu_\mu)/\Gamma(K^-\mu^+\nu_\mu)$ Γ_{17}/Γ_{10}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.043	90	[32] KODAMA 93B	E653	π^- emulsion 600 GeV

[32] KODAMA 93B searched in $K^-\pi^+\pi^-\mu^+\nu_\mu$, but the limit includes other $(\overline{K}^*(892)\pi)^-$ charge states.

$\Gamma(\pi^- e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0037±0.0006 OUR FIT				
$\mathbf{0.0039^{+0.0023}_{-0.0011}\pm0.0004}$	7	[33] ADLER 89	MRK3	e^+e^- 3.77 GeV

[33] This result of ADLER 89 gives $\left|\frac{V_{cd}}{V_{cs}}\cdot\frac{f_+^\pi(0)}{f_+^K(0)}\right|^2 = 0.057^{+0.038}_{-0.015}\pm0.005$.

$\Gamma(\pi^- e^+ \nu_e)/\Gamma(K^- e^+ \nu_e)$ Γ_{18}/Γ_9

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.102±0.017 OUR FIT					
0.101±0.018 OUR AVERAGE					
0.101±0.020±0.003	91	[34] FRABETTI	96B	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV
0.103±0.039±0.013	87	[35] BUTLER	95	CLE2	< 0.156 (90% CL)

[34] FRABETTI 96B uses both e and μ events, and makes a small correction to the μ events to make them effectively e events. This result gives $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}|^2 = 0.050 \pm 0.011 \pm 0.002$.

[35] BUTLER 95 has 87 ± 33 $\pi^- e^+ \nu_e$ events. The result gives $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}|^2 = 0.052 \pm 0.020 \pm 0.007$.

Hadronic modes with a $\overline{K}$ or $\overline{K}K\overline{K}$

$\Gamma(K^-\pi^+)/\Gamma_{total}$ Γ_{20}/Γ

We list measurements *before* radiative corrections are made.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0383±0.0009 OUR FIT					
0.0385±0.0009 OUR AVERAGE					
0.0382±0.0007±0.0012		[36] ARTUSO	98	CLE2	CLEO average
0.0390±0.0009±0.0012	5392	[37] BARATE	97C	ALEP	From Z decays
0.045 ±0.006 ±0.004		[38] ALBRECHT	94	ARG	$e^+e^- \approx \Upsilon(4S)$
0.0341±0.0012±0.0028	1173	[37] ALBRECHT	94F	ARG	$e^+e^- \approx \Upsilon(4S)$
0.0362±0.0034±0.0044		[37] DECAMP	91J	ALEP	From Z decays
0.045 ±0.008 ±0.005	56	[37] ABACHI	88	HRS	e^+e^- 29 GeV
0.042 ±0.004 ±0.004	930	ADLER	88C	MRK3	e^+e^- 3.77 GeV
0.041 ±0.006	263	[39] SCHINDLER	81	MRK2	e^+e^- 3.771 GeV
0.043 ±0.010	130	[40] PERUZZI	77	MRK1	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0381±0.0015±0.0016	1165	[41] ARTUSO	98	CLE2	e^+e^- at $\Upsilon(4S)$
0.0369±0.0011±0.0016		[42] COAN	98	CLE2	
0.0391±0.0008±0.0017	4208	[37,43] AKERIB	93	CLE2	$e^+e^- \approx \Upsilon(4S)$

[36] This combines the CLEO results of ARTUSO 98, COAN 98, and AKERIB 93.

[37] ABACHI 88, DECAMP 91J, AKERIB 93, ALBRECHT 94F, and BARATE 97C use $D^*(2010)^+ \to D^0\pi^+$ decays. The π^+ is both slow and of low p_T with respect to the event thrust axis or nearest jet ($\approx D^{*+}$ direction). The excess number of such π^+'s over background gives the number of $D^*(2010)^+ \to D^0\pi^+$ events, and the fraction with $D^0 \to K^-\pi^+$ gives the $D^0 \to K^-\pi^+$ branching fraction.

[38] ALBRECHT 94 uses D^0 mesons from $\overline{B}^0 \to D^{*+}\ell^-\overline{\nu}_\ell$ decays. This is a different set of events than used by ALBRECHT 94F.

[39] SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.24 ± 0.02 nb. We use the MARK-3 (ADLER 88C) value of σ = 5.8 ± 0.5 ± 0.6 nb.

[40] PERUZZI 77 (MARK-1) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.25 ± 0.05 nb. We use the MARK-3 (ADLER 88C) value of σ = 5.8 ± 0.5 ± 0.6 nb.

[41] ARTUSO 98, following ALBRECHT 94, uses D^0 mesons from $\overline{B}^0 \to D^*(2010)^+ X\ell^-\overline{\nu}_\ell$ decays. Our average uses the CLEO average of this value with the values of COAN 98 and AKERIB 93.

[42] COAN 98 assumes that $\Gamma(B \to \overline{D} X \ell^+ \nu)/\Gamma(B \to X \ell^+ \nu) = 1.0 - 3|V_{ub}/V_{cb}|^2 - 0.010 \pm 0.005$, the last term accounting for $\overline{B} \to D_s^+ K X \ell^- \overline{\nu}$. COAN 98 is included in the CLEO average in ARTUSO 98.

[43] This AKERIB 93 value does not include radiative corrections; with them, the value is $0.0395 \pm 0.0008 \pm 0.0017$. AKERIB 93 is included in the CLEO average in ARTUSO 98.

$\Gamma(\overline{K}^0\pi^0)/\Gamma(K^-\pi^+)$ Γ_{21}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.55±0.06 OUR FIT	Error includes scale factor of 1.1.				
1.36±0.23±0.22	119	ANJOS	92B	E691	γ Be 80–240 GeV

$\Gamma(\overline{K}^0\pi^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{21}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.390±0.031 OUR FIT					
0.378±0.033 OUR AVERAGE					
0.44 ±0.02 ±0.05	1942	PROCARIO	93B	CLE2	e^+e^- 10.36–10.7 GeV
0.34 ±0.04 ±0.02	92	[44] ALBRECHT	92P	ARG	$e^+e^- \approx$ 10 GeV
0.36 ±0.04 ±0.08	104	KINOSHITA	91	CLEO	$e^+e^- \sim$ 10.7 GeV

[44] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^0\pi^+\pi^-)/\Gamma_{total}$ Γ_{22}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.054 ±0.004 OUR FIT	Error includes scale factor of 1.2.				
0.055 ±0.005 OUR AVERAGE					
0.0503±0.0039±0.0049	284	[45] ALBRECHT	94F	ARG	$e^+e^- \approx \Upsilon(4S)$
0.064 ±0.005 ±0.010		ADLER	87	MRK3	e^+e^- 3.77 GeV
0.052 ±0.016	32	[46] SCHINDLER	81	MRK2	e^+e^- 3.771 GeV
0.079 ±0.023	28	[47] PERUZZI	77	MRK1	e^+e^- 3.77 GeV

[45] See the footnote on the ALBRECHT 94F measurement of $\Gamma(K^-\pi^+)/\Gamma_{total}$ for the method used.

[46] SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.30 ± 0.08 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

[47] PERUZZI 77 (MARK-1) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.46 ± 0.12 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

$\Gamma(\overline{K}^0\pi^+\pi^-)/\Gamma(K^-\pi^+)$ Γ_{22}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.42±0.10 OUR FIT	Error includes scale factor of 1.2.				
1.65±0.17 OUR AVERAGE					
1.61±0.10±0.15	856	FRABETTI	94J	E687	γ Be $\overline{E}_\gamma$=220 GeV
1.7 ±0.8	35	AVERY	80	SPEC	$\gamma N \to D^{*+}$
2.8 ±1.0	116	PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV

$\Gamma(\overline{K}^0\rho^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{23}/Γ_{22}

VALUE	DOCUMENT ID		TECN	COMMENT
0.223±0.027 OUR AVERAGE	Error includes scale factor of 1.2.			
0.350±0.028±0.067	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.227±0.032±0.009	ALBRECHT	93D	ARG	$e^+e^- \approx$ 10 GeV
0.215±0.051±0.037	ANJOS	93	E691	γ Be 90–260 GeV
0.20 ±0.06 ±0.03	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.12 ±0.01 ±0.07	ADLER	87	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0 f_0(980))/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{74}/Γ_{22}

Unseen decay modes of the $f_0(980)$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.105±0.029 OUR AVERAGE				
0.131±0.031±0.034	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.088±0.035±0.012	ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(\overline{K}^0 f_2(1270))/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{78}/Γ_{22}

Unseen decay modes of the $f_2(1270)$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.076±0.028 OUR AVERAGE				
0.065±0.025±0.030	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.088±0.037±0.014	ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(\overline{K}^0 f_0(1370))/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{80}/Γ_{22}

Unseen decay modes of the $f_0(1370)$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.13 ±0.04 OUR AVERAGE				
0.123±0.035±0.049	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.131±0.045±0.021	ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(K^*(892)^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{81}/Γ_{22}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.93 ±0.04 OUR FIT Error includes scale factor of 1.1.					
0.96 ±0.04 OUR AVERAGE					
0.938±0.054±0.038		FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
1.08 ±0.063±0.045		ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV
0.720±0.145±0.185		ANJOS	93	E691	γ Be 90–260 GeV
0.96 ±0.12 ±0.075		FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.84 ±0.06 ±0.08		ADLER	87	MRK3	e^+e^- 3.77 GeV
$1.05\ ^{+0.23}_{-0.26}\ ^{+0.07}_{-0.09}$	25	SCHINDLER	81	MRK2	e^+e^- 3.771 GeV

$\Gamma(K_0^*(1430)^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{103}/Γ_{22}

Unseen decay modes of the $\overline{K}_0^*(1430)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.19 ±0.05 OUR AVERAGE				
0.176±0.044±0.047	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
0.208±0.055±0.034	ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(K_2^*(1430)^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{104}/Γ_{22}

Unseen decay modes of the $\overline{K}_2^*(1430)^-$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.15	90	ALBRECHT	93D	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(\overline{K^0}\pi^+\pi^-$ nonresonant$)/\Gamma(\overline{K^0}\pi^+\pi^-)$ Γ_{29}/Γ_{22}

VALUE	DOCUMENT ID		TECN	COMMENT
0.27 ±0.04 OUR AVERAGE				
0.263±0.024±0.041	ANJOS	93	E691	γ Be 90–260 GeV
0.26 ±0.08 ±0.05	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$ = 221 GeV
0.33 ±0.05 ±0.10	ADLER	87	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.139±0.009 OUR FIT Error includes scale factor of 1.3.					
0.131±0.016 OUR AVERAGE					
0.133±0.012±0.013	931	ADLER	88C	MRK3	e^+e^- 3.77 GeV
0.117±0.043	37	[48] SCHINDLER	81	MRK2	e^+e^- 3.771 GeV

[48] SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.68 ± 0.23 nb. We use the MARK-3 (ADLER 88C) value of σ = 5.8 ± 0.5 ± 0.6 nb.

$\Gamma(K^-\pi^+\pi^0)/\Gamma(K^-\pi^+)$ Γ_{30}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
3.63±0.23 OUR FIT Error includes scale factor of 1.4.					
3.47±0.30 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.					
3.81±0.07±0.26	10k	BARISH	96	CLE2	$e^+e^- \approx \Upsilon(4S)$
3.04±0.16±0.34	931	[49] ALBRECHT	92P	ARG	$e^+e^- \approx$ 10 GeV
4.0 ±0.9 ±1.0	69	ALVAREZ	91B	NA14	Photoproduction
2.8 ±0.14±0.52	1050	KINOSHITA	91	CLEO	$e^+e^- \sim$ 10.7 GeV
4.2 ±1.4	41	SUMMERS	84	E691	Photoproduction

[49] This value is calculated from numbers in Table 1 of ALBRECHT 92P.



$\Gamma(K^-\pi^+\pi^0)/\Gamma(K^-\pi^+)$

$\Gamma(K^-\rho^+)/\Gamma(K^-\pi^+\pi^0)$ Γ_{31}/Γ_{30}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.78 ±0.05 OUR AVERAGE					
0.765±0.041±0.054		FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx 220$ GeV
0.647±0.039±0.150		ANJOS	93	E691	γ Be 90–260 GeV
0.81 ±0.03 ±0.06		ADLER	87	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.31\ ^{+0.20}_{-0.14}$	13	SUMMERS	84	E691	Photoproduction
$0.85\ ^{+0.11}_{-0.15}\ ^{+0.09}_{-0.10}$	31	SCHINDLER	81	MRK2	e^+e^- 3.771 GeV

$\Gamma(K^*(892)^-\pi^+)/\Gamma(K^-\pi^+\pi^0)$ Γ_{81}/Γ_{30}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.36 ±0.04 OUR FIT Error includes scale factor of 1.3.				
0.28 ±0.04 OUR AVERAGE				
0.444±0.084±0.147	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx 220$ GeV
0.252±0.033±0.035	ANJOS	93	E691	γ Be 90–260 GeV
0.36 ±0.06 ±0.09	ADLER	87	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\pi^0)/\Gamma(K^-\pi^+\pi^0)$ Γ_{82}/Γ_{30}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.227±0.027 OUR FIT				
0.221±0.029 OUR AVERAGE				
0.248±0.047±0.023	FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx 220$ GeV
0.213±0.027±0.035	ANJOS	93	E691	γ Be 90–260 GeV
0.20 ±0.03 ±0.05	ADLER	87	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-\pi^+\pi^0\ \text{nonresonant})/\Gamma(K^-\pi^+\pi^0)$ Γ_{34}/Γ_{30}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.049±0.018 OUR AVERAGE Error includes scale factor of 1.1.					
0.101±0.033±0.040		FRABETTI	94G	E687	γ Be, $\overline{E}_\gamma \approx 220$ GeV
0.036±0.004±0.018		ANJOS	93	E691	γ Be 90–260 GeV
0.09 ±0.02 ±0.04		ADLER	87	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.51 ±0.22	21	SUMMERS	84	E691	Photoproduction

$\Gamma(\overline{K}^*(892)^0\pi^0)/\Gamma(\overline{K}^0\pi^0)$ Γ_{82}/Γ_{21}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.49±0.23 OUR FIT Error includes scale factor of 1.1.					
$\mathbf{1.65^{+0.39}_{-0.31}\pm0.20}$	122	PROCARIO	93B	CLE2	$\overline{K}^0\pi^0\pi^0$ Dalitz plot

$\Gamma(\overline{K}_2^*(1430)^0\pi^0)/\Gamma(\overline{K}^*(892)^0\pi^0)$ Γ_{105}/Γ_{82}

Unseen decay modes of the $\overline{K}_2^*(1430)^0$ and $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.12	90	PROCARIO	93B	CLE2	$\overline{K}^0\pi^0\pi^0$ Dalitz plot

$\Gamma(\overline{K}^0\pi^0\pi^0$ nonresonant$)/\Gamma(\overline{K}^0\pi^0)$ Γ_{37}/Γ_{21}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.37±0.08±0.04	76	PROCARIO	93B	CLE2	$\overline{K}^0\pi^0\pi^0$ Dalitz plot

$\Gamma(K^-\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0749±0.0031 OUR FIT					
0.075 ±0.006 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
0.079 ±0.015 ±0.009		[50] ALBRECHT	94	ARG	$e^+e^- \approx \Upsilon(4S)$
0.0680±0.0027±0.0057	1430	[51] ALBRECHT	94F	ARG	$e^+e^- \approx \Upsilon(4S)$
0.091 ±0.008 ±0.008	992	ADLER	88C	MRK3	e^+e^- 3.77 GeV
0.117 ±0.025	185	[52] SCHINDLER	81	MRK2	e^+e^- 3.771 GeV
0.062 ±0.019	44	[53] PERUZZI	77	MRK1	e^+e^- 3.77 GeV

[50] ALBRECHT 94 uses D^0 mesons from $\overline{B}^0 \to D^{*+}\ell^-\overline{\nu}_\ell$ decays. This is a different set of events than used by ALBRECHT 94F.

[51] See the footnote on the ALBRECHT 94F measurement of $\Gamma(K^-\pi^+)/\Gamma_{\text{total}}$ for the method used.

[52] SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.68 ± 0.11 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

[53] PERUZZI 77 (MARK-1) measures $\sigma(e^+e^- \to \psi(3770)) \times$ branching fraction to be 0.36 ± 0.10 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.



$\Gamma(K^-\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma(K^-\pi^+\pi^+\pi^-)/\Gamma(K^-\pi^+)$ Γ_{38}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.96±0.07 OUR FIT					
1.97±0.09 OUR AVERAGE					
$1.94\pm0.07^{+0.09}_{-0.11}$		JUN	00	SELX	Σ^- nucleus, 600 GeV
1.7 ±0.2 ±0.2	1745	ANJOS	92C	E691	γ Be 90–260 GeV
1.90±0.25±0.20	337	ALVAREZ	91B	NA14	Photoproduction
2.12±0.16±0.09		BORTOLETTO	88	CLEO	e^+e^- 10.55 GeV
2.0 ±0.9	48	BAILEY	86	ACCM	π^- Be fixed target
2.17±0.28±0.23		ALBRECHT	85F	ARG	e^+e^- 10 GeV
2.0 ±1.0	10	BAILEY	83B	SPEC	π^- Be → D^0
2.2 ±0.8	214	PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV

$\Gamma(K^-\pi^+\rho^0\,\text{total})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{39}/Γ_{38}

This includes $K^-a_1(1260)^+$, $\overline{K}^*(892)^0\rho^0$, etc. The next entry gives the specifically 3-body fraction. We rely on the MARK III and E691 full amplitude analyses of the $K^-\pi^+\pi^+\pi^-$ channel for values of the resonant substructure.

VALUE	DOCUMENT ID		TECN	COMMENT
0.835±0.035 OUR AVERAGE				
0.80 ±0.03 ±0.05	ANJOS	92C	E691	γ Be 90–260 GeV
0.855±0.032±0.030	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.98 ±0.12 ±0.10	ALVAREZ	91B	NA14	Photoproduction

$\Gamma(K^-\pi^+\rho^0\,3\text{-body})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{40}/Γ_{38}

We rely on the MARK III and E691 full amplitude analyses of the $K^-\pi^+\pi^+\pi^-$ channel for values of the resonant substructure.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.063±0.028 OUR AVERAGE					
0.05 ±0.03 ±0.02		ANJOS	92C	E691	γ Be 90–260 GeV
0.084±0.022±0.04		COFFMAN	92B	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.77 ±0.06 ±0.06	[54]	ALVAREZ	91B	NA14	Photoproduction
$0.85\ ^{+0.11}_{-0.22}$	180	PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV

[54] This value is for ρ^0 $(K^-\pi^+)$-nonresonant. ALVAREZ 91B cannot determine what fraction of this is $K^-a_1(1260)^+$.

$\Gamma(\overline{K}^*(892)^0\rho^0)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{87}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included. We rely on the MARK III and E691 full amplitude analyses of the $K^-\pi^+\pi^+\pi^-$ channel for values of the resonant substructure.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.195±0.03±0.03		ANJOS	92C	E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.34 ±0.09±0.09		ALVAREZ	91B	NA14	Photoproduction
0.75 ±0.3	5	BAILEY	83B	SPEC	π Be → D^0
$0.15\ ^{+0.16}_{-0.15}$	20	PICCOLO	77	MRK1	e^+e^- 4.03, 4.41 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,\text{transverse})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{88}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.20 ±0.07 OUR FIT			
0.213±0.024±0.075	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,S\text{-wave})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{89}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.375±0.045±0.06	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,S\text{-wave long.})/\Gamma_{\text{total}}$ Γ_{90}/Γ

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,P\text{-wave})/\Gamma_{\text{total}}$ Γ_{91}/Γ

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	COFFMAN	92B MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.009	90	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\overline{K}^*(892)^0\rho^0\,D\text{-wave})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{92}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.255±0.045±0.06	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(K^-\pi^+ f_0(980))/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.011	90	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\overline{K}^*(892)^0 f_0(980))/\Gamma_{\text{total}}$ Γ_{98}/Γ

Unseen decay modes of the $\overline{K}^*(892)^0$ and $f_0(980)$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(K^-a_1(1260)^+)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{76}/Γ_{38}

Unseen decay modes of the $a_1(1260)^+$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.97 ±0.14 OUR AVERAGE			
0.94 ±0.13 ±0.20	ANJOS	92C E691	γ Be 90–260 GeV
0.984±0.048±0.16	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-a_2(1320)^+)/\Gamma_{\text{total}}$ Γ_{79}/Γ

Unseen decay modes of the $a_2(1320)^+$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.002	90	ANJOS	92C E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.006	90	COFFMAN	92B MRK3	e^+e^- 3.77 GeV

$\Gamma(K_1(1270)^-\pi^+)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{99}/Γ_{38}

Unseen decay modes of the $K_1(1270)^-$ are included. The MARK3 and E691 experiments disagree considerably here.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
0.14 ±0.04 OUR FIT					
0.194±0.056±0.088		COFFMAN	92B	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.013	90	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(K_1(1400)^-\pi^+)/\Gamma_{total}$ Γ_{100}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.012	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(1410)^-\pi^+)/\Gamma_{total}$ Γ_{102}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.012	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\pi^+\pi^-\text{total})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{83}/Γ_{38}

This includes $\overline{K}^*(892)^0\rho^0$, *etc.* The next entry gives the specifically 3-body fraction. Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.30±0.06±0.03	ANJOS	92C	E691	γ Be 90–260 GeV

$\Gamma(\overline{K}^*(892)^0\pi^+\pi^-\text{3-body})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{84}/Γ_{38}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.19 ±0.04 OUR FIT				
0.18 ±0.04 OUR AVERAGE				
0.165±0.03 ±0.045	ANJOS	92C	E691	γ Be 90–260 GeV
0.210±0.027±0.06	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-\pi^+\pi^+\pi^-\text{nonresonant})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{46}/Γ_{38}

VALUE	DOCUMENT ID		TECN	COMMENT
0.233±0.032 OUR AVERAGE				
0.23 ±0.02 ±0.03	ANJOS	92C	E691	γ Be 90–260 GeV
0.242±0.025±0.06	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{47}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.100±0.012 OUR FIT					
0.103±0.022±0.025	140	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.134^{+0.032}_{-0.033}$		[55] BARLAG	92C	ACCM	π^- Cu 230 GeV

[55] BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{47}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.84±0.20 OUR FIT					
1.86±0.23 OUR AVERAGE					
1.80±0.20±0.21	190	[56] ALBRECHT	92P	ARG	$e^+e^-\approx$ 10 GeV
2.8 ±0.8 ±0.8	46	ANJOS	92C	E691	γ Be 90–260 GeV
1.85±0.26±0.30	158	KINOSHITA	91	CLEO	$e^+e^-\sim$ 10.7 GeV

[56] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^0\eta)/\Gamma(K^-\pi^+)$ Γ_{69}/Γ_{20}

Unseen decay modes of the η are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.64	90	ALBRECHT	89D	ARG	e^+e^- 10 GeV

$\Gamma(\overline{K}^0\eta)/\Gamma(\overline{K}^0\pi^0)$ Γ_{69}/Γ_{21}

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.33±0.04 OUR FIT					
0.32±0.04±0.03	225	PROCARIO	93B	CLE2	$\eta\to\gamma\gamma$

$\Gamma(\overline{K}^0\eta)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{69}/Γ_{22}

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.130±0.017 OUR FIT					
0.14 ±0.02 ±0.02	80	PROCARIO	93B	CLE2	$\eta\to\pi^+\pi^-\pi^0$

$\Gamma(\overline{K}^0\omega)/\Gamma(K^-\pi^+)$ Γ_{72}/Γ_{20}

Unseen decay modes of the ω are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.54±0.10 OUR FIT				
1.00±0.36±0.20	ALBRECHT	89D	ARG	e^+e^- 10 GeV

$\Gamma(\overline{K}^0\omega)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{72}/Γ_{22}

Unseen decay modes of the ω are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.38±0.07 OUR FIT					
0.33±0.09 OUR AVERAGE Error includes scale factor of 1.1.					
0.29±0.08±0.05	16	[57] ALBRECHT	92P	ARG	$e^+e^-\approx$ 10 GeV
0.54±0.14±0.16	40	KINOSHITA	91	CLEO	$e^+e^-\sim$ 10.7 GeV

[57] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^0\omega)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{72}/Γ_{47}

Unseen decay modes of the ω are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.21 ±0.04 OUR FIT				
0.220±0.048±0.0116	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0\eta'(958))/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{73}/Γ_{22}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.32±0.04 OUR AVERAGE					
0.31±0.02±0.04	594	PROCARIO	93B	CLE2	$\eta' \to \eta\pi^+\pi^-, \rho^0\gamma$
0.37±0.13±0.06	18	[58] ALBRECHT	92P	ARG	$e^+e^- \approx$ 10 GeV

[58] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^*(892)^-\rho^+)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{93}/Γ_{47}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.606±0.188±0.126	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(892)^-\rho^+ \text{longitudinal})/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{94}/Γ_{47}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.290±0.111	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(892)^-\rho^+ \text{transverse})/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{95}/Γ_{47}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.317±0.180	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(892)^-\rho^+ P\text{-wave})/\Gamma_{\text{total}}$ Γ_{96}/Γ

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.015	90	[59] COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

[59] Obtained using other $\overline{K}^*(892)\rho$ P-wave limits and isospin relations.

$\Gamma(\overline{K}^*(892)^0\rho^0 \text{transverse})/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{88}/Γ_{47}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.15 ±0.06 OUR FIT				
0.126±0.111	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0 a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{77}/Γ

Unseen decay modes of the $a_1(1260)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.019	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K_1(1270)^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{99}/Γ_{47}

Unseen decay modes of the $K_1(1270)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.106±0.028 OUR FIT				
0.10 ±0.03	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}_1(1400)^0\pi^0)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.037	90	COFFMAN	92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^*(892)^0\pi^+\pi^-$ 3-body$)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{84}/Γ_{47}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.14 ±0.04 OUR FIT Error includes scale factor of 1.1.			
0.191±0.105	COFFMAN 92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0\pi^+\pi^-\pi^0$ nonresonant$)/\Gamma(\overline{K}^0\pi^+\pi^-\pi^0)$ Γ_{54}/Γ_{47}

VALUE	DOCUMENT ID	TECN	COMMENT
0.210±0.147±0.150	COFFMAN 92B	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^-\pi^+\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.177±0.029		[60] BARLAG 92C	ACCM	π^- Cu 230 GeV
0.149±0.037±0.030	24	[61] ADLER 88C	MRK3	e^+e^- 3.77 GeV
$0.209^{+0.074}_{-0.043}\pm0.012$	9	[60] AGUILAR-... 87F	HYBR	πp, pp 360, 400 GeV

[60] AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction using topological normalization. They do not distinguish the presence of a third π^0, and thus are not included in the average.

[61] ADLER 88C uses an absolute normalization method finding this decay channel opposite a detected $\overline{D}^0 \to K^+\pi^-$ in pure $D\overline{D}$ events.

$\Gamma(K^-\pi^+\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+)$ Γ_{56}/Γ_{20}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.05±0.10 OUR FIT				
0.98±0.11±0.11	225	[62] ALBRECHT 92P	ARG	$e^+e^- \approx$ 10 GeV

[62] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^-\pi^+\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{56}/Γ_{38}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.54±0.05 OUR FIT				
0.56±0.07 OUR AVERAGE				
$0.55\pm0.07^{+0.12}_{-0.09}$	167	KINOSHITA 91	CLEO	$e^+e^- \sim$ 10.7 GeV
0.57±0.06±0.05	180	ANJOS 90D	E691	Photoproduction

$\Gamma(\overline{K}^*(892)^0\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+\pi^+\pi^-\pi^0)$ Γ_{106}/Γ_{56}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.45±0.15±0.15	ANJOS 90D	E691	Photoproduction

$\Gamma(\overline{K}^*(892)^0\eta)/\Gamma(K^-\pi^+)$ Γ_{107}/Γ_{20}

Unseen decay modes of the $\overline{K}^*(892)^0$ and η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.49±0.12 OUR FIT				
$\mathbf{0.58\pm0.19^{+0.24}_{-0.28}}$	46	KINOSHITA 91	CLEO	$e^+e^- \sim$ 10.7 GeV

$\Gamma(\overline{K}^*(892)^0\eta)/\Gamma(K^-\pi^+\pi^0)$ Γ_{107}/Γ_{30}

Unseen decay modes of the $\overline{K}^*(892)^0$ and η are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.134±0.034 OUR FIT					
0.13 ±0.02 ±0.03	214	PROCARIO	93B	CLE2	$\overline{K}^{*0}\eta \rightarrow K^-\pi^+/\gamma\gamma$

$\Gamma(K^-\pi^+\omega)/\Gamma(K^-\pi^+)$ Γ_{108}/Γ_{20}

Unseen decay modes of the ω are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.78±0.12±0.10	99	[63] ALBRECHT	92P	ARG	$e^+e^- \approx$ 10 GeV

[63] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^*(892)^0\omega)/\Gamma(K^-\pi^+)$ Γ_{109}/Γ_{20}

Unseen decay modes of the $\overline{K}^*(892)^0$ and ω are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.28±0.11±0.04	17	[64] ALBRECHT	92P	ARG	$e^+e^- \approx$ 10 GeV

[64] This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(\overline{K}^*(892)^0\omega)/\Gamma(K^-\pi^+\pi^+\pi^-\pi^0)$ Γ_{109}/Γ_{56}

Unseen decay modes of the $\overline{K}^*(892)^0$ and ω are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.44	90	[65] ANJOS	90D	E691	Photoproduction

[65] Recovered from the published limit, $\Gamma(\overline{K}^*(892)^0\omega)/\Gamma_{\text{total}}$, in order to make our normalization consistent.

$\Gamma(K^-\pi^+\eta'(958))/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{110}/Γ_{38}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.093±0.014±0.019	286	PROCARIO	93B	CLE2	$\eta' \rightarrow \eta\pi^+\pi^-$, $\rho^0\gamma$

$\Gamma(\overline{K}^*(892)^0\eta'(958))/\Gamma(K^-\pi^+\eta'(958))$ $\Gamma_{111}/\Gamma_{110}$

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN
<0.15	90	PROCARIO	93B	CLE2

$\Gamma(\overline{K}^0\pi^+\pi^+\pi^-\pi^-)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{61}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.107±0.029 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.			
0.07 ±0.02 ±0.01	11	[66] ALBRECHT	92P	ARG	$e^+e^- \approx$ 10 GeV
0.149±0.026	56	AMMAR	91	CLEO	$e^+e^- \approx$ 10.5 GeV
0.18 ±0.07 ±0.04	6	ANJOS	90D	E691	Photoproduction

[66] This value is calculated from numbers in Table 1 of ALBRECHT 92P.



$\Gamma\left(\overline{K}^0\,\pi^+\,\pi^+\,\pi^-\,\pi^-\right)/\Gamma\left(\overline{K}^0\,\pi^+\,\pi^-\right)$

$\Gamma(\overline{K}^0\pi^+\pi^-\pi^0\pi^0(\pi^0))/\Gamma_{total}$ Γ_{62}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.106\,^{+0.073}_{-0.029}\pm0.006$	4	[67] AGUILAR-...		87F HYBR	πp, pp 360, 400 GeV

[67] AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization, and does not distinguish the presence of a third π^0.

$\Gamma(\overline{K}^0K^+K^-)/\Gamma(\overline{K}^0\pi^+\pi^-)$ $\Gamma_{63}/\Gamma_{22}=(\Gamma_{65}+\frac{1}{2}\Gamma_{75})/\Gamma_{22}$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.172±0.014 OUR FIT					
0.178±0.019 OUR AVERAGE					
0.20 ±0.05 ±0.04	47	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma=$ 221 GeV
0.170±0.022	136	AMMAR	91	CLEO	$e^+e^-\approx$ 10.5 GeV
0.24 ±0.08		BEBEK	86	CLEO	e^+e^- near $\Upsilon(4S)$
0.185±0.055	52	ALBRECHT	85B	ARG	e^+e^- 10 GeV

$\Gamma(\overline{K}^0\phi)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{75}/Γ_{22}

Unseen decay modes of the ϕ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.158±0.016 OUR FIT					
0.156±0.017 OUR AVERAGE					
0.13 ±0.06 ±0.02	13	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma=$ 221 GeV
0.163±0.023	63	AMMAR	91	CLEO	$e^+e^-\approx$ 10.5 GeV
0.155±0.033	56	ALBRECHT	87E	ARG	e^+e^- 10 GeV
0.14 ±0.05	29	BEBEK	86	CLEO	e^+e^- near $\Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.186±0.052	26	ALBRECHT	85B	ARG	See ALBRECHT 87E

$\Gamma(\overline{K}^0 K^+ K^-$ non-$\phi)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ Γ_{65}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.093±0.014 OUR FIT					
0.088±0.019 OUR AVERAGE					
0.11 ±0.04 ±0.03	20	FRABETTI	92B	E687	γ Be $\overline{E}_\gamma$= 221 GeV
0.084±0.020		ALBRECHT	87E	ARG	e^+e^- 10 GeV

$\Gamma(K^0_S K^0_S K^0_S)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ Γ_{66}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0154±0.0025 OUR AVERAGE					
0.0139±0.0019±0.0024	61	ASNER	96B	CLE2	$e^+e^- \approx \Upsilon(4S)$
0.035 ±0.012 ±0.006	10	FRABETTI	94J	E687	γ Be $\overline{E}_\gamma$=220 GeV
0.016 ±0.005	22	AMMAR	91	CLEO	$e^+e^- \approx$ 10.5 GeV
0.017 ±0.007 ±0.005	5	ALBRECHT	90C	ARG	$e^+e^- \approx$ 10 GeV

$\Gamma(K^+ K^- K^- \pi^+)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{67}/Γ_{38}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0028±0.0007±0.0001	20	FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV

$\Gamma(K^+ K^- \overline{K}^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE		DOCUMENT ID		TECN	COMMENT
$\mathbf{0.0072^{+0.0048}_{-0.0035}}$	[68]	BARLAG	92C	ACCM	π^- Cu 230 GeV

[68] BARLAG 92C computes the branching fraction using topological normalization.

Pionic modes

$\Gamma(\pi^+ \pi^-)/\Gamma(K^- \pi^+)$ Γ_{112}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0397±0.0021 OUR AVERAGE					
0.040 ±0.002 ±0.003	2043	AITALA	98C	E791	π^- nucleus, 500 GeV
0.043 ±0.007 ±0.003	177	FRABETTI	94C	E687	γ Be $\overline{E}_\gamma$= 220 GeV
0.0348±0.0030±0.0023	227	SELEN	93	CLE2	$e^+e^- \approx \Upsilon(4S)$
0.048 ±0.013 ±0.008	51	ADAMOVICH	92	OMEG	π^- 340 GeV
0.055 ±0.008 ±0.005	120	ANJOS	91D	E691	Photoproduction
0.040 ±0.007 ±0.006	57	ALBRECHT	90C	ARG	$e^+e^- \approx$ 10 GeV
0.050 ±0.007 ±0.005	110	ALEXANDER	90	CLEO	e^+e^- 10.5–11 GeV
0.033 ±0.010 ±0.006	39	BALTRUSAIT..	85E	MRK3	e^+e^- 3.77 GeV
0.033 ±0.015		ABRAMS	79D	MRK2	e^+e^- 3.77 GeV

$\Gamma(\pi^0 \pi^0)/\Gamma(K^- \pi^+)$ Γ_{113}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.022±0.004±0.004	40	SELEN	93	CLE2	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{114}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.011 ±0.004 ±0.002	10	[69] BALTRUSAIT...85E		MRK3	e^+e^- 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.0390^{+0.0100}_{-0.0095}$	[70] BARLAG	92C	ACCM		π^- Cu 230 GeV

[69] All the BALTRUSAITIS 85E events are consistent with $\rho^0\pi^0$.
[70] BARLAG 92C computes the branching fraction using topological normalization. Possible contamination by extra π^0's may partly explain the unexpectedly large value.

$\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{115}/Γ_{38}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.098±0.006 OUR AVERAGE					
0.095±0.007±0.002	814	FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx 200$ GeV
0.115±0.023±0.016	64	ADAMOVICH	92	OMEG	π^- 340 GeV
0.108±0.024±0.008	79	FRABETTI	92	E687	γ Be
0.102±0.013	345	[71] AMMAR	91	CLEO	$e^+e^- \approx 10.5$ GeV
0.096±0.018±0.007	66	ANJOS	91	E691	γ Be 80–240 GeV

[71] AMMAR 91 finds 1.25 ± 0.25 ± 0.25 ρ^0's per $\pi^+\pi^+\pi^-\pi^-$ decay, but can't untangle the resonant substructure ($\rho^0\rho^0$, $a_1^\pm\pi^\mp$, $\rho^0\pi^+\pi^-$).

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{total}$ Γ_{116}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.0192^{+0.0041}_{-0.0038}}$	[72] BARLAG	92C	ACCM	π^- Cu 230 GeV

[72] BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-\pi^-)/\Gamma_{total}$ Γ_{117}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0004±0.0003	[73] BARLAG	92C	ACCM	π^- Cu 230 GeV

[73] BARLAG 92C computes the branching fraction using topological normalization.

Hadronic modes with a $K\overline{K}$ pair

$\Gamma(K^+K^-)/\Gamma(K^-\pi^+)$ Γ_{118}/Γ_{20}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.1109±0.0033 OUR FIT					
0.1109±0.0033 OUR AVERAGE					
0.109 ±0.003 ±0.003	3317	AITALA	98C	E791	π^- nucleus, 500 GeV
0.116 ±0.007 ±0.007	1102	ASNER	96B	CLE2	$e^+e^- \approx \Upsilon(4S)$
0.109 ±0.007 ±0.009	581	FRABETTI	94C	E687	γ Be $\overline{E}_\gamma = 220$ GeV
0.107 ±0.029 ±0.015	103	ADAMOVICH	92	OMEG	π^- 340 GeV
0.138 ±0.027 ±0.010	155	FRABETTI	92	E687	γ Be
0.16 ±0.05	34	ALVAREZ	91B	NA14	Photoproduction
0.107 ±0.010 ±0.009	193	ANJOS	91D	E691	Photoproduction
0.10 ±0.02 ±0.01	131	ALBRECHT	90C	ARG	$e^+e^- \approx 10$ GeV
0.117 ±0.010 ±0.007	249	ALEXANDER	90	CLEO	e^+e^- 10.5–11 GeV
0.122 ±0.018 ±0.012	118	BALTRUSAIT...85E		MRK3	e^+e^- 3.77 GeV
0.113 ±0.030		ABRAMS	79D	MRK2	e^+e^- 3.77 GeV

$\Gamma(K^+K^-)/\Gamma(\pi^+\pi^-)$ $\Gamma_{118}/\Gamma_{112}$

The unused results here are redundant with $\Gamma(K^+K^-)/\Gamma(K^-\pi^+)$ and $\Gamma(\pi^+\pi^-)/\Gamma(K^-\pi^+)$ measurements by the same experiments.

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.75\pm0.15\pm0.16$	AITALA	98C	E791	π^- nucleus, 500 GeV
$2.53\pm0.46\pm0.19$	FRABETTI	94C	E687	γ Be $\overline{E}_\gamma=$ 220 GeV
$2.23\pm0.81\pm0.46$	ADAMOVICH	92	OMEG	π^- 340 GeV
$1.95\pm0.34\pm0.22$	ANJOS	91D	E691	Photoproduction
$2.5\ \pm0.7$	ALBRECHT	90C	ARG	$e^+e^-\approx$ 10 GeV
$2.35\pm0.37\pm0.28$	ALEXANDER	90	CLEO	e^+e^- 10.5–11 GeV

$\Gamma(K^0\overline{K}^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{119}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0120 ± 0.0033 OUR FIT Error includes scale factor of 1.3.					
0.0117 ± 0.0033 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.					
$0.0101\pm0.0022\pm0.0016$	26	ASNER	96B	CLE2	$e^+e^-\approx\ \Upsilon(4S)$
$0.039\ \pm0.013\ \pm0.013$	20	FRABETTI	94J	E687	γ Be $\overline{E}_\gamma$=220 GeV
$0.021\ ^{+0.011}_{-0.008}\ \pm0.002$	5	ALEXANDER	90	CLEO	e^+e^- 10.5–11 GeV



$\Gamma(K^0\overline{K}^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$

$\Gamma(K^0\overline{K}^0)/\Gamma(K^+K^-)$ $\Gamma_{119}/\Gamma_{118}$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.15±0.04 OUR FIT	Error includes scale factor of 1.2.				
0.24±0.16	4	[74] CUMALAT	88	SPEC	nN 0–800 GeV

[74] Includes a correction communicated to us by the authors of CUMALAT 88.

$\Gamma(K^0K^-\pi^+)/\Gamma(K^-\pi^+)$ Γ_{120}/Γ_{20}

VALUE	DOCUMENT ID		TECN	COMMENT
0.168±0.026 OUR FIT	Error includes scale factor of 1.1.			
0.16 ±0.06	[75] ANJOS	91	E691	γ Be 80–240 GeV

[75] The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(K^0K^-\pi^+)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{120}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.118±0.018 OUR FIT	Error includes scale factor of 1.1.				
0.119±0.021 OUR AVERAGE	Error includes scale factor of 1.3.				
0.108±0.019	61	AMMAR	91	CLEO	$e^+e^- \approx$ 10.5 GeV
0.16 ±0.03 ±0.02	39	ALBRECHT	90C	ARG	$e^+e^- \approx$ 10 GeV

$\Gamma(\overline{K}^*(892)^0K^0)/\Gamma(K^-\pi^+)$ Γ_{140}/Γ_{20}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.00^{+0.03}_{-0.00}$	[76] ANJOS	91	E691	γ Be 80–240 GeV

[76] The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(\overline{K}^*(892)^0K^0)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{140}/Γ_{22}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.029	90	AMMAR	91	CLEO	$e^+e^- \approx$ 10.5 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.03	90	ALBRECHT	90C	ARG	$e^+e^- \approx$ 10 GeV

$\Gamma(K^*(892)^+K^-)/\Gamma(K^-\pi^+)$ Γ_{141}/Γ_{20}

Unseen decay modes of the $K^*(892)^+$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.090±0.020 OUR FIT				
0.16 $^{+0.08}_{=0.06}$	[77] ANJOS	91	E691	γ Be 80–240 GeV

[77] The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(K^*(892)^+K^-)/\Gamma(\overline{K}^0\pi^+\pi^-)$ Γ_{141}/Γ_{22}

Unseen decay modes of the $K^*(892)^+$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.064±0.014 OUR FIT	Error includes scale factor of 1.1.				
0.058±0.014 OUR AVERAGE					
0.064±0.018	23	AMMAR	91	CLEO	$e^+e^- \approx$ 10.5 GeV
0.05 ±0.02 ±0.01	15	ALBRECHT	90C	ARG	$e^+e^- \approx$ 10 GeV

$\Gamma(K^0 K^- \pi^+ \text{ nonresonant})/\Gamma(K^- \pi^+)$ Γ_{123}/Γ_{20}

VALUE	DOCUMENT ID		TECN	COMMENT
0.06 ± 0.06	[78] ANJOS	91	E691	γ Be 80–240 GeV

[78] The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(\overline{K^0} K^+ \pi^-)/\Gamma(K^- \pi^+)$ Γ_{124}/Γ_{20}

VALUE	DOCUMENT ID		TECN	COMMENT
0.129 ± 0.025 OUR FIT				
0.10 ± 0.05	[79] ANJOS	91	E691	γ Be 80–240 GeV

[79] The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(\overline{K^0} K^+ \pi^-)/\Gamma(\overline{K^0} \pi^+ \pi^-)$ Γ_{124}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.091 ± 0.018 OUR FIT					
0.098 ± 0.020	55	AMMAR	91	CLEO	$e^+ e^- \approx$ 10.5 GeV

$\Gamma(K^*(892)^0 \overline{K^0})/\Gamma(K^- \pi^+)$ Γ_{142}/Γ_{20}

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.00^{+0.04}_{-0.00}$	[80] ANJOS	91	E691	γ Be 80–240 GeV

[80] The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(K^*(892)^0 \overline{K^0})/\Gamma(\overline{K^0} \pi^+ \pi^-)$ Γ_{142}/Γ_{22}

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.015	90	AMMAR	91	CLEO	$e^+ e^- \approx$ 10.5 GeV

$\Gamma(K^*(892)^- K^+)/\Gamma(K^- \pi^+)$ Γ_{143}/Γ_{20}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.00^{+0.03}_{-0.00}$	[81] ANJOS	91	E691	γ Be 80–240 GeV

[81] The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(K^*(892)^- K^+)/\Gamma(\overline{K^0} \pi^+ \pi^-)$ Γ_{143}/Γ_{22}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.034 ± 0.019	12	AMMAR	91	CLEO	$e^+ e^- \approx$ 10.5 GeV

$\Gamma(\overline{K^0} K^+ \pi^- \text{ nonresonant})/\Gamma(K^- \pi^+)$ Γ_{127}/Γ_{20}

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.10^{+0.06}_{-0.05}}$	[82] ANJOS	91	E691	γ Be 80–240 GeV

[82] The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

$\Gamma(K^+K^-\pi^0)/\Gamma(K^-\pi^+\pi^0)$ Γ_{128}/Γ_{30}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0095±0.0026	151	ASNER	96B	CLE2	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(K^0_S K^0_S \pi^0)/\Gamma_{total}$ Γ_{129}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
<0.00059	ASNER	96B	CLE2	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(\phi\pi^0)/\Gamma_{total}$ Γ_{144}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0014	90	ALBRECHT	94I	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(\phi\eta)/\Gamma_{total}$ Γ_{145}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0028	90	ALBRECHT	94I	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(\phi\omega)/\Gamma_{total}$ Γ_{146}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0021	90	ALBRECHT	94I	ARG	$e^+e^-\approx$ 10 GeV

$\Gamma(K^+K^-\pi^+\pi^-)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{130}/Γ_{38}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0334±0.0028 OUR AVERAGE					
0.0313±0.0037±0.0036	136	AITALA	98D	E791	π^- nucleus, 500 GeV
0.035 ±0.004 ±0.002	244	FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV
0.041 ±0.007 ±0.005	114	ALBRECHT	94I	ARG	$e^+e^-\approx$ 10 GeV
0.0314±0.010	89	AMMAR	91	CLEO	$e^+e^- \approx$ 10.5 GeV
$0.028\ ^{+0.008}_{-0.007}$		ANJOS	91	E691	γ Be 80–240 GeV

$\Gamma(\phi\pi^+\pi^-)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{147}/Γ_{38}

Unseen decay modes of the ϕ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.014 ±0.004 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.			
0.011 ±0.003		FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV
0.020 ±0.006 ±0.005	28	ALBRECHT	94I	ARG	$e^+e^-\approx$ 10 GeV
0.024 ±0.006	34	[83] AMMAR	91	CLEO	$e^+e^- \approx$ 10.5 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.0076\ ^{+0.0066}_{-0.0049}$	3	ANJOS	91	E691	γ Be 80–240 GeV

[83] AMMAR 91 measures $\phi\rho^0$, but notes that $\phi\rho^0$ dominates $\phi\pi^+\pi^-$. We put the measurement here to keep from having more $\phi\rho^0$ than $\phi\pi^+\pi^-$.



$\Gamma\left(\phi\,\pi^+\,\pi^-\right)/\Gamma\left(K^-\,\pi^+\,\pi^+\,\pi^-\right)$

$\Gamma\left(\phi\rho^0\right)/\Gamma\left(K^-\pi^+\pi^+\pi^-\right)$ Γ_{148}/Γ_{38}

Unseen decay modes of the ϕ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.008±0.004 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.			
0.02 ±0.009±0.008		AITALA	98D	E791	π^- nucleus, 500 GeV
0.005±0.003		FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx 200$ GeV
0.020±0.006±0.005	28	ALBRECHT	94I	ARG	$e^+e^- \approx 10$ GeV



$\Gamma\left(\phi\rho^0\right)/\Gamma\left(K^-\pi^+\pi^+\pi^-\right)$

$\Gamma\left(\phi\pi^+\pi^-\text{3-body}\right)/\Gamma\left(K^-\pi^+\pi^+\pi^-\right)$ Γ_{149}/Γ_{38}

Unseen decay modes of the ϕ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
0.009±0.004±0.005		AITALA	98D	E791	π^- nucleus, 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.006	90	FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV

$\Gamma\left(K^+K^-\rho^0\text{3-body}\right)/\Gamma\left(K^-\pi^+\pi^+\pi^-\right)$ Γ_{133}/Γ_{38}

VALUE	DOCUMENT ID		TECN	COMMENT
0.012 ±0.003	FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV

$\Gamma\left(K^*(892)^0K^-\pi^+ + \text{c.c.}\right)/\Gamma\left(K^-\pi^+\pi^+\pi^-\right)$ Γ_{150}/Γ_{38}

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
<0.01	90	[84]	AITALA	98D	E791	π^- nucleus, 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.017	90	[84]	FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV
$0.010^{+0.016}_{-0.010}$			ANJOS	91	E691	γ Be 80–240 GeV

[84] These upper limits are in conflict with values in the next two data blocks.

$\Gamma(K^*(892)^0 K^- \pi^+)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{151}/Γ_{38}

The $K^{*0} K^- \pi^+$ and $\overline{K}^{*0} K^+ \pi^-$ modes are distinguished by the charge of the pion in $D^*(2010)^\pm \to D^0 \pi^\pm$ decays. Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.043 \pm 0.014 \pm 0.009$	55	[85] ALBRECHT	94I	ARG	$e^+ e^- \approx 10$ GeV

[85] This ALBRECHT 94I value is in conflict with upper limits given above.

$\Gamma(\overline{K}^*(892)^0 K^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{152}/Γ_{38}

The $K^{*0} K^- \pi^+$ and $\overline{K}^{*0} K^+ \pi^-$ modes are distinguished by the charge of the pion in $D^*(2010)^\pm \to D^0 \pi^\pm$ decays. Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.023 \pm 0.013 \pm 0.009$	30	[86] ALBRECHT	94I	ARG	$e^+ e^- \approx 10$ GeV

[86] This ALBRECHT 94I value is in conflict with upper limits given above.

$\Gamma(K^*(892)^0 \overline{K}^*(892)^0)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{153}/Γ_{38}

Unseen decay modes of the $K^*(892)^0$ and $\overline{K}^*(892)^0$ are included.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
0.018 ± 0.007 OUR AVERAGE			Error includes scale factor of 1.2.			
0.016 ± 0.006			FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx 200$ GeV
$0.036 \,^{+0.020}_{-0.016}$		11	ANJOS	91	E691	γ Be 80–240 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.02	90		AITALA	98D	E791	π^- nucleus, 500 GeV
<0.033	90	[87]	AMMAR	91	CLEO	$e^+ e^- \approx 10.5$ GeV

[87] A corrected value (G. Moneti, private communication).

$\Gamma(K^+ K^- \pi^+ \pi^-$ non-$\phi)/\Gamma_{\text{total}}$ Γ_{136}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0017 ± 0.0005	[88] BARLAG	92C	ACCM	π^- Cu 230 GeV

[88] BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^+ K^- \pi^+ \pi^-$ nonresonant$)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{137}/Γ_{38}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.011	90	FRABETTI	95C	E687	γ Be, $\overline{E}_\gamma \approx 200$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.001 \,^{+0.011}_{-0.001}$		ANJOS	91	E691	γ Be 80–240 GeV

$\Gamma(K^0 \overline{K}^0 \pi^+ \pi^-)/\Gamma(\overline{K}^0 \pi^+ \pi^-)$ Γ_{138}/Γ_{22}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$0.126 \pm 0.038 \pm 0.030$	25	ALBRECHT	94I	ARG	$e^+ e^- \approx 10$ GeV

$\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{139}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0031±0.0020	[89] BARLAG	92C	ACCM	π^- Cu 230 GeV

[89] BARLAG 92C computes the branching fraction using topological normalization.

Radiative modes

$\Gamma(\rho^0\gamma)/\Gamma_{total}$ Γ_{154}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<2.4\times10^{-4}$	90	ASNER	98	CLE2

$\Gamma(\omega\gamma)/\Gamma_{total}$ Γ_{155}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<2.4\times10^{-4}$	90	ASNER	98	CLE2

$\Gamma(\phi\gamma)/\Gamma_{total}$ Γ_{156}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<1.9\times10^{-4}$	90	ASNER	98	CLE2

$\Gamma(\overline{K}^*(892)^0\gamma)/\Gamma_{total}$ Γ_{157}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<7.6\times10^{-4}$	90	ASNER	98	CLE2

Rare or forbidden modes

$\Gamma(K^+\ell^-\overline{\nu}_\ell(\text{via }\overline{D}^0))/\Gamma(K^-\ell^+\nu_\ell)$ Γ_{158}/Γ_8

This is a D^0-$\overline{D}^0$ mixing limit without the complications of possible doubly-Cabibbo-suppressed decays that occur when using hadronic modes. For the limits on $|m_{D_1^0} - m_{D_2^0}|$ and $(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma_{D^0}$ that come from the best mixing limit, see near the beginning of these D^0 Listings.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.005	90	[90] AITALA	96C	E791	π^- nucleus, 500 GeV

[90] AITALA 96C uses $D^{*+} \to D^0\pi^+$ (and charge conjugate) decays to identify the charm at production and $D^0 \to K^-\ell^+\nu_\ell$ (and charge conjugate) decays to identify the charm at decay.

$\Gamma(K^+\pi^-)/\Gamma(K^-\pi^+)$ Γ_{159}/Γ_{20}

The $D^0 \to K^+\pi^-$ decay can occur directly by doubly Cabibbo-suppressed (DCS) decay, or indirectly by $D^0 \to \overline{D}^0$ mixing followed by $\overline{D}^0 \to K^+\pi^-$ decay. The experiments here use the charge of the pion in $D^*(2010)^\pm \to (D^0 \text{ or } \overline{D}^0)\,\pi^\pm$ decay to tell whether a D^0 or a $\overline{D}^0$ was born. Some of the experiments can use the decay-time information to disentagle the two modes. Here, we list the DCS branching ratio; in the next data block we give the limits on the mixing ratio.

Some early limits have been omitted from this Listing; see our 1998 (EPJ **C3** 1) edition.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
0.0038 ±0.0008 OUR AVERAGE Error includes scale factor of 1.1.						
$0.00332\,^{+0.00063}_{-0.00065} \pm 0.00040$		45	[91] GODANG	00	CLE2	e^+e^-
$0.0068\ ^{+0.0034}_{-0.0033} \pm 0.0007$			[92] AITALA	98	E791	π^- nucleus, 500 GeV
$0.0184 \pm 0.0059 \pm 0.0034$		19	[93] BARATE	98W	ALEP	e^+e^- at Z^0
$0.0077 \pm 0.0025 \pm 0.0025$		19	[94] CINABRO	94	CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.011	90		[94] AMMAR	91	CLEO	$e^+e^- \approx$ 10.5 GeV
<0.015	90	1 ± 6	[95] ANJOS	88C	E691	Photoproduction
<0.014	90		[94] ALBRECHT	87K	ARG	e^+e^- 10 GeV

[91] This GODANG 00 result assumes no D^0-$\overline{D}^0$ mixing; the DCS ratio becomes $0.0048 \pm 0.0012 \pm 0.0004$ when mixing is allowed.

[92] This AITALA 98 result assumes no D^0-$\overline{D}^0$ mixing; the DCS ratio becomes $0.0090\,^{+0.0120}_{-0.0109} \pm 0.0044$ when mixing is allowed.

[93] BARATE 98W gets $0.0177\,^{+0.0060}_{-0.0056} \pm 0.0031$ for the DCS ratio when mixing is allowed, assuming no interference between the DCS and mixing amplitudes.

[94] CINABRO 94, AMMAR 91, and ALBRECHT 87K cannot distinguish between doubly Cabibbo-suppressed decay and D^0-$\overline{D}^0$ mixing.

[95] ANJOS 88C allows mixing but assumes no interference between the DCS and mixing amplitudes. When interference is allowed, the limit degrades to 0.049.

$\Gamma(K^+\pi^-$ (via $\overline{D}^0$)$)/\Gamma(K^-\pi^+)$ Γ_{160}/Γ_{20}

This is a D^0-$\overline{D}^0$ mixing limit. The experiments here (1) use the charge of the pion in $D^*(2010)^\pm \to (D^0$ or $\overline{D}^0)\ \pi^\pm$ decay to tell whether a D^0 or a $\overline{D}^0$ was born; and (2) use the decay-time distribution to disentangle doubly Cabibbo-suppressed decay and mixing. For the limits on $\left|m_{D_1^0} - m_{D_2^0}\right|$ and $(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma_{D^0}$ that come from the best mixing limit, see near the beginning of these D^0 Listings.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
<0.00041	95		[96] GODANG	00	CLE2	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.0092	95		[97] BARATE	98W	ALEP	e^+e^- at Z^0
<0.005	90	1 ± 4	[98] ANJOS	88C	E691	Photoproduction

[96] This GODANG 00 result assumes that the strong phase between $D^0 \to K^+\pi^-$ and $\overline{D}^0 \to K^+\pi^-$ is small, and limits only $D^0 \to \overline{D}^0$ transitions via off-shell intermediate states. The limit on transitions via on-shell intermediate states is 0.0017.

[97] This BARATE 98W result assumes no interference between the DCS and mixing amplitudes. When interference is allowed, the limit degrades to 0.036 (95%CL).

[98] This ANJOS 88C result assumes no interference between the DCS and mixing amplitudes. When interference is allowed, the limit degrades to 0.019. Combined with results on $K^\pm\pi^\mp\pi^+\pi^-$, the limit is, assuming no interference, 0.0037.

$\Gamma(K^+\pi^-\pi^+\pi^-)/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{161}/Γ_{38}

The $D^0 \to K^+\pi^-\pi^+\pi^-$ decay can occur directly by doubly Cabibbo-suppressed (DCS) decay, or indirectly by $D^0 \to \overline{D}^0$ mixing followed by $\overline{D}^0 \to K^+\pi^-\pi^+\pi^-$ decay. The experiments here use the charge of the pion in $D^*(2010)^\pm \to (D^0$ or $\overline{D}^0)\,\pi^\pm$ decay to tell whether a D^0 or a $\overline{D}^0$ was born. Some of the experiments can use the decay-time information to disentagle the two modes. Here, we list the DCS branching ratio; in the next data block we give the limits on the mixing ratio.

Some early limits have been omitted from this Listing; see our 1998 (EPJ **C3** 1) edition.

VALUE	*CL%*	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
$\mathbf{0.0025^{+0.0036}_{-0.0034} \pm 0.0003}$			[99] AITALA	98	E791	π^- nucleus, 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.018	90		[100] AMMAR	91	CLEO	$e^+e^- \approx 10.5$ GeV
<0.018	90	5 ± 12	[101] ANJOS	88C	E691	Photoproduction

[99] AITALA 98 uses the charge of the pion in $D^{*\pm} \to (D^0$ or $\overline{D}^0)\,\pi^\pm$ to tell whether a D^0 or a $\overline{D}^0$ was born. This result assumes no D^0-$\overline{D}^0$ mixing; it becomes $-0.0020^{+0.0117}_{-0.0106} \pm 0.0035$ when mixing is allowed and decay-time information is used to distinguish doubly Cabibbo-suppressed decays from mixing.

[100] AMMAR 91 cannot distinguish between doubly Cabibbo-suppressed decay and D^0-$\overline{D}^0$ mixing.

[101] ANJOS 88C uses decay-time information to distinguish doubly Cabibbo-suppressed (DCS) decays from D^0-$\overline{D}^0$ mixing. However, the result assumes no interference between the DCS and mixing amplitudes. When interference is allowed, the limit degrades to 0.033.

$\Gamma(K^+\pi^-\pi^+\pi^-\text{(via }\overline{D}^0\text{)})/\Gamma(K^-\pi^+\pi^+\pi^-)$ Γ_{162}/Γ_{38}

This is a D^0-$\overline{D}^0$ mixing limit. The experiments here (1) use the charge of the pion in $D^*(2010)^\pm \to (D^0$ or $\overline{D}^0)\,\pi^\pm$ decay to tell whether a D^0 or a $\overline{D}^0$ was born; and (2) use the decay-time distribution to disentangle doubly Cabibbo-suppressed decay and mixing. For the limits on $|m_{D_1^0} - m_{D_2^0}|$ and $(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma_{D^0}$ that come from the best mixing limit, see near the beginning of these D^0 Listings.

VALUE	*CL%*	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
<0.005	90	0 ± 4	[102] ANJOS	88C	E691	Photoproduction

[102] ANJOS 88C uses decay-time information to distinguish doubly Cabibbo-suppressed (DCS) decays from D^0-$\overline{D}^0$ mixing. However, the result assumes no interference between the DCS and mixing amplitudes. When interference is allowed, the limit degrades to 0.007. Combined with results on $K^\pm\pi^\mp$, the limit is, assuming no interference, 0.0037.

$\Gamma(K^+\pi^-\text{ or }K^+\pi^-\pi^+\pi^-\text{(via }\overline{D}^0\text{)})/\Gamma(K^-\pi^+\text{ or }K^-\pi^+\pi^+\pi^-)$ Γ_{163}/Γ_0

This is a D^0-$\overline{D}^0$ mixing limit. For the limits on $|m_{D_1^0} - m_{D_2^0}|$ and $(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma_{D^0}$ that come from the best mixing limit, see near the beginning of these D^0 Listings.

VALUE	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<0.0085	90	[103] AITALA	98	E791	π^- nucleus, 500 GeV

[103] AITALA 98 uses decay-time information to distinguish doubly Cabibbo-suppressed decays from D^0-$\overline{D}^0$ mixing. The fit allows interference between the two amplitudes, and also allows *CP* violation in this term. The central value obtained is $0.0039^{+0.0036}_{-0.0032} \pm 0.0016$. When interference is disallowed, the result becomes $0.0021 \pm 0.0009 \pm 0.0002$.

$\Gamma(\mu^-$ anything (via $\overline{D}^0))/\Gamma(\mu^+$ anything) Γ_{164}/Γ_2

This is a D^0-$\overline{D}^0$ mixing limit. See the somewhat better limits above.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0056	90	LOUIS	86	SPEC	π^- W 225 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.012	90	BENVENUTI	85	CNTR	μ C, 200 GeV
<0.044	90	BODEK	82	SPEC	π^-, p Fe $\rightarrow$ D^0

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_{165}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<6.2 \times 10^{-6}$	90		AITALA	99G	E791	π^- N 500 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8.19 \times 10^{-6}$	90		PRIPSTEIN	00	E789	p nucleus, 800 GeV
$<1.3 \times 10^{-5}$	90	0	FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$
$<1.3 \times 10^{-4}$	90		ADLER	88	MRK3	$e^+ e^-$ 3.77 GeV
$<1.7 \times 10^{-4}$	90	7	ALBRECHT	88G	ARG	$e^+ e^-$ 10 GeV
$<2.2 \times 10^{-4}$	90	8	HAAS	88	CLEO	$e^+ e^-$ 10 GeV

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{166}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<4.1 \times 10^{-6}$	90		ADAMOVICH	97	BEAT	π^- Cu, W 350 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.56 \times 10^{-5}$	90		PRIPSTEIN	00	E789	p nucleus, 800 GeV
$<5.2 \times 10^{-6}$	90		AITALA	99G	E791	π^- N 500 GeV
$<4.2 \times 10^{-6}$	90		ALEXOPOU...	96	E771	p Si, 800 GeV
$<3.4 \times 10^{-5}$	90	1	FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$
$<7.6 \times 10^{-6}$	90	0	ADAMOVICH	95	BEAT	See ADAMOVICH 97
$<4.4 \times 10^{-5}$	90	0	KODAMA	95	E653	π^- emulsion 600 GeV
$<3.1 \times 10^{-5}$	90	[104]	MISHRA	94	E789	-4.1 ± 4.8 events
$<7.0 \times 10^{-5}$	90	3	ALBRECHT	88G	ARG	$e^+ e^-$ 10 GeV
$<1.1 \times 10^{-5}$	90		LOUIS	86	SPEC	π^- W 225 GeV
$<3.4 \times 10^{-4}$	90		AUBERT	85	EMC	Deep inelast. μ^- N

[104] Here MISHRA 94 uses "the statistical approach advocated by the PDG." For an alternate approach, giving a limit of 9×10^{-6} at 90% confidence level, see the paper.

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{167}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<4.5 \times 10^{-5}$	90	0	FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma_{total}$ Γ_{168}/Γ

A test for the ΔC=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.8 \times 10^{-4}$	90	2	KODAMA	95	E653	π^- emulsion 600 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<5.4 \times 10^{-4}$	90	3	FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\eta e^+ e^-)/\Gamma_{total}$ Γ_{169}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.1 \times 10^{-4}$	90	0	FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\eta \mu^+ \mu^-)/\Gamma_{total}$ Γ_{170}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<5.3 \times 10^{-4}$	90	0	FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\rho^0 e^+ e^-)/\Gamma_{total}$ Γ_{171}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.0 \times 10^{-4}$	90	2	[105] FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4.5 \times 10^{-4}$	90	2	HAAS	88	CLEO	$e^+ e^-$ 10 GeV

[105] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 1.8 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(\rho^0 \mu^+ \mu^-)/\Gamma_{total}$ Γ_{172}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<2.3 \times 10^{-4}$	90	0	KODAMA	95	E653	π^- emulsion 600 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4.9 \times 10^{-4}$	90	1	[106] FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$
$<8.1 \times 10^{-4}$	90	5	HAAS	88	CLEO	$e^+ e^-$ 10 GeV

[106] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 4.5 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(\omega e^+ e^-)/\Gamma_{total}$ Γ_{173}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.8 \times 10^{-4}$	90	1	[107] FREYBERGER	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$

[107] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 2.7 \times 10^{-4}$ using a photon pole amplitude model.

$\Gamma(\omega\mu^+\mu^-)/\Gamma_{total}$ Γ_{174}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<8.3\times10^{-4}$	90	0	[108] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[108] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $<6.5\times10^{-4}$ using a photon pole amplitude model.

$\Gamma(\phi e^+e^-)/\Gamma_{total}$ Γ_{175}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<5.2\times10^{-5}$	90	2	[109] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[109] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $<7.6\times10^{-5}$ using a photon pole amplitude model.

$\Gamma(\phi\mu^+\mu^-)/\Gamma_{total}$ Γ_{176}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<4.1\times10^{-4}$	90	0	[110] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[110] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $<2.4\times10^{-4}$ using a photon pole amplitude model.

$\Gamma(\overline{K}^0e^+e^-)/\Gamma_{total}$ Γ_{177}/Γ

Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.1\times10^{-4}$	90	0	FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<1.7\times10^{-3}$	90		ADLER	89C	MRK3	e^+e^- 3.77 GeV

$\Gamma(\overline{K}^0\mu^+\mu^-)/\Gamma_{total}$ Γ_{178}/Γ

Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<2.6\times10^{-4}$	90	2	KODAMA	95	E653	π^- emulsion 600 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<6.7\times10^{-4}$	90	1	FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

$\Gamma(\overline{K}^*(892)^0e^+e^-)/\Gamma_{total}$ Γ_{179}/Γ

Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.4\times10^{-4}$	90	1	[111] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[111] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $<2.0\times10^{-4}$ using a photon pole amplitude model.

$\Gamma(\overline{K}^*(892)^0\mu^+\mu^-)/\Gamma_{total}$ Γ_{180}/Γ

Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.18\times10^{-3}$	90	1	[112] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[112] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $<1.0\times10^{-3}$ using a photon pole amplitude model.

$\Gamma(\pi^+\pi^-\pi^0\mu^+\mu^-)/\Gamma_{total}$ Γ_{181}/Γ

A test for the ΔC=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<8.1\times10^{-4}$	90	1	KODAMA	95	E653	π^- emulsion 600 GeV

$\Gamma(\mu^\pm e^\mp)/\Gamma_{total}$ Γ_{182}/Γ

A test of lepton family number conservation.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$< 8.1\times10^{-6}$	90		AITALA	99G	E791	π^- N 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$< 1.72\times10^{-5}$	90		PRIPSTEIN	00	E789	p nucleus, 800 GeV
$< 1.9\times10^{-5}$	90	2	[113] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$
$< 1.0\times10^{-4}$	90	4	ALBRECHT	88G	ARG	e^+e^- 10 GeV
$< 2.7\times10^{-4}$	90	9	HAAS	88	CLEO	e^+e^- 10 GeV
$< 1.2\times10^{-4}$	90		BECKER	87C	MRK3	e^+e^- 3.77 GeV
$< 9\times10^{-4}$	90		PALKA	87	SILI	200 GeV πp
$<21\times10^{-4}$	90	0	[114] RILES	87	MRK2	e^+e^- 29 GeV

[113] This is the corrected result given in the erratum to FREYBERGER 96.

[114] RILES 87 assumes B($D\rightarrow\ K\pi$) = 3.0% and has production model dependency.

$\Gamma(\pi^0 e^\pm\mu^\mp)/\Gamma_{total}$ Γ_{183}/Γ

A test of lepton family number conservation. The value is for the sum of the two charge states.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<8.6\times10^{-5}$	90	2	FREYBERGER 96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

$\Gamma(\eta e^\pm\mu^\mp)/\Gamma_{total}$ Γ_{184}/Γ

A test of lepton family number conservation. The value is for the sum of the two charge states.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.0\times10^{-4}$	90	0	FREYBERGER 96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

$\Gamma(\rho^0 e^\pm\mu^\mp)/\Gamma_{total}$ Γ_{185}/Γ

A test of lepton family number conservation. The value is for the sum of the two charge states.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<4.9\times10^{-5}$	90	0	[115] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[115] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $< 5.0\times10^{-5}$ using a photon pole amplitude model.

$\Gamma(\omega e^\pm\mu^\mp)/\Gamma_{total}$ Γ_{186}/Γ

A test of lepton family number conservation. The value is for the sum of the two charge states.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.2\times10^{-4}$	90	0	[116] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[116] This FREYBERGER 96 limit is obtained using a phase-space model. The same limit is obtained using a photon pole amplitude model.

$\Gamma(\phi e^{\pm}\mu^{\mp})/\Gamma_{total}$ Γ_{187}/Γ

A test of lepton family number conservation. The value is for the sum of the two charge states.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<3.4\times10^{-5}$	90	0	[117] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[117] This FREYBERGER 96 limit is obtained using a phase-space model. The limit changes to $<3.3\times10^{-5}$ using a photon pole amplitude model.

$\Gamma(\overline{K}^0 e^{\pm}\mu^{\mp})/\Gamma_{total}$ Γ_{188}/Γ

A test of lepton family number conservation. The value is for the sum of the two charge states.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.0\times10^{-4}$	90	0	FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

$\Gamma(\overline{K}^*(892)^0 e^{\pm}\mu^{\mp})/\Gamma_{total}$ Γ_{189}/Γ

A test of lepton family number conservation. The value is for the sum of the two charge states.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.0\times10^{-4}$	90	0	[118] FREYBERGER	96	CLE2	$e^+e^-\approx\ \Upsilon(4S)$

[118] This FREYBERGER 96 limit is obtained using a phase-space model. The same limit is obtained using a photon pole amplitude model.

D^0 *CP*-VIOLATING DECAY-RATE ASYMMETRIES

$A_{CP}(K^+K^-)$ in $D^0, \overline{D}^0 \to K^+K^-$

This is the difference between D^0 and $\overline{D}^0$ partial widths for these modes divided by the sum of the widths. The D^0 and $\overline{D}^0$ are distinguished by the charge of the parent D^*: $D^{*+} \to D^0\pi^+$ and $D^{*-} \to D^0\pi^-$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.009 ± 0.021 OUR AVERAGE					
$-0.001\pm0.022\pm0.015$	3330	[119] LINK	00B	FOCS	
$-0.010\pm0.049\pm0.012$	609	[119] AITALA	98C	E791	$-0.093 < A_{CP} < +0.073$ (90% CL)
$+0.080\pm0.061$		BARTELT	95	CLE2	$-0.022 < A_{CP} < +0.18$ (90%CL)
$+0.024\pm0.084$		[119] FRABETTI	94I	E687	$-0.11 < A_{CP} < +0.16$ (90% CL)

[119] FRABETTI 94I, AITALA 98C, and LINK 00B measure $N(D^0 \to K^+K^-)/N(D^0 \to K^-\pi^+)$, the ratio of numbers of events observed, and similarly for the $\overline{D}^0$.

$A_{CP}(\pi^+\pi^-)$ in $D^0, \overline{D}^0 \to \pi^+\pi^-$

This is the difference between D^0 and $\overline{D}^0$ partial widths for these modes divided by the sum of the widths. The D^0 and $\overline{D}^0$ are distinguished by the charge of the parent D^*: $D^{*+} \to D^0\pi^+$ and $D^{*-} \to D^0\pi^-$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$0.03\ \pm0.04$ OUR AVERAGE					
$+0.048\pm0.039\pm0.025$	1177	[120] LINK	00B	FOCS	
$-0.049\pm0.078\pm0.030$	343	[120] AITALA	98C	E791	$-0.186 < A_{CP} < +0.088$ (90% CL)

[120] AITALA 98C and LINK 00B measure $N(D^0 \to \pi^+\pi^-)/N(D^0 \to K^-\pi^+)$, the ratio of numbers of events observed, and similarly for the $\overline{D}^0$.

$A_{CP}(K^0_S\phi)$ in $D^0, \overline{D}^0 \to K^0_S\phi$

This is the difference between D^0 and $\overline{D}^0$ partial widths for these modes divided by the sum of the widths. The D^0 and $\overline{D}^0$ are distinguished by the charge of the parent D^*: $D^{*+} \to D^0\pi^+$ and $D^{*-} \to D^0\pi^-$.

VALUE	DOCUMENT ID		TECN	COMMENT
$=$ **0.028 ± 0.094**	BARTELT	95	CLE2	$-0.182 < A_{CP} < +0.126$ (90%CL)

$A_{CP}(K^0_S\pi^0)$ in $D^0, \overline{D}^0 \to K^0_S\pi^0$

This is the difference between D^0 and $\overline{D}^0$ partial widths for these modes divided by the sum of the widths. The D^0 and $\overline{D}^0$ are distinguished by the charge of the parent D^*: $D^{*+} \to D^0\pi^+$ and $D^{*-} \to D^0\pi^-$.

VALUE	DOCUMENT ID		TECN	COMMENT
-0.018 ± 0.030	BARTELT	95	CLE2	$-0.067 < A_{CP} < +0.031$ (90%CL)

$A_{CP}(K^\pm\pi^\mp)$ in $D^0 \to K^+\pi^-, \overline{D}^0 \to K^-\pi^+$

This is the difference between D^0 and $\overline{D}^0$ partial widths for these modes divided by the sum of the widths. The D^0 and $\overline{D}^0$ are distinguished by the charge of the parent D^*: $D^{*+} \to D^0\pi^+$ and $D^{*-} \to D^0\pi^-$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$+0.02^{+0.19}_{-0.20} \pm 0.01$	45	[121] GODANG	00	CLE2	$-0.43 < A_{CP} < +0.34$ (95%CL)

[121] This GODANG 00 result assumes no D^0-$\overline{D}^0$ mixing; it becomes $-0.01^{+0.16}_{-0.17} \pm 0.01$ when mixing is allowed.

D^0 PRODUCTION CROSS SECTION AT $\psi(3770)$

A compilation of the cross sections for the direct production of D^0 mesons at or near the $\psi(3770)$ peak in e^+e^- production.

VALUE (nanobarns)	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.8 ±0.5 ±0.6	[122] ADLER	88C	MRK3	e^+e^- 3.768 GeV
7.3 ±1.3	[123] PARTRIDGE	84	CBAL	e^+e^- 3.771 GeV
8.00 ± 0.95 ± 1.21	[124] SCHINDLER	80	MRK2	e^+e^- 3.771 GeV
11.5 ±2.5	[125] PERUZZI	77	MRK1	e^+e^- 3.774 GeV

[122] This measurement compares events with one detected D to those with two detected D mesons, to determine the the absolute cross section. ADLER 88C find the ratio of cross sections (neutral to charged) to be 1.36 ± 0.23 ± 0.14.

[123] This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. PARTRIDGE 84 measures 6.4 ± 1.15 nb for the cross section. We take the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and we assume that the $\psi(3770)$ is an isosinglet to evaluate the cross sections. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

[124] This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. SCHINDLER 80 assume the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and that the $\psi(3770)$ is an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

[125] This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. The phase space division of neutral and charged D mesons in $\psi(3770)$ decay

is taken to be 1.33, and $\psi(3770)$ is assumed to be an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction. We exclude this measurement from the average because of uncertainties in the contamination from τ lepton pairs. Also see RAPIDIS 77.

D^0 REFERENCES

BAI	00C	PR D62 052001	J.Z. Bai *et al.*	(BEPC BES Collab.)
GODANG	00	PRL 84 5038	R. Godang *et al.*	(CLEO Collab.)
JUN	00	PRL 84 1857	S.Y. Jun *et al.*	(FNAL SELEX Collab.)
LINK	00	PL B485 62	J.M. Link *et al.*	(FNAL FOCUS Collab.)
LINK	00B	PL B491 232	J.M. Link *et al.*	(FNAL FOCUS Collab.)
Also	00D	PL B495 443 (errata)	J.M. Link *et al.*	(FNAL FOCUS Collab.)
PDG	00	EPJ C15 1	D.E. Groom *et al.*	
PRIPSTEIN	00	PR D61 032005	D. Pripstein *et al.*	(FNAL E789 Collab.)
AITALA	99E	PRL 83 32	E.M. Aitala *et al.*	(FNAL E791 Collab.)
AITALA	99G	PL B462 401	E.M. Aitala *et al.*	(FNAL E791 Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini *et al.*	(CLEO Collab.)
AITALA	98	PR D57 13	E.M. Aitala *et al.*	(FNAL E791 Collab.)
AITALA	98C	PL B421 405	E.M. Aitala *et al.*	(FNAL E791 Collab.)
AITALA	98D	PL B423 185	E.M. Aitala *et al.*	(FNAL E791 Collab.)
ARTUSO	98	PRL 80 3193	M. Artuso *et al.*	(CLEO Collab.)
ASNER	98	PR D58 092001	D.M. Asner *et al.*	(CLEO Collab.)
BARATE	98W	PL B436 211	R. Barate *et al.*	(ALEPH Collab.)
COAN	98	PRL 80 1150	T.E. Coan *et al.*	(CLEO Collab.)
ADAMOVICH	97	PL B408 469	M.I. Adamovich *et al.*	(CERN BEATRICE Collab.)
BARATE	97C	PL B403 367	R. Barate *et al.*	(ALEPH Collab.)
AITALA	96C	PRL 77 2384	E.M. Aitala *et al.*	(FNAL E791 Collab.)
ALBRECHT	96C	PL B374 249	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXOPOU...	96	PRL 77 2380	T. Alexopoulos *et al.*	(FNAL E771 Collab.)
ASNER	96B	PR D54 4211	D.M. Asner *et al.*	(CLEO Collab.)
BARISH	96	PL B373 334	B.C. Barish *et al.*	(CLEO Collab.)
FRABETTI	96B	PL B382 312	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FREYBERGER	96	PRL 76 3065	A. Freyberger *et al.*	(CLEO Collab.)
Also	96B	PRL 77 2147 (errata)	A. Freyberger *et al.*	(CLEO Collab.)
KUBOTA	96B	PR D54 2994	Y. Kubota *et al.*	(CLEO Collab.)
ADAMOVICH	95	PL B353 563	M.I. Adamovich *et al.*	(CERN BEATRICE Collab.)
BARTELT	95	PR D52 4860	J.E. Bartelt *et al.*	(CLEO Collab.)
BUTLER	95	PR D52 2656	F. Butler *et al.*	(CLEO Collab.)
FRABETTI	95C	PL B354 486	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	95G	PL B364 127	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama *et al.*	(FNAL E653 Collab.)
ALBRECHT	94	PL B324 249	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	94F	PL B340 125	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	94I	ZPHY C64 375	H. Albrecht *et al.*	(ARGUS Collab.)
CINABRO	94	PRL 72 1406	D. Cinabro *et al.*	(CLEO Collab.)
FRABETTI	94C	PL B321 295	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	94D	PL B323 459	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	94G	PL B331 217	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	94I	PR D50 R2953	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	94J	PL B340 254	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
KODAMA	94	PL B336 605	K. Kodama *et al.*	(FNAL E653 Collab.)
MISHRA	94	PR D50 R9	C.S. Mishra *et al.*	(FNAL E789 Collab.)
AKERIB	93	PRL 71 3070	D.S. Akerib *et al.*	(CLEO Collab.)
ALBRECHT	93D	PL B308 435	H. Albrecht *et al.*	(ARGUS Collab.)
ANJOS	93	PR D48 56	J.C. Anjos *et al.*	(FNAL E691 Collab.)
BEAN	93C	PL B317 647	A. Bean *et al.*	(CLEO Collab.)
FRABETTI	93I	PL B315 203	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
KODAMA	93B	PL B313 260	K. Kodama *et al.*	(FNAL E653 Collab.)
PROCARIO	93B	PR D48 4007	M. Procario *et al.*	(CLEO Collab.)
SELEN	93	PRL 71 1973	M.A. Selen *et al.*	(CLEO Collab.)
ADAMOVICH	92	PL B280 163	M.I. Adamovich *et al.*	(CERN WA82 Collab.)
ALBRECHT	92P	ZPHY C56 7	H. Albrecht *et al.*	(ARGUS Collab.)
ANJOS	92B	PR D46 R1	J.C. Anjos *et al.*	(FNAL E691 Collab.)
ANJOS	92C	PR D46 1941	J.C. Anjos *et al.*	(FNAL E691 Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag *et al.*	(ACCMOR Collab.)
Also	90D	ZPHY C48 29	S. Barlag *et al.*	(ACCMOR Collab.)
COFFMAN	92B	PR D45 2196	D.M. Coffman *et al.*	(Mark III Collab.)
Also	90	PRL 64 2615	J. Adler *et al.*	(Mark III Collab.)

FRABETTI	92	PL B281 167	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	92B	PL B286 195	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
ALVAREZ	91B	ZPHY C50 11	M.P. Alvarez *et al.*	(CERN NA14/2 Collab.)
AMMAR	91	PR D44 3383	R. Ammar *et al.*	(CLEO Collab.)
ANJOS	91	PR D43 R635	J.C. Anjos *et al.*	(FNAL-TPS Collab.)
ANJOS	91D	PR D44 R3371	J.C. Anjos *et al.*	(FNAL-TPS Collab.)
BAI	91	PRL 66 1011	Z. Bai *et al.*	(Mark III Collab.)
COFFMAN	91	PL B263 135	D.M. Coffman *et al.*	(Mark III Collab.)
CRAWFORD	91B	PR D44 3394	G. Crawford *et al.*	(CLEO Collab.)
DECAMP	91J	PL B266 218	D. Decamp *et al.*	(ALEPH Collab.)
FRABETTI	91	PL B263 584	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
KINOSHITA	91	PR D43 2836	K. Kinoshita *et al.*	(CLEO Collab.)
KODAMA	91	PRL 66 1819	K. Kodama *et al.*	(FNAL E653 Collab.)
ALBRECHT	90C	ZPHY C46 9	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	90	PRL 65 1184	J. Alexander *et al.*	(CLEO Collab.)
ALEXANDER	90B	PRL 65 1531	J. Alexander *et al.*	(CLEO Collab.)
ALVAREZ	90	ZPHY C47 539	M.P. Alvarez *et al.*	(CERN NA14/2 Collab.)
ANJOS	90D	PR D42 2414	J.C. Anjos *et al.*	(FNAL E691 Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag *et al.*	(ACCMOR Collab.)
ADLER	89	PRL 62 1821	J. Adler *et al.*	(Mark III Collab.)
ADLER	89C	PR D40 906	J. Adler *et al.*	(Mark III Collab.)
ALBRECHT	89D	ZPHY C43 181	H. Albrecht *et al.*	(ARGUS Collab.)
ANJOS	89F	PRL 62 1587	J.C. Anjos *et al.*	(FNAL E691 Collab.)
ABACHI	88	PL B205 411	S. Abachi *et al.*	(HRS Collab.)
ADLER	88	PR D37 2023	J. Adler *et al.*	(Mark III Collab.)
ADLER	88C	PRL 60 89	J. Adler *et al.*	(Mark III Collab.)
ALBRECHT	88G	PL B209 380	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	88I	PL B210 267	H. Albrecht *et al.*	(ARGUS Collab.)
AMENDOLIA	88	EPL 5 407	S.R. Amendolia *et al.*	(NA1 Collab.)
ANJOS	88C	PRL 60 1239	J.C. Anjos *et al.*	(FNAL E691 Collab.)
BORTOLETTO	88	PR D37 1719	D. Bortoletto *et al.*	(CLEO Collab.)
Also	89D	PR D39 1471 erratum	D. Bortoletto *et al.*	(CLEO Collab.)
CUMALAT	88	PL B210 253	J.P. Cumalat *et al.*	(E-400 Collab.)
HAAS	88	PRL 60 1614	P. Haas *et al.*	(CLEO Collab.)
RAAB	88	PR D37 2391	J.R. Raab *et al.*	(FNAL E691 Collab.)
ADAMOVICH	87	EPL 4 887	M.I. Adamovich *et al.*	
ADLER	87	PL B196 107	J. Adler *et al.*	(Mark III Collab.)
AGUILAR-...	87D	PL B193 140	M. Aguilar-Benitez *et al.*	(LEBC-EHS Collab.)
Also	88B	ZPHY C40 321	M. Aguilar-Benitez *et al.*	(LEBC-EHS Collab.)
AGUILAR-...	87E	ZPHY C36 551	M. Aguilar-Benitez *et al.*	(LEBC-EHS Collab.)
Also	88B	ZPHY C40 321	M. Aguilar-Benitez *et al.*	(LEBC-EHS Collab.)
AGUILAR-...	87F	ZPHY C36 559	M. Aguilar-Benitez *et al.*	(LEBC-EHS Collab.)
Also	88	ZPHY C38 520 erratum	M. Aguilar-Benitez *et al.*	(LEBC-EHS Collab.)
ALBRECHT	87E	ZPHY C33 359	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87K	PL B199 447	H. Albrecht *et al.*	(ARGUS Collab.)
BARLAG	87B	ZPHY C37 17	S. Barlag *et al.*	(ACCMOR Collab.)
BECKER	87C	PL B193 147	J.J. Becker *et al.*	(Mark III Collab.)
Also	87D	PL B198 590 erratum	J.J. Becker *et al.*	(Mark III Collab.)
CSORNA	87	PL B191 318	S.E. Csorna *et al.*	(CLEO Collab.)
PALKA	87	PL B189 238	H. Palka *et al.*	(ACCMOR Collab.)
RILES	87	PR D35 2914	K. Riles *et al.*	(Mark II Collab.)
ABE	86	PR D33 1	K. Abe *et al.*	
BAILEY	86	ZPHY C30 51	R. Bailey *et al.*	(ACCMOR Collab.)
BEBEK	86	PRL 56 1893	C. Bebek *et al.*	(CLEO Collab.)
GLADNEY	86	PR D34 2601	L. Gladney *et al.*	(Mark II Collab.)
LOUIS	86	PRL 56 1027	W.C. Louis *et al.*	(PRIN, CHIC, ISU)
USHIDA	86B	PRL 56 1771	N. Ushida *et al.*	(AICH, FNAL, KOBE, SEOU+)
ALBRECHT	85B	PL 158B 525	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	85F	PL 150B 235	H. Albrecht *et al.*	(ARGUS Collab.)
AUBERT	85	PL 155B 461	J.J. Aubert *et al.*	(EMC Collab.)
BAILEY	85	ZPHY C28 357	R. Bailey *et al.*	(ABCCMR Collab.)
BALTRUSAIT...	85B	PRL 54 1976	R.M. Baltrusaitis *et al.*	(Mark III Collab.)
BALTRUSAIT...	85E	PRL 55 150	R.M. Baltrusaitis *et al.*	(Mark III Collab.)
BENVENUTI	85	PL 158B 531	A.C. Benvenuti *et al.*	(BCDMS Collab.)
ADAMOVICH	84B	PL 140B 123	M.I. Adamovich *et al.*	(CERN WA58 Collab.)
DERRICK	84	PRL 53 1971	M. Derrick *et al.*	(HRS Collab.)
PARTRIDGE	84	Thesis CALT-68-1150	R.A. Partridge	(Crystal Ball Collab.)
SUMMERS	84	PRL 52 410	D.J. Summers *et al.*	(UCSB, CARL, COLO+)

BAILEY	83B	PL 132B 237	R. Bailey *et al.*	(ACCMOR Collab.)
BODEK	82	PL 113B 82	A. Bodek *et al.*	(ROCH, CIT, CHIC, FNAL+)
FIORINO	81	LNC 30 166	A. Fiorino *et al.*	
SCHINDLER	81	PR D24 78	R.H. Schindler *et al.*	(Mark II Collab.)
TRILLING	81	PRPL 75 57	G.H. Trilling	(LBL, UCB) J
ASTON	80E	PL 94B 113	D. Aston *et al.*	(BONN, CERN, EPOL, GLAS+)
AVERY	80	PRL 44 1309	P. Avery *et al.*	(ILL, FNAL, COLU)
SCHINDLER	80	PR D21 2716	R.H. Schindler *et al.*	(Mark II Collab.)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents *et al.*	(NOVO)
Also	81	SJNP 34 814	A.A. Zholents *et al.*	(NOVO)
		Translated from YAF 34 1471.		
ABRAMS	79D	PRL 43 481	G.S. Abrams *et al.*	(Mark II Collab.)
ATIYA	79	PRL 43 414	M.S. Atiya *et al.*	(COLU, ILL, FNAL)
BALTAY	78C	PRL 41 73	C. Baltay *et al.*	(COLU, BNL)
VUILLEMIN	78	PRL 41 1149	V. Vuillemin *et al.*	(Mark I Collab.)
GOLDHABER	77	PL 69B 503	G. Goldhaber *et al.*	(Mark I Collab.)
PERUZZI	77	PRL 39 1301	I. Peruzzi *et al.*	(Mark I Collab.)
PICCOLO	77	PL 70B 260	M. Piccolo *et al.*	(Mark I Collab.)
RAPIDIS	77	PRL 39 526	P.A. Rapidis *et al.*	(Mark I Collab.)
GOLDHABER	76	PRL 37 255	G. Goldhaber *et al.*	(Mark I Collab.)

OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	J.D. Richman, P.R. Burchat	(UCSB, STAN)
ROSNER	95	CNPP 21 369	J. Rosner	(CHIC)