$D_s^\pm$

was $F^\pm$

$I(J^P) = 0(0^-)$

The angular distributions of the decays of the ϕ and $\overline{K}^*(892)^0$ in the $\phi\pi^+$ and $K^+\overline{K}^*(892)^0$ modes strongly indicate that the spin is zero. The parity given is that expected of a $c\overline{s}$ ground state.

$D_s^\pm$ MASS

The fit includes $D^\pm$, D^0, $D_s^\pm$, $D^{*\pm}$, D^{*0}, and $D_s^{*\pm}$ mass and mass difference measurements. Measurements of the $D_s^\pm$ mass with an error greater than 10 MeV are omitted from the fit and average. A number of early measurements have been omitted altogether.

VALUE (MeV)	*EVTS*	*DOCUMENT ID*	*TECN*	*COMMENT*
1968.6± 0.6 OUR FIT Error includes scale factor of 1.1.				
1969.0± 1.4 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
1967.0± 1.0± 1.0	54	BARLAG	90C ACCM	π^- Cu 230 GeV
1969.3± 1.4± 1.4		ALBRECHT	88 ARG	e^+e^- 9.4–10.6 GeV
1972.7± 1.5± 1.0	21	BECKER	87B SILI	200 GeV π,K,p
1972.4± 3.7± 3.7	27	BLAYLOCK	87 MRK3	e^+e^- 4.14 GeV
1963 ± 3 ± 3	30	DERRICK	85B HRS	e^+e^- 29 GeV
1970 ± 5 ± 5	104	CHEN	83C CLEO	e^+e^- 10.5 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1968.3± 0.7± 0.7	290	[1] ANJOS	88 E691	Photoproduction
1980 ±15	6	USHIDA	86 EMUL	ν wideband
1973.6± 2.6± 3.0	163	ALBRECHT	85D ARG	e^+e^- 10 GeV
1948 ±28 ±10	65	AIHARA	84D TPC	e^+e^- 29 GeV
1975 ± 9 ±10	49	ALTHOFF	84 TASS	e^+e^- 14–25 GeV
1975 ± 4	3	BAILEY	84 ACCM	hadron$^+$ Be $\rightarrow$ $\phi\pi^+$ X

[1] ANJOS 88 enters the fit via $m_{D_s^\pm} - m_{D^\pm}$ (see below).



$D_s^{\pm}$ mass (MeV)

$m_{D_s^{\pm}} - m_{D^{\pm}}$

The fit includes $D^{\pm}$, D^0, $D_s^{\pm}$, $D^{*\pm}$, D^{*0}, and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
99.2±0.5 OUR FIT Error includes scale factor of 1.1.					
99.2±0.5 OUR AVERAGE					
99.5±0.6±0.3		BROWN	94	CLE2	$e^+e^- \approx \Upsilon(4S)$
98.5±1.5	555	CHEN	89	CLEO	e^+e^- 10.5 GeV
99.0±0.8	290	ANJOS	88	E691	Photoproduction

$D_s^{\pm}$ MEAN LIFE

Measurements with an error greater than 0.2×10^{-12} s or with fewer than 100 events are omitted from the average.

VALUE (10^{-12} s)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.496 $^{+0.010}_{-0.009}$ OUR AVERAGE					
0.518 ±0.014 ±0.007	1662	AITALA	99	E791	π^- nucleus, 500 GeV
0.4863±0.0150 $^{+0.0049}_{-0.0051}$	2167	[2] BONVICINI	99	CLE2	$e^+e^- \approx \Upsilon(4S)$
0.475 ±0.020 ±0.007	900	FRABETTI	93F	E687	γ Be, $D_s^+ \rightarrow \phi\pi^+$

0.50	± 0.06	± 0.03	104	FRABETTI	90	E687	γ Be, $\phi\pi^+$
0.56	$^{+0.13}_{-0.12}$	± 0.08	144	ALBRECHT	88I	ARG	e^+e^- 10 GeV
0.47	± 0.04	± 0.02	228	RAAB	88	E691	Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •							
0.33	$^{+0.12}_{-0.08}$	± 0.03	15	ALVAREZ	90	NA14	γ, $D_s^+ \to \phi\pi^+$
0.469	$^{+0.102}_{-0.086}$		54	[3] BARLAG	90C	ACCM	π^- Cu 230 GeV
0.31	$^{+0.24}_{-0.20}$	± 0.05	18	AVERILL	89	HRS	e^+e^- 29 GeV
0.48	$^{+0.06}_{-0.05}$	± 0.02	99	ANJOS	87B	E691	See RAAB 88
0.33	$^{+0.10}_{-0.06}$		21	[4] BECKER	87B	SILI	200 GeV π,K,p
0.57	$^{+0.36}_{-0.26}$	± 0.09	9	BRAUNSCH...	87	TASS	e^+e^- 35–44 GeV
0.47	± 0.22	± 0.05	141	CSORNA	87	CLEO	e^+e^- 10 GeV
0.35	$^{+0.24}_{-0.18}$	± 0.09	17	JUNG	86	HRS	See AVERILL 89
0.26	$^{+0.16}_{-0.09}$		6	USHIDA	86	EMUL	ν wideband
0.32	$^{+0.30}_{-0.13}$		3	BAILEY	84	ACCM	hadron$^+$ Be $\to$ $\phi\pi^+$ X
0.19	$^{+0.13}_{-0.07}$		4	USHIDA	83	EMUL	See USHIDA 86

[2] BONVICINI 99 obtains 1.19 ± 0.04 for the ratio of D_s^+ to D^0 lifetimes.

[3] BARLAG 90C estimates the systematic error to be negligible.

[4] BECKER 87B estimates the systematic error to be negligible.

D_s^+ DECAY MODES

Branching fractions for modes with a resonance in the final state include all the decay modes of the resonance. D_s^- modes are charge conjugates of the modes below.

	Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
	Inclusive modes		
Γ_1	K^- anything	(13 $^{+14}_{-12}$) %	
Γ_2	$\overline{K}^0$ anything + K^0 anything	(39 ± 28) %	
Γ_3	K^+ anything	(20 $^{+18}_{-14}$) %	
Γ_4	non-$K\overline{K}$ anything	(64 ± 17) %	
Γ_5	e^+ anything	(8 $^{+\ 6}_{-\ 5}$) %	
Γ_6	ϕ anything	(18 $^{+15}_{-10}$) %	

Leptonic and semileptonic modes

Γ_7	$\mu^+ \nu_\mu$		(5.1 ± 1.9) $\times 10^{-3}$	S=1.2
Γ_8	$\tau^+ \nu_\tau$		(7 ± 4) %	
Γ_9	$\phi \ell^+ \nu_\ell$	[a]	(2.0 ± 0.5) %	
Γ_{10}	$\eta \ell^+ \nu_\ell$ + $\eta'(958) \ell^+ \nu_\ell$	[a]	(3.5 ± 1.0) %	
Γ_{11}	$\eta \ell^+ \nu_\ell$		(2.6 ± 0.7) %	
Γ_{12}	$\eta'(958) \ell^+ \nu_\ell$		(9.1 ± 3.4) $\times 10^{-3}$	

Hadronic modes with a $K\overline{K}$ pair (including from a ϕ)

Γ_{13}	$K^+ \overline{K}^0$		(3.6 ± 1.1) %	
Γ_{14}	$K^+ K^- \pi^+$	[b]	(4.4 ± 1.2) %	
Γ_{15}	$\phi \pi^+$	[c]	(3.6 ± 0.9) %	
Γ_{16}	$K^+ \overline{K}^*(892)^0$	[c]	(3.3 ± 0.9) %	
Γ_{17}	$f_0(980) \pi^+$	[c]	(1.8 ± 0.8) %	S=1.3
Γ_{18}	$K^+ \overline{K}_0^*(1430)^0$	[c]	(7 ± 4) $\times 10^{-3}$	
Γ_{19}	$f_0(1710) \pi^+ \rightarrow K^+ K^- \pi^+$	[d]	(1.5 ± 1.9) $\times 10^{-3}$	
Γ_{20}	$K^+ K^- \pi^+$ nonresonant		(9 ± 4) $\times 10^{-3}$	
Γ_{21}	$K^0 \overline{K}^0 \pi^+$		—	
Γ_{22}	$K^*(892)^+ \overline{K}^0$	[c]	(4.3 ± 1.4) %	
Γ_{23}	$K^+ K^- \pi^+ \pi^0$		—	
Γ_{24}	$\phi \pi^+ \pi^0$	[c]	(9 ± 5) %	
Γ_{25}	$\phi \rho^+$	[c]	(6.7 ± 2.3) %	
Γ_{26}	$\phi \pi^+ \pi^0$ 3-body	[c]	< 2.6 %	CL=90%
Γ_{27}	$K^+ K^- \pi^+ \pi^0$ non-ϕ		< 9 %	CL=90%
Γ_{28}	$K^+ \overline{K}^0 \pi^+ \pi^-$		< 2.8 %	CL=90%
Γ_{29}	$K^0 K^- \pi^+ \pi^+$		(4.3 ± 1.5) %	
Γ_{30}	$K^*(892)^+ \overline{K}^*(892)^0$	[c]	(5.8 ± 2.5) %	
Γ_{31}	$K^0 K^- \pi^+ \pi^+$ non-$K^{*+} \overline{K}^{*0}$		< 2.9 %	CL=90%
Γ_{32}	$K^+ K^- \pi^+ \pi^+ \pi^-$		(8.4 ± 3.3) $\times 10^{-3}$	
Γ_{33}	$\phi \pi^+ \pi^+ \pi^-$	[c]	(1.18 ± 0.35) %	
Γ_{34}	$K^+ K^- \pi^+ \pi^+ \pi^-$ non-ϕ		(3.0 $^{+\ 3.0}_{-\ 2.0}$) $\times 10^{-3}$	

Hadronic modes without K's

Γ_{35}	$\pi^+ \pi^+ \pi^-$		(1.01 ± 0.28) %	S=1.1
Γ_{36}	$\rho^0 \pi^+$			
Γ_{37}	$f_0(980)^0 \pi^+ \times$ B($f_0 \rightarrow \pi^+ \pi^-$)	[e]	(4.4 ± 2.1) %	
Γ_{38}	$f_2(1270) \pi^+$	[c]	(3.5 ± 1.2) $\times 10^{-3}$	
Γ_{39}	$f_0(1370)^0 \pi^+ \times$B($f_0 \rightarrow \pi^+ \pi^-$)	[e]	(3.3 ± 1.2) $\times 10^{-3}$	
Γ_{40}	$\rho(1450)^0 \pi^+ \times$B($\rho^0 \rightarrow \pi^+ \pi^-$)	[e]	(4.4 ± 2.5) $\times 10^{-4}$	
Γ_{41}	$f_0(1500) \pi^+ \rightarrow \pi^+ \pi^- \pi^+$			
Γ_{42}	$\pi^+ \pi^+ \pi^-$ nonresonant		(5 ±22) $\times 10^{-5}$	
Γ_{43}	$\pi^+ \pi^+ \pi^- \pi^0$		< 12 %	CL=90%
Γ_{44}	$\eta \pi^+$	[c]	(1.7 ± 0.5) %	

Γ_{45}	$\omega\,\pi^+$		[c]	(2.8 ± 1.1)	$\times 10^{-3}$	
Γ_{46}	$\pi^+\,\pi^+\,\pi^+\,\pi^-\,\pi^-$			(7.0 ± 3.0)	$\times 10^{-3}$	
Γ_{47}	$\pi^+\,\pi^+\,\pi^-\,\pi^0\,\pi^0$			—		
Γ_{48}	$\eta\,\rho^+$		[c]	(10.8 ± 3.1)	%	
Γ_{49}	$\eta\,\pi^+\,\pi^0$ 3-body		[c]	< 4	%	CL=90%
Γ_{50}	$\pi^+\,\pi^+\,\pi^+\,\pi^-\,\pi^-\,\pi^0$			(4.9 ± 3.2)	%	
Γ_{51}	$\eta'(958)\,\pi^+$		[c]	(3.9 ± 1.0)	%	
Γ_{52}	$\pi^+\,\pi^+\,\pi^+\,\pi^-\,\pi^-\,\pi^0\,\pi^0$			—		
Γ_{53}	$\eta'(958)\,\rho^+$		[c]	(10.1 ± 2.8)	%	
Γ_{54}	$\eta'(958)\,\pi^+\,\pi^0$ 3-body		[c]	< 1.4	%	CL=90%

Modes with one or three *K*'s

Γ_{55}	$K^0\,\pi^+$			< 8	$\times 10^{-3}$	CL=90%
Γ_{56}	$K^+\,\pi^+\,\pi^-$			(1.0 ± 0.4)	%	
Γ_{57}	$K^+\,\rho^0$			< 2.9	$\times 10^{-3}$	CL=90%
Γ_{58}	$K^*(892)^0\,\pi^+$		[c]	(6.5 ± 2.8)	$\times 10^{-3}$	
Γ_{59}	$K^+\,K^+\,K^-$			< 6	$\times 10^{-4}$	CL=90%
Γ_{60}	$\phi\,K^+$		[c]	< 5	$\times 10^{-4}$	CL=90%

$\Delta C = 1$ weak neutral current (*C1*) modes, or Lepton number (*L*) violating modes

Γ_{61}	$\pi^+\,e^+\,e^-$		[f]	< 2.7	$\times 10^{-4}$	CL=90%
Γ_{62}	$\pi^+\,\mu^+\,\mu^-$		[f]	< 1.4	$\times 10^{-4}$	CL=90%
Γ_{63}	$K^+\,e^+\,e^-$	*C1*		< 1.6	$\times 10^{-3}$	CL=90%
Γ_{64}	$K^+\,\mu^+\,\mu^-$	*C1*		< 1.4	$\times 10^{-4}$	CL=90%
Γ_{65}	$K^*(892)^+\,\mu^+\,\mu^-$	*C1*		< 1.4	$\times 10^{-3}$	CL=90%
Γ_{66}	$\pi^+\,e^\pm\,\mu^\mp$	*LF*	[g]	< 6.1	$\times 10^{-4}$	CL=90%
Γ_{67}	$K^+\,e^\pm\,\mu^\mp$	*LF*	[g]	< 6.3	$\times 10^{-4}$	CL=90%
Γ_{68}	$\pi^-\,e^+\,e^+$	*L*		< 6.9	$\times 10^{-4}$	CL=90%
Γ_{69}	$\pi^-\,\mu^+\,\mu^+$	*L*		< 8.2	$\times 10^{-5}$	CL=90%
Γ_{70}	$\pi^-\,e^+\,\mu^+$	*L*		< 7.3	$\times 10^{-4}$	CL=90%
Γ_{71}	$K^-\,e^+\,e^+$	*L*		< 6.3	$\times 10^{-4}$	CL=90%
Γ_{72}	$K^-\,\mu^+\,\mu^+$	*L*		< 1.8	$\times 10^{-4}$	CL=90%
Γ_{73}	$K^-\,e^+\,\mu^+$	*L*		< 6.8	$\times 10^{-4}$	CL=90%
Γ_{74}	$K^*(892)^-\,\mu^+\,\mu^+$	*L*		< 1.4	$\times 10^{-3}$	CL=90%

Γ_{75}	A dummy mode used by the fit.		(77 ± 6)	%

[*a*] For now, we average together measurements of the $X\,e^+\,\nu_e$ and $X\,\mu^+\,\nu_\mu$ branching fractions. This is the *average*, not the *sum*.

[*b*] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.

[*c*] This branching fraction includes all the decay modes of the final-state resonance.

[*d*] This value includes only K^+K^- decays of the $f_0(1710)$, because branching fractions of this resonance are not known.

[*e*] This value includes only $\pi^+\pi^-$ decays of the intermediate resonance, because branching fractions of this resonance are not known.

[*f*] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

[*g*] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 13 branching ratios uses 24 measurements and one constraint to determine 10 parameters. The overall fit has a $\chi^2 = 12.9$ for 15 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\rm total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_9	70								
x_{11}	60	86							
x_{12}	46	65	56						
x_{14}	67	86	73	56					
x_{15}	73	93	80	60	92				
x_{16}	68	86	74	56	93	93			
x_{35}	64	82	70	53	86	88	84		
x_{37}	36	47	40	30	55	50	51	47	
x_{75}	−69	−89	−80	−61	−93	−93	−91	−85	−76
	x_7	x_9	x_{11}	x_{12}	x_{14}	x_{15}	x_{16}	x_{35}	x_{37}

D_s^+ BRANCHING RATIOS

A few older, now obsolete results have been omitted. They may be found in earlier editions.

Inclusive modes

$\Gamma(K^-\text{anything})/\Gamma_{\rm total}$ **Γ_1/Γ**

VALUE	DOCUMENT ID		TECN	COMMENT
$0.13^{+0.14}_{-0.12}\pm0.02$	COFFMAN	91	MRK3	e^+e^- 4.14 GeV

$[\Gamma(\overline{K^0}\text{anything}) + \Gamma(K^0\text{anything})]/\Gamma_{\rm total}$ **Γ_2/Γ**

VALUE	DOCUMENT ID		TECN	COMMENT
$0.39^{+0.28}_{-0.27}\pm0.04$	COFFMAN	91	MRK3	e^+e^- 4.14 GeV

$\Gamma(K^+ \text{anything})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$0.20^{+0.18}_{-0.13} \pm 0.04$	COFFMAN	91	MRK3	$e^+ e^-$ 4.14 GeV

$\Gamma(\text{non-}K\overline{K}\text{anything})/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$0.64 \pm 0.17 \pm 0.03$	[5] COFFMAN	91	MRK3	$e^+ e^-$ 4.14 GeV

[5] COFFMAN 91 uses the direct measurements of the kaon content to determine this non-$K\overline{K}$ fraction. This number implies that a large fraction of D_s^+ decays involve η, η', and/or non-spectator decays.

$\Gamma(e^+ \text{anything})/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$0.077^{+0.057}_{-0.043}{}^{+0.024}_{-0.021}$		BAI	97	BES	$e^+ e^- \to D_s^+ D_s^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.20	90	[6] BAI	90	MRK3	$e^+ e^-$ 4.14 GeV

[6] Expressed as a value, the BAI 90 result is $\Gamma(e^+ \text{anything})/\Gamma_{\text{total}} = 0.05 \pm 0.05 \pm 0.02$.

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$0.178^{+0.151}_{-0.072}{}^{+0.006}_{-0.063}$	3	BAI	98	BES	$e^+ e^- \to D_s^+ D_s^-$

Leptonic and semileptonic modes

$\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_7/Γ

See the "Note on Pseudoscalar-Meson Decay Constants" in the Listings for the $\pi^\pm$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.015^{+0.013}_{-0.006}{}^{+0.003}_{-0.002}$	3	[7] BAI	95	BES	$e^+ e^- \to D_s^+ D_s^-$
$0.004^{+0.0018}_{-0.0014}{}^{+0.0020}_{-0.0019}$	8	[8] AOKI	93	WA75	π^- emulsion 350 GeV
<0.03	0	[9] AUBERT	83	SPEC	μ^+ Fe, 250 GeV

[7] BAI 95 uses one actual $D_s^+ \to \mu^+ \nu_\mu$ event together with two $D_s^+ \to \tau^+ \nu_\tau$ events and assumes μ-τ universality. This value of $\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ gives a pseudoscalar decay constant of $(430^{+150}_{-130} \pm 40)$ MeV.

[8] AOKI 93 assumes the ratio of production cross sections of the D_s^+ and D^0 is 0.27. The value of $\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ gives a pseudoscalar decay constant $f_{D_s} = (232 \pm 45 \pm 52)$ MeV.

[9] AUBERT 83 assume that the $D_s^\pm$ production rate is 20% of total charm production rate.

$\Gamma(\mu^+\nu_\mu)/\Gamma(\phi\pi^+)$ Γ_7/Γ_{15}

See the "Note on Pseudoscalar-Meson Decay Constants" in the Listings for the $\pi^\pm$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.14 ±0.04 OUR FIT	Error includes scale factor of 1.4.				
0.19 ±0.04 OUR AVERAGE					
0.23 ±0.06 ±0.04	18	[10] ALEXANDROV	00	BEAT	π^- nucleus, 350 GeV
0.173±0.023±0.035	182	[11] CHADA	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.245±0.052±0.074	39	[12] ACOSTA	94	CLE2	See CHADA 98

[10] ALEXANDROV 00 uses $f_D^2/f_{D_s}^2 = 0.82 \pm 0.09$ from a lattice-gauge-theory calculation to get the relative numbers of $D^+ \to \mu^+\nu_\mu$ and $D_s^+ \to \mu^+\nu_\mu$ events. The present result leads to $f_{D_s} = (323 \pm 44 \pm 36)$ MeV.

[11] CHADA 98 obtains $f_{D_s} = (280 \pm 19 \pm 28 \pm 34)$ MeV from this measurement, using $\Gamma(D_s^+ \to \phi\pi^+)/\Gamma(\text{total}) = 0.036 \pm 0.009$.

[12] ACOSTA 94 obtains $f_{D_s} = (344 \pm 37 \pm 52 \pm 42)$ MeV from this measurement, using $\Gamma(D_s^+ \to \phi\pi^+)/\Gamma(\text{total}) = 0.037 \pm 0.009$.

$\Gamma(\mu^+\nu_\mu)/\Gamma(\phi\ell^+\nu_\ell)$ Γ_7/Γ_9

See the "Note on Pseudoscalar-Meson Decay Constants" in the Listings for the $\pi^\pm$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.25±0.07 OUR FIT	Error includes scale factor of 1.4.				
0.16±0.06±0.03	23	[13] KODAMA	96	E653	π^- emulsion, 600 GeV

[13] KODAMA 96 obtains $f_{D_s} = (194 \pm 35 \pm 20 \pm 14)$ MeV from this measurement, using $\Gamma(D_s^+ \to \phi\ell^+\nu)/\Gamma_{\text{total}} = 0.0188 \pm 0.0029$. The third error is from the uncertainty on $\phi\ell^+\nu_\ell$ branching fraction.

$\Gamma(\tau^+\nu_\tau)/\Gamma_{\text{total}}$ Γ_8/Γ

See the "Note on Pseudoscalar-Meson Decay Constants" in the Listings for the $\pi^\pm$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.074±0.028±0.024	16	[14] ACCIARRI	97F	L3	$D_s^{*+} \to \gamma D_s^+$

[14] The second ACCIARRI 97F error here combines in quadrature systematic (0.016) and normalization (0.018) errors. The branching fraction gives $f_{D_s} = (309 \pm 58 \pm 33 \pm 38)$ MeV.

$\Gamma(\phi\ell^+\nu_\ell)/\Gamma(\phi\pi^+)$ Γ_9/Γ_{15}

For now, we average together measurements of the $\Gamma(\phi e^+\nu_e)/\Gamma(\phi\pi^+)$ and $\Gamma(\phi\mu^+\nu_\mu)/\Gamma(\phi\pi^+)$ ratios. See the end of the D_s^+ Listings for measurements of $D_s^+ \to \phi\ell^+\nu_\ell$ form-factor ratios.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.56±0.05 OUR FIT					
0.54±0.05 OUR AVERAGE					
0.54±0.05±0.04	367	[15] BUTLER	94	CLE2	$e^+e^- \approx \Upsilon(4S)$
0.58±0.17±0.07	97	[16] FRABETTI	93G	E687	γ Be $\overline{E}_\gamma = 220$ GeV
0.57±0.15±0.15	104	[17] ALBRECHT	91	ARG	e^+e^- 10.4 GeV
$0.49\pm0.10^{+0.10}_{-0.14}$	54	[18] ALEXANDER	90B	CLEO	e^+e^- 10.5–11 GeV

[15] BUTLER 94 uses both $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$ events, and makes a phase-space adjustment to the latter to use them as $\phi e^+ \nu_e$ events.

[16] FRABETTI 93G measures the $\Gamma(\phi \mu^+ \nu_\mu)/\Gamma(\phi \pi^+)$ ratio.

[17] ALBRECHT 91 measures the $\Gamma(\phi e^+ \nu_e)/\Gamma(\phi \pi^+)$ ratio.

[18] ALEXANDER 90B measures an average of the $\Gamma(\phi e^+ \nu_e)/\Gamma(\phi \pi^+)$ and $\Gamma(\phi \mu^+ \nu_\mu)/\Gamma(\phi \pi^+)$ ratios.

$\Gamma(\eta \ell^+ \nu_\ell)/\Gamma(\phi \ell^+ \nu_\ell)$ Γ_{11}/Γ_9

Unseen decay modes of the η and the ϕ are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.27±0.19 OUR FIT					
1.24±0.12±0.15	440	[19] BRANDENB...	95	CLE2	$e^+ e^- \approx \Upsilon(4S)$

[19] BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

$\Gamma(\eta'(958) \ell^+ \nu_\ell)/\Gamma(\phi \ell^+ \nu_\ell)$ Γ_{12}/Γ_9

Unseen decay modes of the resonances are included.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
0.44±0.13 OUR FIT						
0.43±0.11±0.07		29	[20] BRANDENB...	95	CLE2	$e^+ e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<1.6	90		[21] KODAMA	93B	E653	π^- emulsion 600 GeV

[20] BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

[21] KODAMA 93B uses μ^+ events.

$[\Gamma(\eta \ell^+ \nu_\ell) + \Gamma(\eta'(958) \ell^+ \nu_\ell)]/\Gamma(\phi \ell^+ \nu_\ell)$ $\Gamma_{10}/\Gamma_9 = (\Gamma_{11}+\Gamma_{12})/\Gamma_9$

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.72±0.23 OUR FIT					
3.9 ±1.6	13	[22] KODAMA	93	E653	π^- emulsion 600 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.67±0.17±0.17		[23] BRANDENB...	95	CLE2	$e^+ e^- \approx \Upsilon(4S)$

[22] KODAMA 93 uses μ^+ events.

[23] This BRANDENBURG 95 data is redundant with data in previous blocks.

Hadronic modes with a $K\overline{K}$ pair.

$\Gamma(K^+ \overline{K}^0)/\Gamma(\phi \pi^+)$ Γ_{13}/Γ_{15}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.01±0.16 OUR AVERAGE					
1.15±0.31±0.19	68	ANJOS	90C	E691	γ Be
0.92±0.32±0.20		ADLER	89B	MRK3	$e^+ e^-$ 4.14 GeV
0.99±0.17±0.10		CHEN	89	CLEO	$e^+ e^-$ 10 GeV

$\Gamma(\phi\pi^+)/\Gamma_{total}$ Γ_{15}/Γ

We now have model-independent measurements of this branching fraction, and so we no longer use the earlier, model-dependent results.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
0.036 ±0.009 OUR FIT						
0.036 ±0.009 OUR AVERAGE						
0.0359 ±0.0077 ±0.0048			[24] ARTUSO	96	CLE2	e^+e^- at $\Upsilon(4S)$
0.039 $^{+0.051}_{-0.019}$ $^{+0.018}_{-0.011}$			[25] BAI	95C	BES	e^+e^- 4.03 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.051 ±0.004 ±0.008			[26] BUTLER	94	CLE2	$e^+e^- \approx \Upsilon(4S)$
<0.048	90		MUHEIM	94		
0.046 ±0.015			[27] MUHEIM	94		
0.031 ±0.009			[27] MUHEIM	94		
0.031 ±0.009 ±0.006			[26] FRABETTI	93G	E687	γ Be $\overline{E}_\gamma$= 220 GeV
0.024 ±0.010			[26] ALBRECHT	91	ARG	$e^+e^- \approx$ 10.4 GeV
<0.041	90	0	[25] ADLER	90B	MRK3	e^+e^- 4.14 GeV
0.031 ±0.006 $^{+0.011}_{-0.009}$			[26] ALEXANDER	90B	CLEO	e^+e^- 10.5–11 GeV
0.048 ±0.017 ±0.019			[28] ALVAREZ	90C	NA14	Photoproduction
>0.034	90		[26] ANJOS	90B	E691	γBe, $\overline{E}_\gamma \approx$ 145 GeV
0.02 ±0.01		405	[29] CHEN	89	CLEO	e^+e^- 10 GeV
0.033 ±0.016 ±0.010		9	[29] BRAUNSCH...	87	TASS	e^+e^- 35–44 GeV
0.033 ±0.011		30	[29] DERRICK	85B	HRS	e^+e^- 29 GeV

[24] ARTUSO 96 uses partially reconstructed $\overline{B}^0 \to D^{*+} D_s^{*-}$ decays to get a model-independent value for $\Gamma(D_s^- \to \phi\pi^-)/\Gamma(D^0 \to K^-\pi^+)$ of 0.92 ± 0.20 ± 0.11.

[25] BAI 95C uses $e^+e^- \to D_s^+ D_s^-$ events in which one or both of the $D_s^\pm$ are observed to obtain the first model-independent measurement of the $D_s^+ \to \phi\pi^+$ branching fraction, without assumptions about $\sigma(D_s^\pm)$. However, with only two "doubly-tagged" events, the statistical error is too large for the result to be competitive with indirect measurements. ADLER 90B used the same method to set a limit.

[26] BUTLER 94, FRABETTI 93G, ALBRECHT 91, ALEXANDER 90B, and ANJOS 90B measure the ratio $\Gamma(D_s^+ \to \phi\ell^+\nu_\ell)/\Gamma(D_s^+ \to \phi\pi^+)$, where ℓ = e and/or μ, and then use a theoretical calculation of the ratio of widths $\Gamma(D_s^+ \to \phi\ell^+\nu_\ell)/\Gamma(D^+ \to \overline{K}^{*0}\ell^+\nu)$. Not everyone uses the same value for this ratio.

[27] The two MUHEIM 94 values here are model-dependent calculations based on distinct data sets. The first uses measurements of the $D_2^*(2460)^0$ and $D_{s1}(2536)^+$, the second uses B-decay factorization and $\Gamma(D_s^+ \to \mu^+\nu_\mu)/\Gamma(D_s^+ \to \phi\ell^+\nu_\ell)$. A third calculation using the semileptonic width of $D_s^+ \to \phi\ell^+\nu_\ell$ is not independent of other results listed here. Note also the upper limit, based on the sum of established D_s^+ branching ratios.

[28] ALVAREZ 90C relies on the Lund model to estimate the ratio of D_s^+ to D^+ cross sections.

[29] Values based on crude estimates of the $D_s^\pm$ production level. DERRICK 85B errors are statistical only.

$\Gamma(\phi\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{15}/Γ_{14}

Unseen decay modes of the ϕ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.81 ±0.08 OUR FIT				
0.807±0.067±0.096	FRABETTI	95B	E687	Dalitz plot analysis

$\Gamma(K^+\overline{K}^*(892)^0)/\Gamma(K^+K^-\pi^+)$ Γ_{16}/Γ_{14}

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.75 ±0.07 OUR FIT				
0.717±0.069±0.060	FRABETTI	95B	E687	Dalitz plot analysis

$\Gamma(K^+\overline{K}^*(892)^0)/\Gamma(\phi\pi^+)$ Γ_{16}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.92±0.09 OUR FIT					
0.95±0.10 OUR AVERAGE					
0.85±0.34±0.20	9	ALVAREZ	90C	NA14	Photoproduction
0.84±0.30±0.22		ADLER	89B	MRK3	e^+e^- 4.14 GeV
1.05±0.17±0.12		CHEN	89	CLEO	e^+e^- 10 GeV
0.87±0.13±0.05	117	ANJOS	88	E691	Photoproduction
1.44±0.37	87	ALBRECHT	87F	ARG	e^+e^- 10 GeV

$\Gamma(f_0(980)^0\pi^+\times B(f_0\to\pi^+\pi^-))/\Gamma(K^+K^-\pi^+)$ Γ_{37}/Γ_{14}

Unseen decay modes of the $f_0(980)$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
1.0 ±0.4 OUR FIT				
1.00±0.32±0.24	FRABETTI	95B	E687	Dalitz plot analysis

$\Gamma(f_0(1710)\pi^+\to K^+K^-\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{19}/Γ_{14}

This includes *only* K^+K^- decays of the $f_0(1710)$, because branching fractions of this resonance are not known.

VALUE	DOCUMENT ID		TECN	COMMENT
0.034±0.023±0.035	FRABETTI	95B	E687	Dalitz plot analysis

$\Gamma(K^+\overline{K}_0^*(1430)^0)/\Gamma(K^+K^-\pi^+)$ Γ_{18}/Γ_{14}

Unseen decay modes of the $\overline{K}_0^*(1430)^0$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
0.150±0.052±0.052	FRABETTI	95B	E687	Dalitz plot analysis

$\Gamma(K^+K^-\pi^+\text{nonresonant})/\Gamma(\phi\pi^+)$ Γ_{20}/Γ_{15}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.25±0.07±0.05	48	ANJOS	88	E691	Photoproduction

$\Gamma(K^*(892)^+\overline{K}^0)/\Gamma(\phi\pi^+)$ Γ_{22}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID		TECN	COMMENT
1.20±0.21±0.13	CHEN	89	CLEO	e^+e^- 10 GeV

$\Gamma(K^*(892)^+\overline{K}^0)/\Gamma(K^+\overline{K}^0)$ Γ_{22}/Γ_{13}

Unseen decay modes of the $K^*(892)^+$ are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.9	90	FRABETTI	95	E687	γ Be $\overline{E}_\gamma \approx 200$ GeV

$\Gamma(\phi\pi^+\pi^0)/\Gamma(\phi\pi^+)$ Γ_{24}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
2.4±1.0±0.5		11	ANJOS	89E	E691	Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<2.6	90		ALVAREZ	90C	NA14	Photoproduction

$\Gamma(\phi\rho^+)/\Gamma(\phi\pi^+)$ Γ_{25}/Γ_{15}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.86\pm0.26^{+0.29}_{-0.40}}$	253	AVERY	92	CLE2	$e^+e^- \simeq 10.5$ GeV

$\Gamma(\phi\pi^+\pi^0\ 3\text{-body})/\Gamma(\phi\pi^+)$ Γ_{26}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.71	90	DAOUDI	92	CLE2	$e^+e^- \approx 10.5$ GeV

$\Gamma(K^+K^-\pi^+\pi^0\ \text{non-}\phi)/\Gamma(\phi\pi^+)$ Γ_{27}/Γ_{15}

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
<2.4	90	[30]	ANJOS	89E	E691	Photoproduction

[30] Total minus ϕ component.

$\Gamma(K^+\overline{K}^0\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{28}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.77	90	ALBRECHT	92B	ARG	$e^+e^- \simeq 10.4$ GeV

$\Gamma(K^0K^-\pi^+\pi^+)/\Gamma(\phi\pi^+)$ Γ_{29}/Γ_{15}

VALUE	DOCUMENT ID		TECN	COMMENT
1.2 ±0.2 ±0.2	ALBRECHT	92B	ARG	$e^+e^- \simeq 10.4$ GeV

$\Gamma(K^*(892)^+\overline{K}^*(892)^0)/\Gamma(\phi\pi^+)$ Γ_{30}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID		TECN	COMMENT
1.6±0.4±0.4	ALBRECHT	92B	ARG	$e^+e^- \simeq 10.4$ GeV

$\Gamma(K^0K^-\pi^+\pi^+\ \text{non-}K^{*+}\overline{K}^{*0})/\Gamma(\phi\pi^+)$ Γ_{31}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.80	90	ALBRECHT	92B	ARG	$e^+e^- \simeq 10.4$ GeV

$\Gamma(K^+K^-\pi^+\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{32}/Γ_{14}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.188±0.036±0.040	75	FRABETTI	97C	E687	γ Be, $\overline{E}_\gamma \approx 200$ GeV

$\Gamma(\phi\pi^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{33}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
0.33±0.06 OUR AVERAGE						
0.28±0.06±0.01		40	FRABETTI	97C	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV
0.58±0.21±0.10		21	FRABETTI	92	E687	γ Be
0.42±0.13±0.07		19	ANJOS	88	E691	Photoproduction
1.11±0.37±0.28		62	ALBRECHT	85D	ARG	e^+e^- 10 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.24	90		ALVAREZ	90C	NA14	Photoproduction

$\Gamma(K^+K^-\pi^+\pi^+\pi^-\ \text{non-}\phi)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$0.003\ ^{+0.003}_{-0.002}$	BARLAG	92C	ACCM	π^- 230 GeV

$\Gamma(K^+K^-\pi^+\pi^+\pi^-\ \text{non-}\phi)/\Gamma(\phi\pi^+)$ Γ_{34}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.32	90	10	ANJOS	88	E691	Photoproduction

Hadronic modes without K's

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{35}/Γ_{14}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.227±0.033 OUR FIT	Error includes scale factor of 1.1.				
0.265±0.041±0.031	98	FRABETTI	97D	E687	γ Be ≈ 200 GeV

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{35}/Γ_{15}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.28 ±0.04 OUR FIT	Error includes scale factor of 1.3.				
0.27 ±0.04 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.				
$0.245\pm0.028\ ^{+0.019}_{-0.012}$	848	AITALA	01A	E791	π^- nucleus, 500 GeV
0.33 ±0.10 ±0.04	29	ADAMOVICH	93	WA82	π^- 340 GeV
0.44 ±0.10 ±0.04	68	ANJOS	89	E691	Photoproduction



$\Gamma\left(\pi^+\,\pi^+\,\pi^-\right)/\Gamma\left(\phi\,\pi^+\right)$

$\Gamma\left(\rho^0\,\pi^+\right)/\Gamma\left(\pi^+\,\pi^+\,\pi^-\right)$ Γ_{36}/Γ_{35}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.058\pm0.023\pm0.037$		[31] AITALA	01A	E791	π^- nucleus, 500 GeV
<0.073	90	FRABETTI	97D	E687	γ Be $\approx$ 200 GeV

[31] This AITALA 01B result does not have enough statistical significance to advance it to the Summary Tables.

$\Gamma\left(\rho^0\,\pi^+\right)/\Gamma\left(\phi\pi^+\right)$ Γ_{36}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.08	90	ANJOS	89	E691	Photoproduction
<0.22	90	ALBRECHT	87G	ARG	$e^+\,e^-$ 10 GeV

$\Gamma\left(f_0(980)^0\,\pi^+\times B(f_0\to\pi^+\,\pi^-)\right)/\Gamma\left(\pi^+\,\pi^+\,\pi^-\right)$ Γ_{37}/Γ_{35}

This includes only the $\pi^+\,\pi^-$ decays of the $f_0(980)$, because branching fractions of this resonance are not known.

VALUE	DOCUMENT ID		TECN	COMMENT
$0.565\pm0.043\pm0.047$	AITALA	01A	E791	π^- nucleus, 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.074\pm0.140\pm0.043$	FRABETTI	97D	E687	γ Be $\approx$ 200 GeV

$\Gamma(f_0(980)^0\pi^+ \times B(f_0 \to \pi^+\pi^-))/\Gamma(\phi\pi^+)$ Γ_{37}/Γ_{15}

This includes only the $\pi^+\pi^-$ decays of the $f_0(980)$, because branching fractions of this resonance are not known.

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.28 \pm 0.10 \pm 0.03$	ANJOS	89	E691	Photoproduction

$\Gamma(f_2(1270)\pi^+)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{38}/Γ_{35}

Unseen decay modes of the $f_2(1270)$ are included.

VALUE	DOCUMENT ID		TECN	COMMENT
$0.349 \pm 0.059 \pm 0.011$	[32] AITALA	01A	E791	π^- nucleus, 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.22 \pm 0.10 \pm 0.03$	FRABETTI	97D	E687	γ Be $\approx$ 200 GeV

[32] See AITALA 01A for the magnitude and phase of this amplitude relative to the $f_0(980)\pi^+$ amplitude.

$\Gamma(f_0(1370)^0\pi^+ \times B(f_0 \to \pi^+\pi^-))/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{39}/Γ_{35}

This includes only the $\pi^+\pi^-$ decays of the $f_0(1370)$, because branching fractions of this resonance are not known.

VALUE	DOCUMENT ID		TECN	COMMENT
$0.324 \pm 0.077 \pm 0.017$	[33] AITALA	01A	E791	π^- nucleus, 500 GeV

[33] See AITALA 01A for the magnitude and phase of this amplitude relative to the $f_0(980)\pi^+$ amplitude.

$\Gamma(\rho(1450)^0\pi^+ \times B(\rho^0 \to \pi^+\pi^-))/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{40}/Γ_{35}

This includes only the $\pi^+\pi^-$ decays of the $\rho(1450)^0$, because branching fractions of this resonance are not known.

VALUE	DOCUMENT ID		TECN	COMMENT
$0.044 \pm 0.021 \pm 0.002$	[34] AITALA	01A	E791	π^- nucleus, 500 GeV

[34] See AITALA 01A for the magnitude and phase of this amplitude relative to the $f_0(980)\pi^+$ amplitude.

$\Gamma(f_0(1500)\pi^+ \to \pi^+\pi^-\pi^+)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{41}/Γ_{35}

This includes only $\pi^+\pi^-$ decays of the $f_0(1500)$, because branching fractions of this resonance are not known.

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.274 \pm 0.114 \pm 0.019$	[35] FRABETTI	97D	E687	γ Be $\approx$ 200 GeV

[35] FRABETTI 97D calls this mode $S(1475)\pi^+$, but finds the mass and width of this $S(1475)$ to be in excellent agreement with those of the $f_0(1500)$.

$\Gamma(\pi^+\pi^+\pi^-\ \text{nonresonant})/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{42}/Γ_{35}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$0.005 \pm 0.014 \pm 0.017$		AITALA	01A	E791	π^- nucleus, 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.269	90	[36] FRABETTI	97D	E687	γ Be $\approx$ 200 GeV

[36] See FRABETTI 97D on the difficulty of distengangling the $f_0(1500)\pi^+$ and nonresonant modes.

$\Gamma(\pi^+\pi^+\pi^-$ nonresonant$)/\Gamma(\phi\pi^+)$ Γ_{42}/Γ_{15}

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.29\pm0.09\pm0.03$	ANJOS	89	E691	Photoproduction

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(\phi\pi^+)$ Γ_{43}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<3.3	90	ANJOS	89E	E691	Photoproduction

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+)$ Γ_{44}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$0.48\pm0.03\pm0.04$		920	JESSOP	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$0.54\pm0.09\pm0.06$		165	ALEXANDER	92	CLE2	See JESSOP 98
<1.5	90		ANJOS	89E	E691	Photoproduction

$\Gamma(\omega\pi^+)/\Gamma(\phi\pi^+)$ Γ_{45}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.5	90	ANJOS	89E	E691	Photoproduction

$\Gamma(\omega\pi^+)/\Gamma(\eta\pi^+)$ Γ_{45}/Γ_{44}

VALUE	DOCUMENT ID		TECN	COMMENT
$0.16\pm0.04\pm0.03$	BALEST	97	CLE2	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{46}/Γ_{14}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$0.158\pm0.042\pm0.031$	37	FRABETTI	97C	E687	γ Be, $\overline{E}_\gamma \approx$ 200 GeV

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma(\phi\pi^+)$ Γ_{46}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.29	90	ANJOS	89	E691	Photoproduction

$\Gamma(\eta\rho^+)/\Gamma(\phi\pi^+)$ Γ_{48}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$2.98\pm0.20\pm0.39$	447	JESSOP	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$2.86\pm0.38^{+0.36}_{-0.38}$	217	AVERY	92	CLE2	See JESSOP 98

$\Gamma(\eta\pi^+\pi^0 3\text{-body})/\Gamma(\phi\pi^+)$ Γ_{49}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.1	90	JESSOP	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.82	90	[37] DAOUDI	92	CLE2	See JESSOP 98

[37] We use the JESSOP 98 limit, even though the DAOUDI 92 limit, from the same experiment but with a much smaller data sample, is more restrictive.

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.049^{+0.033}_{-0.030}}$	BARLAG	92C	ACCM	π^- 230 GeV

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$ Γ_{51}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
1.08±0.09 OUR AVERAGE						
1.03±0.06±0.07		537	JESSOP	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
$2.5 \pm 1.0\ ^{+1.5}_{-0.4}$		22	ALVAREZ	91	NA14	Photoproduction
2.5 ±0.5 ±0.3		215	ALBRECHT	90D	ARG	$e^+e^- \approx 10.4$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1.20±0.15±0.11		281	ALEXANDER	92	CLE2	See JESSOP 98
<1.3	90		ANJOS	91B	E691	γ Be, $\overline{E}_\gamma \approx 145$ GeV

$\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$ Γ_{53}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
2.78±0.28±0.30	137	JESSOP	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$3.44 \pm 0.62\ ^{+0.44}_{-0.46}$	68	AVERY	92	CLE2	See JESSOP 98

$\Gamma(\eta'(958)\pi^+\pi^0 3\text{-body})/\Gamma(\phi\pi^+)$ Γ_{54}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.4	90	JESSOP	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.85	90	DAOUDI	92	CLE2	See JESSOP 98

Modes with one or three K's

$\Gamma(K^0\pi^+)/\Gamma(\phi\pi^+)$ Γ_{55}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.21	90	ADLER	89B	MRK3	e^+e^- 4.14 GeV

$\Gamma(K^0\pi^+)/\Gamma(K^+\overline{K}^0)$ Γ_{55}/Γ_{13}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.53	90	FRABETTI	95	E687	γ Be $\overline{E}_\gamma \approx 200$ GeV

$\Gamma(K^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{56}/Γ_{15}

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.28±0.06±0.05	85	FRABETTI	95E	E687	γ Be, $\overline{E}_\gamma$= 220 GeV

$\Gamma(K^+\rho^0)/\Gamma(\phi\pi^+)$ Γ_{57}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.08	90	FRABETTI	95E	E687	γ Be, $\overline{E}_\gamma$= 220 GeV

$\Gamma(K^*(892)^0\pi^+)/\Gamma(\phi\pi^+)$ Γ_{58}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.18±0.05±0.04	25	FRABETTI	95E	E687	γ Be, $\overline{E}_\gamma$= 220 GeV

$\Gamma(K^+K^+K^-)/\Gamma(\phi\pi^+)$ Γ_{59}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.016	90	FRABETTI	95F	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV

$\Gamma(\phi K^+)/\Gamma(\phi\pi^+)$ Γ_{60}/Γ_{15}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.013	90	FRABETTI	95F	E687	γ Be, $\overline{E}_\gamma \approx$ 220 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.071	90	ANJOS	92D	E691	γ Be, $\overline{E}_\gamma$ = 145 GeV

Rare or forbidden modes

$\Gamma(\pi^+e^+e^-)/\Gamma_{total}$ Γ_{61}/Γ

This mode is not a useful test for a ΔC=1 weak neutral current because both quarks must change flavor in this decay.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<2.7 × 10^{-4}	90	AITALA	99G	E791	π^- N 500 GeV

$\Gamma(\pi^+\mu^+\mu^-)/\Gamma_{total}$ Γ_{62}/Γ

This mode is not a useful test for a ΔC=1 weak neutral current because both quarks must change flavor in this decay.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
<1.4 × 10^{-4}	90		AITALA	99G	E791	π^- N 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<4.3 × 10^{-4}	90	0	KODAMA	95	E653	π^- emulsion 600 GeV

$\Gamma(K^+e^+e^-)/\Gamma_{total}$ Γ_{63}/Γ

A test for the ΔC=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.6 × 10^{-3}	90	AITALA	99G	E791	π^- N 500 GeV

$\Gamma(K^+\mu^+\mu^-)/\Gamma_{total}$ Γ_{64}/Γ

A test for the ΔC=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.4\times10^{-4}$	90		AITALA	99G	E791	π^- N 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<5.9\times10^{-4}$	90	0	KODAMA	95	E653	π^- emulsion 600 GeV

$\Gamma(K^*(892)^+\mu^+\mu^-)/\Gamma_{total}$ Γ_{65}/Γ

A test for the ΔC=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.4\times10^{-3}$	90	0	KODAMA	95	E653	π^- emulsion 600 GeV

$\Gamma(\pi^+e^\pm\mu^\mp)/\Gamma_{total}$ Γ_{66}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.1\times10^{-4}$	90	AITALA	99G	E791	π^- N 500 GeV

$\Gamma(K^+e^\pm\mu^\mp)/\Gamma_{total}$ Γ_{67}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.3\times10^{-4}$	90	AITALA	99G	E791	π^- N 500 GeV

$\Gamma(\pi^-e^+e^+)/\Gamma_{total}$ Γ_{68}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.9\times10^{-4}$	90	AITALA	99G	E791	π^- N 500 GeV

$\Gamma(\pi^-\mu^+\mu^+)/\Gamma_{total}$ Γ_{69}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<8.2\times10^{-5}$	90		AITALA	99G	E791	π^- N 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<4.3\times10^{-4}$	90	0	KODAMA	95	E653	π^- emulsion 600 GeV

$\Gamma(\pi^-e^+\mu^+)/\Gamma_{total}$ Γ_{70}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.3\times10^{-4}$	90	AITALA	99G	E791	π^- N 500 GeV

$\Gamma(K^-e^+e^+)/\Gamma_{total}$ Γ_{71}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.3\times10^{-4}$	90	AITALA	99G	E791	π^- N 500 GeV

$\Gamma(K^-\mu^+\mu^+)/\Gamma_{total}$ Γ_{72}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.8\times10^{-4}$	90		AITALA	99G	E791	π^- N 500 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<5.9\times10^{-4}$	90	0	KODAMA	95	E653	π^- emulsion 600 GeV

$\Gamma(K^- e^+ \mu^+)/\Gamma_{total}$ Γ_{73}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.8 \times 10^{-4}$	90	AITALA	99G	E791	π^- *N* 500 GeV

$\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{total}$ Γ_{74}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$<1.4 \times 10^{-3}$	90	0	KODAMA	95	E653	π^- emulsion 600 GeV

$D_s^+ \to \phi \ell^+ \nu_\ell$ FORM FACTORS

$r_2 \equiv A_2(0)/A_1(0)$ in $D_s^+ \to \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.60±0.24 OUR AVERAGE					
1.57±0.25±0.19	271	AITALA	99D	E791	$\phi e^+ \nu_e$, $\phi \mu^+ \nu_\mu$
1.4 ±0.5 ±0.3	308	AVERY	94B	CLE2	$\phi e^+ \nu_e$
1.1 ±0.8 ±0.1	90	FRABETTI	94F	E687	$\phi \mu^+ \nu_\mu$
2.1 $^{+0.6}_{-0.5}$ ±0.2	19	KODAMA	93	E653	$\phi \mu^+ \nu_\mu$

$r_V \equiv V(0)/A_1(0)$ in $D_s^+ \to \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.92±0.32 OUR AVERAGE					
2.27±0.35±0.22	271	AITALA	99D	E791	$\phi e^+ \nu_e$, $\phi \mu^+ \nu_\mu$
0.9 ±0.6 ±0.3	308	AVERY	94B	CLE2	$\phi e^+ \nu_e$
1.8 ±0.9 ±0.2	90	FRABETTI	94F	E687	$\phi \mu^+ \nu_\mu$
2.3 $^{+1.1}_{-0.9}$ ±0.4	19	KODAMA	93	E653	$\phi \mu^+ \nu_\mu$

Γ_L/Γ_T in $D_s^+ \to \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.72±0.18 OUR AVERAGE					
1.0 ±0.3 ±0.2	308	AVERY	94B	CLE2	$\phi e^+ \nu_e$
1.0 ±0.5 ±0.1	90	[38] FRABETTI	94F	E687	$\phi \mu^+ \nu_\mu$
0.54±0.21±0.10	19	[38] KODAMA	93	E653	$\phi \mu^+ \nu_\mu$

[38] FRABETTI 94F and KODAMA 93 evaluate Γ_L/Γ_T for a lepton mass of zero.

$D_s^{\pm}$ REFERENCES

AITALA	01A	PRL 86 765	E.M. Aitala *et al.*	(FNAL E791 Collab.)
AITALA	01B	PRL 86 770	E.M. Aitala *et al.*	(FNAL E791 Collab.)
ALEXANDROV	00	PL B478 31	Y. Alexandrov *et al.*	(CERN BEATRICE Collab.)
AITALA	99	PL B445 449	E.M. Aitala *et al.*	(FNAL E791 Collab.)
AITALA	99D	PL B450 294	E.M. Aitala *et al.*	(FNAL E791 Collab.)
AITALA	99G	PL B462 401	E.M. Aitala *et al.*	(FNAL E791 Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini *et al.*	(CLEO Collab.)
BAI	98	PR D57 28	J.Z. Bai *et al.*	(BEPC BES Collab.)
CHADA	98	PR D58 032002	M. Chada *et al.*	(CLEO Collab.)
JESSOP	98	PR D58 052002	C.P. Jessop *et al.*	(CLEO Collab.)
ACCIARRI	97F	PL B396 327	M. Acciarri *et al.*	(L3 Collab.)
BAI	97	PR D56 3779	J.Z. Bai *et al.*	(BEPC BES Collab.)
BALEST	97	PRL 79 1436	R. Balest *et al.*	(CLEO Collab.)
FRABETTI	97C	PL B401 131	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
ARTUSO	96	PL B378 364	M. Artuso *et al.*	(CLEO Collab.)
KODAMA	96	PL B382 299	K. Kodama *et al.*	(FNAL E653 Collab.)
BAI	95	PRL 74 4599	J.Z. Bai *et al.*	(BES Collab.)
BAI	95C	PR D52 3781	J.Z. Bai *et al.*	(BES Collab.)
BRANDENB...	95	PRL 75 3804	G.W. Brandenburg *et al.*	(CLEO Collab.)
FRABETTI	95	PL B346 199	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	95B	PL B351 591	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	95E	PL B359 403	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	95F	PL B363 259	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama *et al.*	(FNAL E653 Collab.)
ACOSTA	94	PR D49 5690	D. Acosta *et al.*	(CLEO Collab.)
AVERY	94B	PL B337 405	P. Avery *et al.*	(CLEO Collab.)
BROWN	94	PR D50 1884	D. Brown *et al.*	(CLEO Collab.)
BUTLER	94	PL B324 255	F. Butler *et al.*	(CLEO Collab.)
FRABETTI	94F	PL B328 187	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
MUHEIM	94	PR D49 3767	F. Muheim, S. Stone	(SYRA)
ADAMOVICH	93	PL B305 177	M.I. Adamovich *et al.*	(CERN WA82 Collab.)
AOKI	93	PTP 89 131	S. Aoki *et al.*	(CERN WA75 Collab.)
FRABETTI	93F	PRL 71 827	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
FRABETTI	93G	PL B313 253	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
KODAMA	93	PL B309 483	K. Kodama *et al.*	(FNAL E653 Collab.)
KODAMA	93B	PL B313 260	K. Kodama *et al.*	(FNAL E653 Collab.)
ALBRECHT	92B	ZPHY C53 361	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	92	PRL 68 1275	J. Alexander *et al.*	(CLEO Collab.)
ANJOS	92D	PRL 69 2892	J.C. Anjos *et al.*	(FNAL E691 Collab.)
AVERY	92	PRL 68 1279	P. Avery *et al.*	(CLEO Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag *et al.*	(ACCMOR Collab.)
Also	90D	ZPHY C48 29	S. Barlag *et al.*	(ACCMOR Collab.)
DAOUDI	92	PR D45 3965	M. Daoudi *et al.*	(CLEO Collab.)
FRABETTI	92	PL B281 167	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
ALBRECHT	91	PL B255 634	H. Albrecht *et al.*	(ARGUS Collab.)
ALVAREZ	91	PL B255 639	M.P. Alvarez *et al.*	(CERN NA14/2 Collab.)
ANJOS	91B	PR D43 R2063	J.C. Anjos *et al.*	(FNAL E691 Collab.)
COFFMAN	91	PL B263 135	D.M. Coffman *et al.*	(Mark III Collab.)
ADLER	90B	PRL 64 169	J.C. Adler *et al.*	(Mark III Collab.)
ALBRECHT	90D	PL B245 315	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	90B	PRL 65 1531	J. Alexander *et al.*	(CLEO Collab.)
ALVAREZ	90	ZPHY C47 539	M.P. Alvarez *et al.*	(CERN NA14/2 Collab.)
ALVAREZ	90C	PL B246 261	M.P. Alvarez *et al.*	(CERN NA14/2 Collab.)
ANJOS	90B	PRL 64 2885	J.C. Anjos *et al.*	(FNAL E691 Collab.)
ANJOS	90C	PR D41 2705	J.C. Anjos *et al.*	(FNAL E691 Collab.)
BAI	90	PRL 65 686	Z. Bai *et al.*	(Mark III Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag *et al.*	(ACCMOR Collab.)
FRABETTI	90	PL B251 639	P.L. Frabetti *et al.*	(FNAL E687 Collab.)
ADLER	89B	PRL 63 1211	J. Adler *et al.*	(Mark III Collab.)
Also	89D	PRL 63 2858 erratum	J. Adler *et al.*	(Mark III Collab.)
ANJOS	89	PRL 62 125	J.C. Anjos *et al.*	(FNAL E691 Collab.)
ANJOS	89E	PL B223 267	J.C. Anjos *et al.*	(FNAL E691 Collab.)
AVERILL	89	PR D39 123	D.A. Averill *et al.*	(HRS Collab.)
CHEN	89	PL B226 192	W.Y. Chen *et al.*	(CLEO Collab.)

ALBRECHT	88	PL B207 349	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	88I	PL B210 267	H. Albrecht *et al.*	(ARGUS Collab.)
ANJOS	88	PRL 60 897	J.C. Anjos *et al.*	(FNAL E691 Collab.)
RAAB	88	PR D37 2391	J.R. Raab *et al.*	(FNAL E691 Collab.)
ALBRECHT	87F	PL B179 398	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87G	PL B195 102	H. Albrecht *et al.*	(ARGUS Collab.)
ANJOS	87B	PRL 58 1818	J.C. Anjos *et al.*	(FNAL E691 Collab.)
BECKER	87B	PL B184 277	H. Becker *et al.*	(NA11 and NA32 Collab.)
BLAYLOCK	87	PRL 58 2171	G.T. Blaylock *et al.*	(Mark III Collab.)
BRAUNSCH...	87	ZPHY C35 317	W. Braunschweig *et al.*	(TASSO Collab.)
CSORNA	87	PL B191 318	S.E. Csorna *et al.*	(CLEO Collab.)
JUNG	86	PRL 56 1775	C. Jung *et al.*	(HRS Collab.)
USHIDA	86	PRL 56 1767	N. Ushida *et al.*	(FNAL E531 Collab.)
ALBRECHT	85D	PL 153B 343	H. Albrecht *et al.*	(ARGUS Collab.)
DERRICK	85B	PRL 54 2568	M. Derrick *et al.*	(HRS Collab.)
AIHARA	84D	PRL 53 2465	H. Aihara *et al.*	(TPC Collab.)
ALTHOFF	84	PL 136B 130	M. Althoff *et al.*	(TASSO Collab.)
BAILEY	84	PL 139B 320	R. Bailey *et al.*	(ACCMOR Collab.)
AUBERT	83	NP B213 31	J.J. Aubert *et al.*	(EMC Collab.)
CHEN	83C	PRL 51 634	A. Chen *et al.*	(CLEO Collab.)
USHIDA	83	PRL 51 2362	N. Ushida *et al.*	(FNAL E653 Collab.)

OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	J.D. Richman, P.R. Burchat	(UCSB, STAN)