

$\Sigma(1660) P_{11}$ $I(J^P) = 1(\frac{1}{2}^+)$ Status: ***

For results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** (1982).

 $\Sigma(1660)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|--|
| 1630 to 1690 (≈ 1660) OUR ESTIMATE | | | |
| 1665.1 \pm 11.2 | ¹ KOISO | 85 | DPWA $K^- p \rightarrow \Sigma \pi$ |
| 1670 \pm 10 | GOPAL | 80 | DPWA $\bar{K} N \rightarrow \bar{K} N$ |
| 1679 \pm 10 | ALSTON-... | 78 | DPWA $\bar{K} N \rightarrow \bar{K} N$ |
| 1676 \pm 15 | GOPAL | 77 | DPWA $\bar{K} N$ multichannel |
| 1668 \pm 25 | VANHORN | 75 | DPWA $K^- p \rightarrow \Lambda \pi^0$ |
| 1670 \pm 20 | KANE | 74 | DPWA $K^- p \rightarrow \Sigma \pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1565 or 1597 | ² MARTIN | 77 | DPWA $\bar{K} N$ multichannel |
| 1660 \pm 30 | ³ BAILLON | 75 | IPWA $\bar{K} N \rightarrow \Lambda \pi$ |
| 1671 \pm 2 | ⁴ PONTE | 75 | DPWA $K^- p \rightarrow \Lambda \pi^0$ |

 $\Sigma(1660)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|--|
| 40 to 200 (≈ 100) OUR ESTIMATE | | | |
| 81.5 \pm 22.2 | ¹ KOISO | 85 | DPWA $K^- p \rightarrow \Sigma \pi$ |
| 152 \pm 20 | GOPAL | 80 | DPWA $\bar{K} N \rightarrow \bar{K} N$ |
| 38 \pm 10 | ALSTON-... | 78 | DPWA $\bar{K} N \rightarrow \bar{K} N$ |
| 120 \pm 20 | GOPAL | 77 | DPWA $\bar{K} N$ multichannel |
| 230 $\begin{smallmatrix} +165 \\ -60 \end{smallmatrix}$ | VANHORN | 75 | DPWA $K^- p \rightarrow \Lambda \pi^0$ |
| 250 \pm 110 | KANE | 74 | DPWA $K^- p \rightarrow \Sigma \pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 202 or 217 | ² MARTIN | 77 | DPWA $\bar{K} N$ multichannel |
| 80 \pm 40 | ³ BAILLON | 75 | IPWA $\bar{K} N \rightarrow \Lambda \pi$ |
| 81 \pm 10 | ⁴ PONTE | 75 | DPWA $K^- p \rightarrow \Lambda \pi^0$ |

 $\Sigma(1660)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|-------------------------|--------------------------------|
| Γ_1 $N\bar{K}$ | 10–30 % |
| Γ_2 $\Lambda\pi$ | seen |
| Γ_3 $\Sigma\pi$ | seen |

$\Sigma(1660)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

| $\Gamma(N\bar{K})/\Gamma_{\text{total}}$ | Γ_1/Γ |
|---|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| 0.1 to 0.3 OUR ESTIMATE | |
| 0.12±0.03 | GOPAL 80 DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 0.10±0.05 | ALSTON-... 78 DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| <0.04 | GOPAL 77 DPWA See GOPAL 80 |
| 0.27 or 0.29 | ² MARTIN 77 DPWA $\bar{K}N$ multichannel |

| $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1660) \rightarrow \Lambda\pi$ | $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$ |
|--|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| < 0.04 | GOPAL 77 DPWA $\bar{K}N$ multichannel |
| 0.12 ^{+0.12} _{-0.04} | VANHORN 75 DPWA $K^-p \rightarrow \Lambda\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| -0.10 or -0.11 | ² MARTIN 77 DPWA $\bar{K}N$ multichannel |
| -0.04±0.02 | ³ BAILLON 75 IPWA $\bar{K}N \rightarrow \Lambda\pi$ |
| +0.16±0.01 | ⁴ PONTE 75 DPWA $K^-p \rightarrow \Lambda\pi^0$ |

| $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1660) \rightarrow \Sigma\pi$ | $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$ |
|---|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| -0.13±0.04 | ¹ KOISO 85 DPWA $K^-p \rightarrow \Sigma\pi$ |
| -0.16±0.03 | GOPAL 77 DPWA $\bar{K}N$ multichannel |
| -0.11±0.01 | KANE 74 DPWA $K^-p \rightarrow \Sigma\pi$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| -0.34 or -0.37 | ² MARTIN 77 DPWA $\bar{K}N$ multichannel |
| not seen | HEPP 76B DPWA $K^-N \rightarrow \Sigma\pi$ |

$\Sigma(1660)$ FOOTNOTES

- ¹ The evidence of KOISO 85 is weak.
² The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
³ From solution 1 of BAILLON 75; not present in solution 2.
⁴ From solution 2 of PONTE 75; not present in solution 1.

$\Sigma(1660)$ REFERENCES

| | | | | |
|------------|-----|-------------------|---|--------------------|
| KOISO | 85 | NP A433 619 | H. Koiso <i>et al.</i> | (TOKY, MASA) |
| PDG | 82 | PL 111B | M. Roos <i>et al.</i> | (HELSE, CIT, CERN) |
| GOPAL | 80 | Toronto Conf. 159 | G.P. Gopal | (RHEL) IJP |
| ALSTON-... | 78 | PR D18 182 | M. Alston-Garnjost <i>et al.</i> | (LBL, MTHO+) IJP |
| Also | 77 | PRL 38 1007 | M. Alston-Garnjost <i>et al.</i> | (LBL, MTHO+) IJP |
| GOPAL | 77 | NP B119 362 | G.P. Gopal <i>et al.</i> | (LOIC, RHEL) IJP |
| MARTIN | 77 | NP B127 349 | B.R. Martin, M.K. Pidcock, R.G. Moorhouse | (LOUC+) IJP |
| Also | 77B | NP B126 266 | B.R. Martin, M.K. Pidcock | (LOUC) |
| Also | 77C | NP B126 285 | B.R. Martin, M.K. Pidcock | (LOUC) IJP |

| | | | | |
|---------|-----|-------------|-------------------------------|-------------------------|
| HEPP | 76B | PL 65B 487 | V. Hepp <i>et al.</i> | (CERN, HEIDH, MPIM) IJP |
| BAILLON | 75 | NP B94 39 | P.H. Baillon, P.J. Litchfield | (CERN, RHEL) IJP |
| PONTE | 75 | PR D12 2597 | R.A. Ponte <i>et al.</i> | (MASA, TENN, UCR) IJP |
| VANHORN | 75 | NP B87 145 | A.J. van Horn | (LBL) IJP |
| Also | 75B | NP B87 157 | A.J. van Horn | (LBL) IJP |
| KANE | 74 | LBL-2452 | D.F. Kane | (LBL) IJP |
