

$\Upsilon(4S)$   
or  $\Upsilon(10580)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

### $\Upsilon(4S)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
<b>10.5800 ± 0.0035</b>	<sup>1</sup> BEBEK	87 CLEO	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10.5774 ± 0.0010	<sup>2</sup> LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons
<sup>1</sup> Reanalysis of BESSON 85.			
<sup>2</sup> No systematic error given.			

### $\Upsilon(4S)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>14 ± 5 OUR AVERAGE</b>	Error includes scale factor of 1.7.		
10.0 ± 2.8 ± 2.7	<sup>3</sup> ALBRECHT	95E ARG	$e^+e^- \rightarrow$ hadrons
20 ± 2 ± 4	BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
25 ± 2.5	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons
<sup>3</sup> Using LEYAQUANC 77 parametrization of $\Gamma(s)$ .			

### $\Upsilon(4S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $B\bar{B}$	> 96 %	95%
$\Gamma_2$ $B^+B^-$		
$\Gamma_3$ $B^0\bar{B}^0$		
$\Gamma_4$ non- $B\bar{B}$	< 4 %	95%
$\Gamma_5$ $e^+e^-$	$(2.8 \pm 0.7) \times 10^{-5}$	
$\Gamma_6$ $J/\psi(1S)$ anything	< 1.9 $\times 10^{-4}$	95%
$\Gamma_7$ $D^{*+}$ anything + c.c.	< 7.4 %	90%
$\Gamma_8$ $\phi$ anything	< 2.3 $\times 10^{-3}$	90%
$\Gamma_9$ $\Upsilon(1S)$ anything	< 4 $\times 10^{-3}$	90%
$\Gamma_{10}$ $\Upsilon(1S)\pi^+\pi^-$	< 1.2 $\times 10^{-4}$	90%
$\Gamma_{11}$ $\Upsilon(2S)\pi^+\pi^-$	< 3.9 $\times 10^{-4}$	90%


## $\Upsilon(4S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

$\Gamma_5$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.248 \pm 0.031</math> OUR AVERAGE</b>	Error includes scale factor of 1.3. See the ideogram below.		
$0.28 \pm 0.05 \pm 0.01$	<sup>4</sup> ALBRECHT	95E ARG	$e^+e^- \rightarrow$ hadrons
$0.192 \pm 0.007 \pm 0.038$	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons
$0.283 \pm 0.037$	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

<sup>4</sup> Using LEYAOUANC 77 parametrization of  $\Gamma(s)$ .



$\Gamma(e^+e^-)$  (keV)

## $\Upsilon(4S)$ BRANCHING RATIOS

$\Gamma(B^+B^-)/\Gamma(B^0\bar{B}^0)$

$\Gamma_2/\Gamma_3$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.02 \pm 0.14</math></b>	<sup>5</sup> ALEXANDER	01 CLEO	$e^+e^- \rightarrow B\bar{B}$

<sup>5</sup> Assuming isospin conservation with  $(\tau_{B^+})/(\tau_{B^0}) = 1.09^{+0.11}_{-0.10} \pm 0.08$ .

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_5/\Gamma$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.77 \pm 0.50 \pm 0.49</math></b>	<sup>6</sup> ALBRECHT	95E ARG	$e^+e^- \rightarrow$ hadrons

<sup>6</sup> Using LEYAOUANC 77 parametrization of  $\Gamma(s)$ .

**$[\Gamma(D^{*+} \text{ anything}) + \Gamma(\text{c.c.})]/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.074</b>	90	<sup>7</sup> ALEXANDER 90C	CLEO	$e^+ e^-$

<sup>7</sup> For  $x > 0.473$ .

**$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0023</b>	90	<sup>8</sup> ALEXANDER 90C	CLEO	$e^+ e^-$

<sup>8</sup> For  $x > 0.52$ .

**$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$**

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.9</b>	95	<sup>9</sup> ABE 02D	BELL	$e^+ e^- \rightarrow J/\psi X \rightarrow \ell^+ \ell^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.7	90	<sup>9</sup> AUBERT 01C	BABR	$e^+ e^- \rightarrow J/\psi X \rightarrow \ell^+ \ell^- X$
------	----	-------------------------	------	--

<sup>9</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .

**$\Gamma(\Upsilon(1S) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.004</b>	90	ALEXANDER 90C	CLEO	$e^+ e^-$

**$\Gamma(\Upsilon(1S) \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$**

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.2</b>	90	GLENN 99	CLE2	$e^+ e^-$

**$\Gamma(\Upsilon(2S) \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$**

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.9</b>	90	GLENN 99	CLE2	$e^+ e^-$

**$\Gamma(\text{non-}B\bar{B})/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.04</b>	95	BARISH 96B	CLEO	$e^+ e^-$

**$\Upsilon(4S)$  REFERENCES**

ABE	02D	PRL 88 052001	K. Abe <i>et al.</i>	(Belle Collab.)
ALEXANDER	01	PRL 86 2737	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AUBERT	01C	PRL 87 162002	B. Aubert <i>et al.</i>	(BaBar Collab.)
GLENN	99	PR D59 052003	S. Glenn <i>et al.</i>	
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
ALBRECHT	95E	ZPHY C65 619	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	90C	PRL 64 2226	J. Alexander <i>et al.</i>	(CLEO Collab.)
BEBEK	87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)
LEYAOUANC	77	PL B71 397	A. Le Yaouanc <i>et al.</i>	(ORSAY)

————— **OTHER RELATED PAPERS** —————

ABE	01J	PR D64 072001	K. Abe <i>et al.</i>	(Belle Collab.)
HENDERSON	92	PR D45 2212	S. Henderson <i>et al.</i>	(CLEO Collab.)
ANDREWS	80B	PRL 45 219	D. Andrews <i>et al.</i>	(CLEO Collab.)
FINOCCHI...	80	PRL 45 222	G. Finocchiaro <i>et al.</i>	(CUSB Collab.)

---