

$\Upsilon(1S)$

$$J^{PC} = 0^{-}(1^{-}-)$$

$\Upsilon(1S)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9460.30 ± 0.26 OUR AVERAGE	Error includes scale factor of 3.3.		
9460.51 ± 0.09 ± 0.05	¹ ARTAMONOV 00	MD1	$e^+e^- \rightarrow$ hadrons
9459.97 ± 0.11 ± 0.07	MACKAY	84 REDE	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9460.60 ± 0.09 ± 0.05	^{2,3} BARU	92B REDE	$e^+e^- \rightarrow$ hadrons
9460.59 ± 0.12	BARU	86 REDE	$e^+e^- \rightarrow$ hadrons
9460.6 ± 0.4	^{3,4} ARTAMONOV 84	REDE	$e^+e^- \rightarrow$ hadrons
¹ Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).			
² Superseding BARU 86.			
³ Superseded by ARTAMONOV 00.			
⁴ Value includes data of ARTAMONOV 82.			

$\Upsilon(1S)$ WIDTH

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
52.5 ± 1.8 OUR EVALUATION	See the Note on Width Determinations of the Υ states

$\Upsilon(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \quad \tau^+ \tau^-$	$(2.67^{+0.14}_{-0.16}) \%$	
$\Gamma_2 \quad e^+ e^-$	$(2.38 \pm 0.11) \%$	
$\Gamma_3 \quad \mu^+ \mu^-$	$(2.48 \pm 0.06) \%$	
Hadronic decays		
$\Gamma_4 \quad J/\psi(1S)$ anything	$(1.1 \pm 0.4) \times 10^{-3}$	
$\Gamma_5 \quad \rho\pi$	$< 2 \times 10^{-4}$	90%
$\Gamma_6 \quad \pi^+ \pi^-$	$< 5 \times 10^{-4}$	90%
$\Gamma_7 \quad K^+ K^-$	$< 5 \times 10^{-4}$	90%
$\Gamma_8 \quad \rho\bar{\rho}$	$< 5 \times 10^{-4}$	90%
$\Gamma_9 \quad \pi^0 \pi^+ \pi^-$	$< 1.84 \times 10^{-5}$	90%
$\Gamma_{10} \quad D^*(2010)^\pm$ anything		

Radiative decays

Γ_{11}	$\gamma\pi^+\pi^-$	$(6.3 \pm 1.8) \times 10^{-5}$	
Γ_{12}	$\gamma\pi^0\pi^0$	$(1.7 \pm 0.7) \times 10^{-5}$	
Γ_{13}	$\gamma 2h^+ 2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$	
Γ_{14}	$\gamma 3h^+ 3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$	
Γ_{15}	$\gamma 4h^+ 4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$	
Γ_{16}	$\gamma\pi^+\pi^-K^+K^-$	$(2.9 \pm 0.9) \times 10^{-4}$	
Γ_{17}	$\gamma 2\pi^+ 2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$	
Γ_{18}	$\gamma 3\pi^+ 3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$	
Γ_{19}	$\gamma 2\pi^+ 2\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$	
Γ_{20}	$\gamma\pi^+\pi^-p\bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$	
Γ_{21}	$\gamma 2\pi^+ 2\pi^- p\bar{p}$	$(4 \pm 6) \times 10^{-5}$	
Γ_{22}	$\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$	
Γ_{23}	$\gamma\eta'(958)$	$< 1.6 \times 10^{-5}$	90%
Γ_{24}	$\gamma\eta$	$< 3.5 \times 10^{-4}$	90%
Γ_{25}	$\gamma f'_2(1525)$	$< 1.4 \times 10^{-4}$	90%
Γ_{26}	$\gamma f_2(1270)$	$(8 \pm 4) \times 10^{-5}$	
Γ_{27}	$\gamma\eta(1440)$	$< 8.2 \times 10^{-5}$	90%
Γ_{28}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$< 2.6 \times 10^{-4}$	90%
Γ_{29}	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	$< 2 \times 10^{-4}$	90%
Γ_{30}	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	$< 1.5 \times 10^{-5}$	90%
Γ_{31}	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$	$< 3 \times 10^{-3}$	90%
Γ_{32}	γX $X = \text{pseudoscalar with } m < 7.2 \text{ GeV}$	$< 3 \times 10^{-5}$	90%
Γ_{33}	$\gamma X\bar{X}$ $X\bar{X} = \text{vectors with } m < 3.1 \text{ GeV}$	$< 1 \times 10^{-3}$	90%

$\mathcal{R}(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(e^+e^-) \times \Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$				$\Gamma_2\Gamma_3/\Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
31.2±1.6±1.7	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_0\Gamma_2/\Gamma$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.216±0.027 OUR AVERAGE				
1.187±0.023±0.031	⁵ BARU	92B	MD1	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.02 ±0.05	⁵ JAKUBOWSKI	88	CBAL	$e^+e^- \rightarrow \text{hadrons}$
1.37 ±0.06 ±0.09	⁶ GILES	84B	CLEO	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.08 ±0.04	⁶ ALBRECHT	82	DASP	$e^+e^- \rightarrow \text{hadrons}$
1.13 ±0.07 ±0.11	⁶ NICZYPORUK	82	LENA	$e^+e^- \rightarrow \text{hadrons}$
1.09 ±0.25	⁶ BOCK	80	CNTR	$e^+e^- \rightarrow \text{hadrons}$
1.35 ±0.14	⁷ BERGER	79	PLUT	$e^+e^- \rightarrow \text{hadrons}$

⁵ Radiative corrections evaluated following KURAEV 85.

⁶ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

⁷ Radiative corrections reevaluated by ALEXANDER 89 using $B(\mu\mu) = 0.026$.

$\Upsilon(1S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$				Γ_2
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.32 \pm 0.04 \pm 0.03$	⁸ ALBRECHT	95E ARG	$e^+e^- \rightarrow$ hadrons	
⁸ Applying the formula of Kuraev and Fadin.				

$\Upsilon(1S)$ BRANCHING RATIOS

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$				Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0267^{+0.0014}_{-0.0016}$	OUR AVERAGE			
$0.0261 \pm 0.0012^{+0.0009}_{-0.0013}$	25k	CINABRO	94B CLE2	$e^+e^- \rightarrow \tau^+\tau^-$
$0.027 \pm 0.004 \pm 0.002$		⁹ ALBRECHT	85C ARG	$\Upsilon(2S) \rightarrow$ $\pi^+\pi^-\tau^+\tau^-$
$0.034 \pm 0.004 \pm 0.004$		GILES	83 CLEO	$e^+e^- \rightarrow \tau^+\tau^-$
⁹ Using $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.				

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$				Γ_3/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0248 ± 0.0006	OUR AVERAGE			
$0.0249 \pm 0.0008 \pm 0.0013$		ALEXANDER	98 CLE2	$\Upsilon(2S) \rightarrow$ $\pi^+\pi^-\mu^+\mu^-$
$0.0212 \pm 0.0020 \pm 0.0010$		¹⁰ BARU	92 MD1	$e^+e^- \rightarrow$ $\mu^+\mu^-$
$0.0231 \pm 0.0012 \pm 0.0010$		¹⁰ KOBEL	92 CBAL	$e^+e^- \rightarrow$ $\mu^+\mu^-$
$0.0252 \pm 0.0007 \pm 0.0007$		CHEN	89B CLEO	$e^+e^- \rightarrow$ $\mu^+\mu^-$
$0.0261 \pm 0.0009 \pm 0.0011$		KAARSBERG	89 CSB2	$e^+e^- \rightarrow$ $\mu^+\mu^-$
$0.0230 \pm 0.0025 \pm 0.0013$	86	ALBRECHT	87 ARG	$\Upsilon(2S) \rightarrow$ $\pi^+\pi^-\mu^+\mu^-$
$0.029 \pm 0.003 \pm 0.002$	864	BESSON	84 CLEO	$\Upsilon(2S) \rightarrow$ $\pi^+\pi^-\mu^+\mu^-$
$0.027 \pm 0.003 \pm 0.003$		ANDREWS	83 CLEO	$e^+e^- \rightarrow$ $\mu^+\mu^-$
$0.032 \pm 0.013 \pm 0.003$		ALBRECHT	82 DASP	$e^+e^- \rightarrow$ $\mu^+\mu^-$
$0.038 \pm 0.015 \pm 0.002$		NICZYPORUK	82 LENA	$e^+e^- \rightarrow$ $\mu^+\mu^-$
$0.014^{+0.034}_{-0.014}$		BOCK	80 CNTR	$e^+e^- \rightarrow$ $\mu^+\mu^-$
0.022 ± 0.020		BERGER	79 PLUT	$e^+e^- \rightarrow$ $\mu^+\mu^-$

¹⁰ Taking into account interference between the resonance and continuum.

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$					Γ_2/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.0238 ± 0.0011 OUR AVERAGE					
0.0229 ± 0.0008 ± 0.0011		ALEXANDER	98 CLE2	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$	
0.0242 ± 0.0014 ± 0.0014	307	ALBRECHT	87 ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$	
0.028 ± 0.003 ± 0.002	826	BESSION	84 CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$	
0.051 ± 0.030		BERGER	80c PLUT	$e^+e^- \rightarrow e^+e^-$	

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$					Γ_4/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 0.68	90	ALBRECHT	92J ARG	$e^+e^- \rightarrow e^+e^-X$, $e^+e^- \rightarrow \mu^+\mu^-X$	
1.1 ± 0.4 ± 0.2		¹¹ FULTON	89 CLEO	$e^+e^- \rightarrow \mu^+\mu^-X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.7	90	MASCHMANN	90 CBAL	$e^+e^- \rightarrow \text{hadrons}$	
< 20	90	NICZYPORUK	83 LENA		

¹¹ Using $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_6/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 5	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow \pi^+\pi^-$	

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$					Γ_7/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 5	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow K^+K^-$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$					Γ_8/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 5	90	¹² BARU	96 MD1	$\Upsilon(1S) \rightarrow p\bar{p}$	

¹² Supersedes BARU 92 in this node.

$\Gamma(\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_9/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 1.84	90	ANASTASSOV	99 CLE2	$e^+e^- \rightarrow \text{hadrons}$	

$\Gamma(\gamma X)/\Gamma_{\text{total}}$					Γ_{32}/Γ
<i>(X = pseudoscalar with $m < 7.2$ GeV)</i>					
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 3	90	¹³ BALEST	95 CLEO	$e^+e^- \rightarrow \gamma + X$	

¹³ For a noninteracting pseudoscalar X with mass < 7.2 GeV.

$\Gamma(\gamma X \bar{X})/\Gamma_{\text{total}}$ Γ_{33}/Γ

($X \bar{X}$ = vectors with $m < 3.1$ GeV)

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	¹⁴ BALEST	95	CLEO $e^+ e^- \rightarrow \gamma + X \bar{X}$

¹⁴ For a noninteracting vector X with mass < 3.1 GeV.

$\Gamma(\gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$6.3 \pm 1.2 \pm 1.3$	¹⁵ ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow$ hadrons

¹⁵ For $m_{\pi\pi} > 1$ GeV.

$\Gamma(\gamma \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.7 \pm 0.6 \pm 0.3$	¹⁶ ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow$ hadrons

¹⁶ For $m_{\pi\pi} > 1$ GeV.

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 0.7 \pm 0.5$	26 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.9 \pm 0.7 \pm 0.6$	29 ± 8	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma \pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.5 \pm 0.3$	22 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.2 ± 0.2	2 ± 2	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 0.9 \pm 0.8$	17 ± 5	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 0.9 \pm 0.8$	18 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma 2\pi^+ 2\pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.4 \pm 0.4 \pm 0.4$	7 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.0 \pm 1.1 \pm 1.0$	80 ± 12	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.4 \pm 1.5 \pm 1.3$	39 ± 11	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.4 \pm 2.5 \pm 2.5$	36 ± 12	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 10	90	BLINOV	90 MD1	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
< 21	90	NICZYPORUK	83 LENA	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$

$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 19	90	¹⁷ ALBRECHT	92J ARG	$e^+ e^- \rightarrow D^0 \pi^\pm X$
¹⁷ For $x_p > 0.2$.				

$\Gamma(\gamma\eta(1440))/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 8.2	90	¹⁸ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ \pi^\mp K_S^0$
¹⁸ Includes unknown branching ratio of $\eta(1440) \rightarrow K^\pm \pi^\mp K_S^0$.				

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.6	90	RICHICHI	01B CLE2	$\Upsilon(1S) \rightarrow \gamma\eta' \rightarrow \gamma\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 130	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.5	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<14	90	¹⁹ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<19.4	90	¹⁹ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
¹⁹ Assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.				

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.6	90	²⁰ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 6.3	90	²⁰ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<19	90	²⁰ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$
< 8	90	²¹ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<24	90	²² SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$
²⁰ Assuming $B(f_0(1710) \rightarrow K\bar{K}) = 0.38$.				
²¹ Assuming $B(f_0(1710) \rightarrow \pi\pi) = 0.04$.				
²² Assuming $B(f_0(1710) \rightarrow \eta\eta) = 0.18$.				

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$8.1 \pm 2.3^{+2.9}_{-2.7}$		²³ ANASTASSOV	99 CLE2	$e^+ e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<21	90	²³ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<13	90	²³ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<81	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$
²³ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.				

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5	90	²⁴ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 2.9	90	²⁴ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<20	90	²⁴ BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
²⁴ Including unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.				

$\Gamma(\gamma \eta(2225) \rightarrow \gamma \phi\phi)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	²⁵ BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$
²⁵ Assuming that the $\eta(2225)$ decays only into $\phi\phi$.				

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{29} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0002	90	²⁶ BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

²⁶ Assuming that the $f_0(2200)$ decays only into $K^+ K^-$.

 $\Upsilon(1S)$ REFERENCES

RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	95E	ZPHY C65 619	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	E.A. KuraeV, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)

OTHER RELATED PAPERS

KOENIGS...	86	DESY 86/136	K. Koenigsmann	(DESY)
ALBRECHT	84	PL 134B 137	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
BERGER	78	PL 76B 243	C. Berger <i>et al.</i>	(PLUTO Collab.)
BIENLEIN	78	PL 78B 360	J.K. Bienlein <i>et al.</i>	(DESY, HAMB, HEIDP+)
DARDEN	78	PL 76B 246	C.W. Darden <i>et al.</i>	(DESY, DORT, HEIDH+)
GARELICK	78	PR D18 945	D.A. Garelick <i>et al.</i>	(NEAS, WASH, TUFTS)

KAPLAN	78	PRL 40 435	D.M. Kaplan <i>et al.</i>	(STON, FNAL, COLU)
YOH	78	PRL 41 684	J.K. Yoh <i>et al.</i>	(COLU, FNAL, STON)
COBB	77	PL 72B 273	J.H. Cobb <i>et al.</i>	(BNL, CERN, SYRA, YALE)
HERB	77	PRL 39 252	S.W. Herb <i>et al.</i>	(COLU, FNAL, STON)
INNES	77	PRL 39 1240	W.R. Innes <i>et al.</i>	(COLU, FNAL, STON)
