$\chi_{b1}(2P)$

$I^G(J^{PC}) = 0^+(1^{++})$
J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

$\chi_{b1}(2P)$ MASS

VALUE (GeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
10.2552±0.0005 OUR AVERAGE				
10.2547±0.0004±0.0010	[1] HEINTZ	92	CSB2	$e^+e^- \to \gamma X, \ell^+\ell^-\gamma\gamma$
10.2553±0.0005	[2] MORRISON	91	CLE2	$e^+e^- \to \gamma X$

[1] From the average photon energy for inclusive and exclusive events and assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. Supersedes HEINTZ 91 and NARAIN 91.
[2] From γ energy below assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. The error on the $\Upsilon(3S)$ mass is not included in the individual measurements. It is included in the final evaluation.

$m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
23.5±0.7±0.7	[3] HEINTZ	92	CSB2	$e^+e^- \to \gamma X, \ell^+\ell^-\gamma\gamma$

[3] From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
99.90±0.26 OUR AVERAGE					
99 ±1	169	CRAWFORD	92B	CLE2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$
100.1 ±0.4	11147	[4] HEINTZ	92	CSB2	$e^+e^- \to \gamma X$
100.2 ±0.5	223	[5] HEINTZ	92	CSB2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$
99.5 ±0.1 ±0.5	25759	MORRISON	91	CLE2	$e^+e^- \to \gamma X$

[4] A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.
[5] A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

$\chi_{b1}(2P)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1	$\gamma\,\Upsilon(2S)$	(21 ±4) %	1.5
Γ_2	$\gamma\,\Upsilon(1S)$	(8.5±1.3) %	1.3

$\chi_{b1}(2P)$ BRANCHING RATIOS

$\Gamma(\gamma\Upsilon(2S))/\Gamma_{total}$ Γ_1/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.21 ±0.04 OUR AVERAGE	Error includes scale factor of 1.5.			
$0.356 \pm 0.042 \pm 0.092$	[6] CRAWFORD	92B	CLE2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$
$0.199 \pm 0.020 \pm 0.022$	[7] HEINTZ	92	CSB2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$

[6] Using B($\Upsilon(2S) \to \mu^+\mu^-$) = (1.37 ± 0.26)%, B($\Upsilon(3S) \to \gamma\gamma\Upsilon(2S)$)×2 B($\Upsilon(2S) \to \mu^+\mu^-$) = $(10.23 \pm 1.20 \pm 1.26) \times 10^{-4}$, and B($\Upsilon(3S) \to \gamma\chi_{b1}(2P)$) = $0.105^{+0.003}_{-0.002} \pm 0.013$.

[7] Using B($\Upsilon(2S) \to \mu^+\mu^-$) = (1.44 ± 0.10)%, B($\Upsilon(3S) \to \gamma\chi_{b1}(2P)$) = $(11.5 \pm 0.5 \pm 0.5)$% and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma\Upsilon(1S))/\Gamma_{total}$ Γ_2/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.085±0.013 OUR AVERAGE	Error includes scale factor of 1.3.			
$0.120 \pm 0.021 \pm 0.021$	[8] CRAWFORD	92B	CLE2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$
$0.080 \pm 0.009 \pm 0.007$	[9] HEINTZ	92	CSB2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$

[8] Using B($\Upsilon(1S) \to \mu^+\mu^-$) = (2.57 ± 0.07)%, B($\Upsilon(3S) \to \gamma\gamma\Upsilon(1S)$)×2 B($\Upsilon(1S) \to \mu^+\mu^-$) = $(6.47 \pm 1.12 \pm 0.82) \times 10^{-4}$ and B($\Upsilon(3S) \to \gamma\chi_{b1}(2P)$) = $0.105^{+0.003}_{-0.002} \pm 0.013$.

[9] Using B($\Upsilon(1S) \to \mu^+\mu^-$)=$(2.57 \pm 0.07)$%, B($\Upsilon(3S) \to \gamma\chi_{b1}(2P)$) = $(11.5 \pm 0.5 \pm 0.5)$% and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\chi_{b1}(2P)$ REFERENCES

CRAWFORD	92B	PL B294 139	G. Crawford, R. Fulton	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz *et al.*	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz *et al.*	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison *et al.*	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain *et al.*	(CUSB Collab.)

OTHER RELATED PAPERS

EIGEN	82	PRL 49 1616	G. Eigen *et al.*	(CUSB Collab.)
HAN	82	PRL 49 1612	K. Han *et al.*	(CUSB Collab.)