K^0_S

$I(J^P) = \frac{1}{2}(0^-)$

K^0_S MEAN LIFE

For earlier measurements, beginning with BOLDT 58B, see our our 1986 edition, Physics Letters **170B** 130 (1986).

OUR FIT is described in the note on "Fits for K^0_L *CP*-Violation Parameters" in the K^0_L Particle Listings.

VALUE (10^{-10} s)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.8935±0.0008 OUR FIT	Assuming *CPT*				
0.8937±0.0012 OUR FIT	Not assuming *CPT*				
0.8941±0.0009 OUR AVERAGE					
0.8971±0.0021		BERTANZA	97	NA31	
0.8941±0.0014±0.0009		SCHWINGEN...95		E773	Δm free, $\phi_{+-}=\phi_{SW}$
0.8929±0.0016		GIBBONS	93	E731	Δm free, $\phi_{+-}=\phi_{SW}$
0.8920±0.0044	214k	GROSSMAN	87	SPEC	
0.8924±0.0032		[1] CARITHERS	75	SPEC	
0.8937±0.0048	6M	GEWENIGER	74B	ASPK	
0.8958±0.0045	50k	[2] SKJEGGEST...	72	HBC	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.905 ±0.007		[3] ARONSON	82B	SPEC	
0.881 ±0.009	26k	ARONSON	76	SPEC	
0.856 ±0.008	19994	[4] DONALD	68B	HBC	
0.872 ±0.009	20000	[4,5] HILL	68	DBC	

[1] CARITHERS 75 value is for $m_{K^0_L} - m_{K^0_S}$ $\Delta m = 0.5301 \pm 0.0013$. The Δm dependence of the total decay rate (inverse mean life) is $\Gamma(K^0_S) = [(1.122 \pm 0.004) + 0.16(\Delta m - 0.5348)/\Delta m]10^{10}$/s, or, in terms of meanlife $\tau_s = 0.8913 \pm 0.0032 - 0.238(\Delta m - 0.5348)$ where Δm and τ_s are in units of $10^{10}\hbar s^{-1}$ and 10^{-10}s respectively.

[2] HILL 68 has been changed by the authors from the published value (0.865 ± 0.009) because of a correction in the shift due to η_{+-}. SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.

[3] ARONSON 82 find that K^0_S mean life may depend on the kaon energy.

[4] Pre-1971 experiments are excluded from the average because of disagreement with later more precise experiments.

[5] HILL 68 has been changed by the authors from the published value (0.865 ± 0.009) because of a correction in the shift due to η_{+-}. SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.

K^0_S DECAY MODES

	Mode		Fraction (Γ_i/Γ)	Scale factor/ Confidence level
	Hadronic modes			
Γ_1	$\pi^0\pi^0$		(31.40 ± 0.27) %	S=1.2
Γ_2	$\pi^+\pi^-$		(68.60 ± 0.27) %	S=1.2
Γ_3	$\pi^+\pi^-\pi^0$		$(3.2\ ^{+1.2}_{-1.0}) \times 10^{-7}$	
	Modes with photons or $\ell\bar{\ell}$ pairs			
Γ_4	$\pi^+\pi^-\gamma$	[a,b]	$(1.78 \pm 0.05) \times 10^{-3}$	
Γ_5	$\pi^+\pi^- e^+ e^-$		$(4.5 \pm 0.8) \times 10^{-5}$	
Γ_6	$\gamma\gamma$		$(2.5 \pm 0.4) \times 10^{-6}$	
	Semileptonic modes			
Γ_7	$\pi^\pm e^\mp \nu_e$	[c]	$(7.2 \pm 1.4) \times 10^{-4}$	
Γ_8	$\pi^\pm \mu^\mp \nu_\mu$	[c]		
	***CP* violating (*CP*) and $\Delta S = 1$ weak neutral current (*S1*) modes**			
Γ_9	$3\pi^0$	*CP*	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{10}	$\mu^+\mu^-$	*S1*	$< 3.2 \times 10^{-7}$	CL=90%
Γ_{11}	$e^+ e^-$	*S1*	$< 1.4 \times 10^{-7}$	CL=90%
Γ_{12}	$\pi^0 e^+ e^-$	*S1*	$< 1.4 \times 10^{-7}$	CL=90%

[a] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ's.

[b] See the Particle Listings below for the energy limits used in this measurement.

[c] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 15.5$ for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	−100
	x_1

K^0_S DECAY RATES

$\Gamma(\pi^\pm e^\mp \nu_e)$ Γ_7

VALUE (10^6 s^{-1})	EVTS	DOCUMENT ID		TECN	COMMENT
8.1 ±1.6	75	[6] AKHMETSHIN	99	CMD2	Tagged K^0_S using $\phi \to K^0_L K^0_S$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.50±0.08	[7] PDG	98		
seen	BURGUN	72	HBC	$K^+ p \to K^0 p\pi^+$
9.3 ±2.5	AUBERT	65	HLBC	ΔS=ΔQ, *CP* cons. not assumed

[6] AKHMETSHIN 99 is from a measured branching ratio $B(K^0_S \to \pi e \nu_e) = (7.2 \pm 1.4) \times 10^{-4}$ and $\tau_{K^0_S} = (0.8934 \pm 0.0008) \times 10^{-10}$ s. Not independent of measured branching ratio.

[7] PDG 98 from K^0_L measurements, assuming that $\Delta S = \Delta Q$ in K^0 decay so that $\Gamma(K^0_S \to \pi^\pm e^\mp \nu_e) = \Gamma(K^0_L \to \pi^\pm e^\mp \nu_e)$.

$\Gamma(\pi^\pm \mu^\mp \nu_\mu)$ Γ_8

VALUE (10^6 s^{-1})	DOCUMENT ID		

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.25±0.07	[8] PDG	98	

[8] PDG 98 from K^0_L measurements, assuming that $\Delta S = \Delta Q$ in K^0 decay so that $\Gamma(K^0_S \to \pi^\pm \mu^\mp \nu_\mu) = \Gamma(K^0_L \to \pi^\pm \mu^\mp \nu_\mu)$.

K^0_S BRANCHING RATIOS

Hadronic modes

$\Gamma(\pi^0\pi^0)/\Gamma_{total}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID		TECN
0.3140±0.0027 OUR FIT Error includes scale factor of 1.2.				
0.318 ±0.015 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.				
0.335 ±0.014	1066	BROWN	63	HLBC
0.288 ±0.021	198	CHRETIEN	63	HLBC
0.30 ±0.035		BROWN	61	HLBC



$\Gamma\left(\pi^0\,\pi^0\right)/\Gamma_{\text{total}}$

$\Gamma\left(\pi^+\pi^-\right)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.6860±0.0027 OUR FIT Error includes scale factor of 1.2.					
0.670 ±0.010	3447	DOYLE	69	HBC	$\pi^- p \rightarrow \Lambda K^0$

$\Gamma\left(\pi^+\pi^-\right)/\Gamma\left(\pi^0\pi^0\right)$ Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
2.185±0.027 OUR FIT Error includes scale factor of 1.2.					
2.197±0.026 OUR AVERAGE					
2.11 ±0.09	1315	EVERHART	76	WIRE	$\pi^- p \rightarrow \Lambda K^0$
2.169±0.094	16k	COWELL	74	OSPK	$\pi^- p \rightarrow \Lambda K^0$
2.16 ±0.08	4799	HILL	73	DBC	$K^+ d \rightarrow K^0 p p$
2.22 ±0.10	3068	[9] ALITTI	72	HBC	$K^+ p \rightarrow \pi^+ p K^0$
2.22 ±0.08	6380	MORSE	72B	DBC	$K^+ n \rightarrow K^0 p$
2.10 ±0.11	701	[10] NAGY	72	HLBC	$K^+ n \rightarrow K^0 p$
2.22 ±0.095	6150	[11] BALTAY	71	HBC	$K p \rightarrow K^0$ neutrals
2.282±0.043	7944	[12] MOFFETT	70	OSPK	$K^+ n \rightarrow K^0 p$
2.10 ±0.06	3700	MORFIN	69	HLBC	$K^+ n \rightarrow K^0 p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.12 ±0.17	267	[10] BOZOKI	69	HLBC	
2.285±0.055	3016	[12] GOBBI	69	OSPK	$K^+ n \rightarrow K^0 p$

[9] The directly measured quantity is $K^0_S \rightarrow \pi^+\pi^-/\text{all}$ $K^0 = 0.345 \pm 0.005$.
[10] NAGY 72 is a final result which includes BOZOKI 69.
[11] The directly measured quantity is $K^0_S \rightarrow \pi^+\pi^-/\text{all}$ $\overline{K}^0 = 0.345 \pm 0.005$.
[12] MOFFETT 70 is a final result which includes GOBBI 69.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_3/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID		TECN	COMMENT
$3.2^{+1.2}_{-1.0}$ OUR AVERAGE					
$2.5^{+1.3}_{-1.0}{}^{+0.5}_{-0.6}$	500k	[13] ADLER	97B	CPLR	
$4.8^{+2.2}_{-1.6}\pm1.1$		[14] ZOU	96	E621	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$4.1^{+2.5}_{-1.9}{}^{+0.5}_{-0.6}$		[15] ADLER	96E	CPLR	Sup. by ADLER 97B
$3.9^{+5.4}_{-1.8}{}^{+0.9}_{-0.7}$		[16] THOMSON	94	E621	Sup. by ZOU 96

[13] ADLER 97B find the *CP*-conserving parameters $\mathrm{Re}(\lambda) = (28 \pm 7 \pm 3) \times 10^{-3}$, $\mathrm{Im}(\lambda) = (-10 \pm 8 \pm 2) \times 10^{-3}$. They estimate $B(K^0_S \to \pi^+\pi^-\pi^0)$ from $\mathrm{Re}(\lambda)$ and the K^0_L decay parameters. See also ANGELOPOULOS 98C.

[14] ZOU 96 is from the the measured quantities $|\rho_{+-0}| = 0.039^{+0.009}_{-0.006} \pm 0.005$ and $\phi_\rho = (-9 \pm 18)^\circ$.

[15] ADLER 96E is from the measured quantities $\mathrm{Re}(\lambda) = 0.036 \pm 0.010^{+0.002}_{-0.003}$ and $\mathrm{Im}(\lambda)$ consistent with zero. Note that the quantity λ is the same as ρ_{+-0} used in other footnotes.

[16] THOMSON 94 calculates this branching ratio from their measurements $|\rho_{+-0}| = 0.035^{+0.019}_{-0.011} \pm 0.004$ and $\phi_\rho = (-59 \pm 48)^\circ$ where $|\rho_{+-0}|e^{i\phi\rho} = A(K^0_S \to \pi^+\pi^-\pi^0, I = 2)/A(K^0_L \to \pi^+\pi^-\pi^0)$.

Modes with photons or $\ell\bar{\ell}$ pairs

$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-)$ Γ_4/Γ_2

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
2.60 ± 0.08 OUR AVERAGE					
2.56 ± 0.09	1286	RAMBERG	93	E731	p_γ >50 MeV/c
2.68 ± 0.15		[17] TAUREG	76	SPEC	p_γ >50 MeV/c
$2.8\ \pm0.6$		[18] BURGUN	73	HBC	p_γ >50 MeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.10 ± 0.22	3723	RAMBERG	93	E731	p_γ >20 MeV/c
$3.0\ \pm0.6$	29	[19] BOBISUT	74	HLBC	p_γ >40 MeV/c

[17] TAUREG 76 find direct emission contribution <0.06, CL = 90%.

[18] BURGUN 73 estimates that direct emission contribution is 0.3 ± 0.6.

[19] BOBISUT 74 not included in average because p_γ cut differs. Estimates direct emission contribution to be 0.5 or less, CL = 95%.

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{total}$ Γ_5/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID		TECN
$4.5\pm0.7\pm0.4$	56	LAI	00B	NA48

$\Gamma(\gamma\gamma)/\Gamma_{total}$ Γ_6/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID		TECN
2.5 ±0.4 OUR AVERAGE					
2.58±0.36±0.22		149	LAI	00	NA48
2.4 ±0.9		35	[20] BARR	95B	NA31
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.2 ±1.1		16	[21] BARR	95B	NA31
< 13	90		BALATS	89	SPEC
2.4 ±1.2		19	BURKHARDT	87	NA31
<133	90		BARMIN	86B	XEBC

[20] BARR 95B quotes this as the combined BARR 95B + BURKHARDT 87 result after rescaling BURKHARDT 87 to use same branching ratios and lifetimes as BARR 95B.
[21] BARR 95B result is calculated using $B(K_L \to \gamma\gamma) = (5.86 \pm 0.17) \times 10^{-4}$.

Semileptonic modes

$\Gamma(\pi^\pm e^\mp \nu_e)/\Gamma_{total}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2±1.4	75	AKHMETSHIN 99	CMD2	Tagged K^0_S using $\phi \to K^0_L K^0_S$

CP violating (*CP*) and $\Delta S = 1$ weak neutral current (*S1*) modes

$\Gamma(3\pi^0)/\Gamma_{total}$ Γ_9/Γ

Violates *CP* conservation.

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID		TECN
<1.4	90	7M	ACHASOV	99D	SND
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.9	90	17300	[22] ANGELOPO...	98B	CPLR
<3.7	90		BARMIN	83	HLBC

[22] ANGELOPOULOS 98B is from $\mathrm{Im}(\eta_{000}) = -0.05 \pm 0.12 \pm 0.05$, assuming $\mathrm{Re}(\eta_{000}) = \mathrm{Re}(\epsilon) = 1.635 \times 10^{-3}$ and using the value $B(K^0_L \to \pi^0\pi^0\pi^0) = 0.2112 \pm 0.0027$.

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$ Γ_{10}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN
<0.032	90	GJESDAL	73	ASPK
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.7	90	HYAMS	69B	OSPK

$\Gamma(e^+e^-)/\Gamma_{total}$ Γ_{11}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
< 1.4	90		ANGELOPO...	97	CPLR	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 28	90	0	BLICK	94	CNTR	Hyperon facility
<100	90		BARMIN	86	XEBC	

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{total}$ **Γ_{12}/Γ**

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-6})	*CL%*	*EVTS*	*DOCUMENT ID*		*TECN*
< 0.14	90		LAI	01	NA48
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.1	90	0	BARR	93B	NA31
<45	90		GIBBONS	88	E731

CP VIOLATION IN $K_S \to 3\pi$

Written 1996 by T. Nakada (Paul Scherrer Institute) and L. Wolfenstein (Carnegie-Mellon University).

The possible final states for the decay $K^0 \to \pi^+\pi^-\pi^0$ have isospin $I = 0$, 1, 2, and 3. The $I = 0$ and $I = 2$ states have $CP = +1$ and K_S can decay into them without violating CP symmetry, but they are expected to be strongly suppressed by centrifugal barrier effects. The $I = 1$ and $I = 3$ states, which have no centrifugal barrier, have $CP = -1$ so that the K_S decay to these requires CP violation.

In order to see CP violation in $K_S \to \pi^+\pi^-\pi^0$, it is necessary to observe the interference between K_S and K_L decay, which determines the amplitude ratio

$$\eta_{+-0} = \frac{A(K_S \to \pi^+\pi^-\pi^0)}{A(K_L \to \pi^+\pi^-\pi^0)} \ . \qquad (1)$$

If η_{+-0} is obtained from an integration over the whole Dalitz plot, there is no contribution from the $I = 0$ and $I = 2$ final states and a nonzero value of η_{+-0} is entirely due to CP violation.

Only $I = 1$ and $I = 3$ states, which are $CP = -1$, are allowed for $K^0 \to \pi^0\pi^0\pi^0$ decays and the decay of K_S into $3\pi^0$ is an unambiguous sign of CP violation. Similarly to η_{+-0}, η_{000} is defined as

$$\eta_{000} = \frac{A(K_S \to \pi^0\pi^0\pi^0)}{A(K_L \to \pi^0\pi^0\pi^0)} \ . \qquad (2)$$

If one assumes that CPT invariance holds and that there are no transitions to $I = 3$ (or to nonsymmetric $I = 1$ states), it can be shown that

$$\eta_{+-0} = \eta_{000} = \epsilon + i\frac{\text{Im } a_1}{\text{Re } a_1} \ . \tag{3}$$

With the Wu-Yang phase convention, a_1 is the weak decay amplitude for K^0 into $I = 1$ final states; ϵ is determined from CP violation in $K_L \to 2\pi$ decays. The real parts of η_{+-0} and η_{000} are equal to Re(ϵ). Since currently-known upper limits on $|\eta_{+-0}|$ and $|\eta_{000}|$ are much larger than $|\epsilon|$, they can be interpreted as upper limits on Im(η_{+-0}) and Im(η_{000}) and so as limits on the CP-violating phase of the decay amplitude a_1.

CP-VIOLATION PARAMETERS IN K^0_S DECAY

Im(η_{+-0})^2 = $\Gamma(K^0_S \to \pi^+\pi^-\pi^0$, *CP*-violating) / $\Gamma(K^0_L \to \pi^+\pi^-\pi^0)$

CPT assumed valid (i.e. Re(η_{+-0}) $\simeq$ 0).

VALUE	CL%	EVTS	DOCUMENT ID		TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.23	90	601	[23] BARMIN	85	HLBC
<0.12	90	384	METCALF	72	ASPK

[23] BARMIN 85 find Re(η_{+-0}) = (0.05 ± 0.17) and Im(η_{+-0}) = (0.15 ± 0.33). Includes events of BALDO-CEOLIN 75.

Im(η_{+-0}) = Im(A($K^0_S \to \pi^+\pi^-\pi^0$, *CP*-violating) / A($K^0_L \to \pi^+\pi^-\pi^0$))

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$-0.002 \pm 0.009^{+0.002}_{-0.001}$	500k	[24] ADLER	97B	CPLR	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$-0.002 \pm 0.018 \pm 0.003$	137k	[25] ADLER	96D	CPLR	Sup. by ADLER 97B
$-0.015 \pm 0.017 \pm 0.025$	272k	[26] ZOU	94	SPEC	

[24] ADLER 97B also find Re(η_{+-0}) = $-0.002 \pm 0.007^{+0.004}_{-0.001}$. See also ANGELOPOULOS 98C.

[25] The ADLER 96D fit also yields Re(η_{+-0}) = 0.006 ± 0.013 ± 0.001 with a correlation +0.66 between real and imaginary parts. Their results correspond to $|\eta_{+-0}| < 0.037$ with 90% CL.

[26] ZOU 94 use theoretical constraint Re(η_{+-0}) = Re(ϵ) = 0.0016. Without this constraint they find Im(η_{+-0}) = 0.019 ± 0.061 and Re(η_{+-0}) = 0.019 ± 0.027.

$\mathrm{Im}(\eta_{000})^2 = \Gamma(K^0_S \to 3\pi^0) / \Gamma(K^0_L \to 3\pi^0)$

CPT assumed valid (i.e. $\mathrm{Re}(\eta_{000}) \simeq 0$). This limit determines branching ratio $\Gamma(3\pi^0)/\Gamma_{\mathrm{total}}$ above.

VALUE	*CL%*	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.1	90	632	[27] BARMIN	83	HLBC	
<0.28	90		[28] GJESDAL	74B	SPEC	Indirect meas.

[27] BARMIN 83 find $\mathrm{Re}(\eta_{000}) = (-0.08 \pm 0.18)$ and $\mathrm{Im}(\eta_{000}) = (-0.05 \pm 0.27)$. Assuming *CPT* invariance they obtain the limit quoted above.

[28] GJESDAL 74B uses $K2\pi$, $K_{\mu 3}$, and K_{e3} decay results, unitarity, and *CPT*. Calculates $|(\eta_{000})| = 0.26 \pm 0.20$. We convert to upper limit.

$\mathrm{Im}(\eta_{000}) = \mathrm{Im}(A(K^0_S \to \pi^0\pi^0\pi^0)/A(K^0_L \to \pi^0\pi^0\pi^0))$

$K^0_S \to \pi^0\pi^0\pi^0$ violates *CP* conservation, in contrast to $K^0_S \to \pi^+\pi^-\pi^0$ which has a *CP*-conserving part.

VALUE	*EVTS*	*DOCUMENT ID*		*TECN*
$-0.05 \pm 0.12 \pm 0.05$	17300	[29] ANGELOPO...	98B	CPLR

[29] ANGELOPOULOS 98B assumes $\mathrm{Re}(\eta_{000}) = \mathrm{Re}(\epsilon) = 1.635 \times 10^{-3}$. Without assuming *CPT* invariance, they obtain $\mathrm{Re}(\eta_{000}) = 0.18 \pm 0.14 \pm 0.06$ and $\mathrm{Im}(\eta_{000}) = 0.15 \pm 0.20 \pm 0.03$.

K^0_S REFERENCES

LAI	01	PL B514 253	A. Lai *et al.*	(CERN NA48 Collab.)
LAI	00	PL B493 29	A. Lai *et al.*	(CERN NA48 Collab.)
LAI	00B	PL B496 137	A. Lai *et al.*	(CERN NA48 Collab.)
ACHASOV	99D	PL B459 674	M.N. Achasov *et al.*	
AKHMETSHIN	99	PL B456 90	R.R. Akhmetshin *et al.*	(Novosibirsk CMD-2 Collab.)
ANGELOPO...	98B	PL B425 391	A. Angelopoulos *et al.*	(CPLEAR Collab.)
ANGELOPO...	98C	EPJ C5 389	A. Angelopoulos *et al.*	(CPLEAR Collab.)
PDG	98	EPJ C3 1	C. Caso *et al.*	
ADLER	97B	PL B407 193	R. Adler *et al.*	(CPLEAR Collab.)
ANGELOPO...	97	PL B413 232	A. Angelopoulos *et al.*	(CPLEAR Collab.)
BERTANZA	97	ZPHY C73 629	L. Bertanza	(PISA, CERN, EDIN, MANZ, ORSAY+)
ADLER	96D	PL B370 167	R. Adler *et al.*	(CPLEAR Collab.)
ADLER	96E	PL B374 313	R. Adler *et al.*	(CPLEAR Collab.)
ZOU	96	PL B369 362	Y. Zou *et al.*	(RUTG, MINN, MICH)
BARR	95B	PL B351 579	G.D. Barr *et al.*	(CERN, EDIN, MANZ, LALO+)
SCHWINGEN...	95	PRL 74 4376	B. Schwingenheuer *et al.*	(EFI, CHIC+)
BLICK	94	PL B334 234	A.M. Blick *et al.*	(SERP, JINR)
THOMSON	94	PL B337 411	G.B. Thomson *et al.*	(RUTG, MINN, MICH)
ZOU	94	PL B329 519	Y. Zou *et al.*	(RUTG, MINN, MICH)
BARR	93B	PL B304 381	G.D. Barr *et al.*	(CERN, EDIN, MANZ, LALO+)
GIBBONS	93	PRL 70 1199	L.K. Gibbons *et al.*	(FNAL E731 Collab.)
Also	97	PR D55 6625	L.K. Gibbons *et al.*	(FNAL E731 Collab.)
RAMBERG	93	PRL 70 2525	E. Ramberg *et al.*	(FNAL E731 Collab.)
BALATS	89	SJNP 49 828	M.Y. Balats *et al.*	(ITEP)
		Translated from YAF 49 1332.		
GIBBONS	88	PRL 61 2661	L.K. Gibbons *et al.*	(FNAL E731 Collab.)
BURKHARDT	87	PL B199 139	H. Burkhardt *et al.*	(CERN, EDIN, MANZ+)
GROSSMAN	87	PRL 59 18	N. Grossman *et al.*	(MINN, MICH, RUTG)
BARMIN	86	SJNP 44 622	V.V. Barmin *et al.*	(ITEP)
		Translated from YAF 44 965.		
BARMIN	86B	NC 96A 159	V.V. Barmin *et al.*	(ITEP, PADO)
PDG	86B	PL 170B 130	M. Aguilar-Benitez *et al.*	(CERN, CIT+)
BARMIN	85	NC 85A 67	V.V. Barmin *et al.*	(ITEP, PADO)
Also	85B	SJNP 41 759	V.V. Barmin *et al.*	(ITEP)
		Translated from YAF 41 1187.		
BARMIN	83	PL 128B 129	V.V. Barmin *et al.*	(ITEP, PADO)
Also	84	SJNP 39 269	V.V. Barmin *et al.*	(ITEP, PADO)
		Translated from YAF 39 428.		

ARONSON	82	PRL 48 1078	S.H. Aronson *et al.*	(BNL, CHIC, STAN+)
ARONSON	82B	PRL 48 1306	S.H. Aronson *et al.*	(BNL, CHIC, PURD)
Also	82B	PL 116B 73	E. Fischbach *et al.*	(PURD, BNL, CHIC)
Also	83	PR D28 476	S.H. Aronson *et al.*	(BNL, CHIC, PURD)
Also	83B	PR D28 495	S.H. Aronson *et al.*	(BNL, CHIC, PURD)
ARONSON	76	NC 32A 236	S.H. Aronson *et al.*	(WISC, EFI, UCSD+)
EVERHART	76	PR D14 661	G.C. Everhart *et al.*	(PENN)
TAUREG	76	PL 65B 92	H. Taureg *et al.*	(HEIDH, CERN, DORT)
BALDO-...	75	NC 25A 688	M. Baldo-Ceolin *et al.*	(PADO, WISC)
CARITHERS	75	PRL 34 1244	W.C.J. Carithers *et al.*	(COLU, NYU)
BOBISUT	74	LNC 11 646	F. Bobisut *et al.*	(PADO)
COWELL	74	PR D10 2083	P.L. Cowell *et al.*	(STON, COLU)
GEWENIGER	74B	PL 48B 487	C. Geweniger *et al.*	(CERN, HEIDH)
GJESDAL	74B	PL 52B 119	S. Gjesdal *et al.*	(CERN, HEIDH)
BURGUN	73	PL 46B 481	G. Burgun *et al.*	(SACL, CERN)
GJESDAL	73	PL 44B 217	S. Gjesdal *et al.*	(CERN, HEIDH)
HILL	73	PR D8 1290	D.G. Hill *et al.*	(BNL, CMU)
ALITTI	72	PL 39B 568	J. Alitti, E. Lesquoy, A. Muller	(SACL)
BURGUN	72	NP B50 194	G. Burgun *et al.*	(SACL, CERN, OSLO)
METCALF	72	PL 40B 703	M. Metcalf *et al.*	(CERN, IPN, WIEN)
MORSE	72B	PRL 28 388	R. Morse *et al.*	(COLO, PRIN, UMD)
NAGY	72	NP B47 94	E. Nagy, F. Telbisz, G. Vesztergombi	(BUDA)
Also	69	PL 30B 498	G. Bozoki *et al.*	(BUDA)
SKJEGGEST...	72	NP B48 343	O. Skjeggestad *et al.*	(OSLO, CERN, SACL)
BALTAY	71	PRL 27 1678	C. Baltay *et al.*	(COLU)
Also	71	Thesis Nevis 187	W.A. Cooper	(COLU)
MOFFETT	70	BAPS 15 512	R. Moffett *et al.*	(ROCH)
BOZOKI	69	PL 30B 498	G. Bozoki *et al.*	(BUDA)
DOYLE	69	Thesis UCRL 18139	J.C. Doyle	(LRL)
GOBBI	69	PRL 22 682	B. Gobbi *et al.*	(ROCH)
HYAMS	69B	PL 29B 521	B.D. Hyams *et al.*	(CERN, MPIM)
MORFIN	69	PRL 23 660	J.G. Morfin, D. Sinclair	(MICH)
DONALD	68B	PL 27B 58	R.A. Donald *et al.*	(LIVP, CERN, IPNP+)
HILL	68	PR 171 1418	D.G. Hill *et al.*	(BNL, CMU)
AUBERT	65	PL 17 59	B. Aubert *et al.*	(EPOL, ORSAY)
BROWN	63	PR 130 769	J.L. Brown *et al.*	(LRL, MICH)
CHRETIEN	63	PR 131 2208	M. Chretien *et al.*	(BRAN, BROW, HARV+)
BROWN	61	NC 19 1155	J.L. Brown *et al.*	(MICH)
BOLDT	58B	PRL 1 150	E. Boldt, D.O. Caldwell, Y. Pal	(MIT)

OTHER RELATED PAPERS

LITTENBERG	93	ARNPS 43 729	L.S. Littenberg, G. Valencia	(BNL, FNAL)
Rare and Radiative Kaon Decays				
BATTISTON	92	PRPL 214 293	R. Battiston *et al.*	(PGIA, CERN, TRSTT)
Status and Perspectives of K Decay Physics				
TRILLING	65B	UCRL 16473	G.N. Trilling	(LRL)
Updated from 1965 Argonne Conference, page 115.				
CRAWFORD	62	CERN Conf. 827	F.S. Crawford	(LRL)
FITCH	61	NC 22 1160	V.L. Fitch, P.A. Piroue, R.B. Perkins	(PRIN+)
GOOD	61	PR 124 1223	R.H. Good *et al.*	(LRL)
BIRGE	60	Rochester Conf. 601	R.W. Birge *et al.*	(LRL, WISC)
MULLER	60	PRL 4 418	F. Muller *et al.*	(LRL, BNL)