B^0

$I(J^P) = \frac{1}{2}(0^-)$

Quantum numbers not measured. Values shown are quark-model predictions.

See also the $B^{\pm}/B^0$ ADMIXTURE and $B^{\pm}/B^0/B^0_s/b$-baryon ADMIXTURE sections.

See the Note "Production and Decay of b-flavored Hadrons" at the beginning of the $B^{\pm}$ Particle Listings and the Note on "B^0-$\overline{B}^0$ Mixing and CP Violation in B Decay" near the end of the B^0 Particle Listings.

B^0 MASS

The fit uses m_{B^+}, $(m_{B^0} - m_{B^+})$, and m_{B^0} to determine m_{B^+}, m_{B^0}, and the mass difference.

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	COMMENT
5279.4±0.5 OUR FIT					
5279.3±0.7 OUR AVERAGE					
5279.1±0.7 ±0.3	135	[1] CSORNA	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5281.3±2.2 ±1.4	51	ABE	96B	CDF	$p\overline{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5279.2±0.54±2.0	340	ALAM	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5278.0±0.4 ±2.0		BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
5279.6±0.7 ±2.0	40	[2] ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5278.2±1.0 ±3.0	40	ALBRECHT	87C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5279.5±1.6 ±3.0	7	[3] ALBRECHT	87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5280.6±0.8 ±2.0		BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[1] CSORNA 00 uses fully reconstructed 135 $B^0 \rightarrow J/\psi^{(\prime)} K^0_S$ events and invariant masses without beam constraint.

[2] ALBRECHT 90J assumes 10580 for $\Upsilon(4S)$ mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.

[3] Found using fully reconstructed decays with J/ψ. ALBRECHT 87D assume $m_{\Upsilon(4S)}$ = 10577 MeV.

$m_{B^0} - m_{B^+}$

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
0.33±0.28 OUR FIT Error includes scale factor of 1.1.				
0.34±0.32 OUR AVERAGE Error includes scale factor of 1.2.				
0.41±0.25±0.19	ALAM	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
−0.4 ±0.6 ±0.5	BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
−0.9 ±1.2 ±0.5	ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
2.0 ±1.1 ±0.3	[4] BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[4] BEBEK 87 actually measure the difference between half of E_{cm} and the $B^{\pm}$ or B^0 mass, so the $m_{B^0} - m_{B^{\pm}}$ is more accurate. Assume $m_{\Upsilon(4S)}$ = 10580 MeV.

$m_{B^0_H} - m_{B^0_L}$

See the B^0-$\overline{B}^0$ MIXING PARAMETERS section near the end of these B^0 Listings.

B^0 MEAN LIFE

See $B^{\pm}/B^0/B^0_s/b$-baryon ADMIXTURE section for data on B-hadron mean life averaged over species of bottom particles.

"OUR EVALUATION" is an average of the data listed below performed by the LEP B Lifetimes Working Group as described in our review "Production and Decay of b-flavored Hadrons" in the $B^{\pm}$ Section of the Listings. The averaging procedure takes into account correlations between the measurements and asymmetric lifetime errors.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID		TECN	COMMENT
1.542±0.016 OUR EVALUATION					
1.554±0.030±0.019		[5] ABE	02H	BELL	$e^+e^- \to \Upsilon(4S)$
1.529±0.012±0.029		[6] AUBERT	02H	BABR	$e^+e^- \to \Upsilon(4S)$
1.546±0.032±0.022		[5] AUBERT	01F	BABR	$e^+e^- \to \Upsilon(4S)$
1.541±0.028±0.023		[6] ABBIENDI,G	00B	OPAL	$e^+e^- \to Z$
1.518±0.053±0.034		[7] BARATE	00R	ALEP	$e^+e^- \to Z$
1.523±0.057±0.053		[8] ABBIENDI	99J	OPAL	$e^+e^- \to Z$
1.58 ±0.09 ±0.02		[9] ABE	98B	CDF	$p\overline{p}$ at 1.8 TeV
$1.474\pm0.039^{+0.052}_{-0.051}$		[7] ABE	98Q	CDF	$p\overline{p}$ at 1.8 TeV
1.52 ±0.06 ±0.04		[8] ACCIARRI	98S	L3	$e^+e^- \to Z$
1.64 ±0.08 ±0.08		[8] ABE	97J	SLD	$e^+e^- \to Z$
1.532±0.041±0.040		[10] ABREU	97F	DLPH	$e^+e^- \to Z$
$1.25^{+0.15}_{-0.13}\pm0.05$	121	[9] BUSKULIC	96J	ALEP	$e^+e^- \to Z$
$1.49^{+0.17}_{-0.15}{}^{+0.08}_{-0.06}$		[11] BUSKULIC	96J	ALEP	$e^+e^- \to Z$
$1.61^{+0.14}_{-0.13}\pm0.08$		[7,12] ABREU	95Q	DLPH	$e^+e^- \to Z$
1.63 ±0.14 ±0.13		[13] ADAM	95	DLPH	$e^+e^- \to Z$
1.53 ±0.12 ±0.08		[7,14] AKERS	95T	OPAL	$e^+e^- \to Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.54 ±0.08 ±0.06		[7] ABE	96C	CDF	Repl. by ABE 98Q
1.55 ±0.06 ±0.03		[15] BUSKULIC	96J	ALEP	$e^+e^- \to Z$
1.61 ±0.07 ±0.04		[7] BUSKULIC	96J	ALEP	Repl. by BARATE 00R
1.62 ±0.12		[16] ADAM	95	DLPH	$e^+e^- \to Z$
1.57 ±0.18 ±0.08	121	[9] ABE	94D	CDF	Repl. by ABE 98B
$1.17^{+0.29}_{-0.23}\pm0.16$	96	[7] ABREU	93D	DLPH	Sup. by ABREU 95Q
1.55 ±0.25 ±0.18	76	[13] ABREU	93G	DLPH	Sup. by ADAM 95
$1.51^{+0.24}_{-0.23}{}^{+0.12}_{-0.14}$	78	[7] ACTON	93C	OPAL	Sup. by AKERS 95T
$1.52^{+0.20}_{-0.18}{}^{+0.07}_{-0.13}$	77	[7] BUSKULIC	93D	ALEP	Sup. by BUSKULIC 96J
$1.20^{+0.52}_{-0.36}{}^{+0.16}_{-0.14}$	15	[17] WAGNER	90	MRK2	E^{ee}_{cm}= 29 GeV
$0.82^{+0.57}_{-0.37}\pm0.27$		[18] AVERILL	89	HRS	E^{ee}_{cm}= 29 GeV

[5] Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.
[6] Data analyzed using partially reconsturcted $\overline{B}^0 \rightarrow D^{*+} \ell^- \overline{\nu}$ decays.
[7] Data analyzed using $D/D^* \ell X$ event vertices.
[8] Data analyzed using charge of secondary vertex.
[9] Measured mean life using fully reconstructed decays.
[10] Data analyzed using inclusive $D/D^* \ell X$.
[11] Measured mean life using partially reconstructed $D^{*-} \pi^+ X$ vertices.
[12] ABREU 95Q assumes B($B^0 \rightarrow D^{**-} \ell^+ \nu_\ell$) = 3.2 ± 1.7%.
[13] Data analyzed using vertex-charge technique to tag B charge.
[14] AKERS 95T assumes B($B^0 \rightarrow D_s{}^{(*)} D^{0\,(*)}$) = 5.0 ± 0.9% to find B^+/B^0 yield.
[15] Combined result of $D/D^* \ell x$ analysis, fully reconstructed B analysis, and partially reconstructed $D^{*-} \pi^+ X$ analysis.
[16] Combined ABREU 95Q and ADAM 95 result.
[17] WAGNER 90 tagged B^0 mesons by their decays into $D^{*-} e^+ \nu$ and $D^{*-} \mu^+ \nu$ where the D^{*-} is tagged by its decay into $\pi^- \overline{D}^0$.
[18] AVERILL 89 is an estimate of the B^0 mean lifetime assuming that $B^0 \rightarrow D^{*+} + X$ always.

MEAN LIFE RATIO τ_{B+}/τ_{B^0}

τ_{B+}/τ_{B^0} (direct measurements)

"OUR EVALUATION" is an average of the data listed below performed by the LEP B Lifetimes Working Group as described in our review "Production and Decay of b-flavored Hadrons" in the $B^\pm$ Section of the Listings. The averaging procedure takes into account correlations between the measurements and asymmetric lifetime errors.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.					
1.083±0.017 OUR EVALUATION					
1.091±0.023±0.014		[19] ABE	02H	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
1.082±0.026±0.012		[19] AUBERT	01F	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
1.085±0.059±0.018		[20] BARATE	00R	ALEP	$e^+ e^- \rightarrow Z$
1.079±0.064±0.041		[21] ABBIENDI	99J	OPAL	$e^+ e^- \rightarrow Z$
1.06 ±0.07 ±0.02		[22] ABE	98B	CDF	$p\overline{p}$ at 1.8 TeV
$1.110 \pm 0.056 {}^{+0.033}_{-0.030}$		[20] ABE	98Q	CDF	$p\overline{p}$ at 1.8 TeV
1.09 ±0.07 ±0.03		[21] ACCIARRI	98S	L3	$e^+ e^- \rightarrow Z$
1.01 ±0.07 ±0.06		[21] ABE	97J	SLD	$e^+ e^- \rightarrow Z$
$1.27 {}^{+0.23}_{-0.19} {}^{+0.03}_{-0.02}$		[22] BUSKULIC	96J	ALEP	$e^+ e^- \rightarrow Z$
$1.00 {}^{+0.17}_{-0.15} \pm 0.10$		[20,23] ABREU	95Q	DLPH	$e^+ e^- \rightarrow Z$
$1.06 {}^{+0.13}_{-0.10} \pm 0.10$		[24] ADAM	95	DLPH	$e^+ e^- \rightarrow Z$
$0.99 \pm 0.14 {}^{+0.05}_{-0.04}$		[20,25] AKERS	95T	OPAL	$e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.01 \pm 0.11 \pm 0.02$		[20] ABE 96C	CDF	Repl. by ABE 98Q
$1.03 \pm 0.08 \pm 0.02$		[26] BUSKULIC 96J	ALEP	$e^+ e^- \rightarrow Z$
$0.98 \pm 0.08 \pm 0.03$		[20] BUSKULIC 96J	ALEP	Repl. by BARATE 00R
$1.02 \pm 0.16 \pm 0.05$	269	[22] ABE 94D	CDF	Repl. by ABE 98B
$1.11 {}^{+0.51}_{-0.39} \pm 0.11$	188	[20] ABREU 93D	DLPH	Sup. by ABREU 95Q
$1.01 {}^{+0.29}_{-0.22} \pm 0.12$	253	[24] ABREU 93G	DLPH	Sup. by ADAM 95
$1.0 {}^{+0.33}_{-0.25} \pm 0.08$	130	ACTON 93C	OPAL	Sup. by AKERS 95T
$0.96 {}^{+0.19}_{-0.15} {}^{+0.18}_{-0.12}$	154	[20] BUSKULIC 93D	ALEP	Sup. by BUSKULIC 96J

[19] Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.

[20] Data analyzed using $D/D^* \ell X$ vertices.

[21] Data analyzed using charge of secondary vertex.

[22] Measured using fully reconstructed decays.

[23] ABREU 95Q assumes $B(B^0 \rightarrow D^{**-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$.

[24] Data analyzed using vertex-charge technique to tag B charge.

[25] AKERS 95T assumes $B(B^0 \rightarrow D_s{}^{(*)} D^{0\,(*)}) = 5.0 \pm 0.9\%$ to find B^+/B^0 yield.

[26] Combined result of $D/D^* \ell X$ analysis and fully reconstructed B analysis.

τ_{B^+}/τ_{B^0} (inferred from branching fractions)

These measurements are inferred from the branching fractions for semileptonic decay or other spectator-dominated decays by assuming that the rates for such decays are equal for B^0 and B^+. We do not use measurements which assume equal production of B^0 and B^+ because of the large uncertainty in the production ratio.

VALUE	*CL%*	*EVTS*	*DOCUMENT ID*	*TECN*	*COMMENT*

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.95 {}^{+0.117}_{-0.080} \pm 0.091$			[27] ARTUSO 97	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.15 \pm 0.17 \pm 0.06$			[28] JESSOP 97	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.93 \pm 0.18 \pm 0.12$			[29] ATHANAS 94	CLE2	Sup. by ARTUSO 97
$0.91 \pm 0.27 \pm 0.21$			[30] ALBRECHT 92C	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
1.0 ± 0.4		29	[30,31] ALBRECHT 92G	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.89 \pm 0.19 \pm 0.13$			[30] FULTON 91	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.00 \pm 0.23 \pm 0.14$			[30] ALBRECHT 89L	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.49 to 2.3	90		[32] BEAN 87B	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

[27] ARTUSO 97 uses partial reconstruction of $B \rightarrow D^* \ell \nu_\ell$ and independent of B^0 and B^+ production fraction.

[28] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[29] ATHANAS 94 uses events tagged by fully reconstructed B^- decays and partially or fully reconstructed B^0 decays.

[30] Assumes equal production of B^0 and B^+.

[31] ALBRECHT 92G data analyzed using $B \rightarrow D_s \overline{D}$, $D_s \overline{D}^*$, $D_s^* \overline{D}$, $D_s^* \overline{D}^*$ events.

[32] BEAN 87B assume the fraction of $B^0 \overline{B}^0$ events at the $\Upsilon(4S)$ is 0.41.

$\left|\Delta\Gamma_{B^0_d}\right|/\Gamma_{B^0_d}$

$\Gamma_{B^0_d}$ and $\left|\Delta\Gamma_{B^0_d}\right|$ are the decay rate average and difference between two B^0_d *CP* eigenstates.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.80	95	[33,34] BEHRENS	00B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[33] BEHRENS 00B uses high-momentum lepton tags and partially reconstructed $\overline{B}^0 \rightarrow D^{*+}\pi^-$, ρ^- decays to determine the flavor of the B meson.
[34] Assumes Δ_{md}=0.478 ± 0.018 ps^{-1} and τ_{B^0}=1.548 ± 0.032 ps.

B^0 DECAY MODES

$\overline{B}^0$ modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing. Modes which do not identify the charge state of the B are listed in the $B^{\pm}/B^0$ ADMIXTURE section.

The branching fractions listed below assume 50% $B^0\overline{B}^0$ and 50% B^+B^- production at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D, D_s, D^*, and ψ branching ratios to current values whenever this would affect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

	Mode		Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1	$\ell^+\nu_\ell$ anything	[a]	(10.5 ±0.8) %	
Γ_2	$D^-\ell^+\nu_\ell$	[a]	(2.11±0.17) %	
Γ_3	$D^*(2010)^-\ell^+\nu_\ell$	[a]	(4.60±0.21) %	
Γ_4	$\rho^-\ell^+\nu_\ell$	[a]	($2.6^{+0.6}_{-0.7}$) $\times 10^{-4}$	
Γ_5	$\pi^-\ell^+\nu_\ell$		(1.8 ±0.6) $\times 10^{-4}$	
	Inclusive modes			
Γ_6	$\pi^-\mu^+\nu_\mu$			
Γ_7	K^+ anything		(78 ±8) %	

D, D^*, or D_s modes

Γ_{8}	$D^- \pi^+$	$(3.0 \pm 0.4) \times 10^{-3}$	
Γ_{9}	$D^- \rho^+$	$(7.8 \pm 1.4) \times 10^{-3}$	
Γ_{10}	$D^- K^*(892)^+$	$(3.7 \pm 1.8) \times 10^{-4}$	
Γ_{11}	$D^- \omega \pi^+$	$(2.8 \pm 0.6) \times 10^{-3}$	
Γ_{12}	$D^- K^+$	$(2.0 \pm 0.6) \times 10^{-4}$	
Γ_{13}	$\overline{D}^0 \pi^+ \pi^-$	$< 1.6 \times 10^{-3}$	CL=90%
Γ_{14}	$D^*(2010)^- \pi^+$	$(2.76 \pm 0.21) \times 10^{-3}$	
Γ_{15}	$D^- \pi^+ \pi^+ \pi^-$	$(8.0 \pm 2.5) \times 10^{-3}$	
Γ_{16}	$(D^- \pi^+ \pi^+ \pi^-)$ nonresonant	$(3.9 \pm 1.9) \times 10^{-3}$	
Γ_{17}	$D^- \pi^+ \rho^0$	$(1.1 \pm 1.0) \times 10^{-3}$	
Γ_{18}	$D^- a_1(1260)^+$	$(6.0 \pm 3.3) \times 10^{-3}$	
Γ_{19}	$D^*(2010)^- \pi^+ \pi^0$	(1.5 ± 0.5) %	
Γ_{20}	$D^*(2010)^- \rho^+$	$(7.3 \pm 1.5) \times 10^{-3}$	
Γ_{21}	$D^*(2010)^- K^+$	$(2.0 \pm 0.5) \times 10^{-4}$	
Γ_{22}	$D^*(2010)^- K^*(892)^+$	$(3.8 \pm 1.5) \times 10^{-4}$	
Γ_{23}	$D^*(2010)^- \pi^+ \pi^+ \pi^-$	$(7.6 \pm 1.8) \times 10^{-3}$	S=1.4
Γ_{24}	$(D^*(2010)^- \pi^+ \pi^+ \pi^-)$ non-resonant	$(0.0 \pm 2.5) \times 10^{-3}$	
Γ_{25}	$D^*(2010)^- \pi^+ \rho^0$	$(5.7 \pm 3.2) \times 10^{-3}$	
Γ_{26}	$D^*(2010)^- a_1(1260)^+$	(1.30 ± 0.27) %	
Γ_{27}	$D^*(2010)^- \pi^+ \pi^+ \pi^- \pi^0$	(1.76 ± 0.27) %	
Γ_{28}	$D^*(2010)^+ \pi^+ \pi^- \pi^- \pi^0$	(1.8 ± 0.7) %	
Γ_{29}	$D^*(2010)^- p \overline{p} \pi^+$	$(6.5 \pm 1.6) \times 10^{-4}$	
Γ_{30}	$D^*(2010)^- p \overline{n}$	$(1.5 \pm 0.4) \times 10^{-3}$	
Γ_{31}	$\overline{D}^*(2010)^- \omega \pi^+$	$(2.9 \pm 0.5) \times 10^{-3}$	
Γ_{32}	$\overline{D}_2^*(2460)^- \pi^+$	$< 2.2 \times 10^{-3}$	CL=90%
Γ_{33}	$\overline{D}_2^*(2460)^- \rho^+$	$< 4.9 \times 10^{-3}$	CL=90%
Γ_{34}	$D^- D^+$	$< 9.4 \times 10^{-4}$	CL=90%
Γ_{35}	$D^- D_s^+$	$(8.0 \pm 3.0) \times 10^{-3}$	
Γ_{36}	$D^*(2010)^- D_s^+$	(1.11 ± 0.33) %	
Γ_{37}	$D^- D_s^{*+}$	(1.0 ± 0.5) %	
Γ_{38}	$D^*(2010)^- D_s^{*+}$	(1.8 ± 0.6) %	
Γ_{39}	$D_s^+ \pi^-$	$< 2.8 \times 10^{-4}$	CL=90%
Γ_{40}	$D_s^{*+} \pi^-$	$< 5 \times 10^{-4}$	CL=90%
Γ_{41}	$D_s^+ \rho^-$	$< 7 \times 10^{-4}$	CL=90%
Γ_{42}	$D_s^{*+} \rho^-$	$< 8 \times 10^{-4}$	CL=90%
Γ_{43}	$D_s^+ a_1(1260)^-$	$< 2.6 \times 10^{-3}$	CL=90%
Γ_{44}	$D_s^{*+} a_1(1260)^-$	$< 2.2 \times 10^{-3}$	CL=90%
Γ_{45}	$D_s^- K^+$	$< 2.4 \times 10^{-4}$	CL=90%
Γ_{46}	$D_s^{*-} K^+$	$< 1.7 \times 10^{-4}$	CL=90%
Γ_{47}	$D_s^- K^*(892)^+$	$< 9.9 \times 10^{-4}$	CL=90%

Γ_{48}	$D_s^{*-} K^*(892)^+$	< 1.1	$\times 10^{-3}$	CL=90%
Γ_{49}	$D_s^- \pi^+ K^0$	< 5	$\times 10^{-3}$	CL=90%
Γ_{50}	$D_s^{*-} \pi^+ K^0$	< 3.1	$\times 10^{-3}$	CL=90%
Γ_{51}	$D_s^- \pi^+ K^*(892)^0$	< 4	$\times 10^{-3}$	CL=90%
Γ_{52}	$D_s^{*-} \pi^+ K^*(892)^0$	< 2.0	$\times 10^{-3}$	CL=90%
Γ_{53}	$\overline{D}^0 \pi^0$	(2.9 ± 0.5)	$\times 10^{-4}$	
Γ_{54}	$\overline{D}^0 \rho^0$	< 3.9	$\times 10^{-4}$	CL=90%
Γ_{55}	$\overline{D}^0 \eta$	$(1.4 {}^{+0.6}_{-0.5})$	$\times 10^{-4}$	
Γ_{56}	$\overline{D}^0 \eta'$	< 9.4	$\times 10^{-4}$	CL=90%
Γ_{57}	$\overline{D}^0 \omega$	(1.8 ± 0.6)	$\times 10^{-4}$	
Γ_{58}	$\overline{D}^{*0} \gamma$	< 5.0	$\times 10^{-5}$	CL=90%
Γ_{59}	$\overline{D}^*(2007)^0 \pi^0$	(2.5 ± 0.7)	$\times 10^{-4}$	
Γ_{60}	$\overline{D}^*(2007)^0 \rho^0$	< 5.6	$\times 10^{-4}$	CL=90%
Γ_{61}	$\overline{D}^*(2007)^0 \eta$	< 2.6	$\times 10^{-4}$	CL=90%
Γ_{62}	$\overline{D}^*(2007)^0 \eta'$	< 1.4	$\times 10^{-3}$	CL=90%
Γ_{63}	$\overline{D}^*(2007)^0 \omega$	< 7.4	$\times 10^{-4}$	CL=90%
Γ_{64}	$D^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-$	(3.0 ± 0.9)	$\times 10^{-3}$	
Γ_{65}	$D^*(2010)^+ D^*(2010)^-$	$(9.9 {}^{+4.4}_{-3.5})$	$\times 10^{-4}$	
Γ_{66}	$D^*(2010)^+ D^-$	< 6.3	$\times 10^{-4}$	CL=90%
Γ_{67}	$D^{(*)0} \overline{D}^{(*)0}$	< 2.7	%	CL=90%

Charmonium modes

Γ_{68}	$\eta_c K^0$	$(1.1 {}^{+0.6}_{-0.5})$	$\times 10^{-3}$	
Γ_{69}	$J/\psi(1S) K^0$	(8.7 ± 0.5)	$\times 10^{-4}$	
Γ_{70}	$J/\psi(1S) K^+ \pi^-$	(1.2 ± 0.6)	$\times 10^{-3}$	
Γ_{71}	$\quad J/\psi(1S) K^*(892)^0$	(1.31 ± 0.09)	$\times 10^{-3}$	
Γ_{72}	$J/\psi(1S) \phi K^0$	$(8.8 {}^{+3.7}_{-3.3})$	$\times 10^{-5}$	
Γ_{73}	$J/\psi(1S) K(1270)^0$	(1.3 ± 0.5)	$\times 10^{-3}$	
Γ_{74}	$J/\psi(1S) \pi^0$	(2.1 ± 0.5)	$\times 10^{-5}$	
Γ_{75}	$J/\psi(1S) \eta$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{76}	$J/\psi(1S) \rho^0$	< 2.5	$\times 10^{-4}$	CL=90%
Γ_{77}	$J/\psi(1S) \omega$	< 2.7	$\times 10^{-4}$	CL=90%
Γ_{78}	$J/\psi(1S) K^0 \pi^+ \pi^-$	(1.0 ± 0.4)	$\times 10^{-3}$	
Γ_{79}	$\quad J/\psi(1S) K^0 \rho^0$	(5.4 ± 3.0)	$\times 10^{-4}$	
Γ_{80}	$J/\psi(1S) K^*(892)^+ \pi^-$	(8 ± 4)	$\times 10^{-4}$	
Γ_{81}	$J/\psi(1S) K^*(892)^0 \pi^+ \pi^-$	(6.6 ± 2.2)	$\times 10^{-4}$	
Γ_{82}	$\psi(2S) K^0$	(5.7 ± 1.0)	$\times 10^{-4}$	
Γ_{83}	$\psi(2S) K^+ \pi^-$	< 1	$\times 10^{-3}$	CL=90%
Γ_{84}	$\quad \psi(2S) K^*(892)^0$	(8.0 ± 1.3)	$\times 10^{-4}$	

Γ_{85}	$\chi_{c0}(1P)\,K^0$		$< 5.0 \times 10^{-4}$	CL=90%
Γ_{86}	$\chi_{c1}(1P)\,K^0$		$(4.0\ ^{+1.2}_{-1.0}) \times 10^{-4}$	
Γ_{87}	$\chi_{c1}(1P)\,K^*(892)^0$		$(4.1 \pm 1.5) \times 10^{-4}$	
	K or K^* modes			
Γ_{88}	$K^+\pi^-$		$(1.74\pm0.15) \times 10^{-5}$	
Γ_{89}	$K^0\pi^0$		$(1.07\ ^{+0.27}_{-0.25}) \times 10^{-5}$	
Γ_{90}	$\eta' K^0$		$(5.8\ ^{+1.4}_{-1.3}) \times 10^{-5}$	S=1.5
Γ_{91}	$\eta' K^*(892)^0$		$< 2.4 \times 10^{-5}$	CL=90%
Γ_{92}	$\eta\,K^*(892)^0$		$(1.4\ ^{+0.6}_{-0.5}) \times 10^{-5}$	
Γ_{93}	$\eta\,K^0$		$< 9.3 \times 10^{-6}$	CL=90%
Γ_{94}	$\omega\,K^0$		$< 1.3 \times 10^{-5}$	CL=90%
Γ_{95}	$K^0_S X^0$(Familon)		$< 5.3 \times 10^{-5}$	CL=90%
Γ_{96}	$\omega\,K^*(892)^0$		$< 2.3 \times 10^{-5}$	CL=90%
Γ_{97}	K^+K^-		$< 1.9 \times 10^{-6}$	CL=90%
Γ_{98}	$K^0\overline{K}^0$		$< 1.7 \times 10^{-5}$	CL=90%
Γ_{99}	$K^+\rho^-$		$< 3.2 \times 10^{-5}$	CL=90%
Γ_{100}	$K^0\pi^+\pi^-$			
Γ_{101}	$K^0\rho^0$		$< 3.9 \times 10^{-5}$	CL=90%
Γ_{102}	$K^0 f_0(980)$		$< 3.6 \times 10^{-4}$	CL=90%
Γ_{103}	$K^*(892)^+\pi^-$		$< 7.2 \times 10^{-5}$	CL=90%
Γ_{104}	$K^*(892)^0\pi^0$		$< 3.6 \times 10^{-6}$	CL=90%
Γ_{105}	$K^*_2(1430)^+\pi^-$		$< 2.6 \times 10^{-3}$	CL=90%
Γ_{106}	$K^0K^+K^-$		$< 1.3 \times 10^{-3}$	CL=90%
Γ_{107}	$K^0\phi$		$(8.1\ ^{+3.2}_{-2.6}) \times 10^{-6}$	
Γ_{108}	$K^-\pi^+\pi^+\pi^-$	[b]	$< 2.3 \times 10^{-4}$	CL=90%
Γ_{109}	$K^*(892)^0\pi^+\pi^-$		$< 1.4 \times 10^{-3}$	CL=90%
Γ_{110}	$K^*(892)^0\rho^0$		$< 3.4 \times 10^{-5}$	CL=90%
Γ_{111}	$K^*(892)^0 f_0(980)$		$< 1.7 \times 10^{-4}$	CL=90%
Γ_{112}	$K_1(1400)^+\pi^-$		$< 1.1 \times 10^{-3}$	CL=90%
Γ_{113}	$K^- a_1(1260)^+$	[b]	$< 2.3 \times 10^{-4}$	CL=90%
Γ_{114}	$K^*(892)^0K^+K^-$		$< 6.1 \times 10^{-4}$	CL=90%
Γ_{115}	$K^*(892)^0\phi$		$(9.5\ ^{+2.4}_{-2.0}) \times 10^{-6}$	
Γ_{116}	$\overline{K}^*(892)^0 K^*(892)^0$		$< 2.2 \times 10^{-5}$	CL=90%
Γ_{117}	$K^*(892)^0 K^*(892)^0$		$< 3.7 \times 10^{-5}$	CL=90%
Γ_{118}	$K^*(892)^+ K^*(892)^-$		$< 1.41 \times 10^{-4}$	CL=90%
Γ_{119}	$K_1(1400)^0\rho^0$		$< 3.0 \times 10^{-3}$	CL=90%
Γ_{120}	$K_1(1400)^0\phi$		$< 5.0 \times 10^{-3}$	CL=90%
Γ_{121}	$K^*_2(1430)^0\rho^0$		$< 1.1 \times 10^{-3}$	CL=90%
Γ_{122}	$K^*_2(1430)^0\phi$		$< 1.4 \times 10^{-3}$	CL=90%

Γ_{123}	$K^*(892)^0\gamma$		$(4.3 \pm 0.4) \times 10^{-5}$	
Γ_{124}	$K_1(1270)^0\gamma$		$< 7.0 \times 10^{-3}$	CL=90%
Γ_{125}	$K_1(1400)^0\gamma$		$< 4.3 \times 10^{-3}$	CL=90%
Γ_{126}	$K_2^*(1430)^0\gamma$		$< 4.0 \times 10^{-4}$	CL=90%
Γ_{127}	$K^*(1680)^0\gamma$		$< 2.0 \times 10^{-3}$	CL=90%
Γ_{128}	$K_3^*(1780)^0\gamma$		< 1.0 %	CL=90%
Γ_{129}	$K_4^*(2045)^0\gamma$		$< 4.3 \times 10^{-3}$	CL=90%

Light unflavored meson modes

Γ_{130}	$\rho^0\gamma$		$< 1.7 \times 10^{-5}$	CL=90%
Γ_{131}	$\omega\gamma$		$< 9.2 \times 10^{-6}$	CL=90%
Γ_{132}	$\phi\gamma$		$< 3.3 \times 10^{-6}$	CL=90%
Γ_{133}	$\pi^+\pi^-$		$(4.4 \pm 0.9) \times 10^{-6}$	
Γ_{134}	$\pi^0\pi^0$		$< 5.7 \times 10^{-6}$	CL=90%
Γ_{135}	$\eta\pi^0$		$< 2.9 \times 10^{-6}$	CL=90%
Γ_{136}	$\eta\eta$		$< 1.8 \times 10^{-5}$	CL=90%
Γ_{137}	$\eta'\pi^0$		$< 5.7 \times 10^{-6}$	CL=90%
Γ_{138}	$\eta'\eta'$		$< 4.7 \times 10^{-5}$	CL=90%
Γ_{139}	$\eta'\eta$		$< 2.7 \times 10^{-5}$	CL=90%
Γ_{140}	$\eta'\rho^0$		$< 1.2 \times 10^{-5}$	CL=90%
Γ_{141}	$\eta\rho^0$		$< 1.0 \times 10^{-5}$	CL=90%
Γ_{142}	$\omega\eta$		$< 1.2 \times 10^{-5}$	CL=90%
Γ_{143}	$\omega\eta'$		$< 6.0 \times 10^{-5}$	CL=90%
Γ_{144}	$\omega\rho^0$		$< 1.1 \times 10^{-5}$	CL=90%
Γ_{145}	$\omega\omega$		$< 1.9 \times 10^{-5}$	CL=90%
Γ_{146}	$\phi\pi^0$		$< 5 \times 10^{-6}$	CL=90%
Γ_{147}	$\phi\eta$		$< 9 \times 10^{-6}$	CL=90%
Γ_{148}	$\phi\eta'$		$< 3.1 \times 10^{-5}$	CL=90%
Γ_{149}	$\phi\rho^0$		$< 1.3 \times 10^{-5}$	CL=90%
Γ_{150}	$\phi\omega$		$< 2.1 \times 10^{-5}$	CL=90%
Γ_{151}	$\phi\phi$		$< 1.2 \times 10^{-5}$	CL=90%
Γ_{152}	$\pi^+\pi^-\pi^0$		$< 7.2 \times 10^{-4}$	CL=90%
Γ_{153}	$\rho^0\pi^0$		$< 5.5 \times 10^{-6}$	CL=90%
Γ_{154}	$\rho^\mp\pi^\pm$	[c]	$(2.8 \pm 0.9) \times 10^{-5}$	
Γ_{155}	$\pi^+\pi^-\pi^+\pi^-$		$< 2.3 \times 10^{-4}$	CL=90%
Γ_{156}	$\rho^0\rho^0$		$< 1.8 \times 10^{-5}$	CL=90%
Γ_{157}	$a_1(1260)^\mp\pi^\pm$	[c]	$< 4.9 \times 10^{-4}$	CL=90%
Γ_{158}	$a_2(1320)^\mp\pi^\pm$	[c]	$< 3.0 \times 10^{-4}$	CL=90%
Γ_{159}	$\pi^+\pi^-\pi^0\pi^0$		$< 3.1 \times 10^{-3}$	CL=90%
Γ_{160}	$\rho^+\rho^-$		$< 2.2 \times 10^{-3}$	CL=90%
Γ_{161}	$a_1(1260)^0\pi^0$		$< 1.1 \times 10^{-3}$	CL=90%
Γ_{162}	$\omega\pi^0$		$< 3 \times 10^{-6}$	CL=90%
Γ_{163}	$\pi^+\pi^+\pi^-\pi^-\pi^0$		$< 9.0 \times 10^{-3}$	CL=90%
Γ_{164}	$a_1(1260)^+\rho^-$		$< 3.4 \times 10^{-3}$	CL=90%

Γ_{165}	$a_1(1260)^0 \rho^0$			$<$ 2.4	$\times 10^{-3}$	CL=90%
Γ_{166}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-$			$<$ 3.0	$\times 10^{-3}$	CL=90%
Γ_{167}	$a_1(1260)^+ a_1(1260)^-$			$<$ 2.8	$\times 10^{-3}$	CL=90%
Γ_{168}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^- \pi^0$			$<$ 1.1	%	CL=90%

Baryon modes

Γ_{169}	$p \overline{p}$			$<$ 7.0	$\times 10^{-6}$	CL=90%
Γ_{170}	$p \overline{p} \pi^+ \pi^-$			$<$ 2.5	$\times 10^{-4}$	CL=90%
Γ_{171}	$p \overline{\Lambda} \pi^-$			$<$ 1.3	$\times 10^{-5}$	CL=90%
Γ_{172}	$\overline{\Lambda} \Lambda$			$<$ 3.9	$\times 10^{-6}$	CL=90%
Γ_{173}	$\Delta^0 \overline{\Delta}^0$			$<$ 1.5	$\times 10^{-3}$	CL=90%
Γ_{174}	$\Delta^{++} \Delta^{--}$			$<$ 1.1	$\times 10^{-4}$	CL=90%
Γ_{175}	$\overline{\Sigma}_c^{--} \Delta^{++}$			$<$ 1.0	$\times 10^{-3}$	CL=90%
Γ_{176}	$\overline{\Lambda}_c^- p \pi^+ \pi^-$			(1.3 $\pm$0.6)	$\times 10^{-3}$	
Γ_{177}	$\overline{\Lambda}_c^- p$			$<$ 2.1	$\times 10^{-4}$	CL=90%
Γ_{178}	$\overline{\Lambda}_c^- p \pi^0$			$<$ 5.9	$\times 10^{-4}$	CL=90%
Γ_{179}	$\overline{\Lambda}_c^- p \pi^+ \pi^- \pi^0$			$<$ 5.07	$\times 10^{-3}$	CL=90%
Γ_{180}	$\overline{\Lambda}_c^- p \pi^+ \pi^- \pi^+ \pi^-$			$<$ 2.74	$\times 10^{-3}$	CL=90%

Lepton Family number (*LF*) violating modes, or
$\Delta B = 1$ weak neutral current (*B1*) modes

Γ_{181}	$\gamma \gamma$			$<$ 1.7	$\times 10^{-6}$	CL=90%
Γ_{182}	$e^+ e^-$	*B1*		$<$ 8.3	$\times 10^{-7}$	CL=90%
Γ_{183}	$\mu^+ \mu^-$	*B1*		$<$ 6.1	$\times 10^{-7}$	CL=90%
Γ_{184}	$K^0 e^+ e^-$	*B1*		$<$ 2.7	$\times 10^{-6}$	CL=90%
Γ_{185}	$K^0 \mu^+ \mu^-$	*B1*		$<$ 3.3	$\times 10^{-6}$	CL=90%
Γ_{186}	$K^*(892)^0 e^+ e^-$	*B1*		$<$ 6.4	$\times 10^{-6}$	CL=90%
Γ_{187}	$K^*(892)^0 \mu^+ \mu^-$	*B1*		$<$ 4.2	$\times 10^{-6}$	CL=90%
Γ_{188}	$K^*(892)^0 \nu \overline{\nu}$	*B1*		$<$ 1.0	$\times 10^{-3}$	CL=90%
Γ_{189}	$e^{\pm} \mu^{\mp}$	*LF*	[c]	$<$ 1.5	$\times 10^{-6}$	CL=90%
Γ_{190}	$e^{\pm} \tau^{\mp}$	*LF*	[c]	$<$ 5.3	$\times 10^{-4}$	CL=90%
Γ_{191}	$\mu^{\pm} \tau^{\mp}$	*LF*	[c]	$<$ 8.3	$\times 10^{-4}$	CL=90%

[a] An ℓ indicates an e or a μ mode, not a sum over these modes.

[b] B^0 and B_s^0 contributions not separated. Limit is on weighted average of the two decay rates.

[c] The value is for the sum of the charge states or particle/antiparticle states indicated.

B^0 BRANCHING RATIOS

For branching ratios in which the charge of the decaying B is not determined, see the $B^{\pm}$ section.

$\Gamma(\ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.105 ±0.008 OUR AVERAGE				
0.1078±0.0060±0.0069	[35] ARTUSO	97	CLE2	$e^+e^- \to \Upsilon(4S)$
0.093 ±0.011 ±0.015	ALBRECHT	94	ARG	$e^+e^- \to \Upsilon(4S)$
0.099 ±0.030 ±0.009	HENDERSON	92	CLEO	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.109 ±0.007 ±0.011	ATHANAS	94	CLE2	Sup. by ARTUSO 97

[35] ARTUSO 97 uses partial reconstruction of $B \to D^* \ell \nu_\ell$ and inclusive semileptonic branching ratio from BARISH 96B ($0.1049 \pm 0.0017 \pm 0.0043$).

$\Gamma(D^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_2/Γ

ℓ denotes e or μ, not the sum.

VALUE	DOCUMENT ID		TECN	COMMENT
0.0211±0.0017 OUR AVERAGE				
0.0213±0.0012±0.0039	ABE	02E	BELL	$e^+e^- \to \Upsilon(4S)$
0.0209±0.0013±0.0018	[36] BARTELT	99	CLE2	$e^+e^- \to \Upsilon(4S)$
0.0235±0.0020±0.0044	[37] BUSKULIC	97	ALEP	$e^+e^- \to Z$
0.018 ±0.006 ±0.003	[38] FULTON	91	CLEO	$e^+e^- \to \Upsilon(4S)$
0.020 ±0.007 ±0.006	[39] ALBRECHT	89J	ARG	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0187±0.0015±0.0032	[40] ATHANAS	97	CLE2	Repl. by BARTELT 99

[36] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[37] BUSKULIC 97 assumes fraction (B^+) = fraction (B^0) = (37.8 ± 2.2)% and PDG 96 values for B lifetime and branching ratio of D^* and D decays.
[38] FULTON 91 assumes assuming equal production of B^0 and B^+ at the $\Upsilon(4S)$ and uses Mark III D and D^* branching ratios.
[39] ALBRECHT 89J reports $0.018 \pm 0.006 \pm 0.005$. We rescale using the method described in STONE 94 but with the updated PDG 94 B($D^0 \to K^- \pi^+$).
[40] ATHANAS 97 uses missing energy and missing momentum to reconstruct neutrino.

$\Gamma(D^*(2010)^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0460±0.0021 OUR AVERAGE					
0.0459±0.0023±0.0040		[41] ABE	02F	BELL	$e^+e^- \to \Upsilon(4S)$
$0.0470\pm0.0013^{+0.0036}_{-0.0031}$		[42] ABREU	01H	DLPH	$e^+e^- \to Z$
0.0526±0.0020±0.0046		[43] ABBIENDI	00Q	OPAL	$e^+e^- \to Z$
0.0553±0.0026±0.0052		[44] BUSKULIC	97	ALEP	$e^+e^- \to Z$
0.0449±0.0032±0.0039	376	[45] BARISH	95	CLE2	$e^+e^- \to \Upsilon(4S)$
0.045 ±0.003 ±0.004		[46] ALBRECHT	94	ARG	$e^+e^- \to \Upsilon(4S)$
0.047 ±0.005 ±0.005	235	[47] ALBRECHT	93	ARG	$e^+e^- \to \Upsilon(4S)$
0.040 ±0.004 ±0.006		[48] BORTOLETTO	89B	CLEO	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.0508 \pm 0.0021 \pm 0.0066$		[49] ACKERSTAFF	97G	OPAL	Repl. by ABBIENDI 00Q
$0.0552 \pm 0.0017 \pm 0.0068$		[50] ABREU	96P	DLPH	Repl. by ABREU 01H
$0.0518 \pm 0.0030 \pm 0.0062$	410	[51] BUSKULIC	95N	ALEP	Sup. by BUSKULIC 97
seen	398	[52] SANGHERA	93	CLE2	$e^+e^- \to \Upsilon(4S)$
$0.070 \pm 0.018 \pm 0.014$		[53] ANTREASYAN	90B	CBAL	$e^+e^- \to \Upsilon(4S)$
		[54] ALBRECHT	89C	ARG	$e^+e^- \to \Upsilon(4S)$
$0.060 \pm 0.010 \pm 0.014$		[55] ALBRECHT	89J	ARG	$e^+e^- \to \Upsilon(4S)$
$0.070 \pm 0.012 \pm 0.019$	47	[56] ALBRECHT	87J	ARG	$e^+e^- \to \Upsilon(4S)$

[41] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[42] ABREU 01H measured using about 5000 partial reconstructed D^* sample.

[43] ABBIENDI 00Q assumes the fraction $B(b \to B^0) = (39.7^{+1.8}_{-2.2})\%$. This result is an average of two methods using exclusive and partial D^* reconstruction.

[44] BUSKULIC 97 assumes fraction (B^+) = fraction $(B^0) = (37.8 \pm 2.2)\%$ and PDG 96 values for B lifetime and D^* and D branching fractions.

[45] BARISH 95 use $B(D^0 \to K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*+} \to D^0 \pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$.

[46] ALBRECHT 94 assumes $B(D^{*+} \to D^0 \pi^+) = 68.1 \pm 1.0 \pm 1.3\%$. Uses partial reconstruction of D^{*+} and is independent of D^0 branching ratios.

[47] ALBRECHT 93 reports $0.052 \pm 0.005 \pm 0.006$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \to K^- \pi^+)$. We have taken their average e and μ value. They also obtain $\alpha = 2*\Gamma^0/(\Gamma^- + \Gamma^+) - 1 = 1.1 \pm 0.4 \pm 0.2$, $A_{AF} = 3/4*(\Gamma^- - \Gamma^+)/\Gamma = 0.2 \pm 0.08 \pm 0.06$ and a value of $|V_{cb}| = 0.036$–0.045 depending on model assumptions.

[48] We have taken average of the the BORTOLETTO 89B values for electrons and muons, $0.046 \pm 0.005 \pm 0.007$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \to K^- \pi^+)$. The measurement suggests a D^* polarization parameter value $\alpha = 0.65 \pm 0.66 \pm 0.25$.

[49] ACKERSTAFF 97G assumes fraction (B^+) = fraction $(B^0) = (37.8 \pm 2.2)\%$ and PDG 96 values for B lifetime and branching ratio of D^* and D decays.

[50] ABREU 96P result is the average of two methods using exclusive and partial D^* reconstruction.

[51] BUSKULIC 95N assumes fraction (B^+) = fraction $(B^0) = 38.2 \pm 1.3 \pm 2.2\%$ and $\tau_{B^0} = 1.58 \pm 0.06$ ps. $\Gamma(D^{*-} \ell^+ \nu_\ell)$/total = [5.18 − 0.13(fraction(B^0) − 38.2) − 1.5(τ_{B^0} − 1.58)]%.

[52] Combining $\overline{D}^{*0} \ell^+ \nu_\ell$ and $\overline{D}^{*-} \ell^+ \nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4*(\Gamma^- - \Gamma^+)/\Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb}, they measure V, A_1, and A_2, the three form factors for the $D^* \ell \nu_\ell$ decay, where results are slightly dependent on model assumptions.

[53] ANTREASYAN 90B is average over B and $\overline{D}^*(2010)$ charge states.

[54] The measurement of ALBRECHT 89C suggests a D^* polarization γ_L/γ_T of 0.85 ± 0.45. or $\alpha = 0.7 \pm 0.9$.

[55] ALBRECHT 89J is ALBRECHT 87J value rescaled using $B(D^*(2010)^- \to D^0 \pi^-) = 0.57 \pm 0.04 \pm 0.04$. Superseded by ALBRECHT 93.

[56] ALBRECHT 87J assume μ-e universality, the $B(\Upsilon(4S) \to B^0 \overline{B}^0) = 0.45$, the $B(D^0 \to K^- \pi^+) = (0.042 \pm 0.004 \pm 0.004)$, and the $B(D^*(2010)^- \to D^0 \pi^-) = 0.49 \pm 0.08$. Superseded by ALBRECHT 89J.

$\Gamma(\rho^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_4/Γ

$\ell = e$ or μ, not sum over e and μ modes.

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{2.57 \pm 0.29 {}^{+0.53}_{-0.62}}$		[57] BEHRENS	00	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$2.69 \pm 0.41 {}^{+0.61}_{-0.64}$		[58] BEHRENS	00	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$2.5 \pm 0.4 {}^{+0.7}_{-0.9}$		[59] ALEXANDER	96T	CLE2	Repl. by BEHRENS 00
<4.1	90	[60] BEAN	93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[57] Averaging with ALEXANDER 96T results including experimental and theoretical correlations considered, BEHRENS 00 reports systematic errors ${}^{+0.33}_{-0.46} \pm 0.41$, where the second error is theoretical model dependence. We combine these in quadrature.

[58] BEHRENS 00 reports ${}^{+0.35}_{-0.40} \pm 0.50$, where the second error is the theoretical model dependence. We combine these in quadrature. B^+ and B^0 decays combined using isospin symmetry: $\Gamma(B^0 \rightarrow \rho^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \rho^0 \ell^+ \nu) \approx 2\Gamma(B^+ \rightarrow \omega \ell^+ \nu)$. No evidence for $\omega \ell \nu$ is reported.

[59] ALEXANDER 96T reports ${}^{+0.5}_{-0.7} \pm 0.5$ where the second error is the theoretical model dependence. We combine these in quadrature. B^+ and B^0 decays combined using isospin symmetry: $\Gamma(B^0 \rightarrow \rho^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \rho^0 \ell^+ \nu) \approx 2\Gamma(B^+ \rightarrow \omega \ell^+ \nu)$. No evidence for $\omega \ell \nu$ is reported.

[60] BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\omega \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6\text{–}2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow (\omega \text{ or } \rho^0) \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.08\text{–}0.13$ at 90% CL is derived as well.

$\Gamma(\pi^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.8 \pm 0.4 \pm 0.4}$	[61] ALEXANDER	96T	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[61] ALEXANDER 96T gives systematic errors $\pm 0.3 \pm 0.2$ where the second error reflects the estimated model dependence. We combine these in quadrature. Assumes isospin symmetry: $\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2 \times \Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$.

$\Gamma(\pi^- \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID		TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	[62] ALBRECHT	91C	ARG

[62] In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

$\Gamma(K^+ \text{anything})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.78 \pm 0.08}$	[63] ALBRECHT	96D	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[63] Average multiplicity.

$\Gamma(D^- \pi^+)/\Gamma_{total}$ Γ_8/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0030±0.0004 OUR AVERAGE					
0.0029±0.0004±0.0002	81	[64] ALAM	94	CLE2	$e^+ e^- \to \Upsilon(4S)$
0.0027±0.0006±0.0005		[65] BORTOLETTO	92	CLEO	$e^+ e^- \to \Upsilon(4S)$
0.0048±0.0011±0.0011	22	[66] ALBRECHT	90J	ARG	$e^+ e^- \to \Upsilon(4S)$
$0.0051^{+0.0028}_{-0.0025}{}^{+0.0013}_{-0.0012}$	4	[67] BEBEK	87	CLEO	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0031±0.0013±0.0010	7	[66] ALBRECHT	88K	ARG	$e^+ e^- \to \Upsilon(4S)$

[64] ALAM 94 reports $[B(B^0 \to D^- \pi^+) \times B(D^+ \to K^- \pi^+ \pi^+)] = 0.000265 \pm 0.000032 \pm 0.000023$. We divide by our best value $B(D^+ \to K^- \pi^+ \pi^+) = (9.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[65] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[66] ALBRECHT 88K assumes $B^0 \overline{B}^0$:$B^+ B^-$ production ratio is 45:55. Superseded by ALBRECHT 90J which assumes 50:50.

[67] BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

$\Gamma(D^- \rho^+)/\Gamma_{total}$ Γ_9/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0078±0.0014 OUR AVERAGE					
0.0078±0.0013±0.0005	79	[68] ALAM	94	CLE2	$e^+ e^- \to \Upsilon(4S)$
0.009 ±0.005 ±0.003	9	[69] ALBRECHT	90J	ARG	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.022 ±0.012 ±0.009	6	[69] ALBRECHT	88K	ARG	$e^+ e^- \to \Upsilon(4S)$

[68] ALAM 94 reports $[B(B^0 \to D^- \rho^+) \times B(D^+ \to K^- \pi^+ \pi^+)] = 0.000704 \pm 0.000096 \pm 0.000070$. We divide by our best value $B(D^+ \to K^- \pi^+ \pi^+) = (9.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[69] ALBRECHT 88K assumes $B^0 \overline{B}^0$:$B^+ B^-$ production ratio is 45:55. Superseded by ALBRECHT 90J which assumes 50:50.

$\Gamma(D^- K^*(892)^+)/\Gamma_{total}$ Γ_{10}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$(3.7 \pm 1.5 \pm 1.0) \times 10^{-4}$	[70] MAHAPATRA	02	CLE2	$e^+ e^- \to \Upsilon(4S)$

[70] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- \omega \pi^+)/\Gamma_{total}$ Γ_{11}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0028±0.0005±0.0004	[71] ALEXANDER	01B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[71] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(D^- K^+)/\Gamma_{total}$ Γ_{12}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
(2.04±0.50±0.27) × 10^{-4}	[72] ABE	01I	BELL	$e^+e^- \to \Upsilon(4S)$

[72] ABE 01I reports $B(B^0 \to D^- K^+)/B(B^0 \to D^- \pi^+) = 0.068 \pm 0.015 \pm 0.007$. We multiply by our best value $B(B^0 \to D^- \pi^+) = (3.0 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.

$\Gamma(\overline{D}^0 \pi^+ \pi^-)/\Gamma_{total}$ Γ_{13}/Γ

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
<0.0016	90		[73] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.007	90		[74] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
<0.034	90		[75] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$
0.07 ±0.05		5	[76] BEHRENDS	83	CLEO	$e^+e^- \to \Upsilon(4S)$

[73] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[74] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D. The product branching fraction into $D_0^*(2340)\pi$ followed by $D_0^*(2340) \to D^0\pi$ is < 0.0001 at 90% CL and into $D_2^*(2460)$ followed by $D_2^*(2460) \to D^0\pi$ is < 0.0004 at 90% CL.

[75] BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%. $B(D^0 \to K^-\pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ and $B(D^0 \to K^-\pi^+\pi^+\pi^-) = (9.1 \pm 0.8 \pm 0.8)\%$ were used.

[76] Corrected by us using assumptions: $B(D^0 \to K^-\pi^+) = (0.042 \pm 0.006)$ and $B(\Upsilon(4S) \to B^0\overline{B}^0) = 50\%$. The product branching ratio is $B(B^0 \to \overline{D}^0\pi^+\pi^-)B(\overline{D}^0 \to K^+\pi^-) = (0.39 \pm 0.26) \times 10^{-2}$.

$\Gamma(D^*(2010)^- \pi^+)/\Gamma_{total}$ Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.00276±0.00021 OUR AVERAGE					
0.00281±0.00024±0.00005		[77] BRANDENB...	98	CLE2	$e^+e^- \to \Upsilon(4S)$
0.0026 ±0.0003 ±0.0004	82	[78] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
0.00337±0.00096±0.00002		[79] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
0.00236±0.00088±0.00002	12	[80] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$
$0.00236^{+0.00150}_{-0.00110}$±0.00002	5	[81] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.010 ±0.004 ±0.001	8	[82] AKERS	94J	OPAL	$e^+e^- \to Z$
0.0027 ±0.0014 ±0.0010	5	[83] ALBRECHT	87C	ARG	$e^+e^- \to \Upsilon(4S)$
0.0035 ±0.002 ±0.002		[84] ALBRECHT	86F	ARG	$e^+e^- \to \Upsilon(4S)$
0.017 ±0.005 ±0.005	41	[85] GILES	84	CLEO	$e^+e^- \to \Upsilon(4S)$

[77] BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \to D\pi)$.

[78] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \to D^0\pi^+)$ and absolute $B(D^0 \to K^-\pi^+)$ and the PDG 1992 $B(D^0 \to K^-\pi^+\pi^0)/B(D^0 \to K^-\pi^+)$ and $B(D^0 \to K^-\pi^+\pi^+\pi^-)/B(D^0 \to K^-\pi^+)$.

[79] BORTOLETTO 92 reports $0.0040 \pm 0.0010 \pm 0.0007$ for $B(D^*(2010)^+ \to D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[80] ALBRECHT 90J reports $0.0028 \pm 0.0009 \pm 0.0006$ for $B(D^*(2010)^+ \to D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[81] BEBEK 87 reports $0.0028^{+0.0015}_{-0.0012}{}^{+0.0010}_{-0.0006}$ for $B(D^*(2010)^+ \to D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92 and ALBRECHT 90J.

[82] Assumes $B(Z \to b\overline{b}) = 0.217$ and 38% B_d production fraction.

[83] ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \to B^+B^-) = 55\%$ and $B(\Upsilon(4S) \to B^0\overline{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

[84] ALBRECHT 86F uses pseudomass that is independent of D^0 and D^+ branching ratios.

[85] Assumes $B(D^*(2010)^+ \to D^0\pi^+) = 0.60^{+0.08}_{-0.15}$. Assumes $B(\Upsilon(4S) \to B^0\overline{B}^0) = 0.40 \pm 0.02$ Does not depend on D branching ratios.

$\Gamma(D^-\pi^+\pi^+\pi^-)/\Gamma_{total}$ — Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0080±0.0021±0.0014	[86] BORTOLETTO92	CLEO	$e^+e^- \to \Upsilon(4S)$

[86] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma((D^-\pi^+\pi^+\pi^-)\text{ nonresonant})/\Gamma_{total}$ — Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0039±0.0014±0.0013	[87] BORTOLETTO92	CLEO	$e^+e^- \to \Upsilon(4S)$

[87] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(D^-\pi^+\rho^0)/\Gamma_{total}$ — Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0011±0.0009±0.0004	[88] BORTOLETTO92	CLEO	$e^+e^- \to \Upsilon(4S)$

[88] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(D^-a_1(1260)^+)/\Gamma_{total}$ — Γ_{18}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0060±0.0022±0.0024	[89] BORTOLETTO92	CLEO	$e^+e^- \to \Upsilon(4S)$

[89] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(D^*(2010)^- \pi^+ \pi^0)/\Gamma_{total}$ Γ_{19}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0152±0.0052±0.0001	51	[90] ALBRECHT	90J	ARG	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.015 ±0.008 ±0.008	8	[91] ALBRECHT	87C	ARG	$e^+ e^- \to \Upsilon(4S)$

[90] ALBRECHT 90J reports $0.018 \pm 0.004 \pm 0.005$ for $B(D^*(2010)^+ \to D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[91] ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \to B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \to B^0 \overline{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^*(2010)^- \rho^+)/\Gamma_{total}$ Γ_{20}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0073 ±0.0015 OUR AVERAGE					
0.0074 ±0.0010 ±0.0014	76	[92,93] ALAM	94	CLE2	$e^+ e^- \to \Upsilon(4S)$
0.0160 ±0.0113 ±0.0001		[94] BORTOLETTO	92	CLEO	$e^+ e^- \to \Upsilon(4S)$
0.00589±0.00352±0.00004	19	[95] ALBRECHT	90J	ARG	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.081 ±0.029 $^{+0.059}_{-0.024}$	19	[96] CHEN	85	CLEO	$e^+ e^- \to \Upsilon(4S)$

[92] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \to D^0 \pi^+)$ and absolute $B(D^0 \to K^- \pi^+)$ and the PDG 1992 $B(D^0 \to K^- \pi^+ \pi^0)/B(D^0 \to K^- \pi^+)$ and $B(D^0 \to K^- \pi^+ \pi^+ \pi^-)/B(D^0 \to K^- \pi^+)$.

[93] This decay is nearly completely longitudinally polarized, $\Gamma_L/\Gamma = (93 \pm 5 \pm 5)\%$, as expected from the factorization hypothesis (ROSNER 90). The nonresonant $\pi^+ \pi^0$ contribution under the ρ^+ is less than 9% at 90% CL.

[94] BORTOLETTO 92 reports $0.019 \pm 0.008 \pm 0.011$ for $B(D^*(2010)^+ \to D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[95] ALBRECHT 90J reports $0.007 \pm 0.003 \pm 0.003$ for $B(D^*(2010)^+ \to D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[96] Uses $B(D^* \to D^0 \pi^+) = 0.6 \pm 0.15$ and $B(\Upsilon(4S) \to B^0 \overline{B}^0) = 0.4$. Does not depend on D branching ratios.

$\Gamma(D^*(2010)^- K^+)/\Gamma_{total}$ Γ_{21}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
(2.04±0.44±0.16) × 10^{-4}	[97] ABE	01I	BELL	$e^+ e^- \to \Upsilon(4S)$

[97] ABE 01I reports $B(B^0 \to D^*(2010)^- K^+)/B(B^0 \to D^*(2010)^- \pi^+) = 0.074 \pm 0.015 \pm 0.006$. We multiply by our best value $B(B^0 \to D^*(2010)^- \pi^+) = (2.76 \pm 0.21) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.

$\Gamma(D^*(2010)^- K^*(892)^+)/\Gamma_{total}$ Γ_{22}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$(3.8\pm1.3\pm0.8)\times10^{-4}$	[98] MAHAPATRA	02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[98] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and an unpolarized final state.

$\Gamma(D^*(2010)^- \pi^+\pi^+\pi^-)/\Gamma_{total}$ Γ_{23}/Γ

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
0.0076±0.0018 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.						
$0.0063\pm0.0010\pm0.0011$		49	[99,100] ALAM	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$0.0134\pm0.0036\pm0.0001$			[101] BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$0.0101\pm0.0041\pm0.0001$		26	[102] ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$0.033\ \pm0.009\ \pm0.016$		27	[103] ALBRECHT	87C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<0.042	90		[104] BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[99] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2010)^+ \rightarrow D^0\pi^+$) and absolute B($D^0 \rightarrow K^-\pi^+$) and the PDG 1992 B($D^0 \rightarrow K^-\pi^+\pi^0$)/B($D^0 \rightarrow K^-\pi^+$) and B($D^0 \rightarrow K^-\pi^+\pi^+\pi^-$)/B($D^0 \rightarrow K^-\pi^+$).

[100] The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^+, the branching ratio for $\overline{D}^{*-} a_1^+$ is twice that for $\overline{D}^{*-}\pi^+\pi^+\pi^-$.)

[101] BORTOLETTO 92 reports $0.0159 \pm 0.0028 \pm 0.0037$ for B($D^*(2010)^+ \rightarrow D^0\pi^+$) = 0.57 ± 0.06. We rescale to our best value B($D^*(2010)^+ \rightarrow D^0\pi^+$) = $(67.7 \pm 0.5)\times10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[102] ALBRECHT 90J reports $0.012 \pm 0.003 \pm 0.004$ for B($D^*(2010)^+ \rightarrow D^0\pi^+$) = 0.57 ± 0.06. We rescale to our best value B($D^*(2010)^+ \rightarrow D^0\pi^+$) = $(67.7 \pm 0.5)\times10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[103] ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume B($\Upsilon(4S) \rightarrow B^+B^-$) = 55% and B($\Upsilon(4S) \rightarrow B^0\overline{B}^0$) = 45%. Superseded by ALBRECHT 90J.

[104] BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.



$\Gamma\left(D^*(2010)^- \pi^+ \pi^+ \pi^-\right)/\Gamma_{\text{total}}$

$\Gamma\left(\left(D^*(2010)^- \pi^+ \pi^+ \pi^-\right) \text{nonresonant}\right)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE	*DOCUMENT ID*	*TECN*	*COMMENT*
0.0000±0.0019±0.0016	[105] BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

[105] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

$\Gamma\left(D^*(2010)^- \pi^+ \rho^0\right)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	*DOCUMENT ID*	*TECN*	*COMMENT*
0.00573±0.00317±0.00004	[106] BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

[106] BORTOLETTO 92 reports $0.0068 \pm 0.0032 \pm 0.0021$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma\left(D^*(2010)^- a_1(1260)^+\right)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.0130±0.0027 OUR AVERAGE				
0.0126±0.0020±0.0022	[107,108] ALAM	94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0152±0.0070±0.0001	[109] BORTOLETTO	92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

[107] ALAM 94 value is twice their $\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ value based on their observation that the three pions are dominantly in the $a_1(1260)$ mass range 1.0 to 1.6 GeV.

[108] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

[109] BORTOLETTO 92 reports $0.018 \pm 0.006 \pm 0.006$ for $B(D^*(2010)^+ \to D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(D^*(2010)^- \pi^+\pi^+\pi^-\pi^0)/\Gamma_{total}$ — Γ_{27}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0176±0.0027 OUR AVERAGE					
0.0172±0.0014±0.0024		[110] ALEXANDER	01B	CLE2	$e^+e^- \to \Upsilon(4S)$
0.0345±0.0181±0.0003	28	[111] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$

[110] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega\pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

[111] ALBRECHT 90J reports $0.041 \pm 0.015 \pm 0.016$ for $B(D^*(2010)^+ \to D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(D^*(2010)^- p\overline{p}\pi^+)/\Gamma_{total}$ — Γ_{29}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$\mathbf{6.5^{+1.3}_{-1.2}\pm1.0}$	[112] ANDERSON	01	CLE2	$e^+e^- \to \Upsilon(4S)$

[112] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- p\overline{n})/\Gamma_{total}$ — Γ_{30}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$\mathbf{14.5^{+3.4}_{-3.0}\pm2.7}$	[113] ANDERSON	01	CLE2	$e^+e^- \to \Upsilon(4S)$

[113] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^*(2010)^- \omega\pi^+)/\Gamma_{total}$ — Γ_{31}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0029±0.0003±0.0004	[114] ALEXANDER	01B	CLE2	$e^+e^- \to \Upsilon(4S)$

[114] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega\pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(\overline{D}^*_2(2460)^- \pi^+)/\Gamma_{total}$ — Γ_{32}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0022	90	[115] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$

[115] ALAM 94 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \to K^-\pi^+)$ and $B(D^*_2(2460)^+ \to D^0\pi^+) = 30\%$.

$\Gamma(\overline{D}_2^*(2460)^- \rho^+)/\Gamma_{\text{total}}$ — Γ_{33}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0049	90	[116] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$

[116] ALAM 94 assumesequal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute B($D^0 \to K^- \pi^+$) and B($D_2^*(2460)^+ \to D^0 \pi^+$) = 30%.

$\Gamma(D^- D^+)/\Gamma_{\text{total}}$ — Γ_{34}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<9.4 \times 10^{-4}$	90	[117] LIPELES	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<5.9 \times 10^{-3}$	90	BARATE	98Q	ALEP	$e^+e^- \to Z$
$<1.2 \times 10^{-3}$	90	ASNER	97	CLE2	$e^+e^- \to \Upsilon(4S)$

[117] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- D_s^+)/\Gamma_{\text{total}}$ — Γ_{35}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0080±0.0030 OUR AVERAGE					
$0.0084 \pm 0.0030 ^{+0.0020}_{-0.0021}$		[118] GIBAUT	96	CLE2	$e^+e^- \to \Upsilon(4S)$
$0.013 \pm 0.011 \pm 0.003$		[119] ALBRECHT	92G	ARG	$e^+e^- \to \Upsilon(4S)$
$0.007 \pm 0.004 \pm 0.002$		[120] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.012 ± 0.007	3	[121] BORTOLETTO	90	CLEO	$e^+e^- \to \Upsilon(4S)$

[118] GIBAUT 96 reports $0.0087 \pm 0.0024 \pm 0.0020$ for B($D_s^+ \to \phi\pi^+$) = 0.035. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = $(3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[119] ALBRECHT 92G reports $0.017 \pm 0.013 \pm 0.006$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = $(3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ branching ratios, e.g., B($D^+ \to K^- \pi^+ \pi^+$) = 7.7 ± 1.0%.

[120] BORTOLETTO 92 reports $0.0080 \pm 0.0045 \pm 0.0030$ for B($D_s^+ \to \phi\pi^+$) = 0.030 ± 0.011. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = $(3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[121] BORTOLETTO 90 assume B($D_s \to \phi\pi^+$) = 2%. Superseded by BORTOLETTO 92.

$\Gamma(D^*(2010)^- D_s^+)/\Gamma_{\text{total}}$ — Γ_{36}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0111±0.0033 OUR AVERAGE					
$0.0110 \pm 0.0021 ^{+0.0026}_{-0.0027}$		[122] AHMED	00B	CLE2	$e^+e^- \to \Upsilon(4S)$
$0.010 \pm 0.008 \pm 0.003$		[123] ALBRECHT	92G	ARG	$e^+e^- \to \Upsilon(4S)$
$0.013 \pm 0.008 \pm 0.003$		[124] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.0090 \pm 0.0027 \pm 0.0022$		[125] GIBAUT	96	CLE2	Repl. by AHMED 00B
0.024 ± 0.014	3	[126] BORTOLETTO	90	CLEO	$e^+e^- \to \Upsilon(4S)$

[122] AHMED 00B reports $0.0110 \pm 0.0018 \pm 0.0011$ for $B(D_s^+ \to \phi\pi^+) = 0.036$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[123] ALBRECHT 92G reports $0.014 \pm 0.010 \pm 0.003$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ and $D^*(2010)^+$ branching ratios, e.g., $B(D^0 \to K^-\pi^+) = 3.71 \pm 0.25\%$, $B(D^+ \to K^-\pi^+\pi^+) = 7.1 \pm 1.0\%$, and $B(D^*(2010)^+ \to D^0\pi^+) = 55 \pm 4\%$.

[124] BORTOLETTO 92 reports $0.016 \pm 0.009 \pm 0.006$ for $B(D_s^+ \to \phi\pi^+) = 0.030 \pm 0.011$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

[125] GIBAUT 96 reports $0.0093 \pm 0.0023 \pm 0.0016$ for $B(D_s^+ \to \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[126] BORTOLETTO 90 assume $B(D_s \to \phi\pi^+) = 2\%$. Superseded by BORTOLETTO 92.

$\Gamma(D^- D_s^{*+})/\Gamma_{total}$ — Γ_{37}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.010±0.005 OUR AVERAGE				
$0.010 \pm 0.004 \pm 0.002$	[127] GIBAUT	96	CLE2	$e^+e^- \to \Upsilon(4S)$
$0.020 \pm 0.014 \pm 0.005$	[128] ALBRECHT	92G	ARG	$e^+e^- \to \Upsilon(4S)$

[127] GIBAUT 96 reports $0.0100 \pm 0.0035 \pm 0.0022$ for $B(D_s^+ \to \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[128] ALBRECHT 92G reports $0.027 \pm 0.017 \pm 0.009$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ branching ratios, e.g., $B(D^+ \to K^-\pi^+\pi^+) = 7.7 \pm 1.0\%$.

$[\Gamma(D^*(2010)^- D_s^+) + \Gamma(D^*(2010)^- D_s^{*+})]/\Gamma_{total}$ — $(\Gamma_{36}+\Gamma_{38})/\Gamma$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
$\mathbf{4.15 \pm 1.11^{+0.99}_{-1.02}}$	22	[129] BORTOLETTO	90	CLEO	$e^+e^- \to \Upsilon(4S)$

[129] BORTOLETTO 90 reports 7.5 ± 2.0 for $B(D_s^+ \to \phi\pi^+) = 0.02$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^*(2010)^- D_s^{*+})/\Gamma_{total}$ — Γ_{38}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.018±0.006 OUR AVERAGE				
$0.018 \pm 0.004 \pm 0.004$	[130] AHMED	00B	CLE2	$e^+e^- \to \Upsilon(4S)$
$0.019 \pm 0.011 \pm 0.005$	[131] ALBRECHT	92G	ARG	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.020 \pm 0.006 \pm 0.005$	[132] GIBAUT	96	CLE2	Repl. by AHMED 00B

[130] AHMED 00B reports $0.0182 \pm 0.0037 \pm 0.0025$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[131] ALBRECHT 92G reports $0.026 \pm 0.014 \pm 0.006$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ and $D^*(2010)^+$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$, $B(D^+ \rightarrow K^-\pi^+\pi^+) = 7.1 \pm 1.0\%$, and $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 55 \pm 4\%$.

[132] GIBAUT 96 reports $0.0203 \pm 0.0050 \pm 0.0036$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^+\pi^-)/\Gamma_{\text{total}}$ — Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00028	90	[133] ALEXANDER	93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0013	90	[134] BORTOLETTO	90	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[133] ALEXANDER 93B reports $< 2.7 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

[134] BORTOLETTO 90 assume $B(D_s \rightarrow \phi\pi^+) = 2\%$.

$[\Gamma(D_s^+\pi^-) + \Gamma(D_s^- K^+)]/\Gamma_{\text{total}}$ — $(\Gamma_{39}+\Gamma_{45})/\Gamma$

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0013	90	[135] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[135] ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^{*+}\pi^-)/\Gamma_{\text{total}}$ — Γ_{40}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0005	90	[136] ALEXANDER	93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[136] ALEXANDER 93B reports $< 4.4 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$[\Gamma(D_s^{*+}\pi^-) + \Gamma(D_s^{*-} K^+)]/\Gamma_{\text{total}}$ — $(\Gamma_{40}+\Gamma_{46})/\Gamma$

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0009	90	[137] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[137] ALBRECHT 93E reports $< 1.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^+ \rho^-)/\Gamma_{\rm total}$ — Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0007	90	[138] ALEXANDER	93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0016	90	[139] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[138] ALEXANDER 93B reports $< 6.6 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

[139] ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^{*+} \rho^-)/\Gamma_{\rm total}$ — Γ_{42}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0008	90	[140] ALEXANDER	93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0019	90	[141] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[140] ALEXANDER 93B reports $< 7.4 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

[141] ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^+ a_1(1260)^-)/\Gamma_{\rm total}$ — Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0026	90	[142] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[142] ALBRECHT 93E reports $< 3.5 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^{*+} a_1(1260)^-)/\Gamma_{\rm total}$ — Γ_{44}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0022	90	[143] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[143] ALBRECHT 93E reports $< 2.9 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^- K^+)/\Gamma_{\rm total}$ — Γ_{45}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00024	90	[144] ALEXANDER	93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0013	90	[145] BORTOLETTO	90	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[144] ALEXANDER 93B reports $< 2.3 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

[145] BORTOLETTO 90 assume $B(D_s \rightarrow \phi\pi^+) = 2\%$.

$\Gamma(D_s^{*-} K^+)/\Gamma_{total}$ — Γ_{46}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00017	90	[146] ALEXANDER 93B	CLE2	$e^+e^- \to \Upsilon(4S)$

[146] ALEXANDER 93B reports $< 1.7 \times 10^{-4}$ for $B(D_s^+ \to \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

$\Gamma(D_s^- K^*(892)^+)/\Gamma_{total}$ — Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0010	90	[147] ALEXANDER 93B	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0034	90	[148] ALBRECHT 93E	ARG	$e^+e^- \to \Upsilon(4S)$

[147] ALEXANDER 93B reports $< 9.7 \times 10^{-4}$ for $B(D_s^+ \to \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

[148] ALBRECHT 93E reports $< 4.6 \times 10^{-3}$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

$\Gamma(D_s^{*-} K^*(892)^+)/\Gamma_{total}$ — Γ_{48}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0011	90	[149] ALEXANDER 93B	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.004	90	[150] ALBRECHT 93E	ARG	$e^+e^- \to \Upsilon(4S)$

[149] ALEXANDER 93B reports $< 11.0 \times 10^{-4}$ for $B(D_s^+ \to \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

[150] ALBRECHT 93E reports $< 5.8 \times 10^{-3}$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

$\Gamma(D_s^- \pi^+ K^0)/\Gamma_{total}$ — Γ_{49}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.005	90	[151] ALBRECHT 93E	ARG	$e^+e^- \to \Upsilon(4S)$

[151] ALBRECHT 93E reports $< 7.3 \times 10^{-3}$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

$\Gamma(D_s^{*-} \pi^+ K^0)/\Gamma_{total}$ — Γ_{50}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0031	90	[152] ALBRECHT 93E	ARG	$e^+e^- \to \Upsilon(4S)$

[152] ALBRECHT 93E reports $< 4.2 \times 10^{-3}$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

$\Gamma(D_s^- \pi^+ K^*(892)^0)/\Gamma_{total}$ — Γ_{51}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.004	90	[153] ALBRECHT 93E	ARG	$e^+e^- \to \Upsilon(4S)$

[153] ALBRECHT 93E reports $< 5.0 \times 10^{-3}$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

$\Gamma(D_s^{*-}\pi^+ K^*(892)^0)/\Gamma_{total}$ — Γ_{52}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0020	90	[154] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$

[154] ALBRECHT 93E reports $< 2.7 \times 10^{-3}$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.036$.

$\Gamma(\overline{D}^0\pi^0)/\Gamma_{total}$ — Γ_{53}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
2.9 ±0.5 OUR AVERAGE					
3.1 ±0.4 ±0.5		[155] ABE	02J	BELL	$e^+e^- \to \Upsilon(4S)$
$2.74^{+0.36}_{-0.32}\pm0.55$		[155] COAN	02	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.2	90	[156] NEMATI	98	CLE2	Repl. by COAN 02
<4.8	90	[157] ALAM	94	CLE2	Repl. by NEMATI 98

[155] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[156] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.

[157] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \to K^-\pi^+)$ and the PDG 1992 $B(D^0 \to K^-\pi^+\pi^0)/B(D^0 \to K^-\pi^+)$ and $B(D^0 \to K^-\pi^+\pi^+\pi^-)/B(D^0 \to K^-\pi^+)$.

$\Gamma(\overline{D}^0\rho^0)/\Gamma_{total}$ — Γ_{54}/Γ

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
<0.00039	90		[158] NEMATI	98	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.00055	90		[159] ALAM	94	CLE2	Repl. by NEMATI 98
<0.0006	90		[160] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
<0.0027	90	4	[161] ALBRECHT	88K	ARG	$e^+e^- \to \Upsilon(4S)$

[158] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.

[159] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \to K^-\pi^+)$ and the PDG 1992 $B(D^0 \to K^-\pi^+\pi^0)/B(D^0 \to K^-\pi^+)$ and $B(D^0 \to K^-\pi^+\pi^+\pi^-)/B(D^0 \to K^-\pi^+)$.

[160] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[161] ALBRECHT 88K reports < 0.003 assuming $B^0\overline{B}^0$:B^+B^- production ratio is 45:55. We rescale to 50%.

$\Gamma(\overline{D}^0\eta)/\Gamma_{total}$ — Γ_{55}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.4^{+0.5}_{-0.4}\pm0.3}$		[162] ABE	02J	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.3	90	[163] NEMATI	98	CLE2	$e^+e^- \to \Upsilon(4S)$
<6.8	90	[164] ALAM	94	CLE2	Repl. by NEMATI 98

[162] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[163] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.
[164] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute B($D^0 \to K^- \pi^+$) and the PDG 1992 B($D^0 \to K^- \pi^+ \pi^0$)/B($D^0 \to K^- \pi^+$) and B($D^0 \to K^- \pi^+ \pi^+ \pi^-$)/B($D^0 \to K^- \pi^+$).

$\Gamma(\overline{D}^0 \eta')/\Gamma_{total}$ — Γ_{56}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00094	90	[165] NEMATI	98	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.00086	90	[166] ALAM	94	CLE2	Repl. by NEMATI 98

[165] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.
[166] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute B($D^0 \to K^- \pi^+$) and the PDG 1992 B($D^0 \to K^- \pi^+ \pi^0$)/B($D^0 \to K^- \pi^+$) and B($D^0 \to K^- \pi^+ \pi^+ \pi^-$)/B($D^0 \to K^- \pi^+$).

$\Gamma(\overline{D}^0 \omega)/\Gamma_{total}$ — Γ_{57}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
$1.8 \pm 0.5^{+0.4}_{-0.3}$		[167] ABE	02J	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<5.1	90	[168] NEMATI	98	CLE2	$e^+e^- \to \Upsilon(4S)$
<6.3	90	[169] ALAM	94	CLE2	Repl. by NEMATI 98

[167] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[168] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.
[169] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute B($D^0 \to K^- \pi^+$) and the PDG 1992 B($D^0 \to K^- \pi^+ \pi^0$)/B($D^0 \to K^- \pi^+$) and B($D^0 \to K^- \pi^+ \pi^+ \pi^-$)/B($D^0 \to K^- \pi^+$).

$\Gamma(\overline{D}^{*0} \gamma)/\Gamma_{total}$ — Γ_{58}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<5.0 \times 10^{-5}$	90	[170] ARTUSO	00	CLE2	$e^+e^- \to \Upsilon(4S)$

[170] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^*(2007)^0 \pi^0)/\Gamma_{total}$ — Γ_{59}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
2.5 ±0.7 OUR AVERAGE					
$2.7\ {}^{+0.8}_{-0.7}\ {}^{+0.5}_{-0.6}$		[171] ABE	02J	BELL	$e^+e^- \to \Upsilon(4S)$
$2.20^{+0.59}_{-0.52} \pm 0.79$		[171] COAN	02	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.4	90	[172] NEMATI	98	CLE2	Repl. by COAN 02
<9.7	90	[173] ALAM	94	CLE2	Repl. by NEMATI 98

[171] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[172] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.
[173] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2007)^0 \to D^0\pi^0$) and absolute B($D^0 \to K^-\pi^+$) and the PDG 1992 B($D^0 \to K^-\pi^+\pi^0$)/B($D^0 \to K^-\pi^+$) and B($D^0 \to K^-\pi^+\pi^+\pi^-$)/B($D^0 \to K^-\pi^+$).

$\Gamma(\overline{D}^*(2007)^0\rho^0)/\Gamma_{total}$ — Γ_{60}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00056	90	[174] NEMATI	98	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.00117	90	[175] ALAM	94	CLE2	Repl. by NEMATI 98

[174] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.
[175] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2007)^0 \to D^0\pi^0$) and absolute B($D^0 \to K^-\pi^+$) and the PDG 1992 B($D^0 \to K^-\pi^+\pi^0$)/B($D^0 \to K^-\pi^+$) and B($D^0 \to K^-\pi^+\pi^+\pi^-$)/B($D^0 \to K^-\pi^+$).

$\Gamma(\overline{D}^*(2007)^0\eta)/\Gamma_{total}$ — Γ_{61}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00026	90	[176] NEMATI	98	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.00046	90	[177] ABE	02J	BELL	$e^+e^- \to \Upsilon(4S)$
<0.00069	90	[178] ALAM	94	CLE2	Repl. by NEMATI 98

[176] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.
[177] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[178] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2007)^0 \to D^0\pi^0$) and absolute B($D^0 \to K^-\pi^+$) and the PDG 1992 B($D^0 \to K^-\pi^+\pi^0$)/B($D^0 \to K^-\pi^+$) and B($D^0 \to K^-\pi^+\pi^+\pi^-$)/B($D^0 \to K^-\pi^+$).

$\Gamma(\overline{D}^*(2007)^0\eta')/\Gamma_{total}$ — Γ_{62}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0014	90	BRANDENB...	98	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0019	90	[179] NEMATI	98	CLE2	$e^+e^- \to \Upsilon(4S)$
<0.0027	90	[180] ALAM	94	CLE2	Repl. by NEMATI 98

[179] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.
[180] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2007)^0 \to D^0\pi^0$) and absolute B($D^0 \to K^-\pi^+$) and the PDG 1992 B($D^0 \to K^-\pi^+\pi^0$)/B($D^0 \to K^-\pi^+$) and B($D^0 \to K^-\pi^+\pi^+\pi^-$)/B($D^0 \to K^-\pi^+$).

$\Gamma(\overline{D}^*(2007)^0 \omega)/\Gamma_{total}$ — Γ_{63}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00074	90	[181] NEMATI	98	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.00079	90	[182] ABE	02J	BELL	$e^+ e^- \to \Upsilon(4S)$
<0.0021	90	[183] ALAM	94	CLE2	Repl. by NEMATI 98

[181] NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0, D^{*0}, η, η', and ω branching fractions.

[182] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[183] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2007)^0 \to D^0 \pi^0$) and absolute B($D^0 \to K^- \pi^+$) and the PDG 1992 B($D^0 \to K^- \pi^+ \pi^0$)/B($D^0 \to K^- \pi^+$) and B($D^0 \to K^- \pi^+ \pi^+ \pi^-$)/B($D^0 \to K^- \pi^+$).

$\Gamma(D^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-)/\Gamma_{total}$ — Γ_{64}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
3.0±0.7±0.6	[184] EDWARDS	02	CLE2	$e^+ e^- \to \Upsilon(4S)$

[184] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-)/\Gamma(D^*(2010)^+ \pi^+ \pi^- \pi^- \pi^0)$ — Γ_{64}/Γ_{28}

VALUE	DOCUMENT ID		TECN	COMMENT
0.17±0.04±0.02	[185] EDWARDS	02	CLE2	$e^+ e^- \to \Upsilon(4S)$

[185] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{total}$ — Γ_{65}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{(9.9^{+4.2}_{-3.3}\pm1.2) \times 10^{-4}}$		[186] LIPELES	00	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$(6.2^{+4.0}_{-2.9}\pm1.0) \times 10^{-4}$		[187] ARTUSO	99	CLE2	Repl. by LIPELES 00
$< 6.1 \times 10^{-3}$	90	[188] BARATE	98Q	ALEP	$e^+ e^- \to Z$
$< 2.2 \times 10^{-3}$	90	[189] ASNER	97	CLE2	Repl. by ARTUSO 99

[186] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[187] ARTUSO 99 uses B($\Upsilon(4S) \to B^0 \overline{B}^0$)=(48 ± 4)%.

[188] BARATE 98Q (ALEPH) observes 2 events with an expected background of 0.10 ± 0.03 which corresponds to a branching ratio of $(2.3^{+1.9}_{-1.2} \pm 0.4) \times 10^{-3}$.

[189] ASNER 97 at CLEO observes 1 event with an expected background of 0.022 ± 0.011. This correcsponds to a branching ratio of $(5.3^{+7.1}_{-3.7} \pm 1.0) \times 10^{-4}$.

$\Gamma(D^*(2010)^+ D^-)/\Gamma_{total}$ — Γ_{66}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<6.3 \times 10^{-4}}$	90	[190] LIPELES	00	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<5.6 \times 10^{-3}$	90	BARATE	98Q	ALEP	$e^+ e^- \to Z$
$<1.8 \times 10^{-3}$	90	ASNER	97	CLE2	$e^+ e^- \to \Upsilon(4S)$

[190] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^{(*)0}\overline{D}^{(*)0})/\Gamma_{total}$ — Γ_{67}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.027	90	BARATE	98Q	ALEP	$e^+e^- \rightarrow Z$

$\Gamma(\eta_c K^0)/\Gamma_{total}$ — Γ_{68}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.09^{+0.55}_{-0.42}\pm0.33}$	[191] EDWARDS	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[191] EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from B($J/\psi(1S) \rightarrow \gamma\eta_c$) in those modes have been accounted for.

$\Gamma(J/\psi(1S)K^0)/\Gamma_{total}$ — Γ_{69}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
8.7±0.5 OUR AVERAGE						
8.3±0.4±0.5			[192] AUBERT	02	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
9.5±0.8±0.6			[192] AVERY	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
11.5±2.3±1.7			[193] ABE	96H	CDF	$p\overline{p}$ at 1.8 TeV
7.0±4.1±0.1			[194] BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
9.3±7.3±0.2		2	[195] ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$8.5^{+1.4}_{-1.2}\pm0.6$			[192] JESSOP	97	CLE2	Repl. by AVERY 00
7.5±2.4±0.8		10	[194] ALAM	94	CLE2	Sup. by JESSOP 97
<50	90		ALAM	86	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[192] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[193] ABE 96H assumes that B($B^+ \rightarrow J/\psi K^+$) = $(1.02 \pm 0.14) \times 10^{-3}$.
[194] BORTOLETTO 92 reports $6 \pm 3 \pm 2$ for B($J/\psi(1S) \rightarrow e^+e^-$) = 0.069 ± 0.009. We rescale to our best value B($J/\psi(1S) \rightarrow e^+e^-$) = $(5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[195] ALBRECHT 90J reports $8 \pm 6 \pm 2$ for B($J/\psi(1S) \rightarrow e^+e^-$) = 0.069 ± 0.009. We rescale to our best value B($J/\psi(1S) \rightarrow e^+e^-$) = $(5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)K^+\pi^-)/\Gamma_{total}$ — Γ_{70}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
1.16±0.56±0.02			[196] BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<1.3	90		[197] ALBRECHT	87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<6.3	90	2	GILES	84	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[196] BORTOLETTO 92 reports $1.0 \pm 0.4 \pm 0.3$ for B($J/\psi(1S) \rightarrow e^+e^-$) = 0.069 ± 0.009. We rescale to our best value B($J/\psi(1S) \rightarrow e^+e^-$) = $(5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[197] ALBRECHT 87D assume $B^+B^-/B^0\overline{B}^0$ ratio is 55/45. $K\pi$ system is specifically selected as nonresonant.

$\Gamma(J/\psi(1S)K^*(892)^0)/\Gamma_{total}$ Γ_{71}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
1.31±0.09 OUR AVERAGE					
1.24±0.05±0.09		[198] AUBERT	02	BABR	$e^+e^- \to \Upsilon(4S)$
1.74±0.20±0.18		[199] ABE	98O	CDF	$p\overline{p}$ 1.8 TeV
1.32±0.17±0.17		[200] JESSOP	97	CLE2	$e^+e^- \to \Upsilon(4S)$
1.28±0.66±0.02		[201] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
1.28±0.60±0.02	6	[202] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$
4.1 ±1.8 ±0.1	5	[203] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.36±0.27±0.22		[204] ABE	96H	CDF	Sup. by ABE 98O
1.69±0.31±0.18	29	[205] ALAM	94	CLE2	Sup. by JESSOP 97
		[206] ALBRECHT	94G	ARG	$e^+e^- \to \Upsilon(4S)$
4.0 ±0.30		[207] ALBAJAR	91E	UA1	$E_{\rm cm}^{p\overline{p}}$= 630 GeV
3.3 ±0.18	5	[208] ALBRECHT	87D	ARG	$e^+e^- \to \Upsilon(4S)$
4.1 ±0.18	5	[209] ALAM	86	CLEO	Repl. by BEBEK 87

[198] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[199] ABE 98O reports $[B(B^0 \to J/\psi(1S)K^*(892)^0)]/[B(B^+ \to J/\psi(1S)K^+)] = 1.76 \pm 0.14 \pm 0.15$. We multiply by our best value $B(B^+ \to J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[200] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[201] BORTOLETTO 92 reports $1.1 \pm 0.5 \pm 0.3$ for $B(J/\psi(1S) \to e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \to e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[202] ALBRECHT 90J reports $1.1 \pm 0.5 \pm 0.2$ for $B(J/\psi(1S) \to e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \to e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[203] BEBEK 87 reports $3.5 \pm 1.6 \pm 0.3$ for $B(J/\psi(1S) \to e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \to e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Updated in BORTOLETTO 92 to use the same assumptions.

[204] ABE 96H assumes that $B(B^+ \to J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.

[205] The neutral and charged B events together are predominantly longitudinally polarized, $\Gamma_L/\Gamma = 0.080 \pm 0.08 \pm 0.05$. This can be compared with a prediction using HQET, 0.73 (KRAMER 92). This polarization indicates that the $B \to \psi K^*$ decay is dominated by the $CP = -1$ CP eigenstate. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[206] ALBRECHT 94G measures the polarization in the vector-vector decay to be predominantly longitudinal, $\Gamma_T/\Gamma = 0.03 \pm 0.16 \pm 0.15$ making the neutral decay a CP eigenstate when the K^{*0} decays through $K_S^0\pi^0$.

[207] ALBAJAR 91E assumes B_d^0 production fraction of 36%.

[208] ALBRECHT 87D assume $B^+B^-/B^0\overline{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

[209] ALAM 86 assumes $B^\pm/B^0$ ratio is 60/40. The observation of the decay $B^+ \to J/\psi K^*(892)^+$ (HAAS 85) has been retracted in this paper.

$\Gamma(J/\psi(1S)K^*(892)^0)/\Gamma(J/\psi(1S)K^0)$ — Γ_{71}/Γ_{69}

VALUE	DOCUMENT ID		TECN	COMMENT
1.48±0.12 OUR AVERAGE				
1.49±0.10±0.08	[210] AUBERT	02	BABR	$e^+e^- \to \Upsilon(4S)$
1.39±0.36±0.10	ABE	96Q	CDF	$p\overline{p}$

[210] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\phi K^0)/\Gamma_{total}$ — Γ_{72}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$(8.8^{+3.5}_{-3.0}\pm1.3)\times10^{-5}$	[211] ANASTASSOV	00	CLE2	$e^+e^- \to \Upsilon(4S)$

[211] ANASTASSOV 00 finds 10 events on a background of 0.5 ± 0.2. Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$, a uniform Dalitz plot distribution, isotropic $J/\psi(1S)$ and ϕ decays, and B($B^+ \to J/\psi(1S)\phi K^+$)= B($B^0 \to J/\psi(1S)\phi K^0$).

$\Gamma(J/\psi(1S)K(1270)^0)/\Gamma_{total}$ — Γ_{73}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
1.30±0.34±0.32	[212] ABE	01L	BELL	$e^+e^- \to \Upsilon(4S)$

[212] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the PDG value of B($B^+ \to J/\psi(1S)K^+$) = $(1.00 \pm 0.10)\times10^{-3}$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{total}$ — Γ_{74}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
2.1±0.5 OUR AVERAGE						
2.0±0.6±0.2			[213] AUBERT	02	BABR	$e^+e^- \to \Upsilon(4S)$
$2.5^{+1.1}_{-0.9}\pm0.2$			[213] AVERY	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 32	90		[214] ACCIARRI	97C	L3	
< 5.8	90		BISHAI	96	CLE2	$e^+e^- \to \Upsilon(4S)$
<690	90	1	[215] ALEXANDER	95	CLE2	Sup. by BISHAI 96

[213] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[214] ACCIARRI 97C assumes B^0 production fraction (39.5 ± 4.0%) and B_s (12.0 ± 3.0%).
[215] Assumes equal production of B^+B^- and $B^0\overline{B}^0$ on $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$ — Γ_{75}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<1.2\times10^{-3}$	90	[216] ACCIARRI	97C	L3

[216] ACCIARRI 97C assumes B^0 production fraction (39.5 ± 4.0%) and B_s (12.0 ± 3.0%).

$\Gamma(J/\psi(1S)\rho^0)/\Gamma_{total}$ — Γ_{76}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.5\times10^{-4}$	90	BISHAI	96	CLE2	$e^+e^- \to \Upsilon(4S)$

$\Gamma(J/\psi(1S)\omega)/\Gamma_{total}$ — Γ_{77}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.7\times10^{-4}$	90	BISHAI	96	CLE2	$e^+e^- \to \Upsilon(4S)$

$\Gamma(J/\psi(1S)K^0\pi^+\pi^-)/\Gamma_{total}$ — Γ_{78}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
10.3±3.3±1.5	[217] AFFOLDER	02B	CDF	$p\overline{p}$ 1.8 TeV

[217] Uses $B^0 \to J/\psi(1S)K^0_S$ decay as a reference and B($B^0 \to J/\psi(1S)K^0$)= 8.3×10^{-4}.

$\Gamma(J/\psi(1S)K^0\rho^0)/\Gamma_{total}$ — Γ_{79}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
5.4±2.9±0.9	[218] AFFOLDER	02B	CDF	$p\overline{p}$ 1.8 TeV

[218] Uses $B^0 \to J/\psi(1S)K^0_S$ decay as a reference and B($B^0 \to J/\psi(1S)K^0$)= 8.3×10^{-4}.

$\Gamma(J/\psi(1S)K^*(892)^+\pi^-)/\Gamma_{total}$ — Γ_{80}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
7.7±4.1±1.3	[219] AFFOLDER	02B	CDF	$p\overline{p}$ 1.8 TeV

[219] Uses $B^0 \to J/\psi(1S)K^0_S$ decay as a reference and B($B^0 \to J/\psi(1S)K^0$)= 8.3×10^{-4}.

$\Gamma(J/\psi(1S)K^*(892)^0\pi^+\pi^-)/\Gamma_{total}$ — Γ_{81}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
6.6±1.9±1.1	[220] AFFOLDER	02B	CDF	$p\overline{p}$ 1.8 TeV

[220] Uses $B^0 \to J/\psi(1S)K^*(892)^0$ decay as a reference and B($B^0 \to J/\psi(1S)K^0$)= 12.4×10^{-4}.

$\Gamma(\psi(2S)K^0)/\Gamma_{total}$ — Γ_{82}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
5.7±1.0 OUR AVERAGE					
6.9±1.1±1.1		[221] AUBERT	02	BABR	$e^+e^- \to \Upsilon(4S)$
5.0±1.1±0.6		[221] RICHICHI	01	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 8	90	[222] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
<15	90	[222] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
<28	90	[222] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$

[221] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[222] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)K^0)/\Gamma(J/\psi(1S)K^0)$ — Γ_{82}/Γ_{69}

VALUE	DOCUMENT ID		TECN	COMMENT
0.82±0.13±0.12	[223] AUBERT	02	BABR	$e^+e^- \to \Upsilon(4S)$

[223] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)K^+\pi^-)/\Gamma_{total}$ — Γ_{83}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.001	90	[224] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$

[224] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S) K^*(892)^0)/\Gamma_{total}$ — Γ_{84}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
8.0±1.3 OUR AVERAGE					
7.6±1.1±1.0		[225] RICHICHI	01	CLE2	$e^+ e^- \to \Upsilon(4S)$
9.0±2.2±0.9		[226] ABE	98O	CDF	$p\overline{p}$ 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<19	90	[227] ALAM	94	CLE2	Repl. by RICHICHI 01
14 ±8 ±4		[227] BORTOLETTO	92	CLEO	$e^+ e^- \to \Upsilon(4S)$
<23	90	[227] ALBRECHT	90J	ARG	$e^+ e^- \to \Upsilon(4S)$

[225] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[226] ABE 98O reports $[B(B^0 \to \psi(2S) K^*(892)^0)]/[B(B^+ \to J/\psi(1S) K^+)] = 0.908 \pm 0.194 \pm 0.10$. We multiply by our best value $B(B^+ \to J/\psi(1S) K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[227] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c0}(1P) K^0)/\Gamma_{total}$ — Γ_{85}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<5.0 \times 10^{-4}$	90	[228] EDWARDS	01	CLE2	$e^+ e^- \to \Upsilon(4S)$

[228] EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \to \gamma \eta_c)$ in those modes have been accounted for.

$\Gamma(\chi_{c1}(1P) K^0)/\Gamma_{total}$ — Γ_{86}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
$4.0^{+1.2}_{-1.0}$ OUR AVERAGE					
4.7±1.5±0.5		[229] AUBERT	02	BABR	$e^+ e^- \to \Upsilon(4S)$
$3.4^{+1.7}_{-1.2}\pm0.3$		[230] AVERY	00	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<27	90	[231] ALAM	94	CLE2	$e^+ e^- \to \Upsilon(4S)$

[229] AUBERT 02 reports $5.4 \pm 1.4 \pm 1.1$ for $B(\chi_{c1}(1P) \to \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \to \gamma J/\psi(1S)) = (31.6 \pm 3.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[230] AVERY 00 reports $3.9^{+1.9}_{-1.3} \pm 0.4$ for $B(\chi_{c1}(1P) \to \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \to \gamma J/\psi(1S)) = (31.6 \pm 3.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[231] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P) K^*(892)^0)/\Gamma_{total}$ — Γ_{87}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
4.1±1.4±0.4		[232] AUBERT	02	BABR	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<21	90	[233] ALAM	94	CLE2	$e^+ e^- \to \Upsilon(4S)$

[232] AUBERT 02 reports $4.8 \pm 1.4 \pm 0.9$ for $B(\chi_{c1}(1P) \to \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \to \gamma J/\psi(1S)) = (31.6 \pm 3.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[233] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)K^0)/\Gamma(J/\psi(1S)K^0)$ — Γ_{86}/Γ_{69}

VALUE	DOCUMENT ID		TECN	COMMENT
0.57±0.17±0.06	[234] AUBERT	02	BABR	$e^+e^- \to \Upsilon(4S)$

[234] AUBERT 02 reports $0.66 \pm 0.11 \pm 0.17$ for $B(\chi_{c1}(1P) \to \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \to \gamma J/\psi(1S)) = (31.6 \pm 3.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)K^*(892)^0)/\Gamma(\chi_{c1}(1P)K^0)$ — Γ_{87}/Γ_{86}

VALUE	DOCUMENT ID		TECN	COMMENT
0.89±0.34±0.17	[235] AUBERT	02	BABR	$e^+e^- \to \Upsilon(4S)$

[235] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+\pi^-)/\Gamma_{total}$ — Γ_{88}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
1.74±0.15 OUR AVERAGE					
$1.93^{+0.34}_{-0.32}{}^{+0.15}_{-0.06}$		[236] ABE	01H	BELL	$e^+e^- \to \Upsilon(4S)$
$1.67 \pm 0.16 \pm 0.13$		[236] AUBERT	01E	BABR	$e^+e^- \to \Upsilon(4S)$
$1.72^{+0.25}_{-0.24} \pm 0.12$		[236] CRONIN-HEN...	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6.6	90	[237] ABE	00C	SLD	$e^+e^- \to Z$
$1.5 \; {}^{+0.5}_{-0.4} \; \pm 0.14$		GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00
$2.4 \; {}^{+1.7}_{-1.1} \; \pm 0.2$		[238] ADAM	96D	DLPH	$e^+e^- \to Z$
<1.7	90	ASNER	96	CLE2	Sup. by ADAM 96D
<3.0	90	[239] BUSKULIC	96V	ALEP	$e^+e^- \to Z$
<9	90	[240] ABREU	95N	DLPH	Sup. by ADAM 96D
<8.1	90	[241] AKERS	94L	OPAL	$e^+e^- \to Z$
<2.6	90	[242] BATTLE	93	CLE2	$e^+e^- \to \Upsilon(4S)$
<18	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$
<9	90	[243] AVERY	89B	CLEO	$e^+e^- \to \Upsilon(4S)$
<32	90	AVERY	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[236] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[237] ABE 00C assumes $B(Z \to b\overline{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.

[238] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

[239] BUSKULIC 96V assumes PDG 96 production fractions for B^0, B^+, B_s, b baryons.

[240] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s^0 decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.
[241] Assumes B($Z \to b\overline{b}$) = 0.217 and B_d^0 (B_s^0) fraction 39.5% (12%).
[242] BATTLE 93 assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.
[243] Assumes the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$.

$\Gamma(K^+\pi^-)/\Gamma(K^0\pi^0)$ — Γ_{88}/Γ_{89}

VALUE	DOCUMENT ID	TECN	COMMENT
$\mathbf{1.20^{+0.50+0.22}_{-0.58-0.32}}$	[244] ABE 01H	BELL	$e^+e^- \to \Upsilon(4S)$

[244] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$[\Gamma(K^+\pi^-)+\Gamma(\pi^+\pi^-)]/\Gamma_{total}$ — $(\Gamma_{88}+\Gamma_{133})/\Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.9±0.6 OUR AVERAGE				
$2.8^{+1.5}_{-1.0}\pm2.0$		[245] ADAM 96D	DLPH	$e^+e^- \to Z$
$1.8^{+0.6+0.3}_{-0.5-0.4}$	17.2	ASNER 96	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.4^{+0.8}_{-0.7}\pm0.2$		[246] BATTLE 93	CLE2	$e^+e^- \to \Upsilon(4S)$

[245] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.
[246] BATTLE 93 assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(K^0\pi^0)/\Gamma_{total}$ — Γ_{89}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$\mathbf{1.07^{+0.27}_{-0.25}}$ **OUR AVERAGE**				
$1.60^{+0.72+0.25}_{-0.59-0.27}$		[247] ABE 01H	BELL	$e^+e^- \to \Upsilon(4S)$
$0.82^{+0.31}_{-0.27}\pm0.12$		[247] AUBERT 01E	BABR	$e^+e^- \to \Upsilon(4S)$
$1.46^{+0.59+0.24}_{-0.51-0.33}$		[247] CRONIN-HEN...00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<4.1	90	GODANG 98	CLE2	Repl. by CRONIN-HENNESSY 00
<4.0	90	ASNER 96	CLE2	Rep. by GODANG 98

[247] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' K^0)/\Gamma_{total}$ Γ_{90}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$\mathbf{5.8^{+1.4}_{-1.3}}$ **OUR AVERAGE** Error includes scale factor of 1.5. See the ideogram below.			
$5.5^{+1.9}_{-1.6}\pm0.8$	[248] ABE 01M	BELL	$e^+e^- \to \Upsilon(4S)$
$4.2^{+1.3}_{-1.1}\pm0.4$	[248] AUBERT 01G	BABR	$e^+e^- \to \Upsilon(4S)$
$8.9^{+1.8}_{-1.6}\pm0.9$	[248] RICHICHI 00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$4.7^{+2.7}_{-2.0}\pm0.9$	BEHRENS 98	CLE2	Repl. by RICHICHI 00

[248] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.



$\Gamma\left(\eta' K^0\right)/\Gamma_{total}$ (units 10^{-5})

$\Gamma(\eta' K^*(892)^0)/\Gamma_{total}$ Γ_{91}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.4	90	[249] RICHICHI 00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<3.9	90	BEHRENS 98	CLE2	Repl. by RICHICHI 00

[249] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K^*(892)^0)/\Gamma_{total}$ Γ_{92}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
$1.38^{+0.55}_{-0.46}\pm0.16$		[250] RICHICHI	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.0	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[250] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K^0)/\Gamma_{total}$ Γ_{93}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 9.3	90	[251] RICHICHI	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<33	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[251] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega K^0)/\Gamma_{total}$ Γ_{94}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<1.3	90	[252] AUBERT	01G	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.1	90	[252] JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$
<5.7	90	[252] BERGFELD	98	CLE2	Repl. by JESSOP 00

[252] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^0_S X^0(\text{Familon}))/\Gamma_{total}$ Γ_{95}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<5.3	90	[253] AMMAR	01B	CLE2	$e^+e^- \to \Upsilon(4S)$

[253] AMMAR 01B searched for the two-body decay of the B meson to a massless neutral feebly-interacting particle X^0 such as the familon, the Nambu-Goldstone boson associated with a sponateously broken global family symmetry.

$\Gamma(\omega K^*(892)^0)/\Gamma_{total}$ Γ_{96}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<2.3\times10^{-5}$	90	[254] BERGFELD	98	CLE2

[254] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+K^-)/\Gamma_{total}$ Γ_{97}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 1.9	90	[255] CRONIN-HEN...	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 2.7	90	[255] ABE	01H	BELL	$e^+e^- \to \Upsilon(4S)$
< 2.5	90	[255] AUBERT	01E	BABR	$e^+e^- \to \Upsilon(4S)$
< 66	90	[256] ABE	00C	SLD	$e^+e^- \to Z$
< 4.3	90	GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00
< 46		[257] ADAM	96D	DLPH	$e^+e^- \to Z$
< 4	90	ASNER	96	CLE2	Repl. by GODANG 98
< 18	90	[258] BUSKULIC	96V	ALEP	$e^+e^- \to Z$
<120	90	[259] ABREU	95N	DLPH	Sup. by ADAM 96D
< 7	90	[260] BATTLE	93	CLE2	$e^+e^- \to \Upsilon(4S)$

[255] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[256] ABE 00C assumes B($Z \rightarrow b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

[257] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

[258] BUSKULIC 96V assumes PDG 96 production fractions for B^0, B^+, B_s, b baryons.

[259] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s^0 decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

[260] BATTLE 93 assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(K^0\overline{K}^0)/\Gamma_{total}$ — Γ_{98}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.7 × 10^−5	90	GODANG	98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^+\rho^-)/\Gamma_{total}$ — Γ_{99}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<3.2 × 10^−5	90	[261] JESSOP	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.5 × 10^−5	90	ASNER	96	CLE2	Repl. by JESSOP 00

[261] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^0\pi^+\pi^-)/\Gamma_{total}$ — Γ_{100}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.4 × 10^−4	90	ALBRECHT	91E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^0\rho^0)/\Gamma_{total}$ — Γ_{101}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<3.9 × 10^−5	90	ASNER	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.2 × 10^−4	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<5.0 × 10^−4	90	[262] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<0.064	90	[263] AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[262] AVERY 89B reports $< 5.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

[263] AVERY 87 reports < 0.08 assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^0 f_0(980))/\Gamma_{total}$ — Γ_{102}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<3.6 × 10^−4	90	[264] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[264] AVERY 89B reports $< 4.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+\pi^-)/\Gamma_{total}$ Γ_{103}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<7.2 × 10^−5	90	ASNER	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<3.8 × 10^−4	90	[265] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6.2 \times 10^{-4}$	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<5.6 \times 10^{-4}$	90	[266] AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[265] AVERY 89B reports $< 4.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[266] AVERY 87 reports $< 7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^0\pi^0)/\Gamma_{total}$ Γ_{104}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<3.6 × 10^−6	90	[190] JESSOP	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.8 \times 10^{-5}$	90	ASNER	96	CLE2	Repl. by JESSOP 00

$\Gamma(K_2^*(1430)^+\pi^-)/\Gamma_{total}$ Γ_{105}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<2.6 × 10^−3	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^0K^+K^-)/\Gamma_{total}$ Γ_{106}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.3 × 10^−3	90	ALBRECHT	91E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^0\phi)/\Gamma_{total}$ Γ_{107}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{8.1^{+3.1}_{-2.5}\pm0.8}$		[267] AUBERT	01D	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 12.3	90	[267] BRIERE	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 31	90	[267] BERGFELD	98	CLE2	
< 88	90	ASNER	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 720	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 420	90	[268] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<1000	90	[269] AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[267] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[268] AVERY 89B reports $< 4.9 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[269] AVERY 87 reports $< 1.3 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^-\pi^+\pi^+\pi^-)/\Gamma_{total}$ Γ_{108}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<2.3 × 10^−4	90	[270] ADAM	96D	DLPH	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.1 \times 10^{-4}$	90	[271] ABREU	95N	DLPH	Sup. by ADAM 96D

[270] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.
[271] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B^0_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

$\Gamma(K^*(892)^0\pi^+\pi^-)/\Gamma_{\text{total}}$ — Γ_{109}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.4\times10^{-3}$	90	ALBRECHT	91E	ARG	$e^+e^-\to\Upsilon(4S)$

$\Gamma(K^*(892)^0\rho^0)/\Gamma_{\text{total}}$ — Γ_{110}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.4\times10^{-5}$	90	[272] GODANG	02	CLE2	$e^+e^-\to\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.86\times10^{-4}$	90	[273] ABE	00C	SLD	$e^+e^-\to Z$
$<4.6\times10^{-4}$	90	ALBRECHT	91B	ARG	$e^+e^-\to\Upsilon(4S)$
$<5.8\times10^{-4}$	90	[274] AVERY	89B	CLEO	$e^+e^-\to\Upsilon(4S)$
$<9.6\times10^{-4}$	90	[275] AVERY	87	CLEO	$e^+e^-\to\Upsilon(4S)$

[272] Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 2.4×10^{-5}.
[273] ABE 00C assumes $B(Z\to b\overline{b})=(21.7\pm0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.
[274] AVERY 89B reports $<6.7\times10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[275] AVERY 87 reports $<1.2\times10^{-3}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^0 f_0(980))/\Gamma_{\text{total}}$ — Γ_{111}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.7\times10^{-4}$	90	[276] AVERY	89B	CLEO	$e^+e^-\to\Upsilon(4S)$

[276] AVERY 89B reports $<2.0\times10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_1(1400)^+\pi^-)/\Gamma_{\text{total}}$ — Γ_{112}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1\times10^{-3}$	90	ALBRECHT	91B	ARG	$e^+e^-\to\Upsilon(4S)$

$\Gamma(K^- a_1(1260)^+)/\Gamma_{\text{total}}$ — Γ_{113}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.3\times10^{-4}$	90	[277] ADAM	96D	DLPH	$e^+e^-\to Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<3.9\times10^{-4}$	90	[278] ABREU	95N	DLPH	Sup. by ADAM 96D

[277] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.
[278] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B^0_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

$\Gamma(K^*(892)^0 K^+ K^-)/\Gamma_{total}$ — Γ_{114}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.1 \times 10^{-4}$	90	ALBRECHT	91E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^0 \phi)/\Gamma_{total}$ — Γ_{115}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$9.5^{+2.4}_{-2.0}$ OUR AVERAGE					
$8.7^{+2.5}_{-2.1}\pm1.1$		[279] AUBERT	01D	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$11.5^{+4.5}_{-3.7}{}^{+1.8}_{-1.7}$		[279] BRIERE	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<384	90	[280] ABE	00C	SLD	$e^+e^- \rightarrow Z$
<21	90	[279] BERGFELD	98	CLE2	
<43	90	ASNER	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<320	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<380	90	[281] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<380	90	[282] AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[279] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[280] ABE 00C assumes B($Z \rightarrow b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

[281] AVERY 89B reports $< 4.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

[282] AVERY 87 reports $< 4.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\overline{K}^*(892)^0 K^*(892)^0)/\Gamma_{total}$ — Γ_{116}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.2 \times 10^{-5}$	90	[283] GODANG	02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<4.69 \times 10^{-4}$	90	[284] ABE	00C	SLD	$e^+e^- \rightarrow Z$

[283] Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 1.9×10^{-5}.

[284] ABE 00C assumes B($Z \rightarrow b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

$\Gamma(K^*(892)^0 K^*(892)^0)/\Gamma_{total}$ — Γ_{117}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.7 \times 10^{-5}$	90	[285] GODANG	02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[285] Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 2.9×10^{-5}.

$\Gamma(K^*(892)^+ K^*(892)^-)/\Gamma_{total}$ — Γ_{118}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.41 \times 10^{-4}$	90	[286] GODANG	02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[286] Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 8.9×10^{-5}.

$\Gamma(K_1(1400)^0 \rho^0)/\Gamma_{total}$ Γ_{119}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.0 \times 10^{-3}$	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_1(1400)^0 \phi)/\Gamma_{total}$ Γ_{120}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<5.0 \times 10^{-3}$	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2^*(1430)^0 \rho^0)/\Gamma_{total}$ Γ_{121}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1 \times 10^{-3}$	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2^*(1430)^0 \phi)/\Gamma_{total}$ Γ_{122}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.4 \times 10^{-3}$	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^0 \gamma)/\Gamma_{total}$ Γ_{123}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
4.3 ±0.4 OUR AVERAGE						
$4.23 \pm 0.40 \pm 0.22$			[287] AUBERT	02C	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$4.55^{+0.72}_{-0.68} \pm 0.34$			[288] COAN	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<$ 21	90		[289] ADAM	96D	DLPH	$e^+e^- \rightarrow Z$
$4.0 \pm 1.7 \pm 0.8$		8	[290] AMMAR	93	CLE2	Repl. by COAN 00
$<$ 42	90		ALBRECHT	89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 24	90		[291] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<$210	90		AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[287] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[288] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. No evidence for a nonresonant $K\pi\gamma$ contamination was seen; the central value assumes no contamination.

[289] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

[290] AMMAR 93 observed 6.6 ± 2.8 events above background.

[291] AVERY 89B reports $< 2.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_1(1270)^0 \gamma)/\Gamma_{total}$ Γ_{124}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0070	90	[292] ALBRECHT	89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[292] ALBRECHT 89G reports < 0.0078 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_1(1400)^0\gamma)/\Gamma_{\text{total}}$ — Γ_{125}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0043	90	[293] ALBRECHT 89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[293] ALBRECHT 89G reports < 0.0048 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_2^*(1430)^0\gamma)/\Gamma_{\text{total}}$ — Γ_{126}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.0 \times 10^{-4}$	90	[294] ALBRECHT 89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[294] ALBRECHT 89G reports $< 4.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(1680)^0\gamma)/\Gamma_{\text{total}}$ — Γ_{127}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0020	90	[295] ALBRECHT 89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[295] ALBRECHT 89G reports < 0.0022 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_3^*(1780)^0\gamma)/\Gamma_{\text{total}}$ — Γ_{128}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	90	[296] ALBRECHT 89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[296] ALBRECHT 89G reports < 0.011 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_4^*(2045)^0\gamma)/\Gamma_{\text{total}}$ — Γ_{129}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0043	90	[297] ALBRECHT 89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[297] ALBRECHT 89G reports < 0.0048 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\rho^0\gamma)/\Gamma_{\text{total}}$ — Γ_{130}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-5}$	90	[298] COAN 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[298] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega\gamma)/\Gamma_{\text{total}}$ — Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.92 \times 10^{-5}$	90	[299] COAN 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[299] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi\gamma)/\Gamma_{\text{total}}$ — Γ_{132}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.33 \times 10^{-5}$	90	[300] COAN 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[300] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$ — Γ_{133}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
4.4±0.9 OUR AVERAGE						
$5.6^{+2.3}_{-2.0}{}^{+0.4}_{-0.5}$			[301] ABE	01H	BELL	$e^+e^- \to \Upsilon(4S)$
$4.1\pm1.0\pm0.7$			[301] AUBERT	01E	BABR	$e^+e^- \to \Upsilon(4S)$
$4.3^{+1.6}_{-1.4}\pm0.5$			[301] CRONIN-HEN...	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<$ 67	90		[302] ABE	00C	SLD	$e^+e^- \to Z$
$<$ 15	90		GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00
$<$ 45	90		[303] ADAM	96D	DLPH	$e^+e^- \to Z$
$<$ 20	90		ASNER	96	CLE2	Repl. by GODANG 98
$<$ 41	90		[304] BUSKULIC	96V	ALEP	$e^+e^- \to Z$
$<$ 55	90		[305] ABREU	95N	DLPH	Sup. by ADAM 96D
$<$ 47	90		[306] AKERS	94L	OPAL	$e^+e^- \to Z$
$<$ 29	90		[307] BATTLE	93	CLE2	$e^+e^- \to \Upsilon(4S)$
$<$130	90		[307] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$
$<$ 77	90		[308] BORTOLETTO	89	CLEO	$e^+e^- \to \Upsilon(4S)$
$<$260	90		[308] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$
$<$500	90	4	GILES	84	CLEO	$e^+e^- \to \Upsilon(4S)$

[301] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[302] ABE 00C assumes B($Z \to b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

[303] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

[304] BUSKULIC 96V assumes PDG 96 production fractions for B^0, B^+, B_s, b baryons.

[305] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

[306] Assumes B($Z \to b\overline{b}$) = 0.217 and B^0_d (B^0_s) fraction 39.5% (12%).

[307] Assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

[308] Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^-)/\Gamma(K^+\pi^-)$ — Γ_{133}/Γ_{88}

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.29^{+0.13}_{-0.12}{}^{+0.01}_{-0.02}}$	[309] ABE	01H	BELL	$e^+e^- \to \Upsilon(4S)$

[309] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^0\pi^0)/\Gamma_{total}$ — Γ_{134}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<5.7 \times 10^{-6}}$	90	[310] ASNER	02	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<9.3 \times 10^{-6}$	90	GODANG	98	CLE2	Repl. by ASNER 02
$<0.91 \times 10^{-5}$	90	ASNER	96	CLE2	Repl. by GODANG 98
$<6.0 \times 10^{-5}$	90	[311] ACCIARRI	95H	L3	$e^+e^- \to Z$

[310] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[311] ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0$%.

$\Gamma(\eta\pi^0)/\Gamma_{total}$ — Γ_{135}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-6}$	90	[312] RICHICHI 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<8 \times 10^{-6}$	90	BEHRENS 98	CLE2	Repl. by RICHICHI 00
$<2.5 \times 10^{-4}$	90	[313] ACCIARRI 95H	L3	$e^+e^- \rightarrow Z$
$<1.8 \times 10^{-3}$	90	[314] ALBRECHT 90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[312] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[313] ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

[314] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\eta\eta)/\Gamma_{total}$ — Γ_{136}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-5}$	90	BEHRENS 98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<4.1 \times 10^{-4}$	90	[315] ACCIARRI 95H	L3	$e^+e^- \rightarrow Z$

[315] ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

$\Gamma(\eta'\pi^0)/\Gamma_{total}$ — Γ_{137}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.7 \times 10^{-6}$	90	[316] RICHICHI 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.1 \times 10^{-5}$	90	BEHRENS 98	CLE2	Repl. by RICHICHI 00

[316] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta'\eta')/\Gamma_{total}$ — Γ_{138}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.7 \times 10^{-5}$	90	BEHRENS 98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\eta'\eta)/\Gamma_{total}$ — Γ_{139}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.7 \times 10^{-5}$	90	BEHRENS 98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\eta'\rho^0)/\Gamma_{total}$ — Γ_{140}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-5}$	90	[317] RICHICHI 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<2.3 \times 10^{-5}$	90	BEHRENS 98	CLE2	Repl. by RICHICHI 00

[317] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta\rho^0)/\Gamma_{total}$ — Γ_{141}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-5}$	90	[318] RICHICHI 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.3 \times 10^{-5}$	90	BEHRENS 98	CLE2	Repl. by RICHICHI 00

[318] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega\eta)/\Gamma_{total}$ — Γ_{142}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<1.2 \times 10^{-5}$	90	[319] BERGFELD	98	CLE2

[319] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega\eta')/\Gamma_{total}$ — Γ_{143}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<6.0 \times 10^{-5}$	90	[320] BERGFELD	98	CLE2

[320] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega\rho^0)/\Gamma_{total}$ — Γ_{144}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<1.1 \times 10^{-5}$	90	[321] BERGFELD	98	CLE2

[321] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega\omega)/\Gamma_{total}$ — Γ_{145}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<1.9 \times 10^{-5}$	90	[322] BERGFELD	98	CLE2

[322] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi\pi^0)/\Gamma_{total}$ — Γ_{146}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<0.5 \times 10^{-5}$	90	[323] BERGFELD	98	CLE2

[323] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi\eta)/\Gamma_{total}$ — Γ_{147}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<0.9 \times 10^{-5}$	90	[324] BERGFELD	98	CLE2

[324] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi\eta')/\Gamma_{total}$ — Γ_{148}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<3.1 \times 10^{-5}$	90	[325] BERGFELD	98	CLE2

[325] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi\rho^0)/\Gamma_{total}$ — Γ_{149}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.3 \times 10^{-5}$	90	[326] BERGFELD	98	CLE2	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.56 \times 10^{-4}$	90	[327] ABE	00C	SLD	$e^+e^- \rightarrow Z$

[326] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[327] ABE 00C assumes B($Z \rightarrow b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

$\Gamma(\phi\omega)/\Gamma_{total}$ — Γ_{150}/Γ

VALUE	CL%	DOCUMENT ID		TECN
$<2.1 \times 10^{-5}$	90	[328] BERGFELD	98	CLE2

[328] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi\phi)/\Gamma_{total}$ — Γ_{151}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.2 \times 10^{-5}$	90	[329] BERGFELD	98	CLE2	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<3.21 \times 10^{-4}$	90	[330] ABE	00C	SLD	$e^+e^- \to Z$
$<3.9 \times 10^{-5}$	90	ASNER	96	CLE2	$e^+e^- \to \Upsilon(4S)$

[329] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[330] ABE 00C assumes B($Z \to b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ — Γ_{152}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.2 \times 10^{-4}$	90	[331] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[331] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^0\pi^0)/\Gamma_{total}$ — Γ_{153}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<5.5 \times 10^{-6}$	90	[186] JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.4 \times 10^{-5}$	90	ASNER	96	CLE2	Repl. by JESSOP 00
$<4.0 \times 10^{-4}$	90	[332] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[332] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^{\mp}\pi^{\pm})/\Gamma_{total}$ — Γ_{154}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
$2.76^{+0.84}_{-0.74} \pm 0.42$		[333] JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<8.8	90	ASNER	96	CLE2	Repl. by JESSOP 00
<52	90	[334] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$
<520	90	[335] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[333] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[334] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.
[335] BEBEK 87 reports $< 6.1 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{total}$ — Γ_{155}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.3 \times 10^{-4}$	90	[336] ADAM	96D	DLPH	$e^+e^- \to Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.8 \times 10^{-4}$	90	[337] ABREU	95N	DLPH	Sup. by ADAM 96D
$<6.7 \times 10^{-4}$	90	[338] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[336] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

[337] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

[338] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^0\rho^0)/\Gamma_{total}$ — Γ_{156}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.8 \times 10^{-5}$	90	[339] GODANG	02	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.36 \times 10^{-4}$	90	[340] ABE	00C	SLD	$e^+e^- \to Z$
$<2.8 \times 10^{-4}$	90	[341] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$
$<2.9 \times 10^{-4}$	90	[342] BORTOLETTO	89	CLEO	$e^+e^- \to \Upsilon(4S)$
$<4.3 \times 10^{-4}$	90	[342] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[339] Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 1.4×10^{-5}.

[340] ABE 00C assumes B($Z \to b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

[341] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

[342] Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(a_1(1260)^{\mp}\pi^{\pm})/\Gamma_{total}$ — Γ_{157}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<4.9 \times 10^{-4}$	90	[343] BORTOLETTO	89	CLEO	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6.3 \times 10^{-4}$	90	[344] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$
$<1.0 \times 10^{-3}$	90	[343] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[343] Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

[344] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_2(1320)^{\mp}\pi^{\pm})/\Gamma_{total}$ — Γ_{158}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.0 \times 10^{-4}$	90	[345] BORTOLETTO	89	CLEO	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.4 \times 10^{-3}$	90	[345] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[345] Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{total}$ — Γ_{159}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.1 \times 10^{-3}$	90	[346] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[346] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^+\rho^-)/\Gamma_{\text{total}}$ — Γ_{160}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.2\times10^{-3}$	90	[347] ALBRECHT	90B	ARG	$e^+e^-\to\Upsilon(4S)$

[347] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^0\pi^0)/\Gamma_{\text{total}}$ — Γ_{161}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1\times10^{-3}$	90	[348] ALBRECHT	90B	ARG	$e^+e^-\to\Upsilon(4S)$

[348] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ — Γ_{162}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3\times10^{-6}$	90	[349] AUBERT	01G	BABR	$e^+e^-\to\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<5.5\times10^{-6}$	90	[349] JESSOP	00	CLE2	$e^+e^-\to\Upsilon(4S)$
$<1.4\times10^{-5}$	90	[349] BERGFELD	98	CLE2	Repl. by JESSOP 00
$<4.6\times10^{-4}$	90	[350] ALBRECHT	90B	ARG	$e^+e^-\to\Upsilon(4S)$

[349] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[350] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ — Γ_{163}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<9.0\times10^{-3}$	90	[351] ALBRECHT	90B	ARG	$e^+e^-\to\Upsilon(4S)$

[351] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+\rho^-)/\Gamma_{\text{total}}$ — Γ_{164}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.4\times10^{-3}$	90	[352] ALBRECHT	90B	ARG	$e^+e^-\to\Upsilon(4S)$

[352] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^0\rho^0)/\Gamma_{\text{total}}$ — Γ_{165}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.4\times10^{-3}$	90	[353] ALBRECHT	90B	ARG	$e^+e^-\to\Upsilon(4S)$

[353] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-\pi^-)/\Gamma_{\text{total}}$ — Γ_{166}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.0\times10^{-3}$	90	[354] ALBRECHT	90B	ARG	$e^+e^-\to\Upsilon(4S)$

[354] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+a_1(1260)^-)/\Gamma_{\text{total}}$ — Γ_{167}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.8\times10^{-3}$	90	[355] BORTOLETTO	89	CLEO	$e^+e^-\to\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6.0\times10^{-3}$	90	[356] ALBRECHT	90B	ARG	$e^+e^-\to\Upsilon(4S)$

[355] BORTOLETTO 89 reports $<3.2\times10^{-3}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[356] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-\pi^-\pi^0)/\Gamma_{total}$ — Γ_{168}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1 \times 10^{-2}$	90	[357] ALBRECHT	90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[357] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(p\overline{p})/\Gamma_{total}$ — Γ_{169}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.0 \times 10^{-6}$	90	[358] COAN	99	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.8 \times 10^{-5}$	90	[359] BUSKULIC	96V	ALEP	$e^+e^- \rightarrow Z$
$<3.5 \times 10^{-4}$	90	[360] ABREU	95N	DLPH	Sup. by ADAM 96D
$<3.4 \times 10^{-5}$	90	[361] BORTOLETTO	89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<1.2 \times 10^{-4}$	90	[362] ALBRECHT	88F	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<1.7 \times 10^{-4}$	90	[361] BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[358] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[359] BUSKULIC 96V assumes PDG 96 production fractions for B^0, B^+, B_s, b baryons.

[360] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

[361] Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

[362] ALBRECHT 88F reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(p\overline{p}\pi^+\pi^-)/\Gamma_{total}$ — Γ_{170}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2.5	90	[363] BEBEK	89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<9.5	90	[364] ABREU	95N	DLPH	Sup. by ADAM 96D
$5.4\pm1.8\pm2.0$		[365] ALBRECHT	88F	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[363] BEBEK 89 reports $< 2.9 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

[364] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

[365] ALBRECHT 88F reports $6.0 \pm 2.0 \pm 2.2$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(p\overline{\Lambda}\pi^-)/\Gamma_{total}$ — Γ_{171}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.3 \times 10^{-5}$	90	[366] COAN	99	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.8 \times 10^{-4}$	90	[367] ALBRECHT	88F	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[366] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[367] ALBRECHT 88F reports $< 2.0 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\overline{\Lambda}\Lambda)/\Gamma_{total}$ — Γ_{172}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.9 \times 10^{-6}$	90	[368] COAN	99	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[368] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\Delta^0 \overline{\Delta}^0)/\Gamma_{total}$ — Γ_{173}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0015	90	[369] BORTOLETTO	89	CLEO	$e^+ e^- \to \Upsilon(4S)$

[369] BORTOLETTO 89 reports < 0.0018 assuming $\Upsilon(4S)$ decays 43% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(\Delta^{++} \Delta^{--})/\Gamma_{total}$ — Γ_{174}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1 \times 10^{-4}$	90	[370] BORTOLETTO	89	CLEO	$e^+ e^- \to \Upsilon(4S)$

[370] BORTOLETTO 89 reports $< 1.3 \times 10^{-4}$ assuming $\Upsilon(4S)$ decays 43% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(\overline{\Sigma}_c^{--} \Delta^{++})/\Gamma_{total}$ — Γ_{175}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0010	90	[371] PROCARIO	94	CLE2	$e^+ e^- \to \Upsilon(4S)$

[371] PROCARIO 94 reports < 0.0012 for $B(\Lambda_c^+ \to p K^- \pi^+) = 0.043$. We rescale to our best value $B(\Lambda_c^+ \to p K^- \pi^+) = 0.050$.

$\Gamma(\overline{\Lambda}_c^- p \pi^+ \pi^-)/\Gamma_{total}$ — Γ_{176}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
$1.33^{+0.46}_{-0.42} \pm 0.37$	[372] FU	97	CLE2	$e^+ e^- \to \Upsilon(4S)$

[372] FU 97 uses PDG 96 values of Λ_c branching fraction.

$\Gamma(\overline{\Lambda}_c^- p)/\Gamma_{total}$ — Γ_{177}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.1 \times 10^{-4}$	90	[373] FU	97	CLE2	$e^+ e^- \to \Upsilon(4S)$

[373] FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\overline{\Lambda}_c^- p \pi^0)/\Gamma_{total}$ — Γ_{178}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<5.9 \times 10^{-4}$	90	[374] FU	97	CLE2	$e^+ e^- \to \Upsilon(4S)$

[374] FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\overline{\Lambda}_c^- p \pi^+ \pi^- \pi^0)/\Gamma_{total}$ — Γ_{179}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<5.07 \times 10^{-3}$	90	[375] FU	97	CLE2	$e^+ e^- \to \Upsilon(4S)$

[375] FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\overline{\Lambda}_c^- p \pi^+ \pi^- \pi^+ \pi^-)/\Gamma_{total}$ — Γ_{180}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.74 \times 10^{-3}$	90	[376] FU	97	CLE2	$e^+ e^- \to \Upsilon(4S)$

[376] FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\gamma\gamma)/\Gamma_{total}$ Γ_{181}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.7 \times 10^{-6}$	90	[377] AUBERT	01I	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<3.9 \times 10^{-5}$	90	[378] ACCIARRI	95I	L3	$e^+e^- \rightarrow Z$

[377] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[378] ACCIARRI 95I assumes f_{B^0} = 39.5 ± 4.0 and f_{B_s} = 12.0 ± 3.0%.

$\Gamma(e^+e^-)/\Gamma_{total}$ Γ_{182}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<8.3 \times 10^{-7}$	90	[379] BERGFELD	00B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.4 \times 10^{-5}$	90	[380] ACCIARRI	97B	L3	$e^+e^- \rightarrow Z$
$<5.9 \times 10^{-6}$	90	AMMAR	94	CLE2	Repl. by BERGFELD 00B
$<2.6 \times 10^{-5}$	90	[381] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<7.6 \times 10^{-5}$	90	[382] ALBRECHT	87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<6.4 \times 10^{-5}$	90	[383] AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<3 \times 10^{-4}$	90	GILES	84	CLEO	Repl. by AVERY 87

[379] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[380] ACCIARRI 97B assume PDG 96 production fractions for B^+, B^0, B_s, and Λ_b.
[381] AVERY 89B reports $< 3 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[382] ALBRECHT 87D reports $< 8.5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.
[383] AVERY 87 reports $< 8 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$ Γ_{183}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.1 \times 10^{-7}$	90	[384] BERGFELD	00B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<4.0 \times 10^{-5}$	90	ABBOTT	98B	D0	$p\overline{p}$ 1.8 TeV
$<6.8 \times 10^{-7}$	90	[385] ABE	98	CDF	$p\overline{p}$ at 1.8 TeV
$<1.0 \times 10^{-5}$	90	[386] ACCIARRI	97B	L3	$e^+e^- \rightarrow Z$
$<1.6 \times 10^{-6}$	90	[387] ABE	96L	CDF	Repl. by ABE 98
$<5.9 \times 10^{-6}$	90	AMMAR	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$<8.3 \times 10^{-6}$	90	[388] ALBAJAR	91C	UA1	$E_{cm}^{p\overline{p}}$ = 630 GeV
$<1.2 \times 10^{-5}$	90	[389] ALBAJAR	91C	UA1	$E_{cm}^{p\overline{p}}$ = 630 GeV
$<4.3 \times 10^{-5}$	90	[390] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<4.5 \times 10^{-5}$	90	[391] ALBRECHT	87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<7.7 \times 10^{-5}$	90	[392] AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<2 \times 10^{-4}$	90	GILES	84	CLEO	Repl. by AVERY 87

[384] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[385] ABE 98 assumes production of $\sigma(B^0) = \sigma(B^+)$ and $\sigma(B_s)/\sigma(B^0) = 1/3$. They normalize to their measured $\sigma(B^0, p_T(B) > 6, |y| < 1.0) = 2.39 \pm 0.32 \pm 0.44\ \mu$b.
[386] ACCIARRI 97B assume PDG 96 production fractions for B^+, B^0, B_s, and Λ_b.
[387] ABE 96L assumes equal B^0 and B^+ production. They normalize to their measured $\sigma(B^+, p_T(B) > 6$ GeV/c, $|y| < 1) = 2.39 \pm 0.54\ \mu$b.
[388] B^0 and B^0_s are not separated.
[389] Obtained from unseparated B^0 and B^0_s measurement by assuming a B^0:B^0_s ratio 2:1.
[390] AVERY 89B reports $< 5 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[391] ALBRECHT 87D reports $< 5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.
[392] AVERY 87 reports $< 9 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^0 e^+ e^-)/\Gamma_{total}$ — Γ_{184}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<2.7 × 10⁻⁶	90	[393] ABE	02	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<8.45 \times 10^{-6}$	90	[394] ANDERSON	01B	CLE2	$e^+e^- \to \Upsilon(4S)$
$<3.0 \times 10^{-4}$	90	ALBRECHT	91E	ARG	$e^+e^- \to \Upsilon(4S)$
$<5.2 \times 10^{-4}$	90	[395] AVERY	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[393] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[394] The result is for di-lepton masses above 0.5 GeV.
[395] AVERY 87 reports $< 6.5 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^0 \mu^+ \mu^-)/\Gamma_{total}$ — Γ_{185}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.3 \times 10^{-6}$	90	[396] ABE	02	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6.64 \times 10^{-6}$	90	[397] ANDERSON	01B	CLE2	$e^+e^- \to \Upsilon(4S)$
$<5.2 \times 10^{-4}$	90	ALBRECHT	91E	ARG	$e^+e^- \to \Upsilon(4S)$
$<3.6 \times 10^{-4}$	90	[398] AVERY	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[396] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[397] The result is for di-lepton masses above 0.5 GeV.
[398] AVERY 87 reports $< 4.5 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^0 e^+ e^-)/\Gamma_{total}$ — Γ_{186}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.4 \times 10^{-6}$	90	[399] ABE	02	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.9 \times 10^{-4}$	90	ALBRECHT	91E	ARG	$e^+e^- \to \Upsilon(4S)$

[399] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^0 \mu^+ \mu^-)/\Gamma_{total}$ Γ_{187}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<4.2 × 10^{-6}	90	[400] ABE	02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<4.0 \times 10^{-6}$	90	[401] AFFOLDER	99B	CDF	$p\overline{p}$ at 1.8 TeV
$<2.5 \times 10^{-5}$	90	[402] ABE	96L	CDF	Repl. by AFFOLDER 99B
$<2.3 \times 10^{-5}$	90	[403] ALBAJAR	91C	UA1	$E_{cm}^{p\overline{p}} = 630$ GeV
$<3.4 \times 10^{-4}$	90	ALBRECHT	91E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[400] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[401] AFFOLDER 99B measured relative to $B^0 \rightarrow J/\psi(1S) K^*(892)^0$.

[402] ABE 96L measured relative to $B^0 \rightarrow J/\psi(1S) K^*(892)^0$ using PDG 94 branching ratios.

[403] ALBAJAR 91C assumes 36% of $\overline{b}$ quarks give B^0 mesons.

$\Gamma(K^*(892)^0 \nu \overline{\nu})/\Gamma_{total}$ Γ_{188}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.0 × 10^{-3}	90	[404] ADAM	96D	DLPH	$e^+ e^- \rightarrow Z$

[404] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

$\Gamma(e^\pm \mu^\mp)/\Gamma_{total}$ Γ_{189}/Γ

Test of lepton family number conservation. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<15 × 10^{-7}	90	[405] BERGFELD	00B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 3.5 \times 10^{-6}$	90	ABE	98V	CDF	$p\overline{p}$ at 1.8 TeV
$< 1.6 \times 10^{-5}$	90	[406] ACCIARRI	97B	L3	$e^+ e^- \rightarrow Z$
$< 5.9 \times 10^{-6}$	90	AMMAR	94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 3.4 \times 10^{-5}$	90	[407] AVERY	89B	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 4.5 \times 10^{-5}$	90	[408] ALBRECHT	87D	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 7.7 \times 10^{-5}$	90	[409] AVERY	87	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 3 \times 10^{-4}$	90	GILES	84	CLEO	Repl. by AVERY 87

[405] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[406] ACCIARRI 97B assume PDG 96 production fractions for B^+, B^0, B_s, and Λ_b.

[407] Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0 \overline{B}^0$. We rescale to 50%.

[408] ALBRECHT 87D reports $< 5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \overline{B}^0$. We rescale to 50%.

[409] AVERY 87 reports $< 9 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(e^\pm \tau^\mp)/\Gamma_{total}$ Γ_{190}/Γ

Test of lepton family number conservation. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<5.3 × 10^{-4}	90	AMMAR	94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\mu^{\pm}\tau^{\mp})/\Gamma_{total}$ Γ_{191}/Γ

Test of lepton family number conservation. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<8.3\times10^{-4}$	90	AMMAR	94	CLE2	$e^+e^-\to\Upsilon(4S)$

POLARIZATION IN B^0 DECAY

Γ_L/Γ in $B^0\to J/\psi(1S)K^*(892)^0$

$\Gamma_L/\Gamma=1$ would indicate that $B^0\to J/\psi(1S)K^*(892)^0$ followed by $K^*(892)^0\to K^0_S\pi^0$ is a pure CP eigenstate with $CP=-1$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.596±0.028 OUR AVERAGE					
0.597±0.028±0.024		[410] AUBERT	01H	BABR	$e^+e^-\to\Upsilon(4S)$
0.59 ±0.06 ±0.01		[411] AFFOLDER	00N	CDF	$p\overline{p}$ at 1.8 TeV
0.52 ±0.07 ±0.04		[412] JESSOP	97	CLE2	$e^+e^-\to\Upsilon(4S)$
0.65 ±0.10 ±0.04	65	ABE	95Z	CDF	$p\overline{p}$ at 1.8 TeV
0.97 ±0.16 ±0.15	13	[413] ALBRECHT	94G	ARG	$e^+e^-\to\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.80 ±0.08 ±0.05	42	[413] ALAM	94	CLE2	Sup. by JESSOP 97

[410] Averaged over an admixture of B^0 and B^- decays and the P wave fraction is $(16.0\pm3.2\pm1.4)\times10^{-2}$.

[411] AFFOLDER 00N measurements are based on 190 B^0 candidates obtained from a data sample of 89 pb^{-1}. The P-wave fraction is found to be $0.13^{+0.12}_{-0.9}\pm0.06$.

[412] JESSOP 97 is the average over a mixture of B^0 and B^+ decays. The P-wave fraction is found to be $0.16\pm0.08\pm0.04$.

[413] Averaged over an admixture of B^0 and B^+ decays.

Γ_L/Γ in $B^0\to\psi(2S)K^*(892)^0$

VALUE	DOCUMENT ID		TECN	COMMENT
0.45±0.11±0.04	[414] RICHICHI	01	CLE2	$e^+e^-\to\Upsilon(4S)$

[414] Averages between charged and neutral B mesons.

Γ_L/Γ in $B^0\to D_s^{*+}D^{*-}$

VALUE	DOCUMENT ID		TECN	COMMENT
0.506±0.139±0.036	AHMED	00B	CLE2	$e^+e^-\to\Upsilon(4S)$

Γ_L/Γ in $B^0\to D^{*-}\rho^+$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.93±0.05±0.05	76	ALAM	94	CLE2	$e^+e^-\to\Upsilon(4S)$

B^0–$\overline{B}^0$ MIXING

Written March 2000 and revised March 2002 by O. Schneider (University of Lausanne).

Formalism in quantum mechanics

There are two neutral B^0–$\overline{B}^0$ meson systems, B^0_d–$\overline{B}^0_d$ and B^0_s–$\overline{B}^0_s$ (generically denoted B^0_q–$\overline{B}^0_q$, $q = s, d$), which exhibit the phenomenon of particle-antiparticle mixing [1]. Such a system is produced in one of its two possible states of well-defined flavor: $|B^0\rangle$ ($\overline{b}q$) or $|\overline{B}^0\rangle$ ($b\overline{q}$). Due to flavor-changing interactions, this initial state evolves into a time-dependent quantum superposition of the two flavor states, $a(t)|B^0\rangle + b(t)|\overline{B}^0\rangle$, satisfying the equation

$$i\frac{\partial}{\partial t}\begin{pmatrix} a(t) \\ b(t) \end{pmatrix} = \left(\mathbf{M} - \tfrac{i}{2}\mathbf{\Gamma}\right)\begin{pmatrix} a(t) \\ b(t) \end{pmatrix}, \tag{1}$$

where $\mathbf{M}$ and $\mathbf{\Gamma}$, known as the mass and decay matrices, describe the dispersive and absorptive parts of B^0–$\overline{B}^0$ mixing. These matrices are hermitian, and CPT invariance requires $M_{11} = M_{22} \equiv M$ and $\Gamma_{11} = \Gamma_{22} \equiv \Gamma$

The two eigenstates of the effective Hamiltonian matrix $(\mathbf{M} - \frac{i}{2}\mathbf{\Gamma})$ are given by

$$|B_\pm\rangle = p|B^0\rangle \pm q|\overline{B}^0\rangle\,, \tag{2}$$

and correspond to the eigenvalues

$$\lambda_\pm = \left(M - \tfrac{i}{2}\Gamma\right) \pm \frac{q}{p}\left(M_{12} - \tfrac{i}{2}\Gamma_{12}\right)\,, \tag{3}$$

where

$$\frac{q}{p} = \sqrt{\frac{M^*_{12} - \frac{i}{2}\Gamma^*_{12}}{M_{12} - \frac{i}{2}\Gamma_{12}}}\,. \tag{4}$$

We choose a convention where $\mathrm{Re}(q/p) > 0$ and $CP|B^0\rangle = |\overline{B}^0\rangle$.

An alternative notation is

$$|B_\pm\rangle = \frac{(1+\epsilon)|B^0\rangle \pm (1-\epsilon)|\overline{B}^0\rangle}{\sqrt{2(1+|\epsilon|^2)}} \quad \text{with} \quad \frac{1-\epsilon}{1+\epsilon} = \frac{q}{p} . \quad (5)$$

The time dependence of these eigenstates of well-defined masses $M_\pm = \mathrm{Re}(\lambda_\pm)$ and decay widths $\Gamma_\pm = -2\,\mathrm{Im}(\lambda_\pm)$ is given by the phases $e^{-i\lambda_\pm t} = e^{-iM_\pm t} e^{-\frac{1}{2}\Gamma_\pm t}$: the evolution of a pure $|B^0\rangle$ or $|\overline{B}^0\rangle$ state at $t=0$ is thus given by

$$|B^0(t)\rangle = g_+(t)\,|B^0\rangle + \frac{q}{p}\,g_-(t)\,|\overline{B}^0\rangle , \quad (6)$$

$$|\overline{B}^0(t)\rangle = g_+(t)\,|\overline{B}^0\rangle + \frac{p}{q} g_-(t)\,|B^0\rangle , \quad (7)$$

where

$$g_\pm(t) = \frac{1}{2}\left(e^{-i\lambda_+ t} \pm e^{-i\lambda_- t}\right) . \quad (8)$$

This means that the flavor states remain unchanged (+) or oscillate into each other (−) with time-dependent probabilities proportional to

$$|g_\pm(t)|^2 = \frac{e^{-\Gamma t}}{2}\left[\cosh\left(\frac{\Delta\Gamma}{2}\,t\right) \pm \cos(\Delta m\, t)\right] , \quad (9)$$

where

$$\Delta m = |M_+ - M_-| , \qquad \Delta\Gamma = |\Gamma_+ - \Gamma_-| . \quad (10)$$

Time-integrated mixing probabilities are only well defined when considering decays to flavor-specific final states, *i.e.* final states f such that the instantaneous decay amplitudes $A_{\overline{f}} = \langle\overline{f}|H|B^0\rangle$ and $\overline{A}_f = \langle f|H|\overline{B}^0\rangle$, where H is the weak interaction Hamiltonian, are both zero. Due to mixing, a produced B^0 can decay to the final state $\overline{f}$ (mixed event) in addition to the final state

f (unmixed event). Restricting the sample to these two decay channels, the time-integrated mixing probability is given by

$$\chi_f^{B^0\to\overline{B}^0} = \frac{\int_0^\infty |\langle\overline{f}|H|B^0(t)\rangle|^2 dt}{\int_0^\infty |\langle\overline{f}|H|B^0(t)\rangle|^2 dt + \int_0^\infty |\langle f|H|B^0(t)\rangle|^2 dt}$$
$$= \frac{|\xi_f|^2(x^2+y^2)}{|\xi_f|^2(x^2+y^2)+2+x^2-y^2}\,, \tag{11}$$

where we have defined $\xi_f = \frac{q}{p}\frac{\overline{A}_{\overline{f}}}{A_f}$ and

$$x = \frac{\Delta m}{\Gamma}\,, \qquad y = \frac{\Delta\Gamma}{2\Gamma}\,. \tag{12}$$

The mixing probability $\chi_f^{\overline{B}^0\to B^0}$ for the case of a produced $\overline{B}^0$ is obtained by replacing ξ_f with $1/\xi_f$ in Eq. (11). It is different from $\chi_f^{B^0\to\overline{B}^0}$ if $|\xi_f|^2 \neq 1$, a condition reflecting non-invariance under the CP transformation. CP violation in decay amplitudes is discussed elsewhere [2] and we assume $|\overline{A}_{\overline{f}}| = |A_f|$ from now on. The deviation of $|q/p|^2$ from 1, namely the quantity

$$1 - \left|\frac{q}{p}\right|^2 = \frac{4\,\mathrm{Re}(\epsilon)}{1+|\epsilon|^2} + \mathcal{O}\left(\left(\frac{\mathrm{Re}(\epsilon)}{1+|\epsilon|^2}\right)^2\right)\,, \tag{13}$$

describes CP violation in B^0–$\overline{B}^0$ mixing. As can be seen from Eq. (4), this can occur only if $M_{12} \neq 0$, $\Gamma_{12} \neq 0$ and if the phase difference between M_{12} and Γ_{12} is different from 0 or π.

In the absence of CP violation, $|q/p|^2 = 1$, $\mathrm{Re}(\epsilon) = 0$, the mass eigenstates are also CP eigenstates,

$$CP\,|B_\pm\rangle = \pm|B_\pm\rangle\,, \tag{14}$$

the phases $\varphi_{M_{12}} = \arg(M_{12})$ and $\varphi_{\Gamma_{12}} = \arg(\Gamma_{12})$ satisfy

$$\sin(\varphi_{M_{12}} - \varphi_{\Gamma_{12}}) = 0\,, \tag{15}$$

Figure 1: Dominant box diagrams for the $B_q^0 \rightarrow \overline{B}_q^0$ transitions ($q = d$ or s). Similar diagrams exist where one or both t quarks are replaced with c or u quarks.

the mass and decay width differences reduce to

$$\Delta m = 2\,|M_{12}|\,, \qquad \Delta\Gamma = 2\,|\Gamma_{12}|\,, \tag{16}$$

and the time-integrated mixing probabilities $\chi_f^{B^0 \rightarrow \overline{B}^0}$ and $\chi_f^{\overline{B}^0 \rightarrow B^0}$ become both equal to

$$\chi = \frac{x^2 + y^2}{2(x^2 + 1)}\,. \tag{17}$$

Standard Model predictions and phenomenology

In the Standard Model, the transitions $B_q^0 \rightarrow \overline{B}_q^0$ and $\overline{B}_q^0 \rightarrow B_q^0$ are due to the weak interaction. They are described, at the lowest order, by box diagrams involving two W bosons and two up-type quarks (see Fig. **@Fg.box@**), as is the case for K^0–$\overline{K}^0$ mixing. However, the long range interactions arising from intermediate virtual states are negligible for the neutral B meson systems, because the large B mass is off the region of hadronic resonances. The calculation of the dispersive and absorptive parts of the box diagrams yields the following predictions for the off-diagonal element of the mass and decay matrices [3],

$$M_{12} = -\frac{G_F^2 m_W^2 \eta_B m_{B_q} B_{B_q} f_{B_q}^2}{12\pi^2} \, S_0(m_t^2/m_W^2) \, (V_{tq}^* V_{tb})^2 \qquad (18)$$

$$\Gamma_{12} = \frac{G_F^2 m_b^2 \eta'_B m_{B_q} B_{B_q} f_{B_q}^2}{8\pi} \times \left[(V_{tq}^* V_{tb})^2 + V_{tq}^* V_{tb} V_{cq}^* V_{cb} \; \mathcal{O}\left(\frac{m_c^2}{m_b^2}\right) + (V_{cq}^* V_{cb})^2 \; \mathcal{O}\left(\frac{m_c^4}{m_b^4}\right) \right] \qquad (19)$$

where G_F is the Fermi constant, m_W the W boson mass, m_i the mass of quark i, and $m_{B_q} = M$, f_{B_q} and B_{B_q} are the B_q^0 mass, weak decay constant and bag parameter, respectively. The known function $S_0(x_t)$ can be approximated very well with $0.784\, x_t^{0.76}$ [4] and V_{ij} are the elements of the CKM matrix [5]. The QCD corrections η_B and η'_B are of order unity. The only non negligible contributions to M_{12} are from box diagrams involving two top quarks. The phases of M_{12} and Γ_{12} satisfy

$$\varphi_{M_{12}} - \varphi_{\Gamma_{12}} = \pi + \mathcal{O}\left(\frac{m_c^2}{m_b^2}\right) \qquad (20)$$

implying that the mass eigenstates have mass and width differences of opposite signs. This means that, like in the K^0–$\overline{K}^0$ system, the "heavy" state with mass $M_{\rm heavy} = \max(M_+, M_-)$ has a smaller decay width than that of the "light" state with mass $M_{\rm light} = \min(M_+, M_-)$. We thus redefine

$$\Delta m = M_{\rm heavy} - M_{\rm light} \, , \qquad \Delta\Gamma = \Gamma_{\rm light} - \Gamma_{\rm heavy} \, , \qquad (21)$$

where Δm is positive by definition and $\Delta\Gamma$ is expected to be positive in the Standard Model.

Furthermore, the quantity

$$\left|\frac{\Gamma_{12}}{M_{12}}\right| \simeq \frac{3\pi}{2}\frac{m_b^2}{m_W^2}\frac{1}{S_0(m_t^2/m_W^2)} \sim \mathcal{O}\left(\frac{m_b^2}{m_t^2}\right) \tag{22}$$

is small, and a power expansion of $|q/p|^2$ yields

$$\left|\frac{q}{p}\right|^2 = 1 + \left|\frac{\Gamma_{12}}{M_{12}}\right| \sin(\varphi_{M_{12}} - \varphi_{\Gamma_{12}}) + \mathcal{O}\left(\left|\frac{\Gamma_{12}}{M_{12}}\right|^2\right) . \tag{23}$$

Therefore, considering both Eqs. (20) and (22), the CP-violating parameter

$$1 - \left|\frac{q}{p}\right|^2 \simeq \mathrm{Im}\left(\frac{\Gamma_{12}}{M_{12}}\right) \tag{24}$$

is expected to be very small: $\sim \mathcal{O}(10^{-3})$ for the B_d^0–$\overline{B}_d^0$ system and $\lesssim \mathcal{O}(10^{-4})$ for the B_s^0–$\overline{B}_s^0$ system [6].

In the approximation of negligible CP violation in mixing, the ratio $\Delta\Gamma/\Delta m$ is equal to the small quantity $|\Gamma_{12}/M_{12}|$ of Eq. (22); it is hence independent of CKM matrix elements, *i.e.* the same for the B_d^0–$\overline{B}_d^0$ and B_s^0–$\overline{B}_s^0$ systems. It can be calculated with lattice QCD techniques; typical results are $\sim 5 \times 10^{-3}$ with quoted uncertainties of $\sim 30\%$. Given the current experimental knowledge (discussed below) on the mixing parameter x,

$$\begin{cases} x_d = 0.755 \pm 0.015 & (B_d^0\text{–}\overline{B}_d^0 \text{ system}) \\ x_s > 19.0 \text{ at } 95\% \text{ CL} & (B_s^0\text{–}\overline{B}_s^0 \text{ system}) \end{cases} , \tag{25}$$

the Standard Model thus predicts that $\Delta\Gamma/\Gamma$ is very small for the B_d^0–$\overline{B}_d^0$ system (below 1%), but considerably larger for the B_s^0–$\overline{B}_s^0$ system ($\sim 10\%$). This width difference is caused by the existence of final states to which both the B_q^0 and $\overline{B}_q^0$ mesons can decay. Such decays involve $b \to c\overline{c}q$ quark-level transitions, which are Cabibbo-suppressed if $q = d$ and

Cabibbo-allowed if $q = s$. If the final states common to B_s^0 and $\overline{B}_s^0$ are predominantly CP-even as discussed in Ref. 7, then the B_s^0–$\overline{B}_s^0$ mass eigenstate with the largest decay width corresponds to the CP-even eigenstate. Taking Eq. (21) into account, one thus expects $\Gamma_{\rm light} = \Gamma_+$ and

$$\Delta m_s = M_- - M_+ > 0\,, \qquad \Delta\Gamma_s = \Gamma_+ - \Gamma_- > 0\,. \tag{26}$$

Experimental issues and methods for oscillation analyses

Time-integrated measurements of B^0–$\overline{B}^0$ mixing were published for the first time in 1987 by UA1 [8] and ARGUS [9], and since then by many other experiments. These measurements are typically based on counting same-sign and opposite-sign lepton pairs from the semileptonic decay of the produced $b\overline{b}$ pairs. Such analyses cannot easily separate the contributions from the different b-hadron species, therefore the clean environment of $\Upsilon(4S)$ machines (where only B_d^0 and charged B_u mesons are produced) is in principle best suited to measure χ_d.

However, better sensitivity is obtained from time-dependent analyses aimed at the direct measurement of the oscillation frequencies Δm_d and Δm_s, from the proper time distributions of B_d^0 or B_s^0 candidates identified through their decay in (mostly) flavor-specific modes and suitably tagged as mixed or unmixed. (This is particularly true for the B_s^0–$\overline{B}_s^0$ system where the large value of x_s implies maximal mixing, *i.e.* $\chi_s \simeq 1/2$.) In such analyses the B_d^0 or B_s^0 mesons are either fully reconstructed, partially reconstructed from a charm meson, selected from a lepton with the characteristics of a $b \to \ell^-$ decay, or selected from a reconstructed displaced vertex. At high-energy colliders (LEP, SLC, Tevatron), the proper time $t = \dfrac{m_B}{p}L$ is measured from the distance L between the production vertex and the B decay vertex, and from an estimate of the B momentum p. At asymmetric B factories (KEKB, PEP-II), producing $e^+e^- \to \Upsilon(4S) \to B_d^0\overline{B}_d^0$ events with a boost $\beta\gamma$ (= 0.425, 0.55), the proper time difference between the two B candidates is estimated as $\Delta t \simeq \dfrac{\Delta z}{\beta\gamma c}$, where Δz is the spatial separation between the two B decay vertices along the boost direction. In all cases, the good resolution needed on the vertex positions is obtained with silicon detectors.

The statistical significance $\mathcal{S}$ of a B_d^0 or B_s^0 oscillation signal can be approximated as [10]

$$\mathcal{S} \approx \sqrt{N/2}\, f_{\rm sig}\,(1-2\eta)\, e^{-(\Delta m\, \sigma_t)^2/2}\,, \qquad (27)$$

where N and $f_{\rm sig}$ are the number of candidates and the fraction of signal in the selected sample, η is the total mistag probability, and σ_t is the resolution on proper time (or proper time difference). The quantity $\mathcal{S}$ decreases very quickly as Δm increases; this dependence is controlled by σ_t, which is therefore a critical parameter for Δm_s analyses. At high-energy colliders, the proper time resolution $\sigma_t \sim \dfrac{m_B}{\langle p\rangle}\sigma_L \oplus t\dfrac{\sigma_p}{p}$ includes a constant contribution due to the decay length resolution σ_L (typically 0.05–0.3 ps), and a term due to the relative momentum resolution σ_p/p (typically 10–20% for partially reconstructed decays) which increases with proper time. At B factories, the B momentum is reconstructed and/or estimated from the beam energy constraint, and the term due to the spatial resolution dominates (typically 1–1.5 ps because of the much smaller B boost).

In order to tag a B candidate as mixed or unmixed, it is necessary to determine its flavor both in the initial state and in the final state. The initial and final state mistag probabilities, η_i and η_f, degrade $\mathcal{S}$ by a total factor $(1-2\eta) = (1-2\eta_i)(1-2\eta_f)$. In lepton-based analyses, the final state is tagged by the charge of the lepton from $b \to \ell^-$ decays; the biggest contribution to η_f is then due to $\overline{b} \to \overline{c} \to \ell^-$ decays. Alternatively, the charge of a reconstructed charm meson (D^{*-} from B^0_d or D^-_s from B^0_s), or that of a kaon thought to come from a $b \to c \to s$ decay [11], can be used. For fully inclusive analyses based on topological vertexing, final state tagging techniques include jet charge [12] and charge dipole [13,14] methods.

At high-energy colliders, the methods to tag the initial state (i.e. the state at production), can be divided in two groups: the ones that tag the initial charge of the $\overline{b}$ quark contained in the B candidate itself (same-side tag), and the ones that tag the initial charge of the other b quark produced in the event (opposite-side tag). On the same side, the charge of a track from the primary vertex is correlated with the production state of the B if that track is a decay product of a B^{**} state or the first particle in the fragmentation chain [15,16]. Jet- and vertex-charge techniques work on both sides and on the opposite side, respectively. Finally, the charge of a lepton from $b \to \ell^-$ or of a kaon from $b \to c \to s$ can be used as opposite side tags, keeping in mind that their performance is degraded due to integrated mixing. At SLC, the beam polarization produced a sizeable forward-backward asymmetry in the $Z \to b\overline{b}$ decays and provided another very interesting and effective initial state tag based on the polar angle of the B candidate [11]. Initial state tags have also been combined to reach $\eta_i \sim 26\%$ at LEP [16,17], or even 22% at SLD [13] with full efficiency. The equivalent figure at CDF (Tevatron Run I) is $\sim 40\%$ [18].

At B factories, the flavor of a B^0_d meson at production cannot be determined, since the two neutral B mesons produced in a $\Upsilon(4S)$ decay evolve in a coherent P-wave state where they keep opposite flavors at any time. However, as soon as one of them decays, the other follows a time-evolution given by Eqs. (6) or (7), where t is replaced with Δt. Hence, the "initial state" tag of a B can be taken as the final state tag of the other B. Effective mistag probabilities of $\eta_i \sim 24\%$ for full efficiency (corresponding to effective tagging efficiencies of $\sim 27\%$ for perfect tagging) are achieved by BABAR and Belle [19], using different techniques including $b \to \ell^-$ and $b \to c \to s$ tags. It is interesting to note that, in this case, mixing of this other B (i.e. the coherent mixing occurring before the first B decay) does not contribute to the mistag probability.

In the absence of experimental evidence for a width difference, oscillation analyses typically neglect $\Delta\Gamma$ and describe the data with the physics functions $\Gamma e^{-\Gamma t}(1 \pm \cos(\Delta m t))/2$ (high-energy colliders) or $\Gamma e^{-\Gamma|\Delta t|}(1 \pm \cos(\Delta m \Delta t))/4$ (asymmetric $\Upsilon(4S)$ machines). As can be seen from Eq. (9), a non-zero value of $\Delta\Gamma$ would effectively reduce the oscillation amplitude with a small time-dependent factor that would be very difficult to distinguish from time resolution effects. Whereas measurements of Δm_d are usually extracted from the data using a maximum likelihood fit, no significant B_s^0–$\overline{B}_s^0$ oscillations have been seen so far. To extract information useful to set lower limits on Δm_s, B_s^0 analyses follow a method [10] in which a B_s^0 oscillation amplitude $\mathcal{A}$ is measured as a function of a fixed test value of Δm_s, using a maximum likelihood fit based on the functions $\Gamma_s e^{-\Gamma_s t}(1 \pm \mathcal{A}\cos(\Delta m_s t))/2$. To a very good approximation, the statistical uncertainty on $\mathcal{A}$ is Gaussian and equal to $1/\mathcal{S}$ [10]. If $\Delta m_s = \Delta m_s^{\rm true}$, one expects $\mathcal{A} = 1$ within the total uncertainty $\sigma_{\mathcal{A}}$; however, if Δm_s is (far) below its true value, a measurement consistent with $\mathcal{A} = 0$ is expected. A value of Δm_s can be excluded at 95% CL if $\mathcal{A} + 1.645\,\sigma_{\mathcal{A}} \le 1$. If $\Delta m_s^{\rm true}$ is very large, one expects $\mathcal{A} = 0$, and all values of Δm_s such that

$1.645\,\sigma_{\mathcal{A}}(\Delta m_s) < 1$ are expected to be excluded at 95% CL. Because of the proper time resolution, the quantity $\sigma_{\mathcal{A}}(\Delta m_s)$ is an increasing function of Δm_s and one therefore expects to be able to exclude individual Δm_s values up to $\Delta m_s^{\mathrm{sens}}$, where $\Delta m_s^{\mathrm{sens}}$, called here the sensitivity of the analysis, is defined by $1.645\,\sigma_{\mathcal{A}}(\Delta m_s^{\mathrm{sens}}) = 1$.

B_d^0 *mixing studies*

Many B_d^0–$\overline{B}_d^0$ oscillations analyses have been performed by the ALEPH [12,20], BABAR [21], Belle [22,23], CDF [15,24], DELPHI [14,25], L3 [26], OPAL [27] and SLD [11] collaborations. Although a variety of different techniques have been used, the individual Δm_d results obtained at high-energy colliders have remarkably similar precision. Their average is compatible with the recent and more precise measurements from asymmetric B factories. The systematic uncertainties are not negligible; they are often dominated by sample composition, mistag probability, or b-hadron lifetime contributions. Before being combined, the measurements are adjusted on the basis of a common set of input values, including the b-hadron lifetimes and fractions published in this *Review*. Some measurements are statistically correlated. Systematic correlations arise both from common physics sources (fragmentation fractions, lifetimes, branching ratios of b hadrons), and from purely experimental or algorithmic effects (efficiency, resolution, tagging, background description). Combining all published measurements [15,20–22,25–27] and accounting for all identified correlations as described in Ref. 28 yields $\Delta m_d = 0.489 \pm 0.005(\mathrm{stat}) \pm 0.007(\mathrm{syst})\ \mathrm{ps}^{-1}$.

On the other hand, ARGUS and CLEO have published time-integrated measurements [29–31], which average to $\chi_d = 0.182 \pm 0.015$. Following Ref. 31, the width difference $\Delta\Gamma_d$ could in principle be extracted from the measured value of Γ_d

and the above averages for Δm_d and χ_d (see Eqs. (12) and (17)), provided that $\Delta\Gamma_d$ has a negligible impact on the Δm_d measurements. However, a stronger constraint, $\Delta\Gamma_d/\Gamma_d < 20\%$ at 90% CL, has been obtained by DELPHI from a direct time-dependent study [14]. Assuming $\Delta\Gamma_d = 0$ and no CP violation in mixing, and using the measured B_d^0 lifetime, the Δm_d and χ_d results are combined to yield the world average

$$\Delta m_d = 0.489 \pm 0.008 \ \text{ps}^{-1} \tag{28}$$

or, equivalently,

$$\chi_d = 0.181 \pm 0.004\,. \tag{29}$$

Evidence for CP violation in B_d^0 mixing has been searched for, both with flavor-specific and inclusive B_d^0 decays, in samples where the initial flavor state is tagged. In the case of semileptonic (or other flavor-specific) decays, where the final state tag is also available, the following asymmetry

$$\begin{aligned} &\frac{N(\overline{B}_d^0(t) \to \ell^+\nu_\ell X) - N(B_d^0(t) \to \ell^-\overline{\nu}_\ell X)}{N(\overline{B}_d^0(t) \to \ell^+\nu_\ell X) + N(B_d^0(t) \to \ell^-\overline{\nu}_\ell X)} \\ &= a_{CP} \simeq 1 - |q/p|_d^2 \simeq \frac{4\text{Re}(\epsilon_d)}{1+|\epsilon_d|^2} \end{aligned} \tag{30}$$

has been measured, either in time-integrated analyses at CLEO [30–32] and CDF [33], or in time-dependent analyses at LEP [34–36] and BABAR [37]. In the inclusive case, also investigated at LEP [35,36,38], no final state tag is used, and the asymmetry [39]

$$\begin{aligned} &\frac{N(B_d^0(t) \to \text{all}) - N(\overline{B}_d^0(t) \to \text{all})}{N(B_d^0(t) \to \text{all}) + N(\overline{B}_d^0(t) \to \text{all})} \\ &\simeq a_{CP}\left[\frac{x_d}{2}\sin(\Delta m_d\, t) - \sin^2\left(\frac{\Delta m_d\, t}{2}\right)\right] \end{aligned} \tag{31}$$

must be measured as a function of the proper time to extract information on CP violation. In all cases asymmetries compatible with zero have been found, with a precision limited by the available statistics. A simple average of all published and preliminary results [30–38] neglecting small possible statistical correlations and assuming half of the systematics to be correlated between measurements performed at the same energy, is $a_{CP} = -0.002 \pm 0.009(\mathrm{stat}) \pm 0.008(\mathrm{syst})$, a result which does not yet constrain the Standard Model.

The Δm_d result of Eq. (28) provides an estimate of $|M_{12}|$ and can be used, together with Eqs. (16) and (18), to extract the magnitude of the CKM matrix element V_{td} within the Standard Model [40]. The main experimental uncertainties on the resulting estimate of $|V_{td}|$ come from m_t and Δm_d; however, the extraction is at present completely dominated by the uncertainty on the hadronic matrix element $f_{B_d}\sqrt{B_{B_d}} = 230 \pm 40$ MeV obtained from lattice QCD calculations [41].

B_s^0 *mixing studies*

B_s^0–$\overline{B}_s^0$ oscillations have been the subject of many studies from ALEPH [16,42], CDF [43], DELPHI [14,17,44,45], OPAL [46] and SLD [13,47]. No oscillation signal has been found so far. The most sensitive analyses appear to be the ones based on inclusive lepton samples at LEP. Because of their better proper time resolution, the small data samples analyzed inclusively at SLD, as well as the few fully reconstructed B_s decays at LEP, turn out to be also very useful to explore the high Δm_s region.

All results are limited by the available statistics. They can easily be combined, since all experiments provide measurements of the B_s^0 oscillation amplitude. The latter are averaged using the procedure of Ref. 28 to yield the combined amplitudes

$\mathcal{A}$ shown in Fig. **@Fg.amplitude@** as a function of Δm_s. The individual results have been adjusted to common physics inputs, and all known correlations have been accounted for; the sensitivities of the inclusive analyses, which depend directly through Eq. (27) on the assumed fraction f_s of B_s^0 mesons in an unbiased sample of weakly-decaying b hadrons, have also been rescaled to a common preliminary average of $f_s = 0.097 \pm 0.011$. The combined sensitivity for 95% CL exclusion of Δm_s values is found to be 19.3 ps^{-1}. All values of Δm_s below 14.9 ps^{-1} are excluded at 95% CL. The values between 14.9 and 22.4 ps^{-1} cannot be excluded, because the data is compatible with a signal in this region. However, no deviation from $\mathcal{A} = 0$ is seen in Fig. **@Fg.amplitude@** that would indicate the observation of a signal.

Some Δm_s analyses are still unpublished [13,14,42,45,47]. Using only published results, the combined Δm_s result is

$$\Delta m_s > 13.1 \ \text{ps}^{-1} \quad \text{at } 95\% \text{ CL}\,, \tag{32}$$

with a sensitivity of 13.3 ps^{-1}.

The information on $|V_{ts}|$ obtained, in the framework of the Standard Model, from the combined amplitude spectrum is hampered by the hadronic uncertainty, as in the B_d^0 case. However, many uncertainties cancel in the frequency ratio

$$\frac{\Delta m_s}{\Delta m_d} = \frac{m_{B_s}}{m_{B_d}} \xi^2 \left| \frac{V_{ts}}{V_{td}} \right|^2 , \tag{33}$$

where $\xi = (f_{B_s}\sqrt{B_{B_s}})/(f_{B_d}\sqrt{B_{B_d}}) = 1.16 \pm 0.05$ is an SU(3) flavor-symmetry breaking factor obtained from lattice QCD calculations [41]. The CKM matrix can be constrained using the experimental results on Δm_d, Δm_s, $|V_{ub}/V_{cb}|$, ϵ_K and $\sin(2\beta)$ together with theoretical inputs and unitarity conditions [40,49].



Figure 2: Combined measurements of the B_s^0 oscillation amplitude as a function of Δm_s, including all preliminary results available at the time of the winter 2002 conferences [48]. The measurements are dominated by statistical uncertainties. Neighboring points are statistically correlated.

Given all measurements other than Δm_d and Δm_s, the constraint from our knowledge on the ratio $\Delta m_d/\Delta m_s$ is presently more effective in limiting the position of the apex of the CKM unitarity triangle than the one obtained from the Δm_d measurements alone, due to the reduced hadronic uncertainty in Eq. (33). We note also that it would be difficult for the Standard Model to accommodate values of Δm_s above $\sim 25\ \text{ps}^{-1}$ [49].

Information on $\Delta\Gamma_s$ can be obtained by studying the proper time distribution of untagged data samples enriched in B_s^0

mesons [50]. In the case of an inclusive B_s^0 selection [51] or a semileptonic B_s^0 decay selection [17,52], both the short- and long-lived components are present, and the proper time distribution is a superposition of two exponentials with decay constants $\Gamma_s \pm \Delta\Gamma_s/2$. In principle, this provides sensitivity to both Γ_s and $(\Delta\Gamma_s/\Gamma_s)^2$. Ignoring $\Delta\Gamma_s$ and fitting for a single exponential leads to an estimate of Γ_s with a relative bias proportional to $(\Delta\Gamma_s/\Gamma_s)^2$. An alternative approach, which is directly sensitive to first order in $\Delta\Gamma_s/\Gamma_s$, is to determine the lifetime of B_s^0 candidates decaying to CP eigenstates; measurements exist for $B_s^0 \to J/\psi\phi$ [53] and $B_s^0 \to D_s^{(*)+}D_s^{(*)-}$ [54], which are mostly CP-even states [7]. An estimate of $\Delta\Gamma_s/\Gamma_s$ has also been obtained directly from a measurement of the $B_s^0 \to D_s^{(*)+}D_s^{(*)-}$ branching ratio [54], under the assumption that these decays practically account for all the CP-even final states.

Present data is not precise enough to efficiently constrain both Γ_s and $\Delta\Gamma_s/\Gamma_s$; since the B_s^0 and B_d^0 lifetimes are predicted to be equal within less than a percent [55], an expectation compatible with the current experimental data [56], the constraint $\Gamma_s = \Gamma_d$ can also be used to improve the extraction of $\Delta\Gamma_s/\Gamma_s$. Applying the combination procedure of Ref. 28 on the published results [17,52–54,57] yields

$$\Delta\Gamma_s/\Gamma_s < 0.52 \quad \text{at } 95\% \text{ CL} \tag{34}$$

without external constraint, or

$$\Delta\Gamma_s/\Gamma_s < 0.31 \quad \text{at } 95\% \text{ CL} \tag{35}$$

when constraining $1/\Gamma_s$ to the measured B_d^0 lifetime. These results are not yet precise enough to test Standard Model predictions.

Average b-hadron mixing and b-hadron production fractions at high energy

Let f_u, f_d, f_s and $f_{\rm baryon}$ be the B_u, B^0_d, B^0_s and b-baryon fractions composing an unbiased sample of weakly-decaying b hadrons produced in high-energy colliders. LEP experiments have measured $f_s \times {\rm BR}(B^0_s \to D^-_s \ell^+ \nu_\ell X)$ [58], ${\rm BR}(b \to \Lambda^0_b) \times {\rm BR}(\Lambda^0_b \to \Lambda^+_c \ell^- \overline{\nu}_\ell X)$ [59] and ${\rm BR}(b \to \Xi^-_b) \times {\rm BR}(\Xi^-_b \to \Xi^- \ell^- \overline{\nu}_\ell X)$ [60] from partially reconstructed final states including a lepton, $f_{\rm baryon}$ from protons identified in b events [61], and the production rate of charged b hadrons [62]. The various b-hadron fractions have also been measured at CDF from electron-charm final states [63]. All the published results have been combined following the procedure and assumptions described in Ref. 28, to yield $f_u = f_d = (37.3 \pm 2.0)\%$, $f_s = (13.9 \pm 3.8)\%$ and $f_{\rm baryon} = (11.5 \pm 2.0)\%$ under the constraints

$$f_u = f_d \qquad \text{and} \qquad f_u + f_d + f_s + f_{\rm baryon} = 1\,. \tag{36}$$

Time-integrated mixing analyses performed with lepton pairs from $b\overline{b}$ events produced at high-energy colliders measure the quantity

$$\overline{\chi} = f'_d\,\chi_d + f'_s\,\chi_s\,, \tag{37}$$

where f'_d and f'_s are the fractions of B^0_d and B^0_s hadrons in a sample of semileptonic b-hadron decays. Assuming that all b hadrons have the same semileptonic decay width implies $f'_q = f_q/(\Gamma_q \tau_b)$ $(q = s, d)$, where τ_b is the average b-hadron lifetime. Hence $\overline{\chi}$ measurements can be used to improve our knowledge on the fractions f_u, f_d, f_s and $f_{\rm baryon}$.

Combining the above estimates of these fractions with the average $\overline{\chi} = 0.1184 \pm 0.0045$ (published in this *Review*), χ_d from

Eq. (29) and $\chi_s = 1/2$ yields, under the constraints of Eq. (36),

$$f_u = f_d = (38.8 \pm 1.3)\% \,, \tag{38}$$

$$f_s = (10.6 \pm 1.3)\% \,, \tag{39}$$

$$f_{\rm baryon} = (11.8 \pm 2.0)\% \,, \tag{40}$$

showing that mixing information substantially reduces the uncertainty on f_s. These results and the averages quoted in Eqs. (28) and (29) for χ_d and Δm_d have been obtained in a consistent way by the B oscillations working group [28], taking into account the fact that many individual measurements of Δm_d depend on the assumed values for the b-hadron fractions.

Summary and prospects

B^0–$\overline{B}^0$ mixing has been and still is a field of intense study. The mass difference in the B^0_d–$\overline{B}^0_d$ system is very well measured (with an accuracy of 1.7%) but, despite an impressive theoretical effort, the hadronic uncertainty still limits the precision of the extracted estimate of $|V_{td}|$. The mass difference in the B^0_s–$\overline{B}^0_s$ system is much larger and still unmeasured. However, the current experimental lower limit on Δm_s already provides, together with Δm_d, a significant constraint on the CKM matrix within the Standard Model. No strong experimental evidence exists yet for the rather large decay width difference expected in the B^0_s–$\overline{B}^0_s$ system. It is interesting to recall that the ratio $\Delta\Gamma_s/\Delta m_s$ does not depend on CKM matrix elements in the Standard Model (see Eq. (22)), and that a measurement of either Δm_s or $\Delta\Gamma_s$ could be turned into a Standard Model prediction of the other one.

The LEP and SLD experiments have still not finalized all their B^0_s oscillation analyses, but a first measurement of Δm_s from data collected at the Z pole is now very unlikely. In the near future, the most promising prospects for B_s^0 mixing are from Run II at the Tevatron, where both Δm_s and $\Delta\Gamma_s$ are expected to be measured with fully reconstructed B_s^0 decays; for example, with 2 fb^{-1} of data, CDF expects to observe B_s^0 oscillations for values of Δm_s up to $\sim 40-50$ ps^{-1} (depending on event yields and signal-to-background ratios) [64], well above the current Standard Model prediction.

CP violation in B mixing, which has not been seen yet, as well as the phases involved in B mixing, will be further investigated with the large statistics that will become available both at the B factories and at the Tevatron.

B mixing may not have delivered all its secrets yet, because it is one of the phenomena where new physics might very well reveal itself (for example new particles involved in the box diagrams). Theoretical calculations in lattice QCD are becoming more reliable and further progress in reducing hadronic uncertainties is expected. In the long term, a stringent check of the consistency, within the Standard Model, of the B_d^0 and B_s^0 mixing measurements with all other measured observables in B physics (including CP asymmetries in B decays) will be possible, allowing to place limits on new physics or, better, discover new physics.

References

1. T.D. Lee and C.S. Wu, Ann. Rev. Nucl. Sci. **16**, 511 (1966);
   I.I. Bigi and A.I. Sanda, "CP violation," Cambridge, Cambridge Univ. Press, 2000;
   G.C. Branco, L. Lavoura, and J.P. Silva, "CP violation," Clarendon Press Oxford, 1999;
   see also the review on B^0–$\overline{B}^0$ mixing by H. Quinn in C. Caso *et al.*, Eur. Phys. J. **C3**, 1 (1998).
2. See the review on CP violation in B decays by H. Quinn and A. Sanda in this publication.

3. A.J. Buras, W. Slominski, and H. Steger, Nucl. Phys. **B245**, 369 (1984).
4. A.J. Buras and R. Fleischer, in "Heavy Flavours II," ed. A.J. Buras and M. Lindner, Singapore World Scientific (1998).
5. M. Kobayashi and K. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
6. I.I. Bigi *et al.*, in "CP violation," ed. C. Jarlskog, Singapore World Scientific, 1989.
7. R. Aleksan *et al.*, Phys. Lett. **B316**, 567 (1993).
8. C. Albajar *et al.* (**UA1**), Phys. Lett. **B186**, 247 (1987).
9. H. Albrecht *et al.* (**ARGUS**), Phys. Lett. **B192**, 245 (1987).
10. H.-G. Moser and A. Roussarie, Nucl. Instrum. Methods **384**, 491 (1997).
11. **SLD** collab., SLAC-PUB-7228, SLAC-PUB-7229 and SLAC-PUB-7230, contrib. to 28th Int. Conf. on High Energy Physics, Warsaw, 1996;
J. Wittlin, PhD thesis, SLAC-R-582, 2001.
12. **ALEPH** collab., contrib. 596 to Int. Europhysics Conf. on High Energy Physics, Jerusalem, 1997.
13. J. Thom, PhD thesis, SLAC-R-585, 2002.
14. **DELPHI** collab., contrib. 303 to Int. Europhysics Conf. on High Energy Physics, Budapest, 2001.
15. F. Abe *et al.* (**CDF**), Phys. Rev. Lett. **80**, 2057 (1998), Phys. Rev. **D59**, 032001 (1999); Phys. Rev. **D60**, 051101 (1999); Phys. Rev. **D60**, 072003 (1999);
T. Affolder *et al.* (**CDF**), Phys. Rev. **D60**, 112004 (1999).
16. R. Barate *et al.* (**ALEPH**), Eur. Phys. J. **C4**, 367 (1998); Eur. Phys. J. **C7**, 553 (1999).
17. P. Abreu *et al.* (**DELPHI**), Eur. Phys. J. **C16**, 555 (2000); Eur. Phys. J. **C18**, 229 (2000).
18. M. Paulini, private communication.
19. B. Aubert *et al.* (**BABAR**), Phys. Rev. Lett. **87**, 091801 (2001);

K. Abe *et al.* (**Belle**), Phys. Rev. Lett. **87**, 091802 (2001).

20. D. Buskulic *et al.* (**ALEPH**), Z. Phys. **C75**, 397 (1997).
21. B. Aubert *et al.* (**BABAR**), hep-ex/0112044, SLAC-PUB-9061, to appear in Phys. Rev. Lett.; hep-ex/0112045, SLAC-PUB-9096, to appear in Phys. Rev. Lett.
22. K. Abe *et al.* (**Belle**), Phys. Rev. Lett. **86**, 3228 (2001).
23. Y. Sakai, hep-ex/0202010, to appear in the proc. of the 5th KEK topical conference, 2001.
24. M. Paulini, Int. J. Mod. Phys. **A14**, 2791 (1999).
25. P. Abreu *et al.* (**DELPHI**), Z. Phys. **C76**, 579 (1997).
26. M. Acciarri *et al.* (**L3**), Eur. Phys. J. **C5**, 195 (1998).
27. G. Alexander *et al.* (**OPAL**), Z. Phys. **C72**, 377 (1996);
K. Ackerstaff *et al.* (**OPAL**), Z. Phys. **C76**, 401 (1997); Z. Phys. **C76**, 417 (1997);
G. Abbiendi *et al.* (**OPAL**), Phys. Lett. **B493**, 266 (2000).
28. **ALEPH**, **CDF**, **DELPHI**, **L3**, **OPAL**, and **SLD** collab., "Combined results on b-hadron production rates and decay properties," CERN-EP/2001-050;
the combined results on B mixing and b-hadron fractions included in the above paper or published in this *Review* have been obtained by the B oscillations working group; see http://www.cern.ch/LEPBOSC/ for more information.
29. H. Albrecht *et al.* (**ARGUS**), Z. Phys. **C55**, 357 (1992); Phys. Lett. **B324**, 249 (1994).
30. J. Bartelt *et al.* (**CLEO**), Phys. Rev. Lett. **71**, 1680 (1993).
31. B.H. Behrens *et al.* (**CLEO**), Phys. Lett. **B490**, 36 (2000).
32. D.E. Jaffe *et al.* (**CLEO**), Phys. Rev. Lett. **86**, 5000 (2001).
33. F. Abe *et al.* (**CDF**), Phys. Rev. **D55**, 2546 (1997).
34. K. Ackerstaff *et al.* (**OPAL**), Z. Phys. **C76**, 401 (1997).
35. **DELPHI** collab., contrib. 449 to Int. Europhysics Conf. on High Energy Physics, Jerusalem, 1997.
36. R. Barate *et al.* (**ALEPH**), Eur. Phys. J. **C20**, 431 (2001).

37. B. Aubert *et al.* (**BABAR**), hep-ex/0202041, SLAC-PUB-9149, submitted to Phys. Rev. Lett.
38. G. Abbiendi *et al.* (**OPAL**), Eur. Phys. J. **C12**, 609 (2000).
39. M. Beneke, G. Buchalla, and I. Dunietz, Phys. Lett. **B393**, 132 (1997);
    I. Dunietz, Eur. Phys. J. **C7**, 197 (1999).
40. See the review on the CKM quark-mixing matrix by F.J. Gilman, K. Kleinknecht and B. Renk in this publication.
41. For a review of lattice QCD results with heavy quarks, see for example C. Bernard, Nucl. Phys. (Proc. Supp.) **B94**, 159 (2001).
42. A. Heister *et al.* (**ALEPH**), CERN/EP-2002-016, submitted to Eur. Phys. J. C.
43. F. Abe *et al.* (**CDF**), Phys. Rev. Lett. **82**, 3576 (1999).
44. W. Adam *et al.* (**DELPHI**), Phys. Lett. **B414**, 382 (1997).
45. **DELPHI** collab., contrib. 304 to Int. Europhysics Conf. on High Energy Physics, Budapest, 2001.
46. G. Abbiendi *et al.* (**OPAL**), Eur. Phys. J. **C11**, 587 (1999); Eur. Phys. J. **C19**, 241 (2001).
47. **SLD** collab., SLAC-PUB-8568 and SLAC-PUB-8598, contrib. to 30th Int. Conf. on High Energy Physics, Osaka, 2000;
    C.-J.S. Lin, PhD thesis, SLAC-R-584, 2001.
48. D. Abbaneo, "Review of B^0_s oscillations," to appear in the proc. of the 16th Rencontres de la Vallée d'Aoste, La Thuile, 2002.
49. Workshop on the CKM unitarity triangle, CERN, February 2002, http://www.cern.ch/ckm-workshop/, CERN yellow report in preparation.
50. K. Hartkorn and H.-G. Moser, Eur. Phys. J. **C8**, 381 (1999).
51. M. Acciarri *et al.* (**L3**), Phys. Lett. **B438**, 417 (1998).
52. F. Abe *et al.* (**CDF**), Phys. Rev. **D59**, 032004 (1999).
53. F. Abe *et al.* (**CDF**), Phys. Rev. **D57**, 5382 (1998).
54. R. Barate *et al.* (**ALEPH**), Phys. Lett. **B486**, 286 (2000).

55. See for example M. Beneke, G. Buchalla, and I. Dunietz, Phys. Rev. **D54**, 4419 (1996).
56. See the review on production and decay of b-hadrons by L. Gibbons and K. Honscheid in this publication.
57. D. Buskulic *et al.* (**ALEPH**), Phys. Lett. **B377**, 205 (1996);
K. Ackerstaff *et al.* (**OPAL**), Phys. Lett. **B426**, 161 (1998).
58. P. Abreu *et al.* (**DELPHI**), Phys. Lett. **B289**, 199 (1992);
P.D. Acton *et al.* (**OPAL**), Phys. Lett. **B295**, 357 (1992);
D. Buskulic *et al.* (**ALEPH**), Phys. Lett. **B361**, 221 (1995).
59. P. Abreu *et al.* (**DELPHI**), Z. Phys. **C68**, 375 (1995);
R. Barate *et al.* (**ALEPH**), Eur. Phys. J. **C2**, 197 (1998).
60. P. Abreu *et al.* (**DELPHI**), Z. Phys. **C68**, 541 (1995);
D. Buskulic *et al.* (**ALEPH**), Phys. Lett. **B384**, 449 (1996).
61. R. Barate *et al.* (**ALEPH**), Eur. Phys. J. **C5**, 205 (1998).
62. **DELPHI** collab., contrib. 5_515 to Int. Europhysics Conf. on High Energy Physics, Tampere, 1999.
63. F. Abe *et al.* (**CDF**), Phys. Rev. **D60**, 092005 (1999);
T. Affolder *et al.* (**CDF**), Phys. Rev. Lett. **84**, 1663 (2000).
64. K. Anikeev *et al.*, "B physics at the Tevatron: Run II and beyond," `hep-ph/0201071`, FERMILAB-PUB-01/97.

B^0-$\overline{B}^0$ MIXING PARAMETERS

For a discussion of B^0-$\overline{B}^0$ mixing see the note on "B^0-$\overline{B}^0$ Mixing" in the B^0 Particle Listings above.

χ_d is a measure of the time-integrated B^0-$\overline{B}^0$ mixing probability that a produced $B^0(\overline{B}^0)$ decays as a $\overline{B}^0(B^0)$. Mixing violates $\Delta B \neq 2$ rule.

$$\chi_d = \frac{x_d^2}{2(1+x_d^2)}$$

$$x_d = \frac{\Delta m_{B^0}}{\Gamma_{B^0}} = (m_{B^0_H} - m_{B^0_L})\,\tau_{B^0}\ ,$$

where H, L stand for heavy and light states of two B^0 CP eigenstates and $\tau_{B^0} = \frac{1}{0.5(\Gamma_{B^0_H}+\Gamma_{B^0_L})}$.

χ_d

This B^0-$\overline{B}^0$ mixing parameter is the probability (integrated over time) that a produced B^0 (or $\overline{B}^0$) decays as a $\overline{B}^0$ (or B^0), e.g. for inclusive lepton decays

$$\chi_d = \Gamma(B^0 \rightarrow \ell^- X \text{ (via } \overline{B}^0))/\Gamma(B^0 \rightarrow \ell^\pm X)$$
$$= \Gamma(\overline{B}^0 \rightarrow \ell^+ X \text{ (via } B^0))/\Gamma(\overline{B}^0 \rightarrow \ell^\pm X)$$

Where experiments have measured the parameter $r = \chi/(1-\chi)$, we have converted to χ. Mixing violates the $\Delta B \neq 2$ rule.

Note that the measurement of χ at energies higher than the $\Upsilon(4S)$ have not separated χ_d from χ_s where the subscripts indicate $B^0(\overline{b}d)$ or $B^0_s(\overline{b}s)$. They are listed in the B^0_s-$\overline{B}^0_s$ MIXING section.

The experiments at $\Upsilon(4S)$ make an assumption about the $B^0\overline{B}^0$ fraction and about the ratio of the $B^\pm$ and B^0 semileptonic branching ratios (usually that it equals one).

OUR EVALUATION, provided by the LEP B Oscillation Working Group, includes χ_d calculated from Δm_{B^0} and τ_{B^0}.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
0.181±0.004 OUR EVALUATION					
0.182±0.015 OUR AVERAGE					
0.198±0.013±0.014		[415] BEHRENS	00B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.16 ±0.04 ±0.04		[416] ALBRECHT	94	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.149±0.023±0.022		[417] BARTELT	93	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.171±0.048		[418] ALBRECHT	92L	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.20 ±0.13 ±0.12		[419] ALBRECHT	96D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.19 ±0.07 ±0.09		[420] ALBRECHT	96D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.24 ±0.12		[421] ELSEN	90	JADE	e^+e^- 35–44 GeV
$0.158^{+0.052}_{-0.059}$		ARTUSO	89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.17 ±0.05		[422] ALBRECHT	87I	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<0.19	90	[423] BEAN	87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<0.27	90	[424] AVERY	84	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[415] BEHRENS 00B uses high-momentum lepton tags and partially reconstructed $\overline{B}^0 \rightarrow D^{*+}\pi^-$, ρ^- decays to determine the flavor of the B meson.

[416] ALBRECHT 94 reports r=0.194 ± 0.062 ± 0.054. We convert to χ for comparison. Uses tagged events (lepton + pion from D^*).

[417] BARTELT 93 analysis performed using tagged events (lepton+pion from D^*). Using dilepton events they obtain $0.157 \pm 0.016^{+0.033}_{-0.028}$.

[418] ALBRECHT 92L is a combined measurement employing several lepton-based techniques. It uses all previous ARGUS data in addition to new data and therefore supersedes ALBRECHT 87I. A value of r = 20.6 ± 7.0% is directly measured. The value can be used to measure $x = \Delta M/\Gamma = 0.72 \pm 0.15$ for the B_d meson. Assumes $f_{+-}/f_0 = 1.0 \pm 0.05$ and uses $\tau_{B^\pm}/\tau_{B^0} = (0.95 \pm 0.14)\ (f_{+-}/f_0)$.

[419] Uses $D^{*+}K^\pm$ correlations.

[420] Uses $(D^{*+}\ell^-)\ K^\pm$ correlations.

[421] These experiments see a combination of B_s and B_d mesons.

[422] ALBRECHT 87I is inclusive measurement with like-sign dileptons, with tagged B decays plus leptons, and one fully reconstructed event. Measures r=0.21 ± 0.08. We convert to χ for comparison. Superseded by ALBRECHT 92L.

[423] BEAN 87B measured $r <$ 0.24; we converted to χ.
[424] Same-sign dilepton events. Limit assumes semileptonic BR for B^+ and B^0 equal. If $B^0/B^\pm$ ratio <0.58, no limit exists. The limit was corrected in BEAN 87B from $r <$ 0.30 to $r <$ 0.37. We converted this limit to χ.

$\Delta m_{B^0} = m_{B^0_H} - m_{B^0_L}$

$\Delta m_{B^0_s}$ is a measure of 2π times the B^0-$\overline{B}^0$ oscillation frequency in time-dependent mixing experiments.

The second "OUR EVALUATION" (0.489 ± 0.009) is an average of the data listed below performed by the LEP B Oscillation Working Group as described in our "Review of B-$\overline{B}$ Mixing" in the B^0 Section of these Listings. The averaging procedure takes into account correlations between the measurements.

The first "OUR EVALUATION" (0.489 ± 0.008), also provided by the LEP B Oscillation Working Group, includes Δm_d calculated from χ_d measured at $\Upsilon(4S)$.

VALUE (10^{12} $\hbar$ s^{-1})	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.489±0.008 OUR EVALUATION					
0.489±0.009 OUR EVALUATION					
0.516±0.016±0.010		[425] AUBERT	02I	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.493±0.012±0.009		[426] AUBERT	02J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.463±0.008±0.016		[426] ABE	01D	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
0.497±0.024±0.025		[427] ABBIENDI,G	00B	OPAL	$e^+e^- \rightarrow Z$
0.503±0.064±0.071		[428] ABE	99K	CDF	$p\overline{p}$ at 1.8 TeV
0.500±0.052±0.043		[429] ABE	99Q	CDF	$p\overline{p}$ at 1.8 TeV
$0.516\pm0.099^{+0.029}_{-0.035}$		[430] AFFOLDER	99C	CDF	$p\overline{p}$ at 1.8 TeV
$0.471^{+0.078}_{-0.068}{}^{+0.033}_{-0.034}$		[431] ABE	98C	CDF	$p\overline{p}$ at 1.8 TeV
0.458±0.046±0.032		[432] ACCIARRI	98D	L3	$e^+e^- \rightarrow Z$
0.437±0.043±0.044		[433] ACCIARRI	98D	L3	$e^+e^- \rightarrow Z$
0.472±0.049±0.053		[434] ACCIARRI	98D	L3	$e^+e^- \rightarrow Z$
0.523±0.072±0.043		[435] ABREU	97N	DLPH	$e^+e^- \rightarrow Z$
0.493±0.042±0.027		[433] ABREU	97N	DLPH	$e^+e^- \rightarrow Z$
0.499±0.053±0.015		[436] ABREU	97N	DLPH	$e^+e^- \rightarrow Z$
0.480±0.040±0.051		[432] ABREU	97N	DLPH	$e^+e^- \rightarrow Z$
$0.444\pm0.029^{+0.020}_{-0.017}$		[433] ACKERSTAFF	97U	OPAL	$e^+e^- \rightarrow Z$
$0.430\pm0.043^{+0.028}_{-0.030}$		[432] ACKERSTAFF	97V	OPAL	$e^+e^- \rightarrow Z$
0.482±0.044±0.024		[437] BUSKULIC	97D	ALEP	$e^+e^- \rightarrow Z$
0.404±0.045±0.027		[433] BUSKULIC	97D	ALEP	$e^+e^- \rightarrow Z$
0.452±0.039±0.044		[432] BUSKULIC	97D	ALEP	$e^+e^- \rightarrow Z$
0.539±0.060±0.024		[438] ALEXANDER	96V	OPAL	$e^+e^- \rightarrow Z$
$0.567\pm0.089^{+0.029}_{-0.023}$		[439] ALEXANDER	96V	OPAL	$e^+e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.444 \pm 0.028 \pm 0.028$		[440] ACCIARRI	98D	L3	$e^+e^- \rightarrow Z$
0.497 ± 0.035		[441] ABREU	97N	DLPH	$e^+e^- \rightarrow Z$
$0.467 \pm 0.022 {}^{+0.017}_{-0.015}$		[442] ACKERSTAFF	97V	OPAL	$e^+e^- \rightarrow Z$
0.446 ± 0.032		[443] BUSKULIC	97D	ALEP	$e^+e^- \rightarrow Z$
$0.531 {}^{+0.050}_{-0.046} \pm 0.078$		[444] ABREU	96Q	DLPH	Sup. by ABREU 97N
$0.496 {}^{+0.055}_{-0.051} \pm 0.043$		[432] ACCIARRI	96E	L3	Repl. by ACCIARRI 98D
$0.548 \pm 0.050 {}^{+0.023}_{-0.019}$		[445] ALEXANDER	96V	OPAL	$e^+e^- \rightarrow Z$
0.496 ± 0.046		[446] AKERS	95J	OPAL	Repl. by ACKERSTAFF 97V
$0.462 {}^{+0.040}_{-0.053} {}^{+0.052}_{-0.035}$		[432] AKERS	95J	OPAL	Repl. by ACKERSTAFF 97V
$0.50 \pm 0.12 \pm 0.06$		[435] ABREU	94M	DLPH	Sup. by ABREU 97N
$0.508 \pm 0.075 \pm 0.025$		[438] AKERS	94C	OPAL	Repl. by ALEXANDER 96V
$0.57 \pm 0.11 \pm 0.02$	153	[439] AKERS	94H	OPAL	Repl. by ALEXANDER 96V
$0.50 {}^{+0.07}_{-0.06} {}^{+0.11}_{-0.10}$		[432] BUSKULIC	94B	ALEP	Sup. by BUSKULIC 97D
$0.52 {}^{+0.10}_{-0.11} {}^{+0.04}_{-0.03}$		[439] BUSKULIC	93K	ALEP	Sup. by BUSKULIC 97D

[425] Uses a tagged sample of fully-reconstructed neutral B decays at $\Upsilon(4S)$.

[426] Measured based on the time evolution of dilepton events in $\Upsilon(4S)$ decays.

[427] Data analyzed using partially reconsturcted $\overline{B}^0 \rightarrow D^{*+}\ell^-\overline{\nu}$ decay and a combination of flavor tags from the rest of the event.

[428] Uses di-muon events.

[429] Uses jet-charge and lepton-flavor tagging.

[430] Uses $\ell^- D^{*+} - \ell$ events.

[431] Uses π-B in the same side.

[432] Uses ℓ-ℓ.

[433] Uses ℓ-Q_{hem}.

[434] Uses ℓ-ℓ with impact parameters.

[435] Uses $D^{*\pm}$-Q_{hem}.

[436] Uses $\pi_s^{\pm}\,\ell$-Q_{hem}.

[437] Uses $D^{*\pm}$-ℓ/Q_{hem}.

[438] Uses $D^{*\pm}\,\ell$-Q_{hem}.

[439] Uses $D^{*\pm}$-ℓ.

[440] ACCIARRI 98D combines results from ℓ-ℓ, ℓ-Q_{hem}, and ℓ-ℓ with impact parameters.

[441] ABREU 97N combines results from $D^{*\pm}$-Q_{hem}, ℓ-Q_{hem}, $\pi_s^{\pm}\,\ell$-Q_{hem}, and ℓ-ℓ.

[442] ACKERSTAFF 97V combines results from ℓ-ℓ, ℓ-Q_{hem}, D^*-ℓ, and $D^{*\pm}$-Q_{hem}.

[443] BUSKULIC 97D combines results from $D^{*\pm}$-ℓ/Q_{hem}, ℓ-Q_{hem}, and ℓ-ℓ.

[444] ABREU 96Q analysis performed using lepton, kaon, and jet-charge tags.

[445] ALEXANDER 96V combines results from $D^{*\pm}$-ℓ and $D^{*\pm}\,\ell$-Q_{hem}.

[446] AKERS 95J combines results fromt charge measurement, $D^{*\pm}\,\ell$-Q_{hem} and ℓ-ℓ.

$x_d = \Delta m_{B^0}/\Gamma_{B^0}$

The second "OUR EVALUATION" (0.755 ± 0.015) is an average of the data listed in Δm_{B^0} section performed by the LEP B Oscillation Working Group as described in our "Review of B-$\overline{B}$ Mixing" in the B^0 Section of these Listings. The averaging procedure takes into account correlations between the measurements.

The first "OUR EVALUATION" (0.755 ± 0.015), also provided by the LEP B Oscillation Working Group, includes χ_d measured at $\Upsilon(4S)$.

VALUE	*DOCUMENT ID*
0.755±0.015 OUR EVALUATION	
0.755±0.015 OUR EVALUATION	

CP VIOLATION IN *B* DECAY – STANDARD MODEL PREDICTIONS

Revised January 2002 by H. Quinn (SLAC) and A.I. Sanda (Nagoya University).

With the commissioning of the asymmetric B Factories at KEKB and PEP II, and of CESR III and with the completion of the main ring injector at Fermilab, we are headed into an exciting time for the study of CP violation in B meson decays. This review outlines the basic ideas of such studies. For the most part, we follow the discussions given in Refs. [1–3].

Time evolution of neutral B meson states

Neutral B mesons, like neutral K mesons, have mass eigenstates which are not flavor eigenstates. This subject is reviewed separately [4]. Here we give some formulae to establish the notation used in this review. The mass eigenstates are given by:

$$|B_1\rangle = p|B^0\rangle + q|\overline{B}^0\rangle \ ,$$
$$|B_2\rangle = p|B^0\rangle - q|\overline{B}^0\rangle \ , \tag{1}$$

where B^0 and $\overline{B}^0$ are flavor eigenstates containing the $\overline{b}$ and b quarks respectively. The ratio

$$\frac{q}{p} = +\sqrt{\frac{M_{12}^* - \frac{i}{2}\Gamma_{12}^*}{M_{12} - \frac{i}{2}\Gamma_{12}}} \ . \tag{2}$$

Here, the CP operator is defined so that $CP|B^0\rangle = |\overline{B}^0\rangle$, and CPT symmetry is assumed. We define $M_{12} = \overline{M}_{12}e^{i\xi}$, where the phase ξ is restricted to $-\frac{1}{2}\pi < \xi < \frac{1}{2}\pi$, and $\overline{M}_{12}$ is taken to be real but not necessarily positive; and similarly (with a different phase) for Γ_{12}. The convention used here is that the real part of q/p is positive.

The differences in the eigenvalues $\Delta M = M_2 - M_1$ and $\Delta\Gamma = \Gamma_1 - \Gamma_2$ are given by

$$\begin{aligned}
\Delta M &= -2\mathrm{Re}\left(\frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12})\right) \\
&\simeq -2\overline{M}_{12} \\
\Delta\Gamma &= -4\mathrm{Im}\left(\frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12})\right) \\
&\simeq 2\overline{\Gamma}_{12}\cos\zeta \ .
\end{aligned} \tag{3}$$

Here we denoted $\dfrac{\Gamma_{12}}{M_{12}} = re^{i\zeta}$. As we expect $r \sim 10^{-3}$ in the Standard Model for B_d, we kept only the leading order term in r. In the Standard Model, with these conventions and given that all models give a positive value for the parameter B_B, ΔM is positive, so that B_2 is heavier than B_1; this is unlikely to be tested soon. (Note that a common alternative convention is to name the two states B_L and B_H for light and heavy respectively; then the sign of q/p becomes the quantity to be tested.)

This review focuses on the B_d system, but also mentions some possibly interesting studies for CP violation in B_s decays, which may be pursued at hadron colliders. Much of the discussion here can be applied directly for B_s decays with the appropriate replacement of the spectator quark type.

The time evolution of states starting out at time $t = 0$ as pure B^0 or $\overline{B}^0$ is given by:

$$|B^0(t)\rangle = g_+(t)|B^0\rangle + \frac{q}{p}\,g_-(t)|\overline{B}^0\rangle$$
$$|\overline{B}^0(t)\rangle = g_+(t)|\overline{B}^0\rangle + \frac{p}{q}\,g_-(t)|B^0\rangle, \tag{4}$$

where

$$g_\pm(t) = \frac{1}{2}e^{-iM_1t}e^{-\frac{1}{2}\Gamma_1 t}\left[1 \pm e^{-i\Delta Mt}e^{\frac{1}{2}\Delta\Gamma t}\right] . \tag{5}$$

We define

$$A(f) = \langle f|H|B^0\rangle \ ,$$
$$\overline{A}(f) = \langle f|H|\overline{B}^0\rangle \ ,$$
$$\overline{\rho}(f) = \frac{\overline{A}(f)}{A(f)} = \rho(f)^{-1} \ , \tag{6}$$

where f is a final state that is possible for both B^0 and $\overline{B}^0$ decays. The time-dependent decay rates are thus given by

$$\Gamma(B^0(t) \to f)$$
$$\propto e^{-\Gamma_1 t}|A(f)|^2\Bigg[K_+(t) + K_-(t)\left|\frac{q}{p}\right|^2|\overline{\rho}(f)|^2$$
$$+\,2\mathrm{Re}\Big[L^*(t)\left(\frac{q}{p}\right)\overline{\rho}(f)\Big]\Bigg] \ , \tag{7}$$

$$\Gamma(\overline{B}^0(t) \to f)$$
$$\propto e^{-\Gamma_1 t}|\overline{A}(f)|^2\Bigg[K_+(t) + K_-(t)\left|\frac{p}{q}\right|^2|\rho(f)|^2$$
$$+\,2\mathrm{Re}\left[L^*(t)\left(\frac{p}{q}\right)\rho(f)\right]\Bigg], \tag{8}$$

where

$$|g_\pm(t)|^2 = \frac{1}{4} e^{-\Gamma_1 t} K_\pm(t) \ ,$$

$$g_-(t) g_+^*(t) = \frac{1}{4} e^{-\Gamma_1 t} L^*(t) \ ,$$

$$K_\pm(t) = 1 + e^{\Delta\Gamma t} \pm 2 e^{\frac{1}{2}\Delta\Gamma t} \cos \Delta M t \ ,$$

$$L^*(t) = 1 - e^{\Delta\Gamma t} + 2 i e^{\frac{1}{2}\Delta\Gamma t} \sin \ \Delta M t \ . \tag{9}$$

For the case of B_d decays the quantity $\Delta\Gamma/\Gamma$ is small and is usually dropped, for B_s decays it may be significant [6] and hence is retained in Eqs. 4–8.

Three classes of CP violation in B decays

When two amplitudes with different phase-structure contribute to a B decay, they may interfere and produce CP-violating effects [5]. There are three distinct types of CP violation: (1) CP violation from nonvanishing relative phase between the mass and the width parts of the mixing matrix which gives $|q/p| \neq 1$, often called "indirect;" (2) Direct CP violation, which is any effect that indicates two decay amplitudes have different weak phases (those arising from Lagrangian couplings), in particular it occurs whenever $|\rho(f)| \neq 1$; (3) Interference between a decays with and without mixing which can occur for decays to CP eigenstates whenever $\mathrm{Arg}((q/p)\overline{\rho}(f)) \neq 0$. This can occur even for modes where both the other types do not, *i.e.* $|q/p|, \ |\rho(f)| = 1$.

(1) Indirect CP violation

In the next few years, experiments will accumulate a large number of semileptonic B decays. Any asymmetry in the wrong-sign semileptonic decays (or in any other wrong-flavor decays) is a clean sign of indirect CP violation.

The semileptonic asymmetry for the wrong sign B_q decay, where $q = d$ or s, is given by

$$a_{SL}(B_q) = \frac{\Gamma(\overline{B}_q(t) \to \ell^+ X) - \Gamma(B_q(t) \to \ell^- X)}{\Gamma(\overline{B}_q(t) \to \ell^+ X) + \Gamma(B_q(t) \to \ell^- X)}$$
$$= \frac{|p/q|^2 - |q/p|^2}{|p/q|^2 + |q/p|^2} = r_{B_q} \sin\zeta_{B_q} \ , \tag{10}$$

where we kept only the leading order term in r_{B_q}. Within the context of the Standard Model, if hadronic rescattering effects are small then $\sin\zeta_{B_q}$ is small because M_{12} and Γ_{12} acquire their phases from the same combination of CKM matrix elements. Since this asymmetry is tiny in the Standard Model, this may be a fruitful area to search for physics beyond the Standard Model.

(2) Direct CP violation

Direct CP violation is the name given to CP violation that arises because there is a difference between the weak phases of any two decay amplitudes for a single decay. Weak phases are those that arise because of a complex coupling constant in the Lagrangian. Note that a single weak phase from a complex coupling constant is never physically meaningful because it can generally be removed by redefining some field by a phase. Only the differences between the phases of couplings which cannot be changed by such redefinitions are physically meaningful. The strong and electromagnetic couplings can always be defined to be real but, as Kobayashi and Maskawa first observed, in the three generation Standard Model one cannot remove all the phases from the CKM matrix by any choice of field redefinitions [7].

There are two distinct ways to observe direct CP-violation effects in B decays:

• $|\overline{A}_{\overline{f}}/A_f| \neq 1$ leading to rate asymmetries for CP-conjugate decays. Here, two amplitudes with different weak phases must contribute to the same decay; they must also have different strong phases, that is, the phases that arise because of absorptive parts (often called final-state interaction effects). When the final state f has different flavor content than its CP conjugate, this gives a rate asymmetry that is directly observable. The asymmetry is given by

$$a = \frac{2A_1A_2\sin(\xi_1 - \xi_2)\sin(\delta_1 - \delta_2)}{A_1^2 + A_2^2 + 2A_1A_2\cos(\xi_1 - \xi_2)\cos(\delta_1 - \delta_2)} \ , \tag{11}$$

where the A_i are the magnitudes, the ξ_i are the weak phases, and the δ_i are the strong phases of the two amplitudes contributing to A_f. The impact of direct CP violation of this type in decays of neutral B's to flavor eigenstates is discussed below.

• Any difference (other than an overall sign) between the CP asymmetries for decays of B_d mesons to flavor eigenstates, or between those of neutral B_s mesons, is an evidence of direct CP violation. As is shown below, such asymmetries arise whenever the decay weak phase is not canceled by the mixing weak phase, hence any two different results imply that there is a difference between the weak phases of the amplitudes for the two decays. Only if the asymmetries are the same can one choose a phase convention which ascribes all CP-violating phases to the mixing amplitude. For example, the expected asymmetries for the $B \to J/\psi K_S$ and $B \to \pi\pi$ decays are different (whether or not penguin graphs add additional direct CP-violating effects of the type $|\overline{A}_{\overline{f}}/A_f| \neq 1$ in the latter channel) because the dominant decay amplitudes have different weak phases in the Standard Model.

(3) Decays of B^0 and $\overline{B}^0$ to CP eigenstates

In decays to CP eigenstates, the time-dependent asymmetry is given by

$$a_f(t) = \frac{\Gamma(\overline{B}^0(t) \to f) - \Gamma(B^0(t) \to f)}{\Gamma(\overline{B}^0(t) \to f) + \Gamma(B^0(t) \to f)} \; . \tag{12}$$

Asymmetry is generated if: (i) both $A(B \to f)$ and $A(\overline{B} \to f)$ are nonzero; and (ii) the mixing weak phase in $\frac{q}{p}$ is different from the weak decay phase in $\overline{\rho}(f)$. To the leading order in r, the Standard Model predicts

$$q/p = \frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*} = e^{-i2\phi_{\text{mixing}}} \; . \tag{13}$$

If there is only one amplitude (or two with the same weak phase) contributing to $A(B \to f)$ and $A(\overline{B} \to f)$ then $|\overline{\rho}(f)| = 1$ and the relationship between the measured asymmetry and the Kobayshi-Maskawa phases is cleanly predicted by

$$\begin{aligned} a_f(t) &= \mathrm{Im}\left(\frac{q}{p}\overline{\rho}(f)\right) \sin \Delta M t \\ &= -\eta_f \sin 2(\phi_{\text{mixing}} + \phi_{\text{decay}}) \sin \Delta M t \; . \end{aligned} \tag{14}$$

Here we have used the fact that in such cases we can write $\overline{\rho}(f) = \eta_f e^{-i2\phi_{\text{decay}}}$ where $\eta_f = \pm$ is the CP eigenvalue of the state f. The weak phases ϕ_{mixing} and ϕ_{decay} are parameterization dependent quantities, but the combination $\phi_{\text{mixing}} + \phi_{\text{decay}}$ is parameterization independent. This is CP violation due to the interference between decays with and without mixing. Note that a single measurement of $\sin(2\phi)$ yields four ambiguous solutions for ϕ.

When more than one amplitude with different weak phases contribute to a decay to a CP eigenstate there can also be direct

CP violation effects $|\lambda_f = (q/p)\,\rho(f)| \neq 1$ and the asymmetry takes the more complicated form

$$a_f(t) = \frac{(|\lambda_f|^2 - 1)\cos(\Delta Mt) + 2\mathrm{Im}\lambda_f \sin(\Delta Mt)}{(1 + |\lambda_f|^2)} \ . \qquad (15)$$

The quantity λ_f involves the ratio of the two amplitudes that contribute to A_f as well as their relative strong phases and hence introduces the uncertainties of hadronic physics into the relationship between the measured asymmetry and the K–M phases. However in certain cases such channels can be useful in resolving the ambiguities mentioned above. If $\cos(2\phi)$ can be measured as well as $\sin(\phi)$ only a two-fold ambiguity remains. This can be resolved only by knowledge of the sign of certain strong phase shifts [8].

When a B meson decays to a CP self-conjugate set of quarks the final state is in general a mixture of CP even and CP odd states, which contribute opposite sign and hence partially canceling asymmetries. In two special cases, namely the decay to two spin zero particles, or one spin zero and one non-zero spin particle there is a unique CP eigenvalue because there is only one possible relative angular momentum between the two final state particles. Quasi-two-body modes involving two particles with non-zero spin can sometimes be resolved into contributions of definite CP by angular analysis of the decays of the "final-state" particles [9].

There can also be a direct CP violation in these channels from the interference of two contributions to the same decay amplitude, $|\rho(f)| \neq 1$. This introduces dependence on the relative strengths of the two amplitude contributions and on their relative strong phases. Since these cannot be reliably calculated at present, this complicates the attempt to relate the measured asymmetry to the phases of CKM matrix elements.

Standard Model predictions for CP-violating asymmetries

• ***Unitarity Triangles***

The requirement that the CKM matrix be unitary leads to a number of relationships among its entries. The constraints that the product of row i with the complex conjugate of row j is zero are generically referred to as "unitarity triangles" because they each take the form of a sum of three complex numbers equal to zero and hence can be represented by triangles in the complex plane. There are six such relationships, (see for example Ref. 10); the most commonly studied is that with all angles of the same order of magnitude, given by the relationship

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0 \ . \tag{16}$$

This relation can be represented as a triangle on the complex plane, as shown in Fig. **@Fg.unitangle@**, where the signs of all three angles are also defined. When the sides are scaled by $|V_{cd}V_{cb}^*|$, the apex of the triangle is the point ρ, η, where these parameters are defined by the Wolfenstein parameterization of the CKM matrix [11]. If $\eta = 0$, the CKM matrix is real and there is no CP violation in the Standard Model.

The angles of the triangle are

$$\phi_1 = \pi - \arg\left(\frac{-\mathbf{V}_{tb}^* V_{td}}{-V_{cb}^* V_{cd}}\right) = \beta \ ,$$

$$\phi_2 = \arg\left(\frac{V_{tb}^* V_{td}}{-V_{ub}^* V_{ud}}\right) = \alpha \ ,$$

$$\phi_3 = \arg\left(\frac{V_{ub}^* V_{ud}}{-V_{cb}^* V_{cd}}\right) = \gamma \ . \tag{17}$$

Two naming conventions for these angles are commonly used in the literature [12,13]; we provide the translation dictionary



Figure 1: Angles of the unitarity triangle are related to the Kobayashi-Maskawa phases of the CKM matrix. The right-hand rule gives the positive direction of the angle between two vectors. This figure was reproduced from Ref. 1 with permission from Cambridge University Press.

in Eq. (17), but use the ϕ_i notation in the remainder of this review, where ϕ_i is the angle opposite the side $V_{ib}^* V_{id}$ of the unitarity triangle and i represents the i-th up-type quark. As defined here, for consistency with the measured value of ϵ_K, these angles are all positive in the Standard Model, thus a determination of the sign of these angles constitutes a test of the Standard Model [14].

There are two other independent angles of the Standard Model which appear in other triangles. These are denoted

$$\begin{aligned} \chi &= \arg\left(\frac{-V_{cs}^* V_{cb}}{V_{ts}^* V_{tb}}\right) = \beta_s \\ \chi' &= \arg\left(\frac{-V_{ud}^* V_{us}}{V_{cd}^* V_{cs}}\right) = -\beta_K \ . \end{aligned} \tag{18}$$

Again there are two naming conventions in common usage so we give both. These angles are of order λ^2 and λ^4 respectively [15], where $\lambda = V_{us}$. The first of them is the phase of the B_s mixing and thus is in principle measurable, though it will not be easy to achieve a result significantly different from zero for such a small angle. The angle χ' will be even more difficult to measure. Meaningful standard model tests can be defined which use the measured value of λ coupled with χ and any two of the three ϕ_i [16].

A major aim of CP-violation studies of B decays is to make enough independent measurements of the sides and angles that this unitarity triangle is overdetermined, and thereby check the validity of the Standard Model predictions that relate various measurements to aspects of this triangle. Constraints can be made on the basis of present data on the B-meson mixing and lifetime, and on the ratio of charmless decays to decays with charm (V_{ub}/V_{cb}), and on ϵ in K decays [17]. These constraints have been discussed in many places in the literature; for a recent summary of the measurements involved, see Ref. [18]. Note, however, that any given "Standard Model allowed range" cannot be interpreted as a statistically-based error range. The ranges of allowed values depend on matrix element estimates. Improved methods to calculate such quantities, and understand the uncertainties in them, are needed to further sharpen tests of the Standard Model. Recent progress in lattice simulation using dynamical fermions seems encouraging [19]. It can be hoped that reliable computations of f_B, B_B, and B_K will be completed in the next few years. This will reduce the theoretical uncertainties in the relationships between measured mixing effects and the magnitudes of CKM parameters.

In the Standard Model there are only two independent phases in this triangle since, by definition, the three angles add up to π. The literature often discusses tests of whether the angles add up to π; but this really means tests of whether relationships between different measurements, predicted in terms of the two independent parameters in the Standard Model, hold true. For example, many models that go beyond the Standard Model predict an additional contribution to the mixing matrix. Any change in phase of M_{12} will change the measured asymmetries so that $\phi_1(\text{measured}) \to \phi_1 - \phi_{\text{new}}$ and $\phi_2(\text{measured}) \to \phi_2 + \phi_{\text{new}}$. Thus the requirement that the sum of the three angles must add up to π is not sensitive to ϕ_{new} [20]. However, the angles as determined from the sides of the triangle would, in general, no longer coincide with those measured from asymmetries. It is equally important to check the asymmetries in channels for which the Standard model predicts very small or vanishing asymmetries. A new mixing contribution which changes the phase of M_{12} will generate significant asymmetries in such channels. In the Standard Model the CKM matrix must be unitary, this leads to relationships among its entries.

• ***Standard Model decay amplitudes***

In the Standard Model, there are two classes of quark-level diagrams that contribute to hadronic B decays, as shown in Fig. **@Fg.penguin@**. Tree diagrams are those where the W produces an additional quark-antiquark pair. Penguin diagrams are loop diagrams where the W reconnects to the same quark line. Penguin diagrams can further be classified by the nature of the particle emitted from the loop: gluonic or QCD penguins if it is a gluon, and electroweak penguins if it is a photon or a Z boson. In addition, one can label penguin diagrams by the flavor of the up-type quark in the loop; for any process all three flavor types contribute. For some processes, there are additional annihilation-type diagrams; these always contribute to the same CKM structure as the corresponding trees. For a detailed discussion of the status of calculations based on these diagrams, or rather on the more complete operator product approach which also includes higher order QCD corrections see, for example, Ref. 21. Note that the distinction between tree and penguin contributions is a heuristic one, the separation of contributions by the operator that enters is more precise.



Figure 2: Quark level processes for the example of $b \to c\overline{c}s$. (a) Tree diagram; (b) Penguin diagram. In the case of electroweak penguin contributions, the gluon is replaced by a Z or a γ.

To explore possible CP violations, it is useful to tabulate all possible decays by the CKM structure of the various amplitudes. Let us first consider decays $b \to q\overline{q}'s$. The CKM factors for the diagrams for such decays are given in Table 1. Here we have used the fact that, for all such decays, the contribution to the amplitude from penguin graphs has the structure

$$A_P(q\overline{q}s) = V_{tb}V^*_{ts}P_t + V_{cb}V^*_{cs}P_c + V_{ub}V^*_{us}P_u \ , \qquad (19)$$

where the P_i quantities are the amplitudes described by the loop diagram with a flavor i quark apart from the explicitly shown CKM factor (*i.e.*, including strong phases). These are actually divergent quantities, so it is convenient to use a Standard Model unitarity relationship, $V_{tb}V^*_{ts} + V_{cb}V^*_{cs} + V_{ub}V^*_{us} = 0$, to regroup them in the following way

$$A_P(q\overline{q}s) = V_{cb}V^*_{cs}(P_c - P_t) + V_{ub}V^*_{us}(P_u - P_t) \ , \qquad (20)$$

or, equivalently,

$$A_P(q\overline{q}s) = V_{tb}V^*_{ts}(P_t - P_c) + V_{ub}V^*_{us}(P_u - P_c) \ . \qquad (21)$$

The first term is of order λ^2, whereas the second is of order λ^4, and can be ignored in most instances. For modes with $q' \neq q$, there are no penguin contributions. Note also that for the $q\overline{q} = u\overline{u}, d\overline{d}$ cases, the QCD penguin graphs contribute only to the isospin zero combinations, whereas tree graphs contribute only for $u\overline{u}$ and hence have both $\Delta I = 0$ and $\Delta I = 1$ parts, as do electroweak penguins.

The CKM coefficients for $b \to q\overline{q}'d$ are listed in Table 2. A similar exercise to that described above for the penguins yields

$$A_P(q\overline{q}d) = V_{tb}V^*_{td}(P_t - P_c) + V_{ub}V^*_{ud}(P_u - P_c) \ . \qquad (22)$$

Here the two CKM contributions are of the same order of magnitude λ^3, so both must be considered. This grouping is generally preferred over the alternative, because the second term here is somewhat smaller than the first term; it has no top-quark contribution and would vanish if the up and charm quarks were degenerate. In early literature it was often dropped, but, particularly for modes where there is no tree contribution, its effect in generating direct CP violation may be important [22]. Here the $q\overline{q} = u\overline{u}, d\overline{d}$ cases in the penguin graph contribute only to the isospin zero combinations, yielding $\Delta I = 1/2$ for the three-quark combination, whereas tree graphs and electroweak penguins have both $\Delta I = 1/2$ and $\Delta I = 3/2$ parts. For $q\overline{q} = c\overline{c}$, isospin does not distinguish between tree and penguin contributions.

Modes with direct CP violation

The largest direct CP violation is expected when there are two comparable magnitude contributions with different weak phases. Modes where the tree graphs are Cabibbo suppressed, compared to the penguins or modes with two comparable penguin contributions, are thus the best candidates. As can be seen from the tables and expressions for penguin contributions above, there are many possible modes to study. Because strong phases cannot usually be predicted, there is no clean prediction as to which modes will show the largest direct CP-violation effects. One interesting suggestion is to study three-body modes with more than one resonance in the same kinematic region. Then the different amplitudes can have very different, possibly known, strong phase structure because of the resonance (Breit-Wigner) phases [23].

Over the past two years, new information has become available from the CLEO Collaboration which suggests that penguin contributions, at least for some modes, are larger than initial estimates suggested. This is seen by using SU(3) and comparing $B \to K\pi$ and $B \to \pi\pi$ decays. To get an order of magnitude picture, we ignore such details as Clebsch-Gordan coefficients and assume that top penguins dominate the penguin contributions. Thus, we identify the tree and penguin contributions, minus their CKM coefficients, as T and P, the same for both modes. Writing $A_{T,P}(K\pi)$ for the tree and penguin contributions to the $K\pi$ amplitude, and similarly for $\pi\pi$ from the Tables, we see that $|A^T(K\pi)/A^T(\pi\pi)| = \mathcal{O}(\lambda)$. Thus, if the tree graph matrix elements were to dominate both decays, we would expect $\mathrm{Br}(B \to K\pi)/\mathrm{Br}(B \to \pi\pi) \sim \mathcal{O}(\lambda^2)$. Naively, this was expected, since the ratio of tree to penguin contribution was estimated to be $\dfrac{P}{T} = \dfrac{\alpha_S}{12\pi}\log\dfrac{m_t^2}{m_b^2} \sim \mathcal{O}(0.02)$. Experimentally, this is not so [24]; in fact, the $K\pi$ branching ratio is larger. This indicates that $A^P(K\pi) \sim A^T(\pi\pi)$, which suggests that $\dfrac{P}{T} = \mathcal{O}(\lambda)$ or larger, considerably bigger than expected. Note that this is one way that new physics could be hidden in modes with $|\rho(f)| \neq 1$; any new physics contribution can always be written as a sum of two terms with the weak phases of the two Standard Model terms (for example in Eq. (22)), and thus, when added to the Standard Model contributions, appears only as a change in the sizes of P and T from that expected in the Standard Model. However, we cannot calculate these relative sizes well enough to identify such an effect with confidence.

From the point of view of looking for direct CP-violation effects, a large P/T is good news. The largest asymmetry is expected when the interfering amplitudes have comparable magnitudes. This may be so in $B \to K\pi$ decay (or the penguin contribution may even be larger than the tree). There is no reason for the strong phases to be equal (although they could both be small). Therefore, $B^{\pm} \rightarrow K^{\pm}\pi$ is a likely hunting ground for direct CP violation. (Note there is no gluonic penguin contribution to charged $B \rightarrow \pi\pi$, and hence, no significant CP violation expected in the Standard Model.) However, as we will see below, a large P/T complicates the relationship between the measured asymmetry in neutral B decays to $\pi^+\pi^-$ and KM phases.

Studies of CP eigenstates

• $\boldsymbol{f = J/\psi K_S}$

The asymmetry in the "golden mode" $B \rightarrow J/\psi K_S$ has now been measured by both the BaBar and Belle experiments [25]. The Standard Model prediction for this mode is very clean. Since, using Eq. (20), the dominant penguin contribution has the same weak phase as the tree graph, and the remaining term is tiny, there is effectively only one weak phase in the decay amplitude. Hence, in the asymmetry, all dependence on the amplitudes cancel. With about 1% uncertainty,

$$\frac{q}{p}\overline{\rho}(J/\psi K_S) \simeq -\frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*} \cdot \frac{V_{cb} V_{cs}^*}{V_{cb}^* V_{cs}} \cdot \frac{V_{cs} V_{cd}^*}{V_{cs}^* V_{cd}} \equiv -e^{-2i\phi_1} \ , \qquad (23)$$

where the last factor arises from the K^0–$\overline{K}^0$ mixing amplitude and appears because of the K_S in the final state. The asymmetry is thus given by

$$a_{J/\psi K_S} = \sin(2\phi_1)\sin \Delta M t \ , \qquad (24)$$

where the angle ϕ_1 is defined in Fig. 1. The result is consistent within errors with the prediction from the Standard Model, which strongly suggests that the KM ansatz for CP violation is at least one of the sources of this interesting phenomenon.

• $\boldsymbol{B^0 \to \pi^+\pi^-}$

The tree and penguin terms appear at the same order in λ (see Eq. (22) and Table 2.) If penguin decays were negligible the asymmetry would directly measure $\sin(2\phi_2)$. Given the enhanced penguin contribution seen from comparing $\pi\pi$ and $K\pi$ decays, the penguins cannot be ignored, and a treatment that does not assume $|\rho(f)| = 1$ must be made.

If all six modes of $B^+ \to \pi^+\pi^0$, $B^0 \to \pi^+\pi^-$, $B^0 \to \pi^0\pi^0$ and their charge conjugates can be measured with sufficient accuracy, ϕ_2 can be extracted using an isospin analysis [26], up to small corrections from electroweak penguins. However, the branching ratio for the charged modes is less than 10^{-5} [24], and that for the more difficult to measure $B^0 \to \pi^0\pi^0$ is expected to be even smaller. Therefore, further ingenuity is needed to get at this angle cleanly. A future possibility is to study the Dalitz plot of $B \to 3\pi$ decays [27].

To date only upper limits on CP-violating asymmetries in this mode have been reported [25].

Further Measurements

As Tables 1 and 2 suggest there are many more CP-eigenstate modes that are interesting to study, both for B_d and similarly for B_s decays. The latter states are not accessible for the B factories operating at the $\Upsilon(4S)$ resonance, but may be studied at hadronic colliders. The CDF result on the asymmetry in the $J/\psi K_S$ mode is an indication of the capabilities of such facilities for B physics [29]. Upgrades of the Fermilab detectors are in progress and proposals for new detectors with the capability to achieve fast triggers for a larger variety of purely hadronic modes are under development, promising some future improvement in this capability.

In addition to CP-eigenstate modes there are many additional modes for which particular studies have been proposed, in particular those focussed on extracting ϕ_3 (γ). Modes such as DK, DK^* and D^*K where the D mesons decay to CP eigenstates provide theoretically clean extraction of this parameter but have small branching ratios [30]. Other approaches involve the more copious $K\pi$ modes but rely on the use of isospin and SU(3) (U-spin) symmetries, so have larger theoretical uncertainties [31]. This is an active area of current theoretical work.

For a recent review of how predictions for CP-violating effects are affected by Beyond Standard Model effects see Ref. 28. There are also many ways to search for new physics effects in B decays that do not involve just the CP-violation effects. For example searches for isospin breaking effects in $K\pi$ modes have recently been suggested as a likely method to isolate such effects [32].

Table 1: $B \to q\overline{q}s$ decay modes

Quark process	Leading term	Secondary term	Sample B_d modes	B_d angle	Sample B_s modes	B_s angle
$b \to c\overline{c}s$	$V_{cb}V_{cs}^* = A\lambda^2$ tree + penguin$(c-t)$	$V_{ub}V_{us}^* = A\lambda^4(\rho - i\eta)$ penguin only$(u-t)$	$J/\psi\ K_S$	β	$J/\psi\eta$ $D_s\overline{D}_s$	0
$b \to s\overline{s}s$	$V_{cb}V_{cs}^* = A\lambda^2$ penguin only$(c-t)$	$V_{ub}V_{us}^* = A\lambda^4(\rho - i\eta)$ penguin only$(u-t)$	ϕK_S	β	$\phi\eta'$	0
$b \to u\overline{u}s$ $b \to d\overline{d}s$	$V_{cb}V_{cs}^* = A\lambda^2$ penguin only$(c-t)$	$V_{ub}V_{us}^* = A\lambda^4(\rho - i\eta)$ tree + penguin$(u-t)$	$\pi^0\ K_S$ ρK_S	competing terms	$\phi\pi^0$ $K_S\overline{K}_S$	competing terms

Table 2: $B \to q\overline{q}d$ decay modes

Quark process	Leading term	Secondary term	Sample B_d modes	B_d angle	Sample B_s modes	B_s angle
$b \to c\overline{c}d$	$V_{cb}V_{cd}^* = -A\lambda^3$ tree + penguin$(c-u)$	$V_{tb}V_{td}^* = A\lambda^3(1-\rho+i\eta)$ penguin only$(t-u)$	D^+D^-	$^*\beta$	$J/\psi K_S$	$^*\beta_s$
$b \to s\overline{s}d$	$V_{tb}V_{td}^* = A\lambda^3(1-\rho+i\eta)$ penguin only$(t-u)$	$V_{cb}V_{cd}^* = A\lambda^3$ penguin only$(c-u)$	$\phi\pi$ $K_S\overline{K}_S$	competing terms	ϕK_S	competing terms
$b \to u\overline{u}d$ $b \to d\overline{d}d$	$V_{ub}V_{ud}^* = A\lambda^3(\rho - i\eta)$ tree + penguin$(u-c)$	$V_{tb}V_{td}^* = A\lambda^3(1-\rho+i\eta)$ penguin only$(t-c)$	$\pi\pi$; $\pi\rho$ $\pi\, a_1$	$^*\alpha$	π^0K_S ρ^0K_S	competing terms
$b \to c\overline{u}d$	$V_{cb}V_{ud}^* = A\lambda^2$	0	$D^0\pi^0$, $D^0\rho^0$ $\to CP$ eigenstate	β	D^0K_S $\to CP$ eigenstate	0

*Leading terms only, large secondary terms shift asymmetry.

References

1. Much of what is presented is explained in further detail in I.I. Bigi and A.I. Sanda, *CP Violation*, Cambridge University Press, Cambridge (1999).
2. For a review of CP violation in B decays see: I.I. Bigi, V.A. Khoze, N.G. Uraltsev, and A.I. Sanda, in *CP Violation*, ed. C. Jarlskog (World Scientific, Singapore, 1989).
3. Y. Nir and H.R. Quinn, Ann. Rev. Nucl. Sci. **42**, 211 (1992).
4. See our review on "B^0–$\overline{B}^0$ Mixing" by O. Schneider in the B^0 Listings in the full *Review*.
5. A.B. Carter and A.I. Sanda, Phys. Rev. Lett. **45**, 952 (1980); Phys. Rev. **D23**, 1567 (1981).
6. R. Aleksan *et al.*, Phys. Lett. **B316**, 567 (1993); M. Beneke, G. Buchalla, I. Duniety, Phys. Rev. **D54**, 4419 (1996).
7. See our review on the CKM mixing matrix by F.J. Gilman, K. Kleinknecht and B. Renk in the full *Review*.
8. Y. Grossman and H. Quinn, Phys. Rev. **D56**, 7259 (1997).
9. J.R. Dell'Aquila and C.A. Nelson, Phys. Rev. **D33**, 101 (1986); B. Kayser *et al.*, Phys. Lett. **B237**, 3339 (1990); I. Dunietz *et al.*, Phys. Rev. **D43**, 2193 (1991).
10. R. Alexsan, B. Kayser, and D. London, Phys. Rev. Lett. **73**, 18 (1994).
11. L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).
12. The angles of the unitarity triangle ϕ_1, ϕ_2, ϕ_3 are those used in J.L. Rosner, A.I. Sanda, and M. Schmidt in *Proc. Fermilab Workshop on High Sensitivity Beauty Physics at Fermilab*, A.J. Slaughter, N. Lockyer and M. Schmidt, eds., 1987.
13. C.O. Dib *et al.*, Phys. Rev. **D41**, 1522 (1990). See also Y. Nir SLAC Report 412 p. 102 (1992) for careful definition of the signs in the α, β, γ notation.
14. Y. Nir and H. Quinn, Phys. Rev. **D42**, 1473 (1990).

15. I.I. Bigi and A.I. Sanda, Nucl. Phys. **B193**, 85 (1981); I.I. Bigi, in *Proceedings of the Tau-Charm Workshop*, SLAC-Report 343, p. 169 (1989).
16. J.P. Silva and L. Wolfenstein Phys. Rev. **D55**, 5331 (1997).
17. See our review on "CP Violation" by L. Wolfenstein in the full *Review*.
18. A.F. Falk, talk presented at *19th International Symposium on Lepton and Photon Interactions at High-Energies (LP 99)* (Stanford, CA 9–14 Aug 1999) Int. J. Mod. Phys. **A15S1**, 174 (2000).
19. S. Aoki, talk presented at *19th International Symposium on Lepton and Photon Interactions at High-Energies (LP 99)* (Stanford, CA, 9–14 Aug 1999).
20. See for example Y. Nir and D. Silverman, Nucl. Phys. **B345**, 301 (1990).
21. G. Buchalla, A J. Buras, M E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
22. J.M. Gérard and W.S. Hou, Phys. Rev. **D43**, 2909 (1991) and Phys. Lett. **B253**, 478 (1991); H. Simma, G. Eilam, and D. Wyler, Nucl. Phys. **B352**, 367 (1991); R. Fleischer, Phys. Lett. **B341**, 205 (1994).
23. D. Atwood and A. Soni, Z. Phys. **C64**, 241 (1994); Phys. Rev. Lett. **74**, 220 (1995).
24. Y. Kwon *et al.*, CLEO-CONF-99–14, Talk given at *19th International Symposium on Lepton and Photon Interactions at High-Energies (LP 99)* (Stanford, CA, 9–14 Aug 1999) `hep-ex/9908039`.
25. See the Listings for the B^0 in this *Review*.
26. M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990).
27. H.R. Quinn and A.E. Snyder, Phys. Rev. **48**, 2139 (1993).
28. CP Violation beyond the Standard Model. By Y. Grossman, Y. Nir, R. Rattazzi, SLAC-PUB-7379, Jan 1997. In *Heavy flavours II* p. 755, A.J. Buras, M. Lindner, eds., e-Print Archive: `hep-ph/9701231`.

29. CDF Collaboration: T. Affolder *et al.*, Phys. Rev. **D61**, 072005 (2000).
30. I. Dunietz, Phys. Lett. **B270**, 75 (1991);
M. Gronau and D. Wyler, Phys. Lett. **B265**, 172 (1991);
D. Atwood, I. Dunietz, and A. Soni, Phys. Rev. Lett. **78**, 3257 (1997).
31. See for example M. Gronau, J.L. Rosner, and D. London, Phys. Rev. Lett. **73**, 21 (1994);
M. Gronau, O.F. Hernandez, D. London, and J.L. Rosner Phys. Rev. **D50**, 4529 (1994);
M. Gronau and J.L. Rosner, Phys. Rev. Lett. **76**, 1200 (1996);
R. Fleischer and T. Mannel, Phys. Rev. **D57**, 2752 (1998);
M. Neubert and J. L. Rosner, Phys. Lett. **B441**, 403 (1998) and Phys. Rev. Lett. **81**, 5076 (1998);
M. Neubert, Invited talk at *High Energy Physics International Euroconference on Quantum Chromo Dynamics (QCD '99)* (Montpellier, France, 7–13 July 1999) Nucl. Phys. Proc. Supp. **86**, 477 (2000).
32. Y. Grossman, M. Neubert, and A.L. Kagan, JHEP **9910**, 029 (1999).

CP VIOLATION PARAMETERS

$\mathrm{Re}(\epsilon_{B^0})/(1+|\epsilon_{B^0}|^2)$

CP impurity in B^0_d system. It is obtained from either $a_{\ell\ell}$, the charge asymmetry in like-sign dilepton events or a_{cp}, the time-dependent asymmetry of inclusive B^0 and $\overline{B}^0$ decays.

VALUE (units 10^{-3})	*DOCUMENT ID*		*TECN*	*COMMENT*
0 ± 4 OUR AVERAGE				
-3 ± 7	[447] BARATE	01D	ALEP	$e^+e^-\rightarrow Z$
$3.5\pm10.3\pm1.5$	[448] JAFFE	01	CLE2	$e^+e^-\rightarrow \Upsilon(4S)$
$1\pm14\pm3$	[449] ABBIENDI	99J	OPAL	$e^+e^-\rightarrow Z$
$2\pm7\pm3$	[450] ACKERSTAFF	97U	OPAL	$e^+e^-\rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$4\pm18\pm3$	[451] BEHRENS	00B	CLE2	Repl. by JAFFE 01
<45	[452] BARTELT	93	CLE2	$e^+e^-\rightarrow \Upsilon(4S)$

[447] BARATE 01D measured by investigating time-dependent asymmetries in semileptonic and fully inclusive B^0_d decays.

[448] JAFFE 01 finds $a_{\ell\ell} = 0.013 \pm 0.050 \pm 0.005$ and combines with the previous BEHRENS 00B independent measurement.

[449] Data analyzed using the time-dependent asymmetry of inclusive B^0 decay. The production flavor of B^0 mesons is determined using both the jet charge and the charge of secondary vertex in the opposite hemisphere.

[450] ACKERSTAFF 97U assumes CPT and is based on measuring the charge asymmetry in a sample of B^0 decays defined by lepton and Q_{hem} tags. If CPT is not invoked, $\text{Re}(\epsilon_B) = -0.006 \pm 0.010 \pm 0.006$ is found. The indirect CPT violation parameter is determined to $\text{Im}(\delta B) = -0.020 \pm 0.016 \pm 0.006$.

[451] BEHRENS 00B uses high-momentum lepton tags and partially reconstructed $\overline{B}^0 \to D^{*+}\pi^-$, ρ^- decays to determine the flavor of the B meson.

[452] BARTELT 93 finds $a_{\ell\ell} = 0.031 \pm 0.096 \pm 0.032$ which corresponds to $|a_{\ell\ell}| < 0.18$, which yields the above $|\text{Re}(\epsilon_{B^0})/(1+|\epsilon_{B^0}|^2|$.

A_{CP} ($B^0 \to K^+\pi^-$)

A_{CP} is defined as

$$\frac{B(\overline{B}^0 \to \overline{f}) - B(B^0 \to f)}{B(\overline{B}^0 \to \overline{f}) + B(B^0 \to f)},$$

the CP-violation charge asymmetry of inclusive B^0 and $\overline{B}^0$ decay.

VALUE	DOCUMENT ID		TECN	COMMENT
-0.09 ± 0.06 OUR AVERAGE				
$-0.07 \pm 0.08 \pm 0.02$	[453] AUBERT	02D	BABR	$e^+e^- \to \Upsilon(4S)$
$0.044^{+0.186}_{-0.167}{}^{+0.018}_{-0.021}$	[454] ABE	01K	BELL	$e^+e^- \to \Upsilon(4S)$
$-0.19 \pm 0.10 \pm 0.03$	[455] AUBERT	01E	BABR	$e^+e^- \to \Upsilon(4S)$
-0.04 ± 0.16	[456] CHEN	00	CLE2	$e^+e^- \to \Upsilon(4S)$

[453] Corresponds to 90% confidence range $-0.21 < A_{CP} < 0.07$.

[454] Corresponds to 90% confidence range $-0.25 < A_{CP} < 0.37$.

[455] Corresponds to 90% confidence range $-0.35 < A_{CP} < -0.03$.

[456] Corresponds to 90% confidence range $-0.30 < A_{CP} < 0.22$.

$A_{CP}(B^0 \to \phi K^*(892)^0)$

VALUE	DOCUMENT ID		TECN	COMMENT
$0.00 \pm 0.27 \pm 0.03$	[457] AUBERT	02E	BABR	$e^+e^- \to \Upsilon(4S)$

[457] Corresponds to 90% confidence range $-0.44 < A_{CP} < 0.44$.

$C_{\pi\pi}$ ($B^0 \to \pi^+\pi^-$)

$C_{\pi\pi}$ is defined as $(1-|\lambda|^2)/(1+|\lambda|^2)$, where the quantity $\lambda = (q/p)\rho$, involves the ratio of the two amplitudes with different phases that contribute to a deceay to a CP eigenstate. For details, see the note on "CP Violation in B Decay Standard Model Predictions" in the B^0 Particle Listings above.

VALUE	DOCUMENT ID		TECN	COMMENT
$-0.25^{+0.45}_{-0.47} \pm 0.14$	[458] AUBERT	02D	BABR	$e^+e^- \to \Upsilon(4S)$

[458] Corresponds to 90% confidence range $-1.0 < C_{\pi\pi} < 0.47$.

$S_{\pi\pi}$ ($B^0 \rightarrow \pi^+\pi^-$)

$S_{\pi\pi} = 2\mathrm{Im}\lambda/(1+|\lambda|^2)$, see the note in the $C_{\pi\pi}$ datablock above.

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.03^{+0.52}_{-0.56} \pm 0.11}$	[459] AUBERT	02D	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[459] Corresponds to 90% confidence range $-0.89 < S_{\pi\pi} < 0.85$.

sin(2β)

For a discussion of *CP* violation, see the note on "*CP* Violation in *B* Decay Standard Model Predictions" in the B^0 Particle Listings above. $\sin(2\beta)$ is a measure of the *CP*-violating amplitude in the $B^0_d \rightarrow J/\psi(1S)K^0_S$.

VALUE	DOCUMENT ID		TECN	COMMENT
0.79±0.14 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
$0.99 \pm 0.14 \pm 0.06$	[460] ABE	01G	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$0.59 \pm 0.14 \pm 0.05$	[460] AUBERT	01B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.79^{+0.41}_{-0.44}$	[461] AFFOLDER	00C	CDF	$p\overline{p}$ at 1.8 TeV
$0.84^{+0.82}_{-1.04} \pm 0.16$	[462] BARATE	00Q	ALEP	$e^+e^- \rightarrow Z$
$3.2\ ^{+1.8}_{-2.0}\ \pm 0.5$	[463] ACKERSTAFF	98Z	OPAL	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.58^{+0.32}_{-0.34}{}^{+0.09}_{-0.10}$	ABASHIAN	01	BELL	Repl. by ABE 01G
$0.34 \pm 0.20 \pm 0.05$	AUBERT	01	BABR	Repl. by AUBERT 01B
$1.8\ \pm 1.1\ \pm 0.3$	[464] ABE	98U	CDF	Repl. by AFFOLDER 00C

[460] First observation of *CP* violation in B^0 meson system.

[461] AFFOLDER 00C uses about 400 $B^0 \rightarrow J/\psi(1S)K^0_S$ events. The production flavor of B^0 was determined using three tagging algorithms: a same-side tag, a jet-charge tag, and a soft-lepton tag.

[462] BARATE 00Q uses 23 candidates for $B^0 \rightarrow J/\psi(1S)K^0_S$ decays. A combination of jet-charge, vertex-charge, and same-side tagging techniques were used to determine the B^0 production flavor.

[463] ACKERSTAFF 98Z uses 24 candidates for $B^0_d \rightarrow J/\psi(1S)K^0_S$ decay. A combination of jet-charge and vertex-charge techniques were used to tag the B^0_d production flavor.

[464] ABE 98U uses 198 ± 17 $B^0_d \rightarrow J/\psi(1S)K^0$ events. The production flavor of B^0 was determined using the same side tagging technique.



sin(2β)

$B^0 \to D^{*-} \ell^+ \nu_\ell$ FORM FACTORS

R_1 (form factor ratio $\sim V/A_1$)

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
1.18±0.30±0.12	DUBOSCQ	96	CLE2	$e^+ e^- \to \Upsilon(4S)$

R_2 (form factor ratio $\sim A_2/A_1$)

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.71±0.22±0.07	DUBOSCQ	96	CLE2	$e^+ e^- \to \Upsilon(4S)$

$\rho^2_{A_1}$ (form factor slope)

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.91±0.15±0.06	DUBOSCQ	96	CLE2	$e^+ e^- \to \Upsilon(4S)$

B^0 REFERENCES

ABE	02	PRL 88 021801	K. Abe *et al.*	(Belle Collab.)
ABE	02E	PL B526 258	K. Abe *et al.*	(Belle Collab.)
ABE	02F	PL B526 247	K. Abe *et al.*	(Belle Collab.)
ABE	02H	PRL 88 171801	K. Abe *et al.*	(BELLE Collab.)
ABE	02J	PRL 88 052002	K. Abe *et al.*	(BELLE Collab.)
AFFOLDER	02B	PRL 88 071801	T. Affolder *et al.*	(CDF Collab.)
ASNER	02	PR D65 031103R	D.M. Asner *et al.*	(CLEO Collab.)
AUBERT	02	PR D65 032001	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	02D	PR D65 051502	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	02E	PR D65 051101R	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	02H	PRL (to be publ.)	B. Aubert *et al.*	(Babar Collab.)

hep-ex/0202005

AUBERT	02I	PRL (to be publ.)	B. Aubert *et al.*	(BaBar Collab.)
hep-ex/0112044				
AUBERT	02J	PRL (to be publ.)	B. Aubert *et al.*	(BaBar Collab.)
hep-ex/0112045				
COAN	02	PRL 88 062001	T.E. Coan *et al.*	(CLEO Collab.)
EDWARDS	02	PR D65 012002	K.W. Edwards *et al.*	(CLEO Collab.)
GODANG	02	PRL 88 021802	R. Godang *et al.*	(CLEO Collab.)
MAHAPATRA	02	PRL 88 101803	R. Mahapatra *et al.*	(CLEO Collab.)
ABASHIAN	01	PRL 86 2509	A. Abashian *et al.*	(Belle Collab.)
ABE	01D	PRL 86 3228	K. Abe *et al.*	(Belle Collab.)
ABE	01G	PRL 87 091802	K. Abe *et al.*	(Belle Collab.)
ABE	01H	PRL 87 101801	K. Abe *et al.*	(Belle Collab.)
ABE	01I	PRL 87 111801	K. Abe *et al.*	(Belle Collab.)
ABE	01K	PR D64 071101	K. Abe *et al.*	(Belle Collab.)
ABE	01L	PRL 87 161601	K. Abe *et al.*	(Belle Collab.)
ABE	01M	PL B517 309	K. Abe *et al.*	(Belle Collab.)
ABREU	01H	PL B510 55	P. Abreu *et al.*	(DELPHI Collab.)
ALEXANDER	01B	PR D64 092001	J.P. Alexander *et al.*	(CLEO Collab.)
AMMAR	01B	PRL 87 271801	R. Ammar *et al.*	(CLEO Collab.)
ANDERSON	01	PRL 86 2732	S. Anderson *et al.*	(CLEO Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson *et al.*	(CLEO Collab.)
AUBERT	01	PRL 86 2515	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	01B	PRL 87 091801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	01D	PRL 87 151801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	01E	PRL 87 151802	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	01F	PRL 87 201803	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	01G	PRL 87 221802	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	01H	PRL 87 241801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	01I	PRL 87 241803	B. Aubert *et al.*	(BaBar Collab.)
BARATE	01D	EPJ C20 431	R. Barate *et al.*	(ALEPH Collab.)
BRIERE	01	PRL 86 3718	R.A. Biere *et al.*	(CLEO Collab.)
EDWARDS	01	PRL 86 30	K.W. Edwards *et al.*	(CLEO Collab.)
JAFFE	01	PRL 86 5000	D. Jaffe *et al.*	(CLEO Collab.)
RICHICHI	01	PR D63 031103R	S.J. Richichi *et al.*	(CLEO Collab.)
ABBIENDI	00Q	PL B482 15	G. Abbiendi *et al.*	(OPAL Collab.)
ABBIENDI,G	00B	PL B493 266	G. Abbiendi *et al.*	(OPAL Collab.)
ABE	00C	PR D62 071101R	K. Abe *et al.*	(SLD Collab.)
AFFOLDER	00C	PR D61 072005	T. Affolder *et al.*	(CDF Collab.)
AFFOLDER	00N	PRL 85 4668	T. Affolder *et al.*	(CDF Collab.)
AHMED	00B	PR D62 112003	S. Ahmed *et al.*	(CLEO Collab.)
ANASTASSOV	00	PRL 84 1393	A. Anastassov *et al.*	(CLEO Collab.)
ARTUSO	00	PRL 84 4292	M. Artuso *et al.*	(CLEO Collab.)
AVERY	00	PR D62 051101	P. Avery *et al.*	(CLEO Collab.)
BARATE	00Q	PL B492 259	R. Barate *et al.*	(ALEPH Collab.)
BARATE	00R	PL B492 275	R. Barate *et al.*	(ALEPH Collab.)
BEHRENS	00	PR D61 052001	B.H. Behrens *et al.*	(CLEO Collab.)
BEHRENS	00B	PL B490 36	B.H. Behrens *et al.*	(CLEO Collab.)
BERGFELD	00B	PR D62 091102R	T. Bergfeld *et al.*	(CLEO Collab.)
CHEN	00	PRL 85 525	S. Chen *et al.*	(CLEO Collab.)
COAN	00	PRL 84 5283	T.E. Coan *et al.*	(CLEO Collab.)
CRONIN-HEN...	00	PRL 85 515	D. Cronin-Hennessy *et al.*	(CLEO Collab.)
CSORNA	00	PR D61 111101	S.E. Csorna *et al.*	(CLEO Collab.)
JESSOP	00	PRL 85 2881	C.P. Jessop *et al.*	(CLEO Collab.)
LIPELES	00	PR D62 032005	E. Lipeles *et al.*	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi *et al.*	(CLEO Collab.)
ABBIENDI	99J	EPJ C12 609	G. Abbiendi *et al.*	(OPAL Collab.)
ABE	99K	PR D60 051101	F. Abe *et al.*	(CDF Collab.)
ABE	99Q	PR D60 072003	F. Abe *et al.*	(CDF Collab.)
AFFOLDER	99B	PRL 83 3378	T. Affolder *et al.*	(CDF Collab.)
AFFOLDER	99C	PR D60 112004	T. Affolder *et al.*	(CDF Collab.)
ARTUSO	99	PRL 82 3020	M. Artuso *et al.*	(CLEO Collab.)
BARTELT	99	PRL 82 3746	J. Bartelt *et al.*	(CLEO Collab.)
COAN	99	PR D59 111101	T.E. Coan *et al.*	(CLEO Collab.)
ABBOTT	98B	PL B423 419	B. Abbott *et al.*	(D0 Collab.)
ABE	98	PR D57 R3811	F. Abe *et al.*	(CDF Collab.)
ABE	98B	PR D57 5382	F. Abe *et al.*	(CDF Collab.)
ABE	98C	PRL 80 2057	F. Abe *et al.*	(CDF Collab.)
Also	99C	PR D59 032001	F. Abe *et al.*	(CDF Collab.)
ABE	98O	PR D58 072001	F. Abe *et al.*	(CDF Collab.)
ABE	98Q	PR D58 092002	F. Abe *et al.*	(CDF Collab.)
ABE	98U	PRL 81 5513	F. Abe *et al.*	(CDF Collab.)

ABE	98V	PRL 81 5742	F. Abe *et al.*	(CDF Collab.)
ACCIARRI	98D	EPJ C5 195	M. Acciarri *et al.*	(L3 Collab.)
ACCIARRI	98S	PL B438 417	M. Acciarri *et al.*	(L3 Collab.)
ACKERSTAFF	98Z	EPJ C5 379	K. Ackerstaff *et al.*	(OPAL Collab.)
BARATE	98Q	EPJ C4 387	R. Barate *et al.*	(ALEPH Collab.)
BEHRENS	98	PRL 80 3710	B.H. Behrens *et al.*	(CLEO Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld *et al.*	(CLEO Collab.)
BRANDENB...	98	PRL 80 2762	G. Brandenbrug *et al.*	(CLEO Collab.)
GODANG	98	PRL 80 3456	R. Godang *et al.*	(CLEO Collab.)
NEMATI	98	PR D57 5363	B. Nemati *et al.*	(CLEO Collab.)
ABE	97J	PRL 79 590	K. Abe *et al.*	(SLD Collab.)
ABREU	97F	ZPHY C74 19	P. Abreu *et al.*	(DELPHI Collab.)
Also	97K	ZPHY C75 579 erratum	P. Abreu *et al.*	(DELPHI Collab.)
ABREU	97N	ZPHY C76 579	P. Abreu *et al.*	(DELPHI Collab.)
ACCIARRI	97B	PL B391 474	M. Acciarri *et al.*	(L3 Collab.)
ACCIARRI	97C	PL B391 481	M. Acciarri *et al.*	(L3 Collab.)
ACKERSTAFF	97G	PL B395 128	K. Ackerstaff *et al.*	(OPAL Collab.)
ACKERSTAFF	97U	ZPHY C76 401	K. Ackerstaff *et al.*	(OPAL Collab.)
ACKERSTAFF	97V	ZPHY C76 417	K. Ackerstaff *et al.*	(OPAL Collab.)
ARTUSO	97	PL B399 321	M. Artuso *et al.*	(CLEO Collab.)
ASNER	97	PRL 79 799	D. Asner *et al.*	(CLEO Collab.)
ATHANAS	97	PRL 79 2208	M. Athanas *et al.*	(CLEO Collab.)
BUSKULIC	97	PL B395 373	D. Buskulic *et al.*	(ALEPH Collab.)
BUSKULIC	97D	ZPHY C75 397	D. Buskulic *et al.*	(ALEPH Collab.)
FU	97	PRL 79 3125	X. Fu *et al.*	(CLEO Collab.)
JESSOP	97	PRL 79 4533	C.P. Jessop *et al.*	(CLEO Collab.)
ABE	96B	PR D53 3496	F. Abe *et al.*	(CDF Collab.)
ABE	96C	PRL 76 4462	F. Abe *et al.*	(CDF Collab.)
ABE	96H	PRL 76 2015	F. Abe *et al.*	(CDF Collab.)
ABE	96L	PRL 76 4675	F. Abe *et al.*	(CDF Collab.)
ABE	96Q	PR D54 6596	F. Abe *et al.*	(CDF Collab.)
ABREU	96P	ZPHY C71 539	P. Abreu *et al.*	(DELPHI Collab.)
ABREU	96Q	ZPHY C72 17	P. Abreu *et al.*	(DELPHI Collab.)
ACCIARRI	96E	PL B383 487	M. Acciarri *et al.*	(L3 Collab.)
ADAM	96D	ZPHY C72 207	W. Adam *et al.*	(DELPHI Collab.)
ALBRECHT	96D	PL B374 256	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	96T	PRL 77 5000	J.P. Alexander *et al.*	(CLEO Collab.)
ALEXANDER	96V	ZPHY C72 377	G. Alexander *et al.*	(OPAL Collab.)
ASNER	96	PR D53 1039	D.M. Asner *et al.*	(CLEO Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish *et al.*	(CLEO Collab.)
BISHAI	96	PL B369 186	M. Bishai *et al.*	(CLEO Collab.)
BUSKULIC	96J	ZPHY C71 31	D. Buskulic *et al.*	(ALEPH Collab.)
BUSKULIC	96V	PL B384 471	D. Buskulic *et al.*	(ALEPH Collab.)
DUBOSCQ	96	PRL 76 3898	J.E. Duboscq *et al.*	(CLEO Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut *et al.*	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett *et al.*	
ABE	95Z	PRL 75 3068	F. Abe *et al.*	(CDF Collab.)
ABREU	95N	PL B357 255	P. Abreu *et al.*	(DELPHI Collab.)
ABREU	95Q	ZPHY C68 13	P. Abreu *et al.*	(DELPHI Collab.)
ACCIARRI	95H	PL B363 127	M. Acciarri *et al.*	(L3 Collab.)
ACCIARRI	95I	PL B363 137	M. Acciarri *et al.*	(L3 Collab.)
ADAM	95	ZPHY C68 363	W. Adam *et al.*	(DELPHI Collab.)
AKERS	95J	ZPHY C66 555	R. Akers *et al.*	(OPAL Collab.)
AKERS	95T	ZPHY C67 379	R. Akers *et al.*	(OPAL Collab.)
ALEXANDER	95	PL B341 435	J. Alexander *et al.*	(CLEO Collab.)
Also	95C	PL B347 469 (erratum)	J. Alexander *et al.*	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish *et al.*	(CLEO Collab.)
BUSKULIC	95N	PL B359 236	D. Buskulic *et al.*	(ALEPH Collab.)
ABE	94D	PRL 72 3456	F. Abe *et al.*	(CDF Collab.)
ABREU	94M	PL B338 409	P. Abreu *et al.*	(DELPHI Collab.)
AKERS	94C	PL B327 411	R. Akers *et al.*	(OPAL Collab.)
AKERS	94H	PL B336 585	R. Akers *et al.*	(OPAL Collab.)
AKERS	94J	PL B337 196	R. Akers *et al.*	(OPAL Collab.)
AKERS	94L	PL B337 393	R. Akers *et al.*	(OPAL Collab.)
ALAM	94	PR D50 43	M.S. Alam *et al.*	(CLEO Collab.)
ALBRECHT	94	PL B324 249	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	94G	PL B340 217	H. Albrecht *et al.*	(ARGUS Collab.)
AMMAR	94	PR D49 5701	R. Ammar *et al.*	(CLEO Collab.)
ATHANAS	94	PRL 73 3503	M. Athanas *et al.*	(CLEO Collab.)
Also	95	PRL 74 3090 (erratum)	M. Athanas *et al.*	(CLEO Collab.)
BUSKULIC	94B	PL B322 441	D. Buskulic *et al.*	(ALEPH Collab.)

PDG	94	PR D50 1173	L. Montanet *et al.*	(CERN, LBL, BOST+)
PROCARIO	94	PRL 73 1472	M. Procario *et al.*	(CLEO Collab.)
STONE	94	HEPSY 93-11	S. Stone	
Published in B Decays, 2nd Edition, World Scientific, Singapore				
ABREU	93D	ZPHY C57 181	P. Abreu *et al.*	(DELPHI Collab.)
ABREU	93G	PL B312 253	P. Abreu *et al.*	(DELPHI Collab.)
ACTON	93C	PL B307 247	P.D. Acton *et al.*	(OPAL Collab.)
ALBRECHT	93	ZPHY C57 533	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander *et al.*	(CLEO Collab.)
AMMAR	93	PRL 71 674	R. Ammar *et al.*	(CLEO Collab.)
BARTELT	93	PRL 71 1680	J.E. Bartelt *et al.*	(CLEO Collab.)
BATTLE	93	PRL 71 3922	M. Battle *et al.*	(CLEO Collab.)
BEAN	93B	PRL 70 2681	A. Bean *et al.*	(CLEO Collab.)
BUSKULIC	93D	PL B307 194	D. Buskulic *et al.*	(ALEPH Collab.)
Also	94H	PL B325 537 (errata)	D. Buskulic *et al.*	(ALEPH Collab.)
BUSKULIC	93K	PL B313 498	D. Buskulic *et al.*	(ALEPH Collab.)
SANGHERA	93	PR D47 791	S. Sanghera *et al.*	(CLEO Collab.)
ALBRECHT	92C	PL B275 195	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	92L	ZPHY C55 357	H. Albrecht *et al.*	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto *et al.*	(CLEO Collab.)
HENDERSON	92	PR D45 2212	S. Henderson *et al.*	(CLEO Collab.)
KRAMER	92	PL B279 181	G. Kramer, W.F. Palmer	(HAMB, OSU)
ALBAJAR	91C	PL B262 163	C. Albajar *et al.*	(UA1 Collab.)
ALBAJAR	91E	PL B273 540	C. Albajar *et al.*	(UA1 Collab.)
ALBRECHT	91B	PL B254 288	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	91E	PL B262 148	H. Albrecht *et al.*	(ARGUS Collab.)
BERKELMAN	91	ARNPS 41 1	K. Berkelman, S. Stone	(CORN, SYRA)
"Decays of B Mesons"				
FULTON	91	PR D43 651	R. Fulton *et al.*	(CLEO Collab.)
ALBRECHT	90B	PL B241 278	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	90J	ZPHY C48 543	H. Albrecht *et al.*	(ARGUS Collab.)
ANTREASYAN	90B	ZPHY C48 553	D. Antreasyan *et al.*	(Crystal Ball Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto *et al.*	(CLEO Collab.)
ELSEN	90	ZPHY C46 349	E. Elsen *et al.*	(JADE Collab.)
ROSNER	90	PR D42 3732	J.L. Rosner	
WAGNER	90	PRL 64 1095	S.R. Wagner *et al.*	(Mark II Collab.)
ALBRECHT	89C	PL B219 121	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	89G	PL B229 304	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	89J	PL B229 175	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	89L	PL B232 554	H. Albrecht *et al.*	(ARGUS Collab.)
ARTUSO	89	PRL 62 2233	M. Artuso *et al.*	(CLEO Collab.)
AVERILL	89	PR D39 123	D.A. Averill *et al.*	(HRS Collab.)
AVERY	89B	PL B223 470	P. Avery *et al.*	(CLEO Collab.)
BEBEK	89	PRL 62 8	C. Bebek *et al.*	(CLEO Collab.)
BORTOLETTO	89	PRL 62 2436	D. Bortoletto *et al.*	(CLEO Collab.)
BORTOLETTO	89B	PRL 63 1667	D. Bortoletto *et al.*	(CLEO Collab.)
ALBRECHT	88F	PL B209 119	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	88K	PL B215 424	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87C	PL B185 218	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87I	PL B192 245	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87J	PL B197 452	H. Albrecht *et al.*	(ARGUS Collab.)
AVERY	87	PL B183 429	P. Avery *et al.*	(CLEO Collab.)
BEAN	87B	PRL 58 183	A. Bean *et al.*	(CLEO Collab.)
BEBEK	87	PR D36 1289	C. Bebek *et al.*	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam *et al.*	(CLEO Collab.)
ALBRECHT	86F	PL B182 95	H. Albrecht *et al.*	(ARGUS Collab.)
PDG	86	PL 170B	M. Aguilar-Benitez *et al.*	(CERN, CIT+)
CHEN	85	PR D31 2386	A. Chen *et al.*	(CLEO Collab.)
HAAS	85	PRL 55 1248	J. Haas *et al.*	(CLEO Collab.)
AVERY	84	PRL 53 1309	P. Avery *et al.*	(CLEO Collab.)
GILES	84	PR D30 2279	R. Giles *et al.*	(CLEO Collab.)
BEHRENDS	83	PRL 50 881	S. Behrends *et al.*	(CLEO Collab.)