



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

According to the quark model, the Ξ_c^0 (quark content dsc) and Ξ_c^+ form an isospin doublet, and the spin-parity ought to be $J^P = 1/2^+$. None of I , J , or P has actually been measured.

Ξ_c^0 MASS

The fit uses the Ξ_c^0 and Ξ_c^+ mass and mass-difference measurements.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-----------------------|----------|---|
| 2471.8±1.4 OUR FIT | | | | |
| 2471.8±1.4 OUR AVERAGE | | | | |
| 2470.0±2.8±2.6 | 85 | FRABETTI | 98B E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |
| 2469 ±2 ±3 | 9 | HENDERSON | 92B CLEO | $\Omega^- K^+$ |
| 2472.1±2.7±1.6 | 54 | ALBRECHT | 90F ARG | $e^+ e^-$ at $\Upsilon(4S)$ |
| 2473.3±1.9±1.2 | 4 | BARLAG | 90 ACCM | $\pi^- (K^-)$ Cu 230 GeV |
| 2472 ±3 ±4 | 19 | ALAM | 89 CLEO | $e^+ e^-$ 10.6 GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 2462.1±3.1±1.4 | 42 | ¹ FRABETTI | 93C E687 | See FRABETTI 98B |
| 2471 ±3 ±4 | 14 | VERY | 89 CLEO | See ALAM 89 |

¹The FRABETTI 93C mass is well below the other measurements.

$\Xi_c^0 - \Xi_c^+$ MASS DIFFERENCE

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------|---------|--|
| 5.5±1.8 OUR FIT | | | |
| 6.3±2.3 OUR AVERAGE | | | |
| +7.0±4.5±2.2 | ALBRECHT | 90F ARG | $e^+ e^-$ at $\Upsilon(4S)$ |
| +6.8±3.3±0.5 | BARLAG | 90 ACCM | $\pi^- (K^-)$ Cu 230 GeV |
| +5 ±4 ±1 | ALAM | 89 CLEO | $\Xi_c^0 \rightarrow \Xi^- \pi^+, \Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+$ |

Ξ_c^0 MEAN LIFE

| VALUE (10^{-15} s) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|----------|---|
| 98^{+23}_{-15} OUR AVERAGE | | | | |
| $101^{+25}_{-17} \pm 5$ | 42 | FRABETTI | 93C E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |
| 82^{+59}_{-30} | 4 | BARLAG | 90 ACCM | $\pi^- (K^-)$ Cu 230 GeV |

Ξ_c^0 DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| Γ_1 $\Lambda \bar{K}^0$ | seen |
| Γ_2 $\Lambda \bar{K}^0 \pi^+ \pi^-$ | seen |
| Γ_3 $\Lambda K^- \pi^+ \pi^+ \pi^-$ | seen |
| Γ_4 $\Xi^- \pi^+$ | seen |
| Γ_5 $\Xi^- \pi^+ \pi^+ \pi^-$ | seen |
| Γ_6 $p K^- \bar{K}^*(892)^0$ | seen |
| Γ_7 $\Omega^- K^+$ | seen |
| Γ_8 $\Xi^- e^+ \nu_e$ | seen |
| Γ_9 $\Xi^- \ell^+$ anything | seen |

Ξ_c^0 BRANCHING RATIOS

| | |
|---|---|
| $\Gamma(\Lambda \bar{K}^0)/\Gamma_{\text{total}}$ | Γ_1/Γ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| seen | ALBRECHT 95B ARG $e^+ e^- \approx 10.4$ GeV |
| $\Gamma(\Lambda \bar{K}^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ | Γ_2/Γ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| seen | FRABETTI 98B E687 γ Be, $\bar{E}_\gamma = 220$ GeV |
| $\Gamma(\Lambda K^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ | Γ_3/Γ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| seen | FRABETTI 98B E687 γ Be, $\bar{E}_\gamma = 220$ GeV |
| $\Gamma(\Xi^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+ \pi^-)$ | Γ_4/Γ_5 |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| $0.30 \pm 0.12 \pm 0.05$ | ALBRECHT 90F ARG $e^+ e^-$ at $\Upsilon(4S)$ |
| $\Gamma(p K^- \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ | Γ_6/Γ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| seen | BARLAG 90 ACCM $\pi^- (K^-)$ Cu 230 GeV |
| $\Gamma(\Omega^- K^+)/\Gamma(\Xi^- \pi^+)$ | Γ_7/Γ_4 |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| $0.50 \pm 0.21 \pm 0.05$ | HENDERSON 92B CLEO $e^+ e^- \approx 10.6$ GeV |
| $\Gamma(\Xi^- e^+ \nu_e)/\Gamma(\Xi^- \pi^+)$ | Γ_8/Γ_4 |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| $3.1 \pm 1.0^{+0.3}_{-0.5}$ | ALEXANDER 95B CLE2 $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Xi^- \ell^+ \text{anything})/\Gamma(\Xi^- \pi^+)$ Γ_9/Γ_4

The ratio is for the *average* (not the sum) of the $\Xi^- e^+$ anything and $\Xi^- \mu^+$ anything modes.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|---------|----------------------------|
| 0.96±0.43±0.18 | 18 | ALBRECHT | 93B ARG | $e^+ e^- \approx 10.4$ GeV |

$\Gamma(\Xi^- \ell^+ \text{anything})/\Gamma(\Xi^- \pi^+ \pi^+ \pi^-)$ Γ_9/Γ_5

The ratio is for the *average* (not the sum) of the $\Xi^- e^+$ anything and $\Xi^- \mu^+$ anything modes.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|---------|----------------------------|
| 0.29±0.12±0.04 | 18 | ALBRECHT | 93B ARG | $e^+ e^- \approx 10.4$ GeV |

Ξ_c^0 DECAY PARAMETERS

See the note on "Baryon Decay Parameters" in the neutron Listings.

α FOR $\Xi_c^0 \rightarrow \Xi^- \pi^+$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|--------------------------------|
| -0.56±0.39^{+0.10}_{-0.09} | 138 | CHAN | 01 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

Ξ_c^0 REFERENCES

| | | | | |
|-----------|-----|-----------------------|-----------------------------|---------------------|
| CHAN | 01 | PR D63 111102R | S. Chan <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 98B | PL B426 403 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| ALBRECHT | 95B | PL B342 397 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEXANDER | 95B | PRL 74 3113 | J. Alexander <i>et al.</i> | (CLEO Collab.) |
| Also | 95E | PRL 75 4155 (erratum) | J. Alexander <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 93B | PL B303 368 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| FRABETTI | 93C | PRL 70 2058 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| HENDERSON | 92B | PL B283 161 | S. Henderson <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 90F | PL B247 121 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BARLAG | 90 | PL B236 495 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| ALAM | 89 | PL B226 401 | M.S. Alam <i>et al.</i> | (CLEO Collab.) |
| AVERY | 89 | PRL 62 863 | P. Avery <i>et al.</i> | (CLEO Collab.) |