

# $\rho(2150)$

$I^G(J^{PC}) = 1^+(1^{--})$

## OMMITTED FROM SUMMARY TABLE

This entry was previously called  $T_1(2190)$ .

### $\rho(2150)$ MASS

$$e^+ e^- \rightarrow \pi^+ \pi^-, K^+ K^-, 6\pi$$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>2149±17 OUR AVERAGE</b>	Includes data from the datablock that follows this one.			
2153±37	BIAGINI	91	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-, K^+ K^-$
2110±50	<sup>2</sup> CLEGG	90	RVUE	$e^+ e^- \rightarrow 3(\pi^+ \pi^-), 2(\pi^+ \pi^- \pi^0)$

### $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
~ 2191	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 1988	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 2070	<sup>1</sup> OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2170	<sup>3</sup> MARTIN	80B	RVUE
~ 2100	<sup>3</sup> MARTIN	80C	RVUE

<sup>1</sup> See however KLOET 96 who fit  $\pi^+ \pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

### S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
2110±35	<sup>14</sup> ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2190	<sup>4</sup> CUTTS	78B	CNTR	0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2155±15	<sup>4,5</sup> COUPLAND	77	CNTR	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2193± 2	<sup>4,6</sup> ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
2190±10	<sup>7</sup> ABRAMS	70	CNTR	S channel $\bar{p}N$

### $\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

### 2155±21 OUR AVERAGE

2140±30	ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega\pi^0 n$
2170±30	ALDE	92C	GAM4	100 $\pi^- p \rightarrow \omega\pi^0 n$

<sup>2</sup> Includes ATKINSON 85.<sup>3</sup>  $I(J^P) = 1(1^-)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .<sup>4</sup> Isospins 0 and 1 not separated.<sup>5</sup> From a fit to the total elastic cross section.<sup>6</sup> Referred to as  $T$  or  $T'$  region by ALSPECTOR 73.<sup>7</sup> Seen as bump in  $I = 1$  state. See also COOPER 68. PEASLEE 75 confirm  $\bar{p}p$  results of ABRAMS 70, no narrow structure.

## $\rho(2150)$ WIDTH

### $e^+ e^- \rightarrow \pi^+ \pi^-, K^+ K^-, 6\pi$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>363 ± 50 OUR AVERAGE</b>	Includes data from the datablock that follows this one.			
389 ± 79	BIAGINI	91	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-, K^+ K^-$
410 ± 100	9 CLEGG	90	RVUE	$e^+ e^- \rightarrow 3(\pi^+ \pi^-), 2(\pi^+ \pi^- \pi^0)$

### $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 296	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 244	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 40	8 OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	10 MARTIN	80B	RVUE
~ 200	10 MARTIN	80C	RVUE

<sup>8</sup> See however KLOET 96 who fit  $\pi^+ \pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

### S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
230 ± 50	15 ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+ \pi^-$
135 ± 75	11,12 COUPLAND	77	CNTR	0 0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98 ± 8	12 ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
~ 85	13 ABRAMS	70	CNTR	S channel $\bar{p}N$

### $\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

**320 ± 70** ALDE 95 GAM2 38  $\pi^- p \rightarrow \omega\pi^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 300 ALDE 92C GAM4 100  $\pi^- p \rightarrow \omega\pi^0 n$

<sup>9</sup> Includes ATKINSON 85.<sup>10</sup>  $I(J^P) = 1(1^-)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .<sup>11</sup> From a fit to the total elastic cross section.<sup>12</sup> Isospins 0 and 1 not separated.<sup>13</sup> Seen as bump in  $I = 1$  state. See also COOPER 68. PEASLEE 75 confirm  $\bar{p}p$  results of ABRAMS 70, no narrow structure.<sup>14</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.<sup>15</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

## $\rho(2150)$ REFERENCES

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>
BIAGINI	91	NC 104A 363	M.E. Biagini <i>et al.</i>
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>
PEASLEE	75	PL 57B 189	D.C. Peaslee <i>et al.</i>
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>
COOPER	68	PRL 20 1059	W.A. Cooper <i>et al.</i>

(RUTG, NORD)  
(GAMS Collab.) JP  
(LOQM)  
(DURH)  
(BELG, SERP, KEK, LANL+)  
(FRAS, PRAG)  
(LANC, MCHS)  
(BONN, CERN, GLAS+)  
(LOUC, RHEL) JP  
(DURH) JP  
(STON, WISC)  
(LOQM, RHEL)  
(CANB, BARI, BROW+)  
(RUTG, UPNJ)  
(BNL)  
(ANL)

## OTHER RELATED PAPERS

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
EISENHAND...	75	NP B96 109	E. Eisenhandler <i>et al.</i>	(LOQM, LIVP, DARE+)
BRICMAN	69	PL 29B 451	C. Bricman <i>et al.</i>	(CERN, CAEN, SACL)
ABRAMS	67C	PRL 18 1209	R.J. Abrams <i>et al.</i>	(BNL)