

# $K_5^*(2380)$

$$I(J^P) = \frac{1}{2}(5^-)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

## $K_5^*(2380)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
<b>2382±14±19</b>	<sup>1</sup> ASTON	86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

<sup>1</sup> From a fit to all the moments.

## $K_5^*(2380)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
<b>178±37±32</b>	<sup>2</sup> ASTON	86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

<sup>2</sup> From a fit to all the moments.

## $K_5^*(2380)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad K\pi$	(6.1±1.2) %

## $K_5^*(2380)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	$\Gamma_1/\Gamma$	
	<b>0.061±0.012</b>	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	

## $K_5^*(2380)$ REFERENCES

ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)