



$$J = \frac{1}{2}$$

τ discovery paper was PERL 75. $e^+e^- \rightarrow \tau^+\tau^-$ cross-section threshold behavior and magnitude are consistent with pointlike spin-1/2 Dirac particle. BRANDELIK 78 ruled out pointlike spin-0 or spin-1 particle. FELDMAN 78 ruled out $J = 3/2$. KIRKBY 79 also ruled out $J=\text{integer}$, $J = 3/2$.

τ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1776.99^{+0.29}_{-0.26} OUR AVERAGE				
1775.1 ±1.6 ±1.0	13.3k	¹ ABBIENDI	00A OPAL	1990–1995 LEP runs
1778.2 ±0.8 ±1.2		ANASTASSOV 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
1776.96 ^{+0.18} _{-0.21} ^{+0.25} _{-0.17}	65	² BAI	96 BES	$E_{\text{cm}}^{ee} = 3.54\text{--}3.57$ GeV
1776.3 ±2.4 ±1.4	11k	³ ALBRECHT	92M ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
1783 ⁺³ ₋₄	692	⁴ BACINO	78B DLCO	$E_{\text{cm}}^{ee} = 3.1\text{--}7.4$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1777.8 ±0.7 ±1.7	35k	⁵ BALEST	93 CLEO	Repl. by ANASTASSOV 97
1776.9 ^{+0.4} _{-0.5} ±0.2	14	⁶ BAI	92 BES	Repl. by BAI 96

¹ ABBIENDI 00A fit τ pseudomass spectrum in $\tau \rightarrow \pi^\pm \leq 2\pi^0 \nu_\tau$ and $\tau \rightarrow \pi^\pm \pi^+ \pi^- \leq 1\pi^0 \nu_\tau$ decays. Result assumes $m_{\nu_\tau} = 0$.

² BAI 96 fit $\sigma(e^+e^- \rightarrow \tau^+\tau^-)$ at different energies near threshold.

³ ALBRECHT 92M fit τ pseudomass spectrum in $\tau^- \rightarrow 2\pi^- \pi^+ \nu_\tau$ decays. Result assumes $m_{\nu_\tau} = 0$.

⁴ BACINO 78B value comes from $e^\pm X^\mp$ threshold. Published mass 1782 MeV increased by 1 MeV using the high precision $\psi(2S)$ mass measurement of ZHOLENTZ 80 to eliminate the absolute SPEAR energy calibration uncertainty.

⁵ BALEST 93 fit spectra of minimum kinematically allowed τ mass in events of the type $e^+e^- \rightarrow \tau^+\tau^- \rightarrow (\pi^+ n\pi^0 \nu_\tau)(\pi^- m\pi^0 \nu_\tau)$ $n \leq 2, m \leq 2, 1 \leq n+m \leq 3$. If $m_{\nu_\tau} \neq 0$, result increases by $(m_{\nu_\tau}^2 / 1100 \text{ MeV})$.

⁶ BAI 92 fit $\sigma(e^+e^- \rightarrow \tau^+\tau^-)$ near threshold using $e\mu$ events.

$$(m_{\tau^+} - m_{\tau^-})/m_{\text{average}}$$

A test of CPT invariance.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.0 × 10⁻³	90	ABBIENDI	00A OPAL	1990–1995 LEP runs

τ MEAN LIFE

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
290.6 ± 1.1	OUR AVERAGE			
293.2 ± 2.0 ± 1.5		ACCIARRI	00B L3	1991–1995 LEP runs
290.1 ± 1.5 ± 1.1		BARATE	97R ALEP	1989–1994 LEP runs
291.4 ± 3.0		ABREU	96B DLPH	1991–1993 LEP runs
289.2 ± 1.7 ± 1.2		ALEXANDER	96E OPAL	1990–1994 LEP runs
289.0 ± 2.8 ± 4.0	57.4k	BALEST	96 CLEO	$E_{cm}^{ee} = 10.6$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
291.2 ± 2.0 ± 1.2		BARATE	97I ALEP	Repl. by BARATE 97R
290.1 ± 4.0	34k	ACCIARRI	96K L3	Repl. by ACCIARRI 00B
297 ± 9 ± 5	1671	ABE	95Y SLD	1992–1993 SLC runs
304 ± 14 ± 7	4100	BATTLE	92 CLEO	$E_{cm}^{ee} = 10.6$ GeV
301 ± 29	3780	KLEINWORT	89 JADE	$E_{cm}^{ee} = 35$ –46 GeV
288 ± 16 ± 17	807	AMIDEI	88 MRK2	$E_{cm}^{ee} = 29$ GeV
306 ± 20 ± 14	695	BRAUNSCH...	88C TASS	$E_{cm}^{ee} = 36$ GeV
299 ± 15 ± 10	1311	ABACHI	87C HRS	$E_{cm}^{ee} = 29$ GeV
295 ± 14 ± 11	5696	ALBRECHT	87P ARG	$E_{cm}^{ee} = 9.3$ –10.6 GeV
309 ± 17 ± 7	3788	BAND	87B MAC	$E_{cm}^{ee} = 29$ GeV
325 ± 14 ± 18	8470	BEBEK	87C CLEO	$E_{cm}^{ee} = 10.5$ GeV
460 ± 190	102	FELDMAN	82 MRK2	$E_{cm}^{ee} = 29$ GeV

τ MAGNETIC MOMENT ANOMALY

The q^2 dependence is expected to be small providing no thresholds are nearby.

$$\mu_\tau / (e\hbar/2m_\tau) - 1 = (g_\tau - 2)/2$$

For a theoretical calculation [$(g_\tau - 2)/2 = 11773(3) \times 10^{-7}$], see SAMUEL 91B.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
> -0.052 and < 0.058 (CL = 95%) OUR LIMIT				
> -0.052 and < 0.058	95	ACCIARRI	98E L3	1991–1995 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> -0.007 and < 0.005	95	⁷ GONZALEZ-S..00	RVUE	$e^+e^- \rightarrow \tau^+\tau^-$ and $W \rightarrow \tau\nu_\tau$
> -0.068 and < 0.065	95	⁸ ACKERSTAFF	98N OPAL	1990–1995 LEP runs
> -0.004 and < 0.006	95	⁹ ESCRIBANO	97 RVUE	$Z \rightarrow \tau^+\tau^-$ at LEP
< 0.01	95	¹⁰ ESCRIBANO	93 RVUE	$Z \rightarrow \tau^+\tau^-$ at LEP
< 0.12	90	GRIFOLS	91 RVUE	$Z \rightarrow \tau\tau\gamma$ at LEP
< 0.023	95	¹¹ SILVERMAN	83 RVUE	$e^+e^- \rightarrow \tau^+\tau^-$ at PETRA

⁷ GONZALEZ-SPRINBERG 00 use data on tau lepton production at LEP1, SLC, and LEP2, and data from colliders and LEP2 to determine limits. Assume imaginary component is zero.

⁸ ACKERSTAFF 98N use $Z \rightarrow \tau^+\tau^-\gamma$ events. The limit applies to an average of the form factor for off-shell τ 's having p^2 ranging from m_τ^2 to $(M_Z - m_\tau)^2$.

⁹ ESCRIBANO 97 use preliminary experimental results.

¹⁰ ESCRIBANO 93 limit derived from $\Gamma(Z \rightarrow \tau^+ \tau^-)$, and is on the absolute value of the magnetic moment anomaly.

¹¹ SILVERMAN 83 limit is derived from $e^+ e^- \rightarrow \tau^+ \tau^-$ total cross-section measurements for q^2 up to $(37 \text{ GeV})^2$.

τ ELECTRIC DIPOLE MOMENT (d_τ)

A nonzero value is forbidden by both T invariance and P invariance.

The q^2 dependence is expected to be small providing no thresholds are nearby.

Re(d_τ)

VALUE (10^{-16} e cm)	CL%	DOCUMENT ID	TECN	COMMENT
(> -0.22 and < 0.45)	95	¹² INAMI	03 BELL	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<4.6	95	¹³ ALBRECHT	00 ARG	$E_{\text{cm}}^{ee} = 10.4 \text{ GeV}$
> -3.1 and < 3.1	95	ACCIARRI	98E L3	1991-1995 LEP runs
> -3.8 and < 3.6	95	¹⁴ ACKERSTAFF	98N OPAL	1990-1995 LEP runs
<0.11	95	^{15,16} ESCRIBANO	97 RVUE	$Z \rightarrow \tau^+ \tau^-$ at LEP
<0.5	95	¹⁷ ESCRIBANO	93 RVUE	$Z \rightarrow \tau^+ \tau^-$ at LEP
<7	90	GRIFOLS	91 RVUE	$Z \rightarrow \tau \tau \gamma$ at LEP
<1.6	90	DELAGUILA	90 RVUE	$e^+ e^- \rightarrow \tau^+ \tau^-$ $E_{\text{cm}}^{ee} = 35 \text{ GeV}$

¹² INAMI 03 use $e^+ e^- \rightarrow \tau^+ \tau^-$ events.

¹³ ALBRECHT 00 use $e^+ e^- \rightarrow \tau^+ \tau^-$ events. Limit is on the absolute value of Re(d_τ).

¹⁴ ACKERSTAFF 98N use $Z \rightarrow \tau^+ \tau^- \gamma$ events. The limit applies to an average of the form factor for off-shell τ 's having p^2 ranging from m_τ^2 to $(M_Z - m_\tau)^2$.

¹⁵ ESCRIBANO 97 derive the relationship $|d_\tau| = \cot \theta_W |d_\tau^W|$ using effective Lagrangian methods, and use a conference result $|d_\tau^W| < 5.8 \times 10^{-18} \text{ e cm}$ at 95% CL (L. Silvestris, ICHEP96) to obtain this result.

¹⁶ ESCRIBANO 97 use preliminary experimental results.

¹⁷ ESCRIBANO 93 limit derived from $\Gamma(Z \rightarrow \tau^+ \tau^-)$, and is on the absolute value of the electric dipole moment.

Im(d_τ)

VALUE (10^{-16} e cm)	CL%	DOCUMENT ID	TECN	COMMENT
(> -0.25 and < 0.008)	95	¹⁸ INAMI	03 BELL	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<1.8	95	¹⁹ ALBRECHT	00 ARG	$E_{\text{cm}}^{ee} = 10.4 \text{ GeV}$
¹⁸ INAMI 03 use $e^+ e^- \rightarrow \tau^+ \tau^-$ events.				
¹⁹ ALBRECHT 00 use $e^+ e^- \rightarrow \tau^+ \tau^-$ events. Limit is on the absolute value of Im(d_τ).				

τ WEAK DIPOLE MOMENT (d_{τ}^W)

A nonzero value is forbidden by CP invariance.

The q^2 dependence is expected to be small providing no thresholds are nearby.

$\text{Re}(d_{\tau}^W)$

VALUE (10^{-17} ecm)	CL%	DOCUMENT ID	TECN	COMMENT
<0.56	95	ACKERSTAFF 97L	OPAL	1991–1995 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<3.0	90	²⁰ ACCIARRI	98C L3	1991–1995 LEP runs
<0.78	95	²¹ AKERS	95F OPAL	Repl. by ACKER-STAFF 97L
<1.5	95	²¹ BUSKULIC	95C ALEP	1990–1992 LEP runs
<7.0	95	²¹ ACTON	92F OPAL	$Z \rightarrow \tau^+ \tau^-$ at LEP
<3.7	95	²¹ BUSKULIC	92J ALEP	Repl. by BUSKULIC 95C

²⁰ ACCIARRI 98C limit is on the absolute value of the real part of the weak dipole moment.

²¹ Limit is on the absolute value of the real part of the weak dipole moment, and applies for $q^2 = m_Z^2$.

$\text{Im}(d_{\tau}^W)$

VALUE (10^{-17} ecm)	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	95	ACKERSTAFF 97L	OPAL	1991–1995 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<4.5	95	²² AKERS	95F OPAL	Repl. by ACKER-STAFF 97L

²² Limit is on the absolute value of the imaginary part of the weak dipole moment, and applies for $q^2 = m_Z^2$.

τ WEAK ANOMALOUS MAGNETIC DIPOLE MOMENT (α_{τ}^W)

Electroweak radiative corrections are expected to contribute at the 10^{-6} level. See BERNABEU 95.

The q^2 dependence is expected to be small providing no thresholds are nearby.

$\text{Re}(\alpha_{\tau}^W)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.5 × 10⁻³	90	²³ ACCIARRI	98C L3	1991–1995 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> -0.0024 and < 0.0025	95	²⁴ GONZALEZ-S..00	RVUE	$e^+ e^- \rightarrow \tau^+ \tau^-$ and $W \rightarrow \tau \nu_{\tau}$

²³ ACCIARRI 98C limit is on the absolute value of the real part of the weak anomalous magnetic dipole moment.

²⁴ GONZALEZ-SPRINBERG 00 use data on tau lepton production at LEP1, SLC, and LEP2, and data from colliders and LEP2 to determine limits. Assume imaginary component is zero.

$\text{Im}(\alpha_{\tau}^w)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.9 \times 10^{-3}$	90	²⁵ ACCIARRI	98C L3	1991–1995 LEP runs

²⁵ ACCIARRI 98C limit is on the absolute value of the imaginary part of the weak anomalous magnetic dipole moment.

τ^- DECAY MODES

τ^+ modes are charge conjugates of the modes below. “ h^\pm ” stands for π^\pm or K^\pm . “ ℓ ” stands for e or μ . “Neutrals” stands for γ 's and/or π^0 's.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Modes with one charged particle		
Γ_1 particle ⁻ ≥ 0 neutrals $\geq 0K^0\nu_\tau$ (“1-prong”)	(85.35±0.07) %	S=1.1
Γ_2 particle ⁻ ≥ 0 neutrals $\geq 0K_L^0\nu_\tau$	(84.71±0.07) %	S=1.1
Γ_3 $\mu^- \bar{\nu}_\mu \nu_\tau$	[a] (17.36±0.05) %	
Γ_4 $\mu^- \bar{\nu}_\mu \nu_\tau \gamma$	[b] (3.6 ±0.4) × 10 ⁻³	
Γ_5 $e^- \bar{\nu}_e \nu_\tau$	[a] (17.84±0.06) %	
Γ_6 $e^- \bar{\nu}_e \nu_\tau \gamma$	[b] (1.75±0.18) %	
Γ_7 $h^- \geq 0K_L^0 \nu_\tau$	(12.30±0.10) %	S=1.4
Γ_8 $h^- \nu_\tau$	(11.75±0.10) %	S=1.4
Γ_9 $\pi^- \nu_\tau$	[a] (11.06±0.11) %	S=1.4
Γ_{10} $K^- \nu_\tau$	[a] (6.86±0.23) × 10 ⁻³	
Γ_{11} $h^- \geq 1$ neutrals ν_τ	(36.92±0.14) %	S=1.1
Γ_{12} $h^- \pi^0 \nu_\tau$	(25.86±0.13) %	S=1.1
Γ_{13} $\pi^- \pi^0 \nu_\tau$	[a] (25.41±0.14) %	S=1.1
Γ_{14} $\pi^- \pi^0$ non- $\rho(770) \nu_\tau$	(3.0 ±3.2) × 10 ⁻³	
Γ_{15} $K^- \pi^0 \nu_\tau$	[a] (4.50±0.30) × 10 ⁻³	
Γ_{16} $h^- \geq 2\pi^0 \nu_\tau$	(10.76±0.15) %	S=1.1
Γ_{17} $h^- 2\pi^0 \nu_\tau$	(9.39±0.14) %	S=1.1
Γ_{18} $h^- 2\pi^0 \nu_\tau$ (ex. K^0)	(9.23±0.14) %	S=1.1
Γ_{19} $\pi^- 2\pi^0 \nu_\tau$ (ex. K^0)	[a] (9.17±0.14) %	S=1.1
Γ_{20} $\pi^- 2\pi^0 \nu_\tau$ (ex. K^0),	< 9 × 10 ⁻³	CL=95%
Γ_{21} scalar $\pi^- 2\pi^0 \nu_\tau$ (ex. K^0),	< 7 × 10 ⁻³	CL=95%
Γ_{22} vector $K^- 2\pi^0 \nu_\tau$ (ex. K^0)	[a] (5.8 ±2.3) × 10 ⁻⁴	
Γ_{23} $h^- \geq 3\pi^0 \nu_\tau$	(1.37±0.11) %	S=1.1
Γ_{24} $h^- 3\pi^0 \nu_\tau$	(1.21±0.10) %	
Γ_{25} $\pi^- 3\pi^0 \nu_\tau$ (ex. K^0)	[a] (1.08±0.10) %	
Γ_{26} $K^- 3\pi^0 \nu_\tau$ (ex. K^0, η)	[a] (3.7 $\begin{smallmatrix} +2.2 \\ -2.0 \end{smallmatrix}$) × 10 ⁻⁴	
Γ_{27} $h^- 4\pi^0 \nu_\tau$ (ex. K^0)	(1.6 ±0.6) × 10 ⁻³	

Γ_{28}	$h^- 4\pi^0 \nu_\tau$ (ex. K^0, η)	[a]	$(1.0^{+0.6}_{-0.5}) \times 10^{-3}$	
Γ_{29}	$K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma \nu_\tau$		$(1.56 \pm 0.04) \%$	
Γ_{30}	$K^- \geq 1 (\pi^0 \text{ or } K^0 \text{ or } \gamma) \nu_\tau$		$(8.74 \pm 0.35) \times 10^{-3}$	

Modes with K^0 's

Γ_{31}	K_S^0 (particles) $^- \nu_\tau$		$(9.2 \pm 0.4) \times 10^{-3}$	S=1.1
Γ_{32}	$h^- \bar{K}^0 \nu_\tau$		$(1.05 \pm 0.04) \%$	S=1.1
Γ_{33}	$\pi^- \bar{K}^0 \nu_\tau$	[a]	$(8.9 \pm 0.4) \times 10^{-3}$	S=1.1
Γ_{34}	$\pi^- \bar{K}^0$ (non- $K^*(892)^-$) ν_τ		$< 1.7 \times 10^{-3}$	CL=95%
Γ_{35}	$K^- K^0 \nu_\tau$	[a]	$(1.54 \pm 0.16) \times 10^{-3}$	
Γ_{36}	$K^- K^0 \geq 0\pi^0 \nu_\tau$		$(3.09 \pm 0.24) \times 10^{-3}$	
Γ_{37}	$h^- \bar{K}^0 \pi^0 \nu_\tau$		$(5.2 \pm 0.4) \times 10^{-3}$	
Γ_{38}	$\pi^- \bar{K}^0 \pi^0 \nu_\tau$	[a]	$(3.7 \pm 0.4) \times 10^{-3}$	
Γ_{39}	$\bar{K}^0 \rho^- \nu_\tau$		$(2.2 \pm 0.5) \times 10^{-3}$	
Γ_{40}	$K^- K^0 \pi^0 \nu_\tau$	[a]	$(1.55 \pm 0.20) \times 10^{-3}$	
Γ_{41}	$\pi^- \bar{K}^0 \geq 1\pi^0 \nu_\tau$		$(3.2 \pm 1.0) \times 10^{-3}$	
Γ_{42}	$\pi^- \bar{K}^0 \pi^0 \pi^0 \nu_\tau$		$(2.6 \pm 2.4) \times 10^{-4}$	
Γ_{43}	$K^- K^0 \pi^0 \pi^0 \nu_\tau$		$< 1.6 \times 10^{-4}$	CL=95%
Γ_{44}	$\pi^- K^0 \bar{K}^0 \nu_\tau$		$(1.59 \pm 0.29) \times 10^{-3}$	S=1.1
Γ_{45}	$\pi^- K_S^0 K_S^0 \nu_\tau$	[a]	$(2.4 \pm 0.5) \times 10^{-4}$	
Γ_{46}	$\pi^- K_S^0 K_L^0 \nu_\tau$	[a]	$(1.10 \pm 0.28) \times 10^{-3}$	S=1.1
Γ_{47}	$\pi^- K^0 \bar{K}^0 \pi^0 \nu_\tau$		$(3.1 \pm 2.3) \times 10^{-4}$	
Γ_{48}	$\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau$		$< 2.0 \times 10^{-4}$	CL=95%
Γ_{49}	$\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau$		$(3.1 \pm 1.2) \times 10^{-4}$	
Γ_{50}	$K^0 h^+ h^- h^- \geq 0$ neutrals ν_τ		$< 1.7 \times 10^{-3}$	CL=95%
Γ_{51}	$K^0 h^+ h^- h^- \nu_\tau$		$(2.3 \pm 2.0) \times 10^{-4}$	

Modes with three charged particles

Γ_{52}	$h^- h^- h^+ \geq 0$ neutrals $\geq 0K_L^0 \nu_\tau$		$(15.20 \pm 0.07) \%$	S=1.1
Γ_{53}	$h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$) ("3-prong")		$(14.57 \pm 0.07) \%$	S=1.1
Γ_{54}	$h^- h^- h^+ \nu_\tau$		$(10.01 \pm 0.09) \%$	S=1.2
Γ_{55}	$h^- h^- h^+ \nu_\tau$ (ex. K^0)		$(9.65 \pm 0.09) \%$	S=1.2
Γ_{56}	$h^- h^- h^+ \nu_\tau$ (ex. K^0, ω)		$(9.61 \pm 0.09) \%$	S=1.2
Γ_{57}	$\pi^- \pi^+ \pi^- \nu_\tau$		$(9.52 \pm 0.10) \%$	S=1.2
Γ_{58}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0)		$(9.22 \pm 0.10) \%$	S=1.2
Γ_{59}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0), non-axial vector		$< 2.4 \%$	CL=95%
Γ_{60}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0, ω)	[a]	$(9.18 \pm 0.10) \%$	S=1.2
Γ_{61}	$h^- h^- h^+ \geq 1$ neutrals ν_τ		$(5.19 \pm 0.10) \%$	S=1.3
Γ_{62}	$h^- h^- h^+ \geq 1$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$)		$(4.92 \pm 0.10) \%$	S=1.3

Γ ₆₃	$h^- h^- h^+ \pi^0 \nu_\tau$	(4.53 ± 0.09) %	S=1.3
Γ ₆₄	$h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0)	(4.35 ± 0.09) %	S=1.3
Γ ₆₅	$h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0, ω)	(2.62 ± 0.09) %	S=1.2
Γ ₆₆	$\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$	(4.37 ± 0.10) %	S=1.3
Γ ₆₇	$\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0)	(4.24 ± 0.10) %	S=1.3
Γ ₆₈	$\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0, ω)	[a] (2.51 ± 0.09) %	S=1.2
Γ ₆₉	$h^- \rho \pi^0 \nu_\tau$		
Γ ₇₀	$h^- \rho^+ h^- \nu_\tau$		
Γ ₇₁	$h^- \rho^- h^+ \nu_\tau$		
Γ ₇₂	$h^- h^- h^+ 2\pi^0 \nu_\tau$	(5.5 ± 0.4) × 10 ⁻³	
Γ ₇₃	$h^- h^- h^+ 2\pi^0 \nu_\tau$ (ex. K^0)	(5.4 ± 0.4) × 10 ⁻³	
Γ ₇₄	$h^- h^- h^+ 2\pi^0 \nu_\tau$ (ex. K^0, ω, η)	[a] (1.1 ± 0.4) × 10 ⁻³	
Γ ₇₅	$h^- h^- h^+ 3\pi^0 \nu_\tau$	[a] (2.3 ± 0.8) × 10 ⁻⁴	S=1.6
Γ ₇₆	$K^- h^+ h^- \geq 0$ neutrals ν_τ	(6.5 ± 0.5) × 10 ⁻³	S=1.4
Γ ₇₇	$K^- h^+ \pi^- \nu_\tau$ (ex. K^0)	(4.4 ± 0.5) × 10 ⁻³	S=1.5
Γ ₇₈	$K^- h^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0)	(1.10 ± 0.22) × 10 ⁻³	
Γ ₇₉	$K^- \pi^+ \pi^- \geq 0$ neutrals ν_τ	(4.5 ± 0.5) × 10 ⁻³	S=1.4
Γ ₈₀	$K^- \pi^+ \pi^- \geq 0 \pi^0 \nu_\tau$ (ex. K^0)	(3.5 ± 0.5) × 10 ⁻³	S=1.4
Γ ₈₁	$K^- \pi^+ \pi^- \nu_\tau$	(3.3 ± 0.5) × 10 ⁻³	S=1.5
Γ ₈₂	$K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0)	[a] (2.8 ± 0.5) × 10 ⁻³	S=1.5
Γ ₈₃	$K^- \rho^0 \nu_\tau \rightarrow$ $K^- \pi^+ \pi^- \nu_\tau$	(1.3 ± 0.5) × 10 ⁻³	
Γ ₈₄	$K^- \pi^+ \pi^- \pi^0 \nu_\tau$	(1.24 ± 0.25) × 10 ⁻³	
Γ ₈₅	$K^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0)	(7.0 ± 2.4) × 10 ⁻⁴	
Γ ₈₆	$K^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0, η)	[a] (6.4 ± 2.4) × 10 ⁻⁴	
Γ ₈₇	$K^- \pi^+ K^- \geq 0$ neut. ν_τ	< 9 × 10 ⁻⁴	CL=95%
Γ ₈₈	$K^- K^+ \pi^- \geq 0$ neut. ν_τ	(2.00 ± 0.23) × 10 ⁻³	
Γ ₈₉	$K^- K^+ \pi^- \nu_\tau$	[a] (1.60 ± 0.19) × 10 ⁻³	
Γ ₉₀	$K^- K^+ \pi^- \pi^0 \nu_\tau$	[a] (4.0 ± 1.6) × 10 ⁻⁴	
Γ ₉₁	$K^- K^+ K^- \geq 0$ neut. ν_τ	< 2.1 × 10 ⁻³	CL=95%
Γ ₉₂	$K^- K^+ K^- \nu_\tau$	< 1.9 × 10 ⁻⁴	CL=90%
Γ ₉₃	$\pi^- K^+ \pi^- \geq 0$ neut. ν_τ	< 2.5 × 10 ⁻³	CL=95%
Γ ₉₄	$e^- e^- e^+ \bar{\nu}_e \nu_\tau$	(2.8 ± 1.5) × 10 ⁻⁵	
Γ ₉₅	$\mu^- e^- e^+ \bar{\nu}_\mu \nu_\tau$	< 3.6 × 10 ⁻⁵	CL=90%

Modes with five charged particles

Γ ₉₆	$3h^- 2h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^- \pi^+$) ("5-prong")	(1.00 ± 0.06) × 10 ⁻³	
Γ ₉₇	$3h^- 2h^+ \nu_\tau$ (ex. K^0)	[a] (8.2 ± 0.6) × 10 ⁻⁴	
Γ ₉₈	$3h^- 2h^+ \pi^0 \nu_\tau$ (ex. K^0)	[a] (1.81 ± 0.27) × 10 ⁻⁴	
Γ ₉₉	$3h^- 2h^+ 2\pi^0 \nu_\tau$	< 1.1 × 10 ⁻⁴	CL=90%

Miscellaneous other allowed modes

Γ_{100}	$(5\pi)^- \nu_\tau$	$(8.0 \pm 0.7) \times 10^{-3}$	
Γ_{101}	$4h^- 3h^+ \geq 0$ neutrals ν_τ ("7-prong")	$< 2.4 \times 10^{-6}$	CL=90%
Γ_{102}	$X^-(S=-1)\nu_\tau$	$(2.86 \pm 0.09) \%$	S=1.1
Γ_{103}	$K^*(892)^- \geq 0$ neutrals \geq $0K_L^0 \nu_\tau$	$(1.42 \pm 0.18) \%$	S=1.4
Γ_{104}	$K^*(892)^- \nu_\tau$	$(1.29 \pm 0.05) \%$	
Γ_{105}	$K^*(892)^0 K^- \geq 0$ neutrals ν_τ	$(3.2 \pm 1.4) \times 10^{-3}$	
Γ_{106}	$K^*(892)^0 K^- \nu_\tau$	$(2.1 \pm 0.4) \times 10^{-3}$	
Γ_{107}	$\bar{K}^*(892)^0 \pi^- \geq 0$ neutrals ν_τ	$(3.8 \pm 1.7) \times 10^{-3}$	
Γ_{108}	$\bar{K}^*(892)^0 \pi^- \nu_\tau$	$(2.2 \pm 0.5) \times 10^{-3}$	
Γ_{109}	$(\bar{K}^*(892)\pi)^- \nu_\tau \rightarrow$ $\pi^- \bar{K}^0 \pi^0 \nu_\tau$	$(1.0 \pm 0.4) \times 10^{-3}$	
Γ_{110}	$K_1(1270)^- \nu_\tau$	$(4.7 \pm 1.1) \times 10^{-3}$	
Γ_{111}	$K_1(1400)^- \nu_\tau$	$(1.7 \pm 2.6) \times 10^{-3}$	S=1.7
Γ_{112}	$K^*(1410)^- \nu_\tau$	$(1.5 \begin{smallmatrix} +1.4 \\ -1.0 \end{smallmatrix}) \times 10^{-3}$	
Γ_{113}	$K_0^*(1430)^- \nu_\tau$	$< 5 \times 10^{-4}$	CL=95%
Γ_{114}	$K_2^*(1430)^- \nu_\tau$	$< 3 \times 10^{-3}$	CL=95%
Γ_{115}	$a_0(980)^- \geq 0$ neutrals ν_τ		
Γ_{116}	$\eta \pi^- \nu_\tau$	$< 1.4 \times 10^{-4}$	CL=95%
Γ_{117}	$\eta \pi^- \pi^0 \nu_\tau$	[a] $(1.74 \pm 0.24) \times 10^{-3}$	
Γ_{118}	$\eta \pi^- \pi^0 \pi^0 \nu_\tau$	$(1.5 \pm 0.5) \times 10^{-4}$	
Γ_{119}	$\eta K^- \nu_\tau$	[a] $(2.7 \pm 0.6) \times 10^{-4}$	
Γ_{120}	$\eta K^*(892)^- \nu_\tau$	$(2.9 \pm 0.9) \times 10^{-4}$	
Γ_{121}	$\eta K^- \pi^0 \nu_\tau$	$(1.8 \pm 0.9) \times 10^{-4}$	
Γ_{122}	$\eta \bar{K}^0 \pi^- \nu_\tau$	$(2.2 \pm 0.7) \times 10^{-4}$	
Γ_{123}	$\eta \pi^+ \pi^- \pi^- \geq 0$ neutrals ν_τ	$< 3 \times 10^{-3}$	CL=90%
Γ_{124}	$\eta \pi^- \pi^+ \pi^- \nu_\tau$	$(2.3 \pm 0.5) \times 10^{-4}$	
Γ_{125}	$\eta a_1(1260)^- \nu_\tau \rightarrow \eta \pi^- \rho^0 \nu_\tau$	$< 3.9 \times 10^{-4}$	CL=90%
Γ_{126}	$\eta \eta \pi^- \nu_\tau$	$< 1.1 \times 10^{-4}$	CL=95%
Γ_{127}	$\eta \eta \pi^- \pi^0 \nu_\tau$	$< 2.0 \times 10^{-4}$	CL=95%
Γ_{128}	$\eta'(958) \pi^- \nu_\tau$	$< 7.4 \times 10^{-5}$	CL=90%
Γ_{129}	$\eta'(958) \pi^- \pi^0 \nu_\tau$	$< 8.0 \times 10^{-5}$	CL=90%
Γ_{130}	$\phi \pi^- \nu_\tau$	$< 2.0 \times 10^{-4}$	CL=90%
Γ_{131}	$\phi K^- \nu_\tau$	$< 6.7 \times 10^{-5}$	CL=90%
Γ_{132}	$f_1(1285) \pi^- \nu_\tau$	$(5.8 \pm 2.3) \times 10^{-4}$	
Γ_{133}	$f_1(1285) \pi^- \nu_\tau \rightarrow$ $\eta \pi^- \pi^+ \pi^- \nu_\tau$	$(1.3 \pm 0.4) \times 10^{-4}$	
Γ_{134}	$\pi(1300)^- \nu_\tau \rightarrow (\rho \pi)^- \nu_\tau \rightarrow$ $(3\pi)^- \nu_\tau$	$< 1.0 \times 10^{-4}$	CL=90%
Γ_{135}	$\pi(1300)^- \nu_\tau \rightarrow$ $((\pi \pi)_{S\text{-wave}} \pi)^- \nu_\tau \rightarrow$ $(3\pi)^- \nu_\tau$	$< 1.9 \times 10^{-4}$	CL=90%

Γ_{136}	$h^- \omega \geq 0$ neutrals ν_τ	(2.37 ± 0.08) %
Γ_{137}	$h^- \omega \nu_\tau$	[a] (1.94 ± 0.07) %
Γ_{138}	$h^- \omega \pi^0 \nu_\tau$	[a] (4.3 ± 0.5) $\times 10^{-3}$
Γ_{139}	$h^- \omega 2\pi^0 \nu_\tau$	(1.4 ± 0.5) $\times 10^{-4}$
Γ_{140}	$2h^- h^+ \omega \nu_\tau$	(1.20 ± 0.22) $\times 10^{-4}$

**Lepton Family number (LF), Lepton number (L),
or Baryon number (B) violating modes**

L means lepton number violation (e.g. $\tau^- \rightarrow e^+ \pi^- \pi^-$). Following common usage, *LF* means lepton family violation *and not* lepton number violation (e.g. $\tau^- \rightarrow e^- \pi^+ \pi^-$). *B* means baryon number violation.

Γ_{141}	$e^- \gamma$	<i>LF</i>	< 2.7	$\times 10^{-6}$	CL=90%
Γ_{142}	$\mu^- \gamma$	<i>LF</i>	< 1.1	$\times 10^{-6}$	CL=90%
Γ_{143}	$e^- \pi^0$	<i>LF</i>	< 3.7	$\times 10^{-6}$	CL=90%
Γ_{144}	$\mu^- \pi^0$	<i>LF</i>	< 4.0	$\times 10^{-6}$	CL=90%
Γ_{145}	$e^- K_S^0$	<i>LF</i>	< 9.1	$\times 10^{-7}$	CL=90%
Γ_{146}	$\mu^- K_S^0$	<i>LF</i>	< 9.5	$\times 10^{-7}$	CL=90%
Γ_{147}	$e^- \eta$	<i>LF</i>	< 8.2	$\times 10^{-6}$	CL=90%
Γ_{148}	$\mu^- \eta$	<i>LF</i>	< 9.6	$\times 10^{-6}$	CL=90%
Γ_{149}	$e^- \rho^0$	<i>LF</i>	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{150}	$\mu^- \rho^0$	<i>LF</i>	< 6.3	$\times 10^{-6}$	CL=90%
Γ_{151}	$e^- K^*(892)^0$	<i>LF</i>	< 5.1	$\times 10^{-6}$	CL=90%
Γ_{152}	$\mu^- K^*(892)^0$	<i>LF</i>	< 7.5	$\times 10^{-6}$	CL=90%
Γ_{153}	$e^- \bar{K}^*(892)^0$	<i>LF</i>	< 7.4	$\times 10^{-6}$	CL=90%
Γ_{154}	$\mu^- \bar{K}^*(892)^0$	<i>LF</i>	< 7.5	$\times 10^{-6}$	CL=90%
Γ_{155}	$e^- \phi$	<i>LF</i>	< 6.9	$\times 10^{-6}$	CL=90%
Γ_{156}	$\mu^- \phi$	<i>LF</i>	< 7.0	$\times 10^{-6}$	CL=90%
Γ_{157}	$e^- e^+ e^-$	<i>LF</i>	< 2.9	$\times 10^{-6}$	CL=90%
Γ_{158}	$e^- \mu^+ \mu^-$	<i>LF</i>	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{159}	$e^+ \mu^- \mu^-$	<i>LF</i>	< 1.5	$\times 10^{-6}$	CL=90%
Γ_{160}	$\mu^- e^+ e^-$	<i>LF</i>	< 1.7	$\times 10^{-6}$	CL=90%
Γ_{161}	$\mu^+ e^- e^-$	<i>LF</i>	< 1.5	$\times 10^{-6}$	CL=90%
Γ_{162}	$\mu^- \mu^+ \mu^-$	<i>LF</i>	< 1.9	$\times 10^{-6}$	CL=90%
Γ_{163}	$e^- \pi^+ \pi^-$	<i>LF</i>	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{164}	$e^+ \pi^- \pi^-$	<i>L</i>	< 1.9	$\times 10^{-6}$	CL=90%
Γ_{165}	$\mu^- \pi^+ \pi^-$	<i>LF</i>	< 8.2	$\times 10^{-6}$	CL=90%
Γ_{166}	$\mu^+ \pi^- \pi^-$	<i>L</i>	< 3.4	$\times 10^{-6}$	CL=90%
Γ_{167}	$e^- \pi^+ K^-$	<i>LF</i>	< 6.4	$\times 10^{-6}$	CL=90%
Γ_{168}	$e^- \pi^- K^+$	<i>LF</i>	< 3.8	$\times 10^{-6}$	CL=90%
Γ_{169}	$e^+ \pi^- K^-$	<i>L</i>	< 2.1	$\times 10^{-6}$	CL=90%
Γ_{170}	$e^- K_S^0 K_S^0$	<i>LF</i>	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{171}	$e^- K^+ K^-$	<i>LF</i>	< 6.0	$\times 10^{-6}$	CL=90%
Γ_{172}	$e^+ K^- K^-$	<i>L</i>	< 3.8	$\times 10^{-6}$	CL=90%
Γ_{173}	$\mu^- \pi^+ K^-$	<i>LF</i>	< 7.5	$\times 10^{-6}$	CL=90%

Γ_{174}	$\mu^- \pi^- K^+$	<i>LF</i>	< 7.4	$\times 10^{-6}$	CL=90%
Γ_{175}	$\mu^+ \pi^- K^-$	<i>L</i>	< 7.0	$\times 10^{-6}$	CL=90%
Γ_{176}	$\mu^- K_S^0 K_S^0$	<i>LF</i>	< 3.4	$\times 10^{-6}$	CL=90%
Γ_{177}	$\mu^- K^+ K^-$	<i>LF</i>	< 1.5	$\times 10^{-5}$	CL=90%
Γ_{178}	$\mu^+ K^- K^-$	<i>L</i>	< 6.0	$\times 10^{-6}$	CL=90%
Γ_{179}	$e^- \pi^0 \pi^0$	<i>LF</i>	< 6.5	$\times 10^{-6}$	CL=90%
Γ_{180}	$\mu^- \pi^0 \pi^0$	<i>LF</i>	< 1.4	$\times 10^{-5}$	CL=90%
Γ_{181}	$e^- \eta \eta$	<i>LF</i>	< 3.5	$\times 10^{-5}$	CL=90%
Γ_{182}	$\mu^- \eta \eta$	<i>LF</i>	< 6.0	$\times 10^{-5}$	CL=90%
Γ_{183}	$e^- \pi^0 \eta$	<i>LF</i>	< 2.4	$\times 10^{-5}$	CL=90%
Γ_{184}	$\mu^- \pi^0 \eta$	<i>LF</i>	< 2.2	$\times 10^{-5}$	CL=90%
Γ_{185}	$\bar{p} \gamma$	<i>L,B</i>	< 3.5	$\times 10^{-6}$	CL=90%
Γ_{186}	$\bar{p} \pi^0$	<i>L,B</i>	< 1.5	$\times 10^{-5}$	CL=90%
Γ_{187}	$\bar{p} 2\pi^0$	<i>L,B</i>	< 3.3	$\times 10^{-5}$	CL=90%
Γ_{188}	$\bar{p} \eta$	<i>L,B</i>	< 8.9	$\times 10^{-6}$	CL=90%
Γ_{189}	$\bar{p} \pi^0 \eta$	<i>L,B</i>	< 2.7	$\times 10^{-5}$	CL=90%
Γ_{190}	e^- light boson	<i>LF</i>	< 2.7	$\times 10^{-3}$	CL=95%
Γ_{191}	μ^- light boson	<i>LF</i>	< 5	$\times 10^{-3}$	CL=95%

[a] Basis mode for the τ .

[b] See the Particle Listings below for the energy limits used in this measurement.

CONSTRAINED FIT INFORMATION

An overall fit to 64 branching ratios uses 126 measurements and one constraint to determine 31 parameters. The overall fit has a $\chi^2 = 57.3$ for 96 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_5	1									
x_9	-1	-1								
x_{10}	0	0	-21							
x_{13}	-14	-14	-25	1						
x_{15}	0	0	1	-3	-22					
x_{19}	-14	-15	-25	0	-36	4				
x_{22}	0	0	1	-2	5	-21	-16			
x_{25}	-8	-8	-15	0	-18	5	-24	3		
x_{26}	0	0	1	-2	5	-19	3	-13	-22	
x_{28}	-4	-4	-7	0	-10	0	-12	0	-7	0
x_{33}	-2	-2	-12	0	-3	1	-8	1	-2	1
x_{35}	-1	-1	-4	-1	2	-12	-1	-8	2	-8
x_{38}	-2	-2	-3	0	-5	3	-5	2	-9	2
x_{40}	-1	-1	0	-1	2	-16	1	-11	-1	-10
x_{45}	0	0	-2	0	0	0	0	0	0	0
x_{46}	-2	-2	-2	0	-4	1	-4	1	-2	1
x_{60}	-1	-1	-2	0	-3	0	-3	0	-2	0
x_{68}	-1	-1	-3	0	-3	0	-4	0	-2	0
x_{74}	0	0	0	0	0	0	0	0	0	0
x_{75}	0	0	0	0	0	0	0	0	0	0
x_{82}	0	0	0	0	0	0	0	0	0	0
x_{86}	0	0	0	0	0	0	0	0	0	0
x_{89}	0	0	0	0	0	0	0	0	0	0
x_{90}	0	0	0	0	0	0	0	0	0	0
x_{97}	0	0	-1	0	-1	0	-1	0	0	0
x_{98}	0	0	0	0	0	0	0	0	0	0
x_{117}	-1	-1	-1	0	-2	0	-2	0	-1	0
x_{119}	0	0	0	0	0	-3	0	-2	-2	-2
x_{137}	-1	-1	-2	0	-3	0	-3	0	-2	0
x_{138}	-1	-1	-1	0	-2	0	-2	0	-1	0
	x_3	x_5	x_9	x_{10}	x_{13}	x_{15}	x_{19}	x_{22}	x_{25}	x_{26}

x33	-1									
x35	0	-5								
x38	-1	-7	0							
x40	0	-2	-15	-19						
x45	0	-2	-1	-2	0					
x46	-1	-12	-4	-10	-3	-3				
x60	-1	-7	-3	3	2	0	0			
x68	-1	4	2	-5	-2	0	0	-45		
x74	2	1	0	1	0	0	0	-7	-7	
x75	0	0	0	0	0	0	0	-3	-3	0
x82	0	0	0	0	0	0	0	-43	2	0
x86	0	0	0	0	0	0	0	5	-15	0
x89	0	0	0	0	0	0	0	-8	-2	0
x90	0	0	0	0	0	0	0	-6	-1	0
x97	0	0	0	0	0	0	0	-1	-1	0
x98	0	0	0	0	0	0	0	0	0	0
x117	-14	0	0	0	0	0	0	-1	-1	-14
x119	0	0	-1	0	-2	0	0	0	0	0
x137	-1	2	1	-2	-1	0	0	-22	-28	-3
x138	-1	1	0	1	0	0	0	-8	-10	-44
	x28	x33	x35	x38	x40	x45	x46	x60	x68	x74

x82	0									
x86	0	-19								
x89	0	-14	8							
x90	0	10	-46	-14						
x97	0	0	0	0	0					
x98	0	0	0	0	0	-19				
x117	0	0	0	0	0	0	0			
x119	0	0	-6	0	0	0	0	0		
x137	-1	-1	2	-1	2	0	0	0	0	
x138	-1	0	0	0	-1	0	0	0	0	-4
	x75	x82	x86	x89	x90	x97	x98	x117	x119	x137

A REVIEW GOES HERE – Check our WWW List of Reviews

τ^- BRANCHING RATIOS

$$\Gamma(\text{particle}^- \geq 0 \text{ neutrals} \geq 0 K^0 \nu_\tau \text{ ("1-prong")}) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$


$$\Gamma_1 / \Gamma = (\Gamma_3 + \Gamma_5 + \Gamma_9 + \Gamma_{10} + \Gamma_{13} + \Gamma_{15} + \Gamma_{19} + \Gamma_{22} + \Gamma_{25} + \Gamma_{26} + \Gamma_{28} + \Gamma_{33} + \Gamma_{35} + \Gamma_{38} + \Gamma_{40} + 2\Gamma_{45} + \Gamma_{46} + 0.708\Gamma_{117} + 0.715\Gamma_{119} + 0.09\Gamma_{137} + 0.09\Gamma_{138}) / \Gamma$$

The charged particle here can be e , μ , or hadron. In many analyses, the sum of the topological branching fractions (1, 3, and 5 prongs) is constrained to be unity. Since the 5-prong fraction is very small, the measured 1-prong and 3-prong fractions are highly correlated and cannot be treated as independent quantities in our overall fit. We arbitrarily choose to use the 3-prong fraction in our fit, and leave the 1-prong fraction out. We do, however, use these 1-prong measurements in our average below. The measurements used only for the average are marked "avg," whereas "f&a" marks a result used for the fit and the average.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
85.35 ± 0.07 OUR FIT				Error includes scale factor of 1.1.
85.26 ± 0.13 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
85.316 ± 0.093 ± 0.049	avg	78k	²⁶ ABREU	01M DLPH 1992–1995 LEP runs
85.274 ± 0.105 ± 0.073	avg		²⁷ ACHARD	01D L3 1992–1995 LEP runs
84.48 ± 0.27 ± 0.23	avg		ACTON	92H OPAL 1990–1991 LEP runs
85.45 $\begin{smallmatrix} +0.69 \\ -0.73 \end{smallmatrix}$ ± 0.65	f&a		DECAMP	92C ALEP 1989–1990 LEP runs

²⁶ The correlation coefficients between this measurement and the ABREU 01M measurements of $B(\tau \rightarrow 3\text{-prong})$ and $B(\tau \rightarrow 5\text{-prong})$ are -0.98 and -0.08 respectively.

²⁷ The correlation coefficients between this measurement and the ACHARD 01D measurements of $B(\tau \rightarrow 3\text{-prong})$ and $B(\tau \rightarrow 5\text{-prong})$ are -0.978 and -0.082 respectively.



$$\Gamma(\text{particle}^- \geq 0 \text{ neutrals} \geq 0 K^0 \nu_\tau \text{ ("1-prong")}) / \Gamma_{\text{total}}$$

$$\Gamma_2 / \Gamma = (\Gamma_3 + \Gamma_5 + \Gamma_9 + \Gamma_{10} + \Gamma_{13} + \Gamma_{15} + \Gamma_{19} + \Gamma_{22} + \Gamma_{25} + \Gamma_{26} + \Gamma_{28} + 0.6569\Gamma_{33} + 0.6569\Gamma_{35} + 0.6569\Gamma_{38} + 0.6569\Gamma_{40} + 1.0985\Gamma_{45} + 0.3139\Gamma_{46} + 0.708\Gamma_{117} + 0.715\Gamma_{119} + 0.09\Gamma_{137} + 0.09\Gamma_{138}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
84.71±0.07 OUR FIT	Error includes scale factor of 1.1.			
85.1 ±0.4 OUR AVERAGE				
85.6 ±0.6 ±0.3 avg	3300	²⁸ ADEVA	91F L3	$E_{\text{cm}}^{ee} = 88.3\text{--}94.3 \text{ GeV}$
84.9 ±0.4 ±0.3 avg		BEHREND	89B CELL	$E_{\text{cm}}^{ee} = 14\text{--}47 \text{ GeV}$
84.7 ±0.8 ±0.6 avg		²⁹ AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
86.4 ±0.3 ±0.3		ABACHI	89B HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
87.1 ±1.0 ±0.7		³⁰ BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
87.2 ±0.5 ±0.8		SCHMIDKE	86 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
84.7 ±1.1 $\begin{smallmatrix} +1.6 \\ -1.3 \end{smallmatrix}$	169	³¹ ALTHOFF	85 TASS	$E_{\text{cm}}^{ee} = 34.5 \text{ GeV}$
86.1 ±0.5 ±0.9		BARTEL	85F JADE	$E_{\text{cm}}^{ee} = 34.6 \text{ GeV}$
87.8 ±1.3 ±3.9		³² BERGER	85 PLUT	$E_{\text{cm}}^{ee} = 34.6 \text{ GeV}$
86.7 ±0.3 ±0.6		FERNANDEZ	85 MAC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

²⁸ Not independent of ADEVA 91F $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ value.

²⁹ Not independent of AIHARA 87B $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}$, and Γ_{total} values.

³⁰ Not independent of SCHMIDKE 86 value (also not independent of BURCHAT 87 value for $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$).

³¹ Not independent of ALTHOFF 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$, Γ_{total} , and $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ values.

³² Not independent of (1-prong + $0\pi^0$) and (1-prong + $\geq 1\pi^0$) values.

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$

Γ_3/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

To minimize the effect of experiments with large systematic errors, we exclude experiments which together would contribute 5% of the weight in the average.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
17.36 ±0.05				OUR FIT
17.33 ±0.06				OUR AVERAGE
17.34 ±0.09 ±0.06	f&a 31.4k	ABBIENDI	03 OPAL	1990–1995 LEP runs
17.342±0.110±0.067	f&a 21.5k	³³ ACCIARRI	01F L3	1991–1995 LEP runs
17.325±0.095±0.077	f&a 27.7k	ABREU	99X DLPH	1991–1995 LEP runs
17.37 ±0.08 ±0.18	avg	³⁴ ANASTASSOV	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
17.31 ±0.11 ±0.05	f&a 20.7k	BUSKULIC	96C ALEP	1991–1993 LEP runs
17.35 ±0.41 ±0.37	f&a	DECAMP	92C ALEP	1989–1990 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
17.02 ±0.19 ±0.24	6586	ABREU	95T DLPH	Repl. by ABREU 99X
17.36 ±0.27	7941	AKERS	95I OPAL	Repl. by ABBIENDI 03
17.6 ±0.4 ±0.4	2148	ADRIANI	93M L3	Repl. by ACCIARRI 01F
17.4 ±0.3 ±0.5		³⁵ ALBRECHT	93G ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
17.7 ±0.8 ±0.4	568	BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35$ GeV
17.4 ±1.0	2197	ADEVA	88 MRKJ	$E_{\text{cm}}^{ee} = 14\text{--}16$ GeV
17.7 ±1.2 ±0.7		AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29$ GeV
18.3 ±0.9 ±0.8		BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
18.6 ±0.8 ±0.7	558	³⁶ BARTEL	86D JADE	$E_{\text{cm}}^{ee} = 34.6$ GeV
12.9 ±1.7 $\begin{smallmatrix} +0.7 \\ -0.5 \end{smallmatrix}$		ALTHOFF	85 TASS	$E_{\text{cm}}^{ee} = 34.5$ GeV
18.0 ±0.9 ±0.5	473	³⁶ ASH	85B MAC	$E_{\text{cm}}^{ee} = 29$ GeV
18.0 ±1.0 ±0.6		³⁷ BALTRUSAIT..	85 MRK3	$E_{\text{cm}}^{ee} = 3.77$ GeV
19.4 ±1.6 ±1.7	153	BERGER	85 PLUT	$E_{\text{cm}}^{ee} = 34.6$ GeV
17.6 ±2.6 ±2.1	47	BEHREND	83C CELL	$E_{\text{cm}}^{ee} = 34$ GeV
17.8 ±2.0 ±1.8		BERGER	81B PLUT	$E_{\text{cm}}^{ee} = 9\text{--}32$ GeV

³³ The correlation coefficient between this measurement and the ACCIARRI 01F measurement of $B(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)$ is 0.08.

³⁴ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(e^- \bar{\nu}_e \nu_\tau)$, $B(\mu^- \bar{\nu}_\mu \nu_\tau)/B(e^- \bar{\nu}_e \nu_\tau)$, $B(h^- \nu_\tau)$, and $B(h^- \nu_\tau)/B(e^- \bar{\nu}_e \nu_\tau)$ are 0.50, 0.58, 0.50, and 0.08 respectively.

³⁵ Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ and ALBRECHT 93G $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}^2$ values.

³⁶ Modified using $B(e^- \bar{\nu}_e \nu_\tau)/B(\text{"1 prong"})$ and $B(\text{"1 prong"}) = 0.855$.

³⁷ Error correlated with BALTRUSAITIS 85 $e\nu\bar{\nu}$ value.

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.361 ± 0.016 ± 0.035		³⁸ BERGFELD 00	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
0.30 ± 0.04 ± 0.05	116	³⁹ ALEXANDER 96S	OPAL	1991–1994 LEP runs
0.23 ± 0.10	10	⁴⁰ WU 90	MRK2	$E_{\text{cm}}^{ee} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

³⁸ BERGFELD 00 impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_\gamma^* > 10$ MeV. For $E_\gamma^* > 20$ MeV, they quote $(3.04 \pm 0.14 \pm 0.30) \times 10^{-3}$.

³⁹ ALEXANDER 96S impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_\gamma > 20$ MeV.

⁴⁰ WU 90 reports $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) = 0.013 \pm 0.006$, which is converted to $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma_{\text{total}}$ using $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma_{\text{total}} = 17.35\%$. Requirements on detected γ 's correspond to a τ rest frame energy cutoff $E_\gamma > 37$ MeV.

$\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$ Γ_5/Γ

To minimize the effect of experiments with large systematic errors, we exclude experiments which together would contribute 5% of the weight in the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
17.84 ± 0.06 OUR FIT				
17.81 ± 0.06 OUR AVERAGE				
17.806 ± 0.104 ± 0.076	24.7k	⁴¹ ACCIARRI 01F	L3	1991–1995 LEP runs
17.81 ± 0.09 ± 0.06	33.1k	ABBIENDI 99H	OPAL	1991–1995 LEP runs
17.877 ± 0.109 ± 0.110	23.3k	ABREU 99X	DLPH	1991–1995 LEP runs
17.76 ± 0.06 ± 0.17		⁴² ANASTASSOV 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
17.79 ± 0.12 ± 0.06	20.6k	BUSKULIC 96C	ALEP	1991–1993 LEP runs
18.09 ± 0.45 ± 0.45		DECAMP 92C	ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •				
17.78 ± 0.10 ± 0.09	25.3k	ALEXANDER 96D	OPAL	Repl. by ABBI- ENDI 99H
17.51 ± 0.23 ± 0.31	5059	ABREU 95T	DLPH	Repl. by ABREU 99X
17.9 ± 0.4 ± 0.4	2892	ADRIANI 93M	L3	Repl. by ACCIA- RRI 01F
17.5 ± 0.3 ± 0.5		⁴³ ALBRECHT 93G	ARG	$E_{\text{cm}}^{ee} = 9.4$ – 10.6 GeV
17.97 ± 0.14 ± 0.23	3970	AKERIB 92	CLEO	Repl. by ANAS- TASSOV 97
19.1 ± 0.4 ± 0.6	2960	⁴⁴ AMMAR 92	CLEO	$E_{\text{cm}}^{ee} = 10.5$ – 10.9 GeV
17.0 ± 0.5 ± 0.6	1.7k	ABACHI 90	HRS	$E_{\text{cm}}^{ee} = 29$ GeV
18.4 ± 0.8 ± 0.4	644	BEHREND 90	CELL	$E_{\text{cm}}^{ee} = 35$ GeV
16.3 ± 0.3 ± 3.2		JANSSEN 89	CBAL	$E_{\text{cm}}^{ee} = 9.4$ – 10.6 GeV
18.4 ± 1.2 ± 1.0		AIHARA 87B	TPC	$E_{\text{cm}}^{ee} = 29$ GeV
19.1 ± 0.8 ± 1.1		BURCHAT 87	MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
16.8 ± 0.7 ± 0.9	515	⁴⁴ BARTEL 86D	JADE	$E_{\text{cm}}^{ee} = 34.6$ GeV
20.4 ± 3.0 $\begin{smallmatrix} +1.4 \\ -0.9 \end{smallmatrix}$		ALTHOFF 85	TASS	$E_{\text{cm}}^{ee} = 34.5$ GeV

17.8	± 0.9	± 0.6	390	44 ASH	85B MAC	$E_{cm}^{ee} = 29$ GeV
18.2	± 0.7	± 0.5		45 BALTRUSAIT..85	MRK3	$E_{cm}^{ee} = 3.77$ GeV
13.0	± 1.9	± 2.9		BERGER	85 PLUT	$E_{cm}^{ee} = 34.6$ GeV
18.3	± 2.4	± 1.9	60	BEHREND	83C CELL	$E_{cm}^{ee} = 34$ GeV
16.0	± 1.3		459	46 BACINO	78B DLCO	$E_{cm}^{ee} = 3.1-7.4$ GeV

⁴¹ The correlation coefficient between this measurement and the ACCIARRI 01F measurement of $B(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)$ is 0.08.

⁴² The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu \bar{\nu}_\mu \nu_\tau)$, $B(\mu \bar{\nu}_\mu \nu_\tau)/B(e \bar{\nu}_e \nu_\tau)$, $B(h^- \nu_\tau)$, and $B(h^- \nu_\tau)/B(e \bar{\nu}_e \nu_\tau)$ are 0.50, -0.42, 0.48, and -0.39 respectively.

⁴³ Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ and ALBRECHT 93G $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{total}^2$ values.

⁴⁴ Modified using $B(e^- \bar{\nu}_e \nu_\tau)/B(\text{"1 prong"})$ and $B(\text{"1 prong"})$, = 0.855.

⁴⁵ Error correlated with BALTRUSAITIS 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{total}$.

⁴⁶ BACINO 78B value comes from fit to events with e^\pm and one other nonelectron charged prong.

$\Gamma(e^- \bar{\nu}_e \nu_\tau \gamma)/\Gamma_{total}$ Γ_6/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.75\pm0.06\pm0.17	47 BERGFELD	00	CLEO $E_{cm}^{ee} = 10.6$ GeV

⁴⁷ BERGFELD 00 impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_\gamma^* > 10$ MeV.

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ Γ_3/Γ_5

Standard Model prediction including mass effects is 0.9726.

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.973 \pm 0.004 OUR FIT			
0.978 \pm 0.011 OUR AVERAGE			

0.9777 \pm 0.0063 \pm 0.0087 f&a ⁴⁸ ANASTASSOV 97 CLEO $E_{cm}^{ee} = 10.6$ GeV

0.997 \pm 0.035 \pm 0.040 f&a ALBRECHT 92D ARG $E_{cm}^{ee} = 9.4-10.6$ GeV

⁴⁸ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu \bar{\nu}_\mu \nu_\tau)$, $B(e \bar{\nu}_e \nu_\tau)$, $B(h^- \nu_\tau)$, and $B(h^- \nu_\tau)/B(e \bar{\nu}_e \nu_\tau)$ are 0.58, -0.42, 0.07, and 0.45 respectively.

$\Gamma(h^- \geq 0K_L^0 \nu_\tau)/\Gamma_{total}$ Γ_7/Γ

$$\Gamma_7/\Gamma = (\Gamma_9 + \Gamma_{10} + \frac{1}{2}\Gamma_{33} + \frac{1}{2}\Gamma_{35} + \Gamma_{45})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.30\pm0.10 OUR FIT				Error includes scale factor of 1.4.
12.44\pm0.14 OUR AVERAGE				
12.44 \pm 0.11 \pm 0.11	f&a 15k	49 BUSKULIC	96	ALEP 1991-1993 LEP run
12.47 \pm 0.26 \pm 0.43	f&a 2967	50 ACCIARRI	95	L3 1992 LEP run
12.4 \pm 0.7 \pm 0.7	f&a 283	51 ABREU	92N	DLPH 1990 LEP run

12.98 ± 0.44 ± 0.33	f&a		52 DECAMP	92C ALEP	1989–1990 LEP runs
12.1 ± 0.7 ± 0.5	f&a	309	ALEXANDER	91D OPAL	1990 LEP run
11.3 ± 0.5 ± 0.8	avg	798	53 FORD	87 MAC	$E_{cm}^{ee} = 29$ GeV
• • •	We do not use the following data for averages, fits, limits, etc. • • •				
11.7 ± 0.6 ± 0.8			54 ALBRECHT	92D ARG	$E_{cm}^{ee} = 9.4–10.6$ GeV
12.3 ± 0.9 ± 0.5		1338	BEHREND	90 CELL	$E_{cm}^{ee} = 35$ GeV
11.1 ± 1.1 ± 1.4			55 BURCHAT	87 MRK2	$E_{cm}^{ee} = 29$ GeV
12.3 ± 0.6 ± 1.1		328	56 BARTEL	86D JADE	$E_{cm}^{ee} = 34.6$ GeV
13.0 ± 2.0 ± 4.0			BERGER	85 PLUT	$E_{cm}^{ee} = 34.6$ GeV
11.2 ± 1.7 ± 1.2		34	57 BEHREND	83C CELL	$E_{cm}^{ee} = 34$ GeV

⁴⁹ BUSKULIC 96 quote $11.78 \pm 0.11 \pm 0.13$ We add 0.66 to undo their correction for unseen K_L^0 and modify the systematic error accordingly.

⁵⁰ ACCIARRI 95 with 0.65% added to remove their correction for $\pi^- K_L^0$ backgrounds.

⁵¹ ABREU 92N with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.

⁵² DECAMP 92C quote $B(h^- \geq 0 K_L^0 \geq 0 (K_S^0 \rightarrow \pi^+ \pi^-) \nu_\tau) = 13.32 \pm 0.44 \pm 0.33$.

We subtract 0.35 to correct for their inclusion of the K_S^0 decays.

⁵³ FORD 87 result for $B(\pi^- \nu_\tau)$ with 0.67% added to remove their K^- correction and adjusted for 1992 B (“1 prong”).

⁵⁴ Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma(e^- \bar{\nu}_e \nu_\tau)$, $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)$, and $\Gamma(h^- \geq 0 K_L^0 \nu_\tau) / \Gamma(e^- \bar{\nu}_e \nu_\tau)$ values.

⁵⁵ BURCHAT 87 with 1.1% added to remove their correction for K^- and $K^*(892)^-$ backgrounds.

⁵⁶ BARTEL 86D result for $B(\pi^- \nu_\tau)$ with 0.59% added to remove their K^- correction and adjusted for 1992 B (“1 prong”).

⁵⁷ BEHREND 83C quote $B(\pi^- \nu_\tau) = 9.9 \pm 1.7 \pm 1.3$ after subtracting 1.3 ± 0.5 to correct for $B(K^- \nu_\tau)$.

$\Gamma(h^- \nu_\tau) / \Gamma_{total}$

$$\Gamma_8 / \Gamma = (\Gamma_9 + \Gamma_{10}) / \Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

<u>VALUE (%)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.75 ± 0.10 OUR FIT		Error includes scale factor of 1.4.		
11.65 ± 0.21 OUR AVERAGE		Error includes scale factor of 1.9.		
11.98 ± 0.13 ± 0.16	f&a	ACKERSTAFF	98M OPAL	1991–1995 LEP runs
11.52 ± 0.05 ± 0.12	f&a	58 ANASTASSOV	97 CLEO	$E_{cm}^{ee} = 10.6$ GeV

⁵⁸ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu^- \bar{\nu}_\mu \nu_\tau)$, $B(e^- \bar{\nu}_e \nu_\tau)$, $B(\mu^- \bar{\nu}_\mu \nu_\tau) / B(e^- \bar{\nu}_e \nu_\tau)$, and $B(h^- \nu_\tau) / B(e^- \bar{\nu}_e \nu_\tau)$ are 0.50, 0.48, 0.07, and 0.63 respectively.

$\Gamma(h^- \nu_\tau) / \Gamma(e^- \bar{\nu}_e \nu_\tau)$

$$\Gamma_8 / \Gamma_5 = (\Gamma_9 + \Gamma_{10}) / \Gamma_5$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.659 ± 0.006 OUR FIT		Error includes scale factor of 1.3.		
0.6484 ± 0.0041 ± 0.0060 avg		59 ANASTASSOV	97 CLEO	$E_{cm}^{ee} = 10.6$ GeV

⁵⁹ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu\bar{\nu}_\mu\nu_\tau)$, $B(e\bar{\nu}_e\nu_\tau)$, $B(\mu\bar{\nu}_\mu\nu_\tau)/B(e\bar{\nu}_e\nu_\tau)$, and $B(h^-\nu_\tau)$ are 0.08, -0.39, 0.45, and 0.63 respectively.

$\Gamma(\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_9/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

11.06±0.11 OUR FIT Error includes scale factor of 1.4.

11.07±0.18 OUR AVERAGE

11.06±0.11±0.14	avg	⁶⁰ BUSKULIC	96	ALEP	LEP 1991–1993 data
11.7 ±0.4 ±1.8	f&a	1138	BLOCKER	82D MRK2	$E_{\text{cm}}^{ee} = 3.5\text{--}6.7$ GeV

⁶⁰ Not independent of BUSKULIC 96 $B(h^-\nu_\tau)$ and $B(K^-\nu_\tau)$ values.

$\Gamma(K^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.686±0.023 OUR FIT

0.685±0.023 OUR AVERAGE

0.658±0.027±0.029		⁶¹ ABBIENDI	01J	OPAL	1990–1995 LEP runs
0.696±0.025±0.014	2032	BARATE	99K	ALEP	1991–1995 LEP runs
0.85 ±0.18	27	ABREU	94K	DLPH	LEP 1992 Z data
0.66 ±0.07 ±0.09	99	BATTLE	94	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.72 ±0.04 ±0.04	728	BUSKULIC	96	ALEP	Repl. by BARATE 99K
0.59 ±0.18	16	MILLS	84	DLCO	$E_{\text{cm}}^{ee} = 29$ GeV
1.3 ±0.5	15	BLOCKER	82B	MRK2	$E_{\text{cm}}^{ee} = 3.9\text{--}6.7$ GeV

⁶¹ The correlation coefficient between this measurement and the ABBIENDI 01J $B(\tau^- \rightarrow K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma\nu_\tau)$ is 0.60.

$\Gamma(h^- \geq 1 \text{ neutrals}\nu_\tau)/\Gamma_{\text{total}}$ Γ_{11}/Γ

$\Gamma_{11}/\Gamma = (\Gamma_{13} + \Gamma_{15} + \Gamma_{19} + \Gamma_{22} + \Gamma_{25} + \Gamma_{26} + \Gamma_{28} + 0.157\Gamma_{33} + 0.157\Gamma_{35} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.0985\Gamma_{45} + 0.708\Gamma_{117} + 0.715\Gamma_{119} + 0.09\Gamma_{137} + 0.09\Gamma_{138})/\Gamma$

VALUE (%) DOCUMENT ID TECN COMMENT

36.92±0.14 OUR FIT Error includes scale factor of 1.1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.14±0.33±0.58	⁶² AKERS	94E	OPAL	1991–1992 LEP runs
38.4 ±1.2 ±1.0	⁶³ BURCHAT	87	MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
42.7 ±2.0 ±2.9	BERGER	85	PLUT	$E_{\text{cm}}^{ee} = 34.6$ GeV

⁶² Not independent of ACKERSTAFF 98M $B(h^-\pi^0\nu_\tau)$ and $B(h^-\geq 2\pi^0\nu_\tau)$ values.

⁶³ BURCHAT 87 quote for $B(\pi^\pm \geq 1 \text{ neutral}\nu_\tau) = 0.378 \pm 0.012 \pm 0.010$. We add 0.006 to account for contribution from $(K^{*-}\nu_\tau)$ which they fixed at BR = 0.013.

$\Gamma(h^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma = (\Gamma_{13} + \Gamma_{15})/\Gamma$

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

25.86 ± 0.13 OUR FIT Error includes scale factor of 1.1.

25.76 ± 0.15 OUR AVERAGE

25.89 ± 0.17 ± 0.29		ACKERSTAFF	98M	OPAL	1991–1995 LEP runs
25.76 ± 0.15 ± 0.13	31k	BUSKULIC	96	ALEP	LEP 1991–1993 data
25.05 ± 0.35 ± 0.50	6613	ACCIARRI	95	L3	1992 LEP run
25.87 ± 0.12 ± 0.42	51k	⁶⁴ ARTUSO	94	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

25.98 ± 0.36 ± 0.52		⁶⁵ AKERS	94E	OPAL	Repl. by ACKER-STAFF 98M
22.9 ± 0.8 ± 1.3	283	⁶⁶ ABREU	92N	DLPH	$E_{\text{cm}}^{ee} = 88.2$ – 94.2 GeV
23.1 ± 0.4 ± 0.9	1249	⁶⁷ ALBRECHT	92Q	ARG	$E_{\text{cm}}^{ee} = 10$ GeV
25.02 ± 0.64 ± 0.88	1849	DECAMP	92C	ALEP	1989–1990 LEP runs
22.0 ± 0.8 ± 1.9	779	ANTREASYAN	91	CBAL	$E_{\text{cm}}^{ee} = 9.4$ – 10.6 GeV
22.6 ± 1.5 ± 0.7	1101	BEHREND	90	CELL	$E_{\text{cm}}^{ee} = 35$ GeV
23.1 ± 1.9 ± 1.6		BEHREND	84	CELL	$E_{\text{cm}}^{ee} = 14,22$ GeV

⁶⁴ARTUSO 94 reports the combined result from three independent methods, one of which (23% of the $\tau^- \rightarrow h^- \pi^0 \nu_\tau$) is normalized to the inclusive one-prong branching fraction, taken as 0.854 ± 0.004 . Renormalization to the present value causes negligible change.

⁶⁵AKERS 94E quote $(26.25 \pm 0.36 \pm 0.52) \times 10^{-2}$; we subtract 0.27% from their number to correct for $\tau^- \rightarrow h^- K_L^0 \nu_\tau$.

⁶⁶ABREU 92N with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.

⁶⁷ALBRECHT 92Q with 0.5% added to remove their correction for $\tau^- \rightarrow K^*(892)^- \nu_\tau$ background.

$\Gamma(\pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{13}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

25.41 ± 0.14 OUR FIT Error includes scale factor of 1.1.

25.31 ± 0.18 OUR AVERAGE

25.30 ± 0.15 ± 0.13	avg	⁶⁸ BUSKULIC	96	ALEP	LEP 1991–1993 data
25.36 ± 0.44	avg	⁶⁹ ARTUSO	94	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

21.5 ± 0.4 ± 1.9	4400	^{70,71} ALBRECHT	88L	ARG	$E_{\text{cm}}^{ee} = 10$ GeV
23.0 ± 1.3 ± 1.7	582	ADLER	87B	MRK3	$E_{\text{cm}}^{ee} = 3.77$ GeV
25.8 ± 1.7 ± 2.5		⁷² BURCHAT	87	MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
22.3 ± 0.6 ± 1.4	629	⁷¹ YELTON	86	MRK2	$E_{\text{cm}}^{ee} = 29$ GeV

⁶⁸Not independent of BUSKULIC 96 B($h^- \pi^0 \nu_\tau$) and B($K^- \pi^0 \nu_\tau$) values.

⁶⁹Not independent of ARTUSO 94 B($h^- \pi^0 \nu_\tau$) and BATTLE 94 B($K^- \pi^0 \nu_\tau$) values.

⁷⁰The authors divide by $(\Gamma_3 + \Gamma_5 + \Gamma_9 + \Gamma_{10})/\Gamma = 0.467$ to obtain this result.

⁷¹Experiment had no hadron identification. Kaon corrections were made, but insufficient information is given to permit their removal.

⁷²BURCHAT 87 value is not independent of YELTON 86 value. Nonresonant decays included.

$\Gamma(\pi^- \pi^0 \text{non-}\rho(770) \nu_\tau) / \Gamma_{\text{total}}$ Γ_{14} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.3 ± 0.1 ± 0.3		⁷³ BEHREND	84	CELL $E_{\text{cm}}^{ee} = 14,22 \text{ GeV}$

⁷³ BEHREND 84 assume a flat nonresonant mass distribution down to the $\rho(770)$ mass, using events with mass above 1300 to set the level.

$\Gamma(K^- \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{15} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.450 ± 0.030 OUR FIT

0.449 ± 0.034 OUR AVERAGE

0.444 ± 0.026 ± 0.024	923	BARATE	99K	ALEP 1991–1995 LEP runs
0.51 ± 0.10 ± 0.07	37	BATTLE	94	CLEO $E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.52 ± 0.04 ± 0.05	395	BUSKULIC	96	ALEP Repl. by BARATE 99K
--------------------	-----	----------	----	--------------------------

$\Gamma(h^- \geq 2\pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{16} / Γ

$$\Gamma_{16} / \Gamma = (\Gamma_{19} + \Gamma_{22} + \Gamma_{25} + \Gamma_{26} + \Gamma_{28} + 0.157\Gamma_{33} + 0.157\Gamma_{35} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.0985\Gamma_{45} + 0.319\Gamma_{117} + 0.322\Gamma_{119}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

10.76 ± 0.15 OUR FIT Error includes scale factor of 1.1.

10.0 ± 0.4 OUR AVERAGE

9.91 ± 0.31 ± 0.27 f&a		ACKERSTAFF	98M	OPAL 1991–1995 LEP runs
12.0 ± 1.4 ± 2.5 f&a		⁷⁴ BURCHAT	87	MRK2 $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.89 ± 0.34 ± 0.55		⁷⁵ AKERS	94E	OPAL Repl. by ACKER-STAFF 98M
14.0 ± 1.2 ± 0.6	938	⁷⁶ BEHREND	90	CELL $E_{\text{cm}}^{ee} = 35 \text{ GeV}$
13.9 ± 2.0 $\begin{smallmatrix} +1.9 \\ -2.2 \end{smallmatrix}$		⁷⁷ AIHARA	86E	TPC $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

⁷⁴ Error correlated with BURCHAT 87 $\Gamma(\rho^- \nu_e) / \Gamma(\text{total})$ value.

⁷⁵ AKERS 94E not independent of AKERS 94E $B(h^- \geq 1\pi^0 \nu_\tau)$ and $B(h^- \pi^0 \nu_\tau)$ measurements.

⁷⁶ No independent of BEHREND 90 $\Gamma(h^- 2\pi^0 \nu_\tau (\text{exp. } K^0))$ and $\Gamma(h^- \geq 3\pi^0 \nu_\tau)$.

⁷⁷ AIHARA 86E (TPC) quote $B(2\pi^0 \pi^- \nu_\tau) + 1.6B(3\pi^0 \pi^- \nu_\tau) + 1.1B(\pi^0 \eta \pi^- \nu_\tau)$.

$\Gamma(h^- 2\pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{17} / Γ

$$\Gamma_{17} / \Gamma = (\Gamma_{19} + \Gamma_{22} + 0.157\Gamma_{33} + 0.157\Gamma_{35}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

9.39 ± 0.14 OUR FIT Error includes scale factor of 1.1.

9.48 ± 0.13 ± 0.10 ⁷⁸ BUSKULIC 96 ALEP LEP 1991–1993 data

⁷⁸ BUSKULIC 96 quote $9.29 \pm 0.13 \pm 0.10$. We add 0.19 to undo their correction for $\tau^- \rightarrow h^- K^0 \nu_\tau$.

$\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ Γ_{18}/Γ
 $\Gamma_{18}/\Gamma = (\Gamma_{19} + \Gamma_{22})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. f&a marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
9.23±0.14 OUR FIT					Error includes scale factor of 1.1.
9.08±0.34 OUR AVERAGE					
8.88±0.37±0.42	f&a	1060	ACCIARRI	95 L3	1992 LEP run
8.96±0.16±0.44	avg		⁷⁹ PROCARIO	93 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
10.38±0.66±0.82	f&a	809	⁸⁰ DECAMP	92C ALEP	1989–1990 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
5.7 ±0.5 ^{+1.7} / _{-1.0}		133	⁸¹ ANTREASYAN	91 CBAL	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
10.0 ±1.5 ±1.1		333	⁸² BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35$ GeV
8.7 ±0.4 ±1.1		815	⁸³ BAND	87 MAC	$E_{\text{cm}}^{ee} = 29$ GeV
6.2 ±0.6 ±1.2			⁸⁴ GAN	87 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
6.0 ±3.0 ±1.8			BEHREND	84 CELL	$E_{\text{cm}}^{ee} = 14,22$ GeV

⁷⁹ PROCARIO 93 entry is obtained from $B(h^- 2\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.

⁸⁰ We subtract 0.0015 to account for $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

⁸¹ ANTREASYAN 91 subtract 0.001 to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

⁸² BEHREND 90 subtract 0.002 to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

⁸³ BAND 87 assume $B(\pi^- 3\pi^0 \nu_\tau) = 0.01$ and $B(\pi^- \pi^0 \eta \nu_\tau) = 0.005$.

⁸⁴ GAN 87 analysis use photon multiplicity distribution.

$\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma(h^- \pi^0 \nu_\tau)$ Γ_{18}/Γ_{12}
 $\Gamma_{18}/\Gamma_{12} = (\Gamma_{19} + \Gamma_{22})/(\Gamma_{13} + \Gamma_{15})$

VALUE		DOCUMENT ID	TECN	COMMENT
0.357±0.006 OUR FIT				Error includes scale factor of 1.1.
0.342±0.006±0.016		⁸⁵ PROCARIO	93 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

⁸⁵ PROCARIO 93 quote $0.345 \pm 0.006 \pm 0.016$ after correction for 2 kaon backgrounds assuming $B(K^{*-} \nu_\tau) = 1.42 \pm 0.18\%$ and $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$. We multiply by 0.990 ± 0.010 to remove these corrections to $B(h^- \pi^0 \nu_\tau)$.

$\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ Γ_{19}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		DOCUMENT ID	TECN	COMMENT
9.17±0.14 OUR FIT				Error includes scale factor of 1.1.
9.21±0.13±0.11	avg	⁸⁶ BUSKULIC	96 ALEP	LEP 1991–1993 data

⁸⁶ Not independent of BUSKULIC 96 $B(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ and $B(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ values.

$\Gamma(\pi^- 2\pi^0 \nu_\tau(\text{ex. } K^0), \text{ scalar})/\Gamma(\pi^- 2\pi^0 \nu_\tau(\text{ex. } K^0))$ Γ_{20}/Γ_{19}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.094	95	⁸⁷ BROWDER	00 CLEO	$4.7 \text{ fb}^{-1} E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

⁸⁷ Model-independent limit from structure function analysis on contribution to $B(\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau(\text{ex. } K^0))$ from scalars.

$\Gamma(\pi^- 2\pi^0 \nu_\tau(\text{ex. } K^0), \text{ vector})/\Gamma(\pi^- 2\pi^0 \nu_\tau(\text{ex. } K^0))$ Γ_{21}/Γ_{19}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.073	95	⁸⁸ BROWDER	00 CLEO	$4.7 \text{ fb}^{-1} E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

⁸⁸ Model-independent limit from structure function analysis on contribution to $B(\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau(\text{ex. } K^0))$ from vectors.

$\Gamma(K^- 2\pi^0 \nu_\tau(\text{ex. } K^0))/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.058 ± 0.023 OUR FIT				
0.058 ± 0.024 OUR AVERAGE				
0.056 ± 0.020 ± 0.015	131	BARATE	99K ALEP	1991–1995 LEP runs
0.09 ± 0.10 ± 0.03	3	⁸⁹ BATTLE	94 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.08 ± 0.02 ± 0.02	59	BUSKULIC	96 ALEP	Repl. by BARATE 99K
⁸⁹ BATTLE 94 quote $0.14 \pm 0.10 \pm 0.03$ or $< 0.3\%$ at 90% CL. We subtract $(0.05 \pm 0.02)\%$ to account for $\tau^- \rightarrow K^- (K^0 \rightarrow \pi^0 \pi^0) \nu_\tau$ background.				

$\Gamma(h^- \geq 3\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{23}/Γ

$$\Gamma_{23}/\Gamma = (\Gamma_{25} + \Gamma_{26} + \Gamma_{28} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.0985\Gamma_{45} + 0.319\Gamma_{117} + 0.322\Gamma_{119})/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.37 ± 0.11 OUR FIT	Error includes scale factor of 1.1.			
1.53 ± 0.40 ± 0.46	186	DECAMP	92C ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.2 ± 1.0 ± 1.0		BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35 \text{ GeV}$

$\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{24}/Γ

$$\Gamma_{24}/\Gamma = (\Gamma_{25} + \Gamma_{26} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.322\Gamma_{119})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.21 ± 0.10 OUR FIT				
1.22 ± 0.10 OUR AVERAGE				
1.24 ± 0.09 ± 0.11	f&a 2.3k	⁹⁰ BUSKULIC	96 ALEP	LEP 1991–1993 data
1.70 ± 0.24 ± 0.38	f&a 293	ACCIARRI	95 L3	1992 LEP run
1.15 ± 0.08 ± 0.13	avg	⁹¹ PROCARIO	93 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0 ^{+1.4} _{-0.1} ^{+1.1} _{-0.1}		⁹² GAN	87 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

⁹⁰ BUSKULIC 96 quote $B(h^- 3\pi^0 \nu_\tau (\text{ex. } K^0)) = 1.17 \pm 0.09 \pm 0.11$. We add 0.07 to remove their correction for K^0 backgrounds.

⁹¹ PROCARIO 93 entry is obtained from $B(h^- 3\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.

⁹² Highly correlated with GAN 87 $\Gamma(\eta \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ value. Authors quote $B(\pi^\pm 3\pi^0 \nu_\tau) + 0.67B(\pi^\pm \eta \pi^0 \nu_\tau) = 0.047 \pm 0.010 \pm 0.011$.

$$\frac{\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma(h^- \pi^0 \nu_\tau)}{\Gamma_{24}/\Gamma_{12}} = \frac{\Gamma_{24}/\Gamma_{12}}{(\Gamma_{25} + \Gamma_{26} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.322\Gamma_{119})/(\Gamma_{13} + \Gamma_{15})} \quad \Gamma_{24}/\Gamma_{12}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.047±0.004 OUR FIT			Error includes scale factor of 1.1.
0.044±0.003±0.005	⁹³ PROCARIO 93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

⁹³ PROCARIO 93 quote $0.041 \pm 0.003 \pm 0.005$ after correction for 2 kaon backgrounds assuming $B(K^{*-} \nu_\tau) = 1.42 \pm 0.18\%$ and $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$. We add 0.003 ± 0.003 and multiply the sum by 0.990 ± 0.010 to remove these corrections.

$$\frac{\Gamma(\pi^- 3\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}}{\Gamma_{25}/\Gamma} \quad \Gamma_{25}/\Gamma$$

VALUE (%)	DOCUMENT ID
1.08±0.10 OUR FIT	

$$\frac{\Gamma(K^- 3\pi^0 \nu_\tau (\text{ex. } K^0, \eta))/\Gamma_{\text{total}}}{\Gamma_{26}/\Gamma} \quad \Gamma_{26}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.037^{+0.022}_{-0.020} OUR FIT				
0.037±0.021±0.011	22	BARATE	99K ALEP	1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.05 ± 0.13 ⁹⁴ BUSKULIC 94E ALEP Repl. by BARATE 99K

⁹⁴ BUSKULIC 94E quote $B(K^- \geq 0\pi^0 \geq 0K^0 \nu_\tau) - [B(K^- \nu_\tau) + B(K^- \pi^0 \nu_\tau) + B(K^- K^0 \nu_\tau) + B(K^- \pi^0 \pi^0 \nu_\tau) + B(K^- \pi^0 K^0 \nu_\tau)] = 0.05 \pm 0.13\%$ accounting for common systematic errors in BUSKULIC 94E and BUSKULIC 94F measurements of these modes. We assume $B(K^- \geq 2K^0 \nu_\tau)$ and $B(K^- \geq 4\pi^0 \nu_\tau)$ are negligible.

$$\frac{\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}}{\Gamma_{27}/\Gamma} = \frac{\Gamma_{27}/\Gamma}{(\Gamma_{28} + 0.319\Gamma_{117})/\Gamma} \quad \Gamma_{27}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.16±0.06 OUR FIT				
0.16±0.06 OUR AVERAGE				

$0.16 \pm 0.04 \pm 0.09$ 232 ⁹⁵ BUSKULIC 96 ALEP LEP 1991–1993 data

$0.16 \pm 0.05 \pm 0.05$ ⁹⁶ PROCARIO 93 CLEO $E_{\text{cm}}^{ee} \approx 10.6$ GeV

⁹⁵ BUSKULIC 96 quote result for $\tau^- \rightarrow h^- \geq 4\pi^0 \nu_\tau$. We assume $B(h^- \geq 5\pi^0 \nu_\tau)$ is negligible.

⁹⁶ PROCARIO 93 quotes $B(h^- 4\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau) = 0.006 \pm 0.002 \pm 0.002$. We multiply by the ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$ to obtain $B(h^- 4\pi^0 \nu_\tau)$. PROCARIO 93 assume $B(h^- \geq 5\pi^0 \nu_\tau)$ is small and do not correct for it.

$\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex. } K^0, \eta))/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (%) DOCUMENT ID

0.10^{+0.06}_{-0.05} OUR FIT

$\Gamma(K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma \nu_\tau)/\Gamma_{\text{total}}$ Γ_{29}/Γ

$$\Gamma_{29}/\Gamma = (\Gamma_{10} + \Gamma_{15} + \Gamma_{22} + \Gamma_{26} + \Gamma_{35} + \Gamma_{40} + 0.715\Gamma_{119})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

1.56 ± 0.04 OUR FIT

1.53 ± 0.04 OUR AVERAGE

1.528 ± 0.039 ± 0.040	f&a		⁹⁷ ABBIENDI	01J OPAL	1990–1995 LEP runs
1.520 ± 0.040 ± 0.041	avg	4006	⁹⁸ BARATE	99K ALEP	1991–1995 LEP runs
1.54 ± 0.24	f&a		ABREU	94K DLPH	LEP 1992 Z data
1.70 ± 0.12 ± 0.19	f&a	202	⁹⁹ BATTLE	94 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
1.6 ± 0.4 ± 0.2	f&a	35	AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29$ GeV
1.71 ± 0.29	f&a	53	MILLS	84 DLCO	$E_{\text{cm}}^{ee} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.70 ± 0.05 ± 0.06		1610	¹⁰⁰ BUSKULIC	96 ALEP	Repl. by BARATE 99K
--------------------	--	------	-------------------------	---------	------------------------

⁹⁷ The correlation coefficient between this measurement and the ABBIENDI 01J $B(\tau^- \rightarrow K^- \nu_\tau)$ is 0.60.

⁹⁸ Not independent of BARATE 99K $B(K^- \nu_\tau)$, $B(K^- \pi^0 \nu_\tau)$, $B(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$, $B(K^- 3\pi^0 \nu_\tau (\text{ex. } K^0))$, $B(K^- K^0 \nu_\tau)$, and $B(K^- K^0 \pi^0 \nu_\tau)$ values.

⁹⁹ BATTLE 94 quote $1.60 \pm 0.12 \pm 0.19$. We add 0.10 ± 0.02 to correct for their rejection of $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

¹⁰⁰ Not independent of BUSKULIC 96 $B(K^- \nu_\tau)$, $B(K^- \pi^0 \nu_\tau)$, $B(K^- 2\pi^0 \nu_\tau)$, $B(K^- K^0 \nu_\tau)$, and $B(K^- K^0 \pi^0 \nu_\tau)$ values.

$\Gamma(K^- \geq 1(\pi^0 \text{ or } K^0 \text{ or } \gamma) \nu_\tau)/\Gamma_{\text{total}}$ Γ_{30}/Γ

$$\Gamma_{30}/\Gamma = (\Gamma_{15} + \Gamma_{22} + \Gamma_{26} + \Gamma_{35} + \Gamma_{40} + 0.715\Gamma_{119})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.874 ± 0.035 OUR FIT

0.86 ± 0.05 OUR AVERAGE

0.869 ± 0.031 ± 0.034	avg		¹⁰¹ ABBIENDI	01J OPAL	1990–1995 LEP runs
0.69 ± 0.25	avg		¹⁰² ABREU	94K DLPH	LEP 1992 Z data
1.2 ± 0.5 ^{+0.2} _{-0.4}	f&a	9	AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29$ GeV

¹⁰¹ Not independent of ABBIENDI 01J $B(\tau^- \rightarrow K^- \nu_\tau)$ and $B(\tau^- \rightarrow K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma \nu_\tau)$ values.

¹⁰² Not independent of ABREU 94K $B(K^- \nu_\tau)$ and $B(K^- \geq 0 \text{ neutrals } \nu_\tau)$ measurements.

$$\Gamma(K_S^0(\text{particles})^- \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{31} / \Gamma$$

$$\Gamma_{31} / \Gamma = (\frac{1}{2}\Gamma_{33} + \frac{1}{2}\Gamma_{35} + \frac{1}{2}\Gamma_{38} + \frac{1}{2}\Gamma_{40} + \Gamma_{45} + \Gamma_{46}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.92 ± 0.04 OUR FIT				Error includes scale factor of 1.1.
0.97 ± 0.07 OUR AVERAGE				
0.970 ± 0.058 ± 0.062	929	BARATE	98E ALEP	1991–1995 LEP runs
0.97 ± 0.09 ± 0.06	141	AKERS	94G OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$$\Gamma(h^- \bar{K}^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{32} / \Gamma = (\Gamma_{33} + \Gamma_{35}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.05 ± 0.04 OUR FIT				Error includes scale factor of 1.1.
0.90 ± 0.07 OUR AVERAGE				
1.01 ± 0.11 ± 0.07 avg	555	¹⁰³ BARATE	98E ALEP	1991–1995 LEP runs
0.855 ± 0.036 ± 0.073 f&a	1242	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

¹⁰³ Not independent of BARATE 98E $B(\tau^- \rightarrow \pi^- \bar{K}^0 \nu_\tau)$ and $B(\tau^- \rightarrow K^- K^0 \nu_\tau)$ values.

$$\Gamma(\pi^- \bar{K}^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{33} / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.89 ± 0.04 OUR FIT				Error includes scale factor of 1.1.
0.88 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.2.
0.933 ± 0.068 ± 0.049 f&a	377	ABBIENDI	00C OPAL	1991–1995 LEP runs
0.928 ± 0.045 ± 0.034 f&a	937	¹⁰⁴ BARATE	99K ALEP	1991–1995 LEP runs
0.855 ± 0.117 ± 0.066 avg	509	¹⁰⁵ BARATE	98E ALEP	1991–1995 LEP runs
0.704 ± 0.041 ± 0.072 avg		¹⁰⁶ COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
0.95 ± 0.15 ± 0.06 f&a		¹⁰⁷ ACCIARRI	95F L3	1991–1993 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.79 ± 0.10 ± 0.09	98	¹⁰⁸ BUSKULIC	96 ALEP	Repl. by BARATE 99K
--------------------	----	-------------------------	---------	---------------------

¹⁰⁴ BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

¹⁰⁵ BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays. Not independent of BARATE 98E $B(K^0 \text{ particles}^- \nu_\tau)$ value.

¹⁰⁶ Not independent of COAN 96 $B(h^- K^0 \nu_\tau)$ and $B(K^- K^0 \nu_\tau)$ measurements.

¹⁰⁷ ACCIARRI 95F do not identify π^- / K^- and assume $B(K^- K^0 \nu_\tau) = (0.29 \pm 0.12)\%$.

¹⁰⁸ BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$$\Gamma(\pi^- \bar{K}^0(\text{non-}K^*(892)^-) \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{34} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.17	95	ACCIARRI	95F L3	1991–1993 LEP runs

$\Gamma(K^- K^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.154±0.016 OUR FIT

0.158±0.017 OUR AVERAGE

0.162±0.021±0.011	150	¹⁰⁹ BARATE	99K ALEP	1991–1995 LEP runs
0.158±0.042±0.017	46	¹¹⁰ BARATE	98E ALEP	1991–1995 LEP runs
0.151±0.021±0.022	111	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ±0.09 ±0.02	13	¹¹¹ BUSKULIC	96 ALEP	Repl. by BARATE 99K
------------------	----	-------------------------	---------	---------------------

¹⁰⁹ BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

¹¹⁰ BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

¹¹¹ BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$\Gamma(K^- K^0 \geq 0\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{36}/\Gamma = (\Gamma_{35} + \Gamma_{40})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.309±0.024 OUR FIT

0.330±0.055±0.039 124 ABBIENDI 00C OPAL 1991–1995 LEP runs

$\Gamma(h^- \bar{K}^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{37}/\Gamma = (\Gamma_{38} + \Gamma_{40})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.52 ±0.04 OUR FIT

0.50 ±0.06 OUR AVERAGE Error includes scale factor of 1.2.

0.446±0.052±0.046	avg	¹¹² BARATE	98E ALEP	1991–1995 LEP runs
-------------------	-----	-----------------------	----------	--------------------

0.562±0.050±0.048	f&a	264	COAN	96 CLEO $E_{\text{cm}}^{ee} \approx 10.6$ GeV
-------------------	-----	-----	------	---

¹¹² Not independent of BARATE 98E $B(\tau^- \rightarrow \pi^- \bar{K}^0 \pi^0 \nu_\tau)$ and $B(\tau^- \rightarrow K^- K^0 \pi^0 \nu_\tau)$ values.

$\Gamma(\pi^- \bar{K}^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{38}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.37 ±0.04 OUR FIT

0.36 ±0.04 OUR AVERAGE

0.347±0.053±0.037	f&a	299	¹¹³ BARATE	99K ALEP 1991–1995 LEP runs
-------------------	-----	-----	-----------------------	-----------------------------

0.294±0.073±0.037	f&a	142	¹¹⁴ BARATE	98E ALEP 1991–1995 LEP runs
-------------------	-----	-----	-----------------------	-----------------------------

0.417±0.058±0.044	avg	¹¹⁵ COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
-------------------	-----	---------------------	---------	---------------------------------------

0.41 ±0.12 ±0.03	f&a	¹¹⁶ ACCIARRI	95F L3	1991–1993 LEP runs
------------------	-----	-------------------------	--------	--------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.32 ±0.11 ±0.05	23	¹¹⁷ BUSKULIC	96 ALEP	Repl. by BARATE 99K
------------------	----	-------------------------	---------	---------------------

¹¹³ BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

¹¹⁴ BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

¹¹⁵ Not independent of COAN 96 $B(h^- K^0 \pi^0 \nu_\tau)$ and $B(K^- K^0 \pi^0 \nu_\tau)$ measurements.

¹¹⁶ ACCIARRI 95F do not identify π^-/K^- and assume $B(K^- K^0 \pi^0 \nu_\tau) = (0.05 \pm 0.05)\%$.

¹¹⁷ BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$\Gamma(\bar{K}^0 \rho^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{39}/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.22 ± 0.05 OUR AVERAGE			
0.250 ± 0.057 ± 0.044	118 BARATE	99K ALEP	1991–1995 LEP runs
0.188 ± 0.054 ± 0.038	119 BARATE	98E ALEP	1991–1995 LEP runs

118 BARATE 99K measure K^0 's by detecting K_L^0 's in hadron calorimeter. They determine the $\bar{K}^0 \rho^-$ fraction in $\tau^- \rightarrow \pi^- \bar{K}^0 \pi^0 \nu_\tau$ decays to be $(0.72 \pm 0.12 \pm 0.10)$ and multiply their $B(\pi^- \bar{K}^0 \pi^0 \nu_\tau)$ measurement by this fraction to obtain the quoted result.

119 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays. They determine the $\bar{K}^0 \rho^-$ fraction in $\tau^- \rightarrow \pi^- \bar{K}^0 \pi^0 \nu_\tau$ decays to be $(0.64 \pm 0.09 \pm 0.10)$ and multiply their $B(\pi^- \bar{K}^0 \pi^0 \nu_\tau)$ measurement by this fraction to obtain the quoted result.

$\Gamma(K^- K^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{40}/Γ

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.155 ± 0.020 OUR FIT				
0.144 ± 0.023 OUR AVERAGE				
0.143 ± 0.025 ± 0.015	78	120 BARATE	99K ALEP	1991–1995 LEP runs
0.152 ± 0.076 ± 0.021	15	121 BARATE	98E ALEP	1991–1995 LEP runs
0.145 ± 0.036 ± 0.020	32	COAN	96 CLEO	$E_{\text{cm}}^{e\bar{e}} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.10 ± 0.05 ± 0.03 5 122 BUSKULIC 96 ALEP Repl. by BARATE 99K

120 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

121 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

122 BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$\Gamma(\pi^- \bar{K}^0 \geq 1\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{41}/\Gamma = (\Gamma_{38} + \Gamma_{42})/\Gamma$

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.324 ± 0.074 ± 0.066	148	ABBIENDI	00C OPAL	1991–1995 LEP runs

$\Gamma(\pi^- \bar{K}^0 \pi^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{42}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.26 ± 0.24			123 BARATE	99R ALEP	1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.66 95 17 124 BARATE 99K ALEP 1991–1995 LEP runs

0.58 ± 0.33 ± 0.14 5 125 BARATE 98E ALEP 1991–1995 LEP runs

123 BARATE 99R combine the BARATE 98E and BARATE 99K measurements to obtain this value.

124 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

125 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

$\Gamma(K^- K^0 \pi^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.16 \times 10^{-3}$	95	126 BARATE	99R ALEP	1991–1995 LEP runs
$<0.18 \times 10^{-3}$	95	127 BARATE	99K ALEP	1991–1995 LEP runs
$<0.39 \times 10^{-3}$	95	128 BARATE	98E ALEP	1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •
 126 BARATE 99R combine the BARATE 98E and BARATE 99K bounds to obtain this value.
 127 BARATE 99K measure K^0 's by detecting K_L^0 's in hadron calorimeter.
 128 BARATE 98E reconstruct K^0 's by using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

$\Gamma(\pi^- K^0 \bar{K}^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{44}/\Gamma = (2\Gamma_{45} + \Gamma_{46})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.159 ± 0.029 OUR FIT				Error includes scale factor of 1.1.
0.153 ± 0.030 ± 0.016 avg	74	129 BARATE	98E ALEP	1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •
 0.31 ± 0.12 ± 0.04 130 ACCIARRI 95F L3 1991–1993 LEP runs
 129 BARATE 98E obtain this value by adding twice their $B(\pi^- K_S^0 K_S^0 \nu_\tau)$ value to their $B(\pi^- K_S^0 K_L^0 \nu_\tau)$ value.
 130 ACCIARRI 95F assume $B(\pi^- K_S^0 K_S^0 \nu) = B(\pi^- K_S^0 K_L^0 \nu) = 1/2 B(\pi^- K_S^0 K_L^0 \nu)$.

$\Gamma(\pi^- K_S^0 K_S^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{45}/Γ

Bose-Einstein correlations might make the mixing fraction different than 1/4.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.024 ± 0.005 OUR FIT				
0.024 ± 0.005 OUR AVERAGE				
0.026 ± 0.010 ± 0.005	6	BARATE	98E ALEP	1991–1995 LEP runs
0.023 ± 0.005 ± 0.003	42	COAN	96 CLEO	$E_{\text{cm}}^{e\bar{e}} \approx 10.6$ GeV

$\Gamma(\pi^- K_S^0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.110 ± 0.028 OUR FIT				Error includes scale factor of 1.1.
0.101 ± 0.023 ± 0.013	68	BARATE	98E ALEP	1991–1995 LEP runs

$\Gamma(\pi^- K^0 \bar{K}^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
(0.31 ± 0.23) × 10⁻³	131 BARATE	99R ALEP	1991–1995 LEP runs

131 BARATE 99R combine BARATE 98E $\Gamma(\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ measurements to obtain this value.

$\Gamma(\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.020	95	BARATE	98E ALEP	1991–1995 LEP runs

$\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$					Γ_{49}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.031 ± 0.011 ± 0.005	11	BARATE	98E ALEP	1991–1995 LEP runs	

$\Gamma(K^0 h^+ h^- h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$					Γ_{50}/Γ
VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.17	95	TSCHIRHART	88 HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.27	90	BELTRAMI	85 HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
-------	----	----------	--------	---------------------------------------

$\Gamma(K^0 h^+ h^- h^- \nu_\tau)/\Gamma_{\text{total}}$					Γ_{51}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.023 ± 0.019 ± 0.007	6	¹³² BARATE	98E ALEP	1991–1995 LEP runs	

¹³²BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$					Γ_{52}/Γ
$\Gamma_{52}/\Gamma = (0.3431\Gamma_{33} + 0.3431\Gamma_{35} + 0.3431\Gamma_{38} + 0.3431\Gamma_{40} + 0.4307\Gamma_{45} + 0.6861\Gamma_{46} + \Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.9101\Gamma_{137} + 0.9101\Gamma_{138})/\Gamma$					
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
15.20 ± 0.07 OUR FIT	Error includes scale factor of 1.1.				
14.8 ± 0.4 OUR AVERAGE					
14.4 ± 0.6 ± 0.3		ADEVA	91F L3	$E_{\text{cm}}^{ee} = 88.3\text{--}94.3 \text{ GeV}$	
15.0 ± 0.4 ± 0.3		BEHREND	89B CELL	$E_{\text{cm}}^{ee} = 14\text{--}47 \text{ GeV}$	
15.1 ± 0.8 ± 0.6		AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹³³BURCHAT 87 value is not independent of SCHMIDKE 86 value.

¹³⁴Not independent of BERGER 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$, $\Gamma(h^- \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$, and $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$, and therefore not used in the fit.

¹³⁵Low energy experiments are not in average or fit because the systematic errors in background subtraction are judged to be large.

$$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-) (\text{"3-prong"})) / \Gamma_{\text{total}} \quad \Gamma_{53} / \Gamma$$

$$\Gamma_{53} / \Gamma = (\Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.9101\Gamma_{137} + 0.9101\Gamma_{138}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

14.57 ± 0.07 OUR FIT Error includes scale factor of 1.1.

14.59 ± 0.08 OUR AVERAGE Error includes scale factor of 1.1.

14.569 ± 0.093 ± 0.048	f&a	23k	¹³⁶ ABREU	01M DLPH	1992–1995 LEP runs
14.556 ± 0.105 ± 0.076	f&a		¹³⁷ ACHARD	01D L3	1992–1995 LEP runs
14.96 ± 0.09 ± 0.22	f&a	10.4k	AKERS	95Y OPAL	1991–1994 LEP runs
14.22 ± 0.10 ± 0.37	avg		¹³⁸ BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
15.26 ± 0.26 ± 0.22			ACTON	92H OPAL	Repl. by AK-ERS 95Y
13.3 ± 0.3 ± 0.8			¹³⁹ ALBRECHT	92D ARG	$E_{\text{cm}}^{ee} = 9.4$ – 10.6 GeV
14.35 $\begin{smallmatrix} +0.40 \\ -0.45 \end{smallmatrix}$ ± 0.24			DECAMP	92C ALEP	1989–1990 LEP runs

¹³⁶ The correlation coefficients between this measurement and the ABREU 01M measurements of $B(\tau \rightarrow 1\text{-prong})$ and $B(\tau \rightarrow 5\text{-prong})$ are -0.98 and -0.08 respectively.

¹³⁷ The correlation coefficients between this measurement and the ACHARD 01D measurements of $B(\tau \rightarrow \text{"1-prong"})$ and $B(\tau \rightarrow \text{"5-prong"})$ are -0.978 and -0.19 respectively.

¹³⁸ Not independent of BALEST 95C $B(h^- h^- h^+ \nu_\tau)$ and $B(h^- h^- h^+ \pi^0 \nu_\tau)$ values, and BORTOLETTO 93 $B(h^- h^- h^+ 2\pi^0 \nu_\tau) / B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau)$ value.

¹³⁹ This ALBRECHT 92D value is not independent of their $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}^2$ value.

$$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{54} / \Gamma$$

$$\Gamma_{54} / \Gamma = (0.3431\Gamma_{33} + 0.3431\Gamma_{35} + \Gamma_{60} + \Gamma_{82} + \Gamma_{89} + 0.0221\Gamma_{137}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.


VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

10.01 ± 0.09 OUR FIT Error includes scale factor of 1.2.

9.8 ± 0.6 OUR AVERAGE Error includes scale factor of 4.4. See the ideogram below.

7.6 ± 0.1 ± 0.5	avg	7.5k	¹⁴⁰ ALBRECHT	96E ARG	$E_{\text{cm}}^{ee} = 9.4$ – 10.6 GeV
9.92 ± 0.10 ± 0.09	f&a	11.2k	¹⁴¹ BUSKULIC	96 ALEP	LEP 1991–1993 data
9.49 ± 0.36 ± 0.63	f&a		DECAMP	92C ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •					
8.7 ± 0.7 ± 0.3		694	¹⁴² BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35$ GeV
7.0 ± 0.3 ± 0.7		1566	¹⁴³ BAND	87 MAC	$E_{\text{cm}}^{ee} = 29$ GeV
6.7 ± 0.8 ± 0.9			¹⁴⁴ BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
6.4 ± 0.4 ± 0.9			¹⁴⁵ RUCKSTUHL	86 DLCO	$E_{\text{cm}}^{ee} = 29$ GeV
7.8 ± 0.5 ± 0.8		890	SCHMIDKE	86 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
8.4 ± 0.4 ± 0.7		1255	¹⁴⁵ FERNANDEZ	85 MAC	$E_{\text{cm}}^{ee} = 29$ GeV
9.7 ± 2.0 ± 1.3			BEHREND	84 CELL	$E_{\text{cm}}^{ee} = 14, 22$ GeV

- 140 ALBRECHT 96E not independent of ALBRECHT 93C $\Gamma(h^- h^- h^+ \nu_\tau(\text{ex. } K^0)) \times \Gamma(\text{particle}^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}^2$ value.
- 141 BUSKULIC 96 quote $B(h^- h^- h^+ \nu_\tau(\text{ex. } K^0)) = 9.50 \pm 0.10 \pm 0.11$. We add 0.42 to remove their K^0 correction and reduce the systematic error accordingly.
- 142 BEHREND 90 subtract 0.3% to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution to measured events.
- 143 BAND 87 subtract for charged kaon modes; not independent of FERNANDEZ 85 value.
- 144 BURCHAT 87 value is not independent of SCHMIDKE 86 value.
- 145 Value obtained by multiplying paper's $R = B(h^- h^- h^+ \nu_\tau) / B(3\text{-prong})$ by $B(3\text{-prong}) = 0.143$ and subtracting 0.3% for $K^*(892)$ background.



$$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{\text{total}} (\%)$$

$$\Gamma(h^- h^- h^+ \nu_\tau(\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{55} / \Gamma$$

$$\Gamma_{55} / \Gamma = (\Gamma_{60} + \Gamma_{82} + \Gamma_{89} + 0.0221 \Gamma_{137}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
9.65±0.09 OUR FIT	Error includes scale factor of 1.2.				
9.57±0.11 OUR AVERAGE					
9.50±0.10±0.11	avg	11.2k	146 BUSKULIC	96 ALEP	LEP 1991–1993 data
9.87±0.10±0.24	avg		147 AKERS	95Y OPAL	1991–1994 LEP runs
9.51±0.07±0.20	f&a	37.7k	BALEST	95C CLEO	$E_{\text{cm}}^e \approx 10.6 \text{ GeV}$

- 146 Not independent of BUSKULIC 96 $B(h^- h^- h^+ \nu_\tau)$ value.
- 147 Not independent of AKERS 95Y $B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau(\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$ and $B(h^- h^- h^+ \nu_\tau(\text{ex. } K^0)) / B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau(\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$ values.

$$\Gamma(h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$$

Γ₅₅/Γ₅₃

$$\Gamma_{55}/\Gamma_{53} = (\Gamma_{60} + \Gamma_{82} + \Gamma_{89} + 0.0221\Gamma_{137}) / (\Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.9101\Gamma_{137} + 0.9101\Gamma_{138})$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.662 ± 0.006 OUR FIT			Error includes scale factor of 1.3.
0.660 ± 0.004 ± 0.014	AKERS	95Y OPAL	1991–1994 LEP runs

$$\Gamma(h^- h^- h^+ \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}}$$

Γ₅₆/Γ = (Γ₆₀ + Γ₈₂ + Γ₈₉)/Γ

VALUE (%)	DOCUMENT ID
9.61 ± 0.09 OUR FIT	Error includes scale factor of 1.2.

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}}$$

Γ₅₇/Γ = (0.3431Γ₃₃ + Γ₆₀ + 0.0221Γ₁₃₇)/Γ

VALUE (%)	DOCUMENT ID
9.52 ± 0.10 OUR FIT	Error includes scale factor of 1.2.

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$$

Γ₅₈/Γ = (Γ₆₀ + 0.0221Γ₁₃₇)/Γ

VALUE (%)	DOCUMENT ID
9.22 ± 0.10 OUR FIT	Error includes scale factor of 1.2.

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0), \text{ non-axial vector}) / \Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$$

Γ₅₉/Γ₅₈ = Γ₅₉/(Γ₆₀ + 0.0221Γ₁₃₇)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.261	95	148 ACKERSTAFF	97R OPAL	1992–1994 LEP runs

¹⁴⁸ Model-independent limit from structure function analysis on contribution to $B(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ from non-axial vectors.

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}}$$

Γ₆₀/Γ

VALUE (%)	DOCUMENT ID
9.18 ± 0.10 OUR FIT	Error includes scale factor of 1.2.

$$\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}}$$

Γ₆₁/Γ

$$\Gamma_{61}/\Gamma = (0.3431\Gamma_{38} + 0.3431\Gamma_{40} + 0.4307\Gamma_{45} + 0.6861\Gamma_{46} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{86} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.888\Gamma_{137} + 0.9101\Gamma_{138}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
5.19 ± 0.10 OUR FIT				Error includes scale factor of 1.3.

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.6 ± 0.7 ± 0.3	352	¹⁴⁹ BEHREND	90	CELL	$E_{\text{cm}}^{ee} = 35 \text{ GeV}$
4.2 ± 0.5 ± 0.9	203	¹⁵⁰ ALBRECHT	87L	ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
6.1 ± 0.8 ± 0.9		¹⁵¹ BURCHAT	87	MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
7.6 ± 0.4 ± 0.9		^{152,153} RUCKSTUHL	86	DLCO	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
4.7 ± 0.5 ± 0.8	530	¹⁵⁴ SCHMIDKE	86	MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
5.6 ± 0.4 ± 0.7		¹⁵³ FERNANDEZ	85	MAC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
6.2 ± 2.3 ± 1.7		BEHREND	84	CELL	$E_{\text{cm}}^{ee} = 14, 22 \text{ GeV}$

- 149 BEHREND 90 value is not independent of BEHREND 90 $B(3h\nu_\tau \geq 1 \text{ neutrals}) + B(5\text{-prong})$.
- 150 ALBRECHT 87L measure the product of branching ratios $B(3\pi^\pm \pi^0 \nu_\tau) B((e\bar{\nu} \text{ or } \mu\bar{\nu} \text{ or } \pi \text{ or } K \text{ or } \rho)\nu_\tau) = 0.029$ and use the PDG 86 values for the second branching ratio which sum to 0.69 ± 0.03 to get the quoted value.
- 151 BURCHAT 87 value is not independent of SCHMIDKE 86 value.
- 152 Contributions from kaons and from $>1\pi^0$ are subtracted. Not independent of $(3\text{-prong} + 0\pi^0)$ and $(3\text{-prong} + \geq 0\pi^0)$ values.
- 153 Value obtained using paper's $R = B(h^- h^- h^+ \nu_\tau)/B(3\text{-prong})$ and current $B(3\text{-prong}) = 0.143$.
- 154 Not independent of SCHMIDKE 86 $h^- h^- h^+ \nu_\tau$ and $h^- h^- h^+ (\geq 0\pi^0)\nu_\tau$ values.

$$\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))/\Gamma_{\text{total}} \quad \Gamma_{62}/\Gamma$$

$$\Gamma_{62}/\Gamma = (\Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{86} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.888\Gamma_{137} + 0.9101\Gamma_{138})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.92±0.10 OUR FIT				Error includes scale factor of 1.3.
5.07±0.24 OUR AVERAGE				
5.09±0.10±0.23	avg	155 AKERS	95Y OPAL	1991–1994 LEP runs
4.95±0.29±0.65	f&a	570 DECAMP	92C ALEP	1989–1990 LEP runs

- 155 Not independent of AKERS 95Y $B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$ and $B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K^0))/B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$ values.

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{63}/\Gamma$$

$$\Gamma_{63}/\Gamma = (0.3431\Gamma_{38} + 0.3431\Gamma_{40} + \Gamma_{68} + \Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119} + 0.888\Gamma_{137} + 0.0221\Gamma_{138})/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.53±0.09 OUR FIT				Error includes scale factor of 1.3.
4.45±0.09±0.07	6.1k	156 BUSKULIC	96 ALEP	LEP 1991–1993 data

- 156 BUSKULIC 96 quote $B(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0)) = 4.30 \pm 0.09 \pm 0.09$. We add 0.15 to remove their K^0 correction and reduce the systematic error accordingly.

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}} \quad \Gamma_{64}/\Gamma$$

$$\Gamma_{64}/\Gamma = (\Gamma_{68} + \Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119} + 0.888\Gamma_{137} + 0.0221\Gamma_{138})/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.35±0.09 OUR FIT				Error includes scale factor of 1.3.
4.23±0.06±0.22	7.2k	BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0, \omega))/\Gamma_{\text{total}} \quad \Gamma_{65}/\Gamma = (\Gamma_{68} + \Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119})/\Gamma$$

VALUE (%)	DOCUMENT ID
2.62±0.09 OUR FIT	Error includes scale factor of 1.2.

$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (\text{ex. } K^0, \omega, \eta))/\Gamma_{\text{total}}$ Γ_{74}/Γ
VALUE (%) DOCUMENT ID
0.11 ± 0.04 OUR FIT

$\Gamma(h^- h^- h^+ 3\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{75}/Γ
VALUE (%) EVTS DOCUMENT ID TECN COMMENT
0.023 ± 0.008 OUR FIT Error includes scale factor of 1.6.
0.023 ± 0.005 OUR AVERAGE

0.022 ± 0.003 ± 0.004	139	ANASTASSOV 01	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
0.11 ± 0.04 ± 0.05	440	¹⁵⁸ BUSKULIC 96	ALEP	LEP 1991–1993 data

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0285 ± 0.0056 ± 0.0051	57	ANDERSON 97	CLEO	Repl. by ANASTASSOV 01
--------------------------	----	-------------	------	------------------------

¹⁵⁸BUSKULIC 96 state their measurement is for $B(h^- h^- h^+ \geq 3\pi^0 \nu_\tau)$. We assume that $B(h^- h^- h^+ \geq 4\pi^0 \nu_\tau)$ is very small.

$\Gamma(K^- h^+ h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$
 $\Gamma_{76}/\Gamma = (0.3431\Gamma_{35} + 0.3431\Gamma_{40} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{119})/\Gamma$
VALUE (%) CL% DOCUMENT ID TECN COMMENT
0.65 ± 0.05 OUR FIT Error includes scale factor of 1.4.
< 0.6 90 AIHARA 84C TPC $E_{\text{cm}}^{ee} = 29$ GeV

$\Gamma(K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ $\Gamma_{77}/\Gamma = (\Gamma_{82} + \Gamma_{89})/\Gamma$
VALUE (%) DOCUMENT ID
0.44 ± 0.05 OUR FIT Error includes scale factor of 1.5.

$\Gamma(K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0))/\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$
 $\Gamma_{77}/\Gamma_{58} = (\Gamma_{82} + \Gamma_{89})/(\Gamma_{60} + 0.0221\Gamma_{137})$
VALUE (%) EVTS DOCUMENT ID TECN COMMENT
4.7 ± 0.6 OUR FIT Error includes scale factor of 1.5.
5.44 ± 0.21 ± 0.53 7.9k RICHICHI 99 CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(K^- h^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ $\Gamma_{78}/\Gamma = (\Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119})/\Gamma$
VALUE (%) DOCUMENT ID
0.110 ± 0.022 OUR FIT

$\Gamma(K^- h^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0))$
 $\Gamma_{78}/\Gamma_{67} = (\Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119})/(\Gamma_{68} + 0.888\Gamma_{137} + 0.0221\Gamma_{138})$
VALUE (%) EVTS DOCUMENT ID TECN COMMENT
2.6 ± 0.5 OUR FIT
2.61 ± 0.45 ± 0.42 719 RICHICHI 99 CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(K^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$
 $\Gamma_{79}/\Gamma = (0.3431\Gamma_{35} + 0.3431\Gamma_{40} + \Gamma_{82} + \Gamma_{86} + 0.285\Gamma_{119})/\Gamma$
VALUE (%) EVTS DOCUMENT ID TECN COMMENT
0.45 ± 0.05 OUR FIT Error includes scale factor of 1.4.

0.58^{+0.15}_{-0.13} ± 0.12	20	¹⁵⁹ BAUER	94	TPC	$E_{\text{cm}}^{ee} = 29$ GeV
--	----	----------------------	----	-----	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22 ^{+0.16} _{-0.13} ± 0.05	9	¹⁶⁰ MILLS	85	DLCO	$E_{\text{cm}}^{ee} = 29$ GeV
---	---	----------------------	----	------	-------------------------------

¹⁵⁹We multiply 0.58% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

¹⁶⁰Error correlated with MILLS 85 ($K K \pi \nu$) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain the systematic error.

$$\Gamma(K^- \pi^+ \pi^- \geq 0 \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{80} / \Gamma = (\Gamma_{82} + \Gamma_{86} + 0.231 \Gamma_{119}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		DOCUMENT ID	TECN	COMMENT
-----------	--	-------------	------	---------

0.35 ± 0.05 OUR FIT Error includes scale factor of 1.4.

0.30 ± 0.05 OUR AVERAGE

0.343 ± 0.073 ± 0.031	f&a	ABBIENDI	00D OPAL	1990–1995 LEP runs
0.275 ± 0.064	avg	¹⁶¹ BARATE	98 ALEP	1991–1995 LEP runs

¹⁶¹Not independent of BARATE 98 $\Gamma(\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}}$ and $\Gamma(\tau^- \rightarrow K^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ values.

$$\Gamma(K^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{81} / \Gamma = (0.3431 \Gamma_{35} + \Gamma_{82}) / \Gamma$$

VALUE (%)	DOCUMENT ID
-----------	-------------

0.33 ± 0.05 OUR FIT Error includes scale factor of 1.5.

$$\Gamma(K^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{82} / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------


0.28 ± 0.05 OUR FIT Error includes scale factor of 1.5.

0.28 ± 0.05 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

0.360 ± 0.082 ± 0.048	avg		ABBIENDI	00D OPAL	1990–1995 LEP runs
0.346 ± 0.023 ± 0.056	avg	158	¹⁶² RICHICHI	99 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
0.214 ± 0.037 ± 0.029	f&a		BARATE	98 ALEP	1991–1995 LEP runs

¹⁶²Not independent of RICHICHI 99

$\Gamma(\tau^- \rightarrow K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$, $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau) / \Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ values.



$$\Gamma(K^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} (\%)$$

$$\Gamma(K^- \rho^0 \nu_\tau \rightarrow K^- \pi^+ \pi^- \nu_\tau) / \Gamma(K^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) \quad \Gamma_{83} / \Gamma_{82}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.48±0.14±0.10	163 ASNER	00B CLEO	$E_{\text{cm}}^{\text{e}^{\text{e}}}$ = 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.39±0.14	164 BARATE	99R ALEP	1991–1995 LEP runs
-----------	------------	----------	--------------------

163 ASNER 00B assume $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0) decays proceed only through $K\rho$ and $K^* \pi$ intermediate states. They assume the resonance structure of $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0) decays is dominated by $K_1(1270)^-$ and $K_1(1400)^-$ resonances, and assume $B(K_1(1270) \rightarrow K^*(892)\pi) = (16 \pm 5)\%$, $B(K_1(1270) \rightarrow K\rho) = (42 \pm 6)\%$, and $B(K_1(1400) \rightarrow K\rho) = 0$.

164 BARATE 99R assume $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0) decays proceed only through $K\rho$ and $K^* \pi$ intermediate states. The quoted error is statistical only.

$$\Gamma(K^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{84} / \Gamma = (0.3431\Gamma_{40} + \Gamma_{86} + 0.231\Gamma_{119}) / \Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID
--------------------------	-------------

12.4±2.5 OUR FIT

$$\Gamma(K^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{85} / \Gamma = (\Gamma_{86} + 0.231\Gamma_{119}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.0±2.4 OUR FIT				
7.0±2.5 OUR AVERAGE				
7.5±2.6±1.8	avg	165 RICHICHI	99 CLEO	$E_{cm}^{ee} = 10.6$ GeV
6.1±3.9±1.8	f&a	BARATE	98 ALEP	1991–1995 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<17	95	ABBIENDI	00D OPAL	1990–1995 LEP runs
165 Not independent of RICHICHI 99				
$\Gamma(\tau^- \rightarrow K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0))/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$, $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ values.				

$\Gamma(K^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0, \eta))/\Gamma_{\text{total}}$ **Γ_{86}/Γ**
 Test of lepton family number conservation.

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
6.4±2.4 OUR FIT	

$\Gamma(K^- \pi^+ K^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ **Γ_{87}/Γ**

<u>VALUE (%)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.09	95	BAUER	94 TPC	$E_{cm}^{ee} = 29$ GeV

$\Gamma(K^- K^+ \pi^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ **$\Gamma_{88}/\Gamma = (\Gamma_{89} + \Gamma_{90})/\Gamma$**
 Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.200±0.023 OUR FIT				
0.203±0.031 OUR AVERAGE				
0.159±0.053±0.020	f&a	ABBIENDI	00D OPAL	1990–1995 LEP runs
0.238±0.042	avg	166 BARATE	98 ALEP	1991–1995 LEP runs
0.15 $\begin{smallmatrix} +0.09 \\ -0.07 \end{smallmatrix}$ ±0.03	f&a	4 167 BAUER	94 TPC	$E_{cm}^{ee} = 29$ GeV

166 Not independent of BARATE 98 $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ values.

167 We multiply 0.15% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$\Gamma(K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ **Γ_{89}/Γ**
 Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.160±0.019 OUR FIT				
0.151±0.019 OUR AVERAGE				
0.087±0.056±0.040	avg	ABBIENDI	00D OPAL	1990–1995 LEP runs
0.145±0.013±0.028	avg	2.3k 168 RICHICHI	99 CLEO	$E_{cm}^{ee} = 10.6$ GeV
0.163±0.021±0.017	f&a	BARATE	98 ALEP	1991–1995 LEP runs
0.22 $\begin{smallmatrix} +0.17 \\ -0.11 \end{smallmatrix}$ ±0.05	f&a	9 169 MILLS	85 DLCO	$E_{cm}^{ee} = 29$ GeV

¹⁶⁸ Not independent of RICHICHI 99 $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau) / \Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ values.

¹⁶⁹ Error correlated with MILLS 85 ($K \pi \pi \pi^0 \nu$) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain obtain the systematic error.

$$\Gamma(K^- K^+ \pi^- \nu_\tau) / \Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) \quad \Gamma_{89} / \Gamma_{58} = \Gamma_{89} / (\Gamma_{60} + 0.0221 \Gamma_{137})$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ± 0.20 OUR FIT				
1.60 ± 0.15 ± 0.30	2.3k	RICHICHI	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{90} / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.6 OUR FIT					
4.4 ± 1.8 OUR AVERAGE					Error includes scale factor of 1.1.
3.3 ± 1.8 ± 0.7	avg	158	¹⁷⁰ RICHICHI	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
7.5 ± 2.9 ± 1.5	f&a		BARATE	98	ALEP 1991–1995 LEP runs
• • •					We do not use the following data for averages, fits, limits, etc. • • •
<27	95		ABBIENDI	00D	OPAL 1990–1995 LEP runs

¹⁷⁰ Not independent of RICHICHI 99 $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau) / \Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ values.

$$\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau) / \Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0)) \quad \Gamma_{90} / \Gamma_{67} = \Gamma_{90} / (\Gamma_{68} + 0.888 \Gamma_{137} + 0.0221 \Gamma_{138})$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.9 ± 0.4 OUR FIT				
0.79 ± 0.44 ± 0.16	158	¹⁷¹ RICHICHI	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

¹⁷¹ RICHICHI 99 also quote a 95%CL upper limit of 0.0157 for this measurement.

$$\Gamma(K^- K^+ K^- \geq 0 \text{ neut. } \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{91} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.21	95	BAUER	94	TPC $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$$\Gamma(K^- K^+ K^- \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{92} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.9 × 10⁻⁴	90	BARATE	98	ALEP 1991–1995 LEP runs

$$\Gamma(\pi^- K^+ \pi^- \geq 0 \text{ neut. } \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{93} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.25	95	BAUER	94	TPC $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$\Gamma(e^- e^- e^+ \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$					Γ_{94}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.8 \pm 1.4 \pm 0.4$	5	ALAM	96	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\mu^- e^- e^+ \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$					Γ_{95}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
< 3.6	90	ALAM	96	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(3h^- 2h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^- \pi^+ \text{) ("5-prong")})/\Gamma_{\text{total}}$ Γ_{96}/Γ
 Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average. $\Gamma_{96}/\Gamma = (\Gamma_{97} + \Gamma_{98})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.100 ± 0.006 OUR FIT				
0.111 ± 0.008 OUR AVERAGE				Error includes scale factor of 1.1.
$0.115 \pm 0.013 \pm 0.006$	f&a	112 ¹⁷² ABREU	01M	DLPH 1992–1995 LEP runs
$0.170 \pm 0.022 \pm 0.026$	f&a	173 ACHARD	01D	L3 1992–1995 LEP runs
$0.119 \pm 0.013 \pm 0.008$	avg	119 ¹⁷⁴ ACKERSTAFF	99E	OPAL 1991–1995 LEP runs
$0.097 \pm 0.005 \pm 0.011$	f&a	419 GIBAUT	94B	CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV
0.102 ± 0.029	f&a	13 BYLSMA	87	HRS $E_{\text{cm}}^{ee} = 29$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.26 \pm 0.06 \pm 0.05$		ACTON	92H	OPAL $E_{\text{cm}}^{ee} = 88.2\text{--}94.2$ GeV
$0.10 \begin{smallmatrix} +0.05 \\ -0.04 \end{smallmatrix} \pm 0.03$		DECAMP	92C	ALEP 1989–1990 LEP runs
$0.16 \pm 0.13 \pm 0.04$		BEHREND	89B	CELL $E_{\text{cm}}^{ee} = 14\text{--}47$ GeV
$0.3 \pm 0.1 \pm 0.2$		BARTEL	85F	JADE $E_{\text{cm}}^{ee} = 34.6$ GeV
0.13 ± 0.04	10	BELTRAMI	85	HRS Repl. by BYLSMA 87
$0.16 \pm 0.08 \pm 0.04$	4	BURCHAT	85	MRK2 $E_{\text{cm}}^{ee} = 29$ GeV
1.0 ± 0.4	10	BEHREND	82	CELL Repl. by BEHREND 89B

¹⁷² The correlation coefficients between this measurement and the ABREU 01M measurements of $B(\tau \rightarrow 1\text{-prong})$ and $B(\tau \rightarrow 3\text{-prong})$ are -0.08 and -0.08 respectively.

¹⁷³ The correlation coefficients between this measurement and the ACHARD 01D measurements of $B(\tau \rightarrow "1\text{-prong}")$ and $B(\tau \rightarrow "3\text{-prong}")$ are -0.082 and -0.19 respectively.

¹⁷⁴ Not independent of ACKERSTAFF 99E $B(\tau^- \rightarrow 3h^- 2h^+ \nu_\tau \text{ (ex. } K^0))$ and $B(\tau^- \rightarrow 3h^- 2h^+ \pi^0 \nu_\tau \text{ (ex. } K^0))$ measurements.

$\Gamma(3h^- 2h^+ \nu_\tau \text{ (ex. } K^0))/\Gamma_{\text{total}}$					Γ_{97}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.082 ± 0.006 OUR FIT					
0.076 ± 0.007 OUR AVERAGE					
$0.091 \pm 0.014 \pm 0.006$	97	ACKERSTAFF	99E	OPAL 1991–1995 LEP runs	
$0.080 \pm 0.011 \pm 0.013$	58	BUSKULIC	96	ALEP LEP 1991–1993 data	
$0.077 \pm 0.005 \pm 0.009$	295	GIBAUT	94B	CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV	
$0.064 \pm 0.023 \pm 0.01$	12	ALBRECHT	88B	ARG $E_{\text{cm}}^{ee} = 10$ GeV	
0.051 ± 0.020	7	BYLSMA	87	HRS $E_{\text{cm}}^{ee} = 29$ GeV	


$\Gamma(K^*(892)^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{103} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.42 ± 0.18 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			

1.19 ± 0.15 ^{+0.13} _{-0.18}	104	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$
1.94 ± 0.27 ± 0.15	74	¹⁷⁹ AKERS	94G OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
1.43 ± 0.11 ± 0.13	475	¹⁸⁰ GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9 \text{ GeV}$

¹⁷⁹ AKERS 94G reject events in which a K_S^0 accompanies the $K^*(892)^-$. We do not correct for them.

¹⁸⁰ GOLDBERG 90 estimates that 10% of observed $K^*(892)$ are accompanied by a π^0 .



$\Gamma(K^*(892)^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{103} / Γ

$\Gamma(K^*(892)^- \nu_\tau) / \Gamma_{\text{total}}$ Γ_{104} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.29 ± 0.05 OUR AVERAGE				

1.326 ± 0.063		BARATE	99R ALEP	1991–1995 LEP runs
1.11 ± 0.12	181	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
1.42 ± 0.22 ± 0.09		¹⁸² ACCIARRI	95F L3	1991–1993 LEP runs
1.23 ± 0.21 ^{+0.11} _{-0.21}	54	¹⁸³ ALBRECHT	88L ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
1.9 ± 0.3 ± 0.4	44	¹⁸⁴ TSCHIRHART	88 HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
1.5 ± 0.4 ± 0.4	15	¹⁸⁵ AIHARA	87C TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
1.3 ± 0.3 ± 0.3	31	YELTON	86 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.39 ± 0.09 ± 0.10		¹⁸⁶ BUSKULIC	96 ALEP	Repl. by BARATE 99R
1.45 ± 0.13 ± 0.11	273	¹⁸⁷ BUSKULIC	94F ALEP	Repl. by BUSKULIC 96
1.7 ± 0.7	11	DORFAN	81 MRK2	$E_{\text{cm}}^{ee} = 4.2\text{--}6.7 \text{ GeV}$

- 181 Not independent of COAN 96 $B(\pi^- \bar{K}^0 \nu_\tau)$ and BATTLE 94 $B(K^- \pi^0 \nu_\tau)$ measurements. $K\pi$ final states are consistent with and assumed to originate from $K^*(892)^-$ production.
- 182 This result is obtained from their $B(\pi^- \bar{K}^0 \nu_\tau)$ assuming all those decays originate in $K^*(892)^-$ decays.
- 183 The authors divide by $\Gamma_2/\Gamma = 0.865$ to obtain this result.
- 184 Not independent of TSCHIRHART 88 $\Gamma(\tau^- \rightarrow h^- \bar{K}^0 \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau)/\Gamma(\text{total})$.
- 185 Decay π^- identified in this experiment, is assumed in the others.
- 186 Not independent of BUSKULIC 96 $B(\pi^- \bar{K}^0 \nu_\tau)$ and $B(K^- \pi^0 \nu_\tau)$ measurements.
- 187 BUSKULIC 94F obtain this result from BUSKULIC 94F $B(\bar{K}^0 \pi^- \nu_\tau)$ and BUSKULIC 94E $B(K^- \pi^0 \nu_\tau)$ assuming all of those decays originate in $K^*(892)^-$ decays.

$\Gamma(K^*(892)^- \nu_\tau)/\Gamma(\pi^- \pi^0 \nu_\tau)$		Γ_{104}/Γ_{13}		
VALUE		DOCUMENT ID	TECN	COMMENT
0.075 ± 0.027		188 ABREU	94K DLPH	LEP 1992 Z data

188 ABREU 94K quote $B(\tau^- \rightarrow K^*(892)^- \nu_\tau)B(K^*(892)^- \rightarrow K^- \pi^0)/B(\tau^- \rightarrow \rho^- \nu_\tau) = 0.025 \pm 0.009$. We divide by $B(K^*(892)^- \rightarrow K^- \pi^0) = 0.333$ to obtain this result.

$\Gamma(K^*(892)^0 K^- \geq 0 \text{ neutrals} \nu_\tau)/\Gamma_{\text{total}}$		Γ_{105}/Γ		
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.08 ± 0.12	119	GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9 \text{ GeV}$

$\Gamma(K^*(892)^0 K^- \nu_\tau)/\Gamma_{\text{total}}$		Γ_{106}/Γ		
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.21 ± 0.04 OUR AVERAGE				
0.213 ± 0.048		189 BARATE	98 ALEP	1991–1995 LEP runs
0.20 ± 0.05 ± 0.04	47	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$

189 BARATE 98 measure the $K^- (\rho^0 \rightarrow \pi^+ \pi^-)$ fraction in $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ decays to be $(35 \pm 11)\%$ and derive this result from their measurement of $\Gamma(\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ assuming the intermediate states are all $K^- \rho$ and $K^- K^*(892)^0$.

$\Gamma(\bar{K}^*(892)^0 \pi^- \geq 0 \text{ neutrals} \nu_\tau)/\Gamma_{\text{total}}$		Γ_{107}/Γ		
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.38 ± 0.11 ± 0.13	105	GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9 \text{ GeV}$

$\Gamma(\bar{K}^*(892)^0 \pi^- \nu_\tau)/\Gamma_{\text{total}}$		Γ_{108}/Γ		
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.22 ± 0.05 OUR AVERAGE				
0.209 ± 0.058		190 BARATE	98 ALEP	1991–1995 LEP runs
0.25 ± 0.10 ± 0.05	27	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$

190 BARATE 98 measure the $K^- K^*(892)^0$ fraction in $\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau$ decays to be $(87 \pm 13)\%$ and derive this result from their measurement of $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$.

$\Gamma((\bar{K}^*(892)\pi)^-\nu_\tau \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.10 ± 0.04 OUR AVERAGE			
0.097 ± 0.044 ± 0.036	191 BARATE	99K ALEP	1991–1995 LEP runs
0.106 ± 0.037 ± 0.032	192 BARATE	98E ALEP	1991–1995 LEP runs

191 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter. They determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.72 \pm 0.12 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by one minus this fraction to obtain the quoted result.

192 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+\pi^-$ decays. They determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.64 \pm 0.09 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by one minus this fraction to obtain the quoted result.

$\Gamma(K_1(1270)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.47 ± 0.11 OUR AVERAGE				
0.48 ± 0.11		BARATE	99R ALEP	1991–1995 LEP runs
0.41 ^{+0.41} _{-0.35} ± 0.10	5	193 BAUER	94 TPC	$E_{\text{cm}}^{e^+e^-} = 29$ GeV

193 We multiply 0.41% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$\Gamma(K_1(1400)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.26 OUR AVERAGE				Error includes scale factor of 1.7.
0.05 ± 0.17		BARATE	99R ALEP	1991–1995 LEP runs
0.76 ^{+0.40} _{-0.33} ± 0.20	11	194 BAUER	94 TPC	$E_{\text{cm}}^{e^+e^-} = 29$ GeV

194 We multiply 0.76% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$[\Gamma(K_1(1270)^-\nu_\tau) + \Gamma(K_1(1400)^-\nu_\tau)]/\Gamma_{\text{total}}$ $(\Gamma_{110} + \Gamma_{111})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.17^{+0.41}_{-0.37} ± 0.29	16	195 BAUER	94 TPC	$E_{\text{cm}}^{e^+e^-} = 29$ GeV

195 We multiply 1.17% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error. Not independent of BAUER 94 $B(K_1(1270)^-\nu_\tau)$ and BAUER 94 $B(K_1(1400)^-\nu_\tau)$ measurements.

$\Gamma(K_1(1270)^-\nu_\tau)/[\Gamma(K_1(1270)^-\nu_\tau) + \Gamma(K_1(1400)^-\nu_\tau)]$ $\Gamma_{110}/(\Gamma_{110} + \Gamma_{111})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.69 ± 0.15 OUR AVERAGE			
0.71 ± 0.16 ± 0.11	196 ABBIENDI	00D OPAL	1990–1995 LEP runs
0.66 ± 0.19 ± 0.13	197 ASNER	00B CLEO	$E_{\text{cm}}^{e^+e^-} = 10.6$ GeV

196 ABBIENDI 00D assume the resonance structure of $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ decays is dominated by the $K_1(1270)^-$ and $K_1(1400)^-$ resonances.

197 ASNER 00B assume the resonance structure of $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays is dominated by $K_1(1270)^-$ and $K_1(1400)^-$ resonances.

$\Gamma(K^*(1410)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.5^{+1.4}_{-1.0}$	BARATE	99R ALEP	1991–1995 LEP runs

$\Gamma(K_0^*(1430)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{113}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	95	BARATE	99R ALEP	1991–1995 LEP runs

$\Gamma(K_2^*(1430)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.3	95		TSCHIRHART	88 HRS	$E_{\text{cm}}^{ee} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.33	95		¹⁹⁸ ACCIARRI	95F L3	1991–1993 LEP runs
<0.9	95	0	DORFAN	81 MRK2	$E_{\text{cm}}^{ee} = 4.2\text{--}6.7$ GeV

¹⁹⁸ACCIARRI 95F quote $B(\tau^- \rightarrow K^*(1430)^- \rightarrow \pi^- \bar{K}^0 \nu_\tau) < 0.11\%$. We divide by $B(K^*(1430)^- \rightarrow \pi^- \bar{K}^0) = 0.33$ to obtain the limit shown.

$\Gamma(a_0(980)^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}} \times B(a_0(980) \rightarrow K^0 K^-)$ $\Gamma_{115}/\Gamma \times B$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9$ GeV

$\Gamma(\eta\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 1.4	95	0	BARTELT	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.2	95		BUSKULIC	97C ALEP	1991–1994 LEP runs
< 3.4	95		ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
< 90	95		ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10$ GeV
<140	90		BEHREND	88 CELL	$E_{\text{cm}}^{ee} = 14\text{--}46.8$ GeV
<180	95		BARINGER	87 CLEO	$E_{\text{cm}}^{ee} = 10.5$ GeV
<250	90	0	COFFMAN	87 MRK3	$E_{\text{cm}}^{ee} = 3.77$ GeV
510 $\pm 100 \pm 120$		65	DERRICK	87 HRS	$E_{\text{cm}}^{ee} = 29$ GeV
<100	95		GAN	87B MRK2	$E_{\text{cm}}^{ee} = 29$ GeV

$\Gamma(\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.174 \pm 0.024 OUR FIT					
0.173 \pm 0.024 OUR AVERAGE					
0.18 $\pm 0.04 \pm 0.02$			BUSKULIC	97C ALEP	1991–1994 LEP runs
0.17 $\pm 0.02 \pm 0.02$		125	ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.10	95	ALBRECHT	88M ARG	$E_{cm}^{ee} \approx 10$ GeV
<2.10	95	BARINGER	87 CLEO	$E_{cm}^{ee} = 10.5$ GeV
$4.20^{+0.70}_{-1.20} \pm 1.60$		¹⁹⁹ GAN	87 MRK2	$E_{cm}^{ee} = 29$ GeV

¹⁹⁹ Highly correlated with GAN 87 $\Gamma(\pi^- 3\pi^0 \nu_\tau)/\Gamma(\text{total})$ value.

$\Gamma(\eta \pi^- \pi^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{118}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.5 ± 0.5		30	²⁰⁰ ANASTASSOV 01	CLEO	$E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.4 \pm 0.6 \pm 0.3$		15	²⁰¹ BERGFELD	97 CLEO	Repl. by ANAS-TASSOV 01
< 4.3	95		ARTUSO	92 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV
<120	95		ALBRECHT	88M ARG	$E_{cm}^{ee} \approx 10$ GeV

²⁰⁰ Weighted average of BERGFELD 97 and ANASTASSOV 01 value of $(1.5 \pm 0.6 \pm 0.3) \times 10^{-4}$ obtained using η 's reconstructed from $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

²⁰¹ BERGFELD 97 reconstruct η 's using $\eta \rightarrow \gamma\gamma$ decays.

$\Gamma(\eta K^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.7 ± 0.6 OUR FIT					
2.7 ± 0.6 OUR AVERAGE					

$2.9^{+1.3}_{-1.2} \pm 0.7$			BUSKULIC	97c ALEP	1991–1994 LEP runs
$2.6 \pm 0.5 \pm 0.5$		85	BARTELT	96 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.7	95		ARTUSO	92 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV
------	----	--	--------	---------	--------------------------------

$\Gamma(\eta K^*(892)^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.90 ± 0.80 ± 0.42	25	BISHAI	99 CLEO	$E_{cm}^{ee} = 10.6$ GeV

$\Gamma(\eta K^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.77 ± 0.56 ± 0.71	36	BISHAI	99 CLEO	$E_{cm}^{ee} = 10.6$ GeV

$\Gamma(\eta \bar{K}^0 \pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20 ± 0.70 ± 0.22	15	²⁰² BISHAI	99 CLEO	$E_{cm}^{ee} = 10.6$ GeV

²⁰² We multiply the BISHAI 99 measurement $B(\tau^- \rightarrow \eta K_S^0 \pi^- \nu_\tau) = (1.10 \pm 0.35 \pm 0.11) \times 10^{-4}$ by 2 to obtain the listed value.

$\Gamma(\eta \pi^+ \pi^- \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.3	90	ABACHI	87B HRS	$E_{cm}^{ee} = 29$ GeV

$\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.5	170	203 ANASTASSOV 01	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.4^{+0.6}_{-0.5} \pm 0.6$	89	204 BERGFELD	97 CLEO	Repl. by ANASTASSOV 01
-----------------------------	----	--------------	---------	------------------------

²⁰³ Weighted average of BERGFELD 97 and ANASTASSOV 01 measurements using η 's reconstructed from $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow 3\pi^0$ decays.

²⁰⁴ BERGFELD 97 reconstruct η 's using $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi^0$ decays.

$\Gamma(\eta a_1(1260)^-\nu_\tau \rightarrow \eta\pi^-\rho^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.9 \times 10^{-4}$	90	BERGFELD	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\eta\eta\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.1	95	ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 83	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10$ GeV
--------	----	----------	---------	-------------------------------------

$\Gamma(\eta\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.0	95	ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 90	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10$ GeV
--------	----	----------	---------	-------------------------------------

$\Gamma(\eta'(958)\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.4 \times 10^{-5}$	90	BERGFELD	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\eta'(958)\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.0 \times 10^{-5}$	90	BERGFELD	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\phi\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.0 \times 10^{-4}$	90	²⁰⁵ AVERY	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.5 \times 10^{-4}$	90	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
------------------------	----	----------	---------	---

²⁰⁵ AVERY 97 limit varies from $(1.2\text{--}2.0) \times 10^{-4}$ depending on decay model assumptions.

$\Gamma(\phi K^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.7 \times 10^{-5}$	90	²⁰⁶ AVERY	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

²⁰⁶ AVERY 97 limit varies from $(5.4\text{--}6.7) \times 10^{-5}$ depending on decay model assumptions.

$\Gamma(f_1(1285)\pi^-\nu_\tau)/\Gamma_{\text{total}}$					Γ_{132}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$5.8^{+1.4}_{-1.3} \pm 1.8$	54	BERGFELD	97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(f_1(1285)\pi^-\nu_\tau \rightarrow \eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)$					$\Gamma_{133}/\Gamma_{124}$
VALUE		DOCUMENT ID	TECN	COMMENT	
0.55 ± 0.14		BERGFELD	97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\pi(1300)^-\nu_\tau \rightarrow (\rho\pi)^-\nu_\tau \rightarrow (3\pi)^-\nu_\tau)/\Gamma_{\text{total}}$					Γ_{134}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 1.0 \times 10^{-4}$	90	ASNER	00	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\pi(1300)^-\nu_\tau \rightarrow ((\pi\pi)_{\text{S-wave}}\pi)^-\nu_\tau \rightarrow (3\pi)^-\nu_\tau)/\Gamma_{\text{total}}$					Γ_{135}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 1.9 \times 10^{-4}$	90	ASNER	00	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(h^-\omega \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$					Γ_{136}/Γ
$\Gamma_{136}/\Gamma = (\Gamma_{137} + \Gamma_{138})/\Gamma$					

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
2.37 ± 0.08 OUR FIT					
1.65 ± 0.3 ± 0.2	avg	1513	ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10$ GeV

$\Gamma(h^-\omega\nu_\tau)/\Gamma_{\text{total}}$					Γ_{137}/Γ
Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.					

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
1.94 ± 0.07 OUR FIT					
1.92 ± 0.07 OUR AVERAGE					
1.91 ± 0.07 ± 0.06	f&a	5803	BUSKULIC	97C ALEP	1991–1994 LEP
1.95 ± 0.07 ± 0.11	avg	2223	²⁰⁷ BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
1.60 ± 0.27 ± 0.41	f&a	139	BARINGER	87 CLEO	$E_{\text{cm}}^{ee} = 10.5$ GeV

²⁰⁷ Not independent of BALEST 95C $B(\tau^- \rightarrow h^-\omega\nu_\tau)/B(\tau^- \rightarrow h^-h^+h^+\pi^0\nu_\tau)$ value.

$\Gamma(h^-\omega\nu_\tau)/\Gamma(h^-h^+h^+\pi^0\nu_\tau \text{ (ex. } K^0))$					Γ_{137}/Γ_{64}
$\Gamma_{137}/\Gamma_{64} = \Gamma_{137}/(\Gamma_{68} + \Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119} + 0.888\Gamma_{137} + 0.0221\Gamma_{138})$					

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.446 ± 0.015 OUR FIT				
0.453 ± 0.019 OUR AVERAGE				
0.431 ± 0.033	2350	²⁰⁸ BUSKULIC	96	ALEP LEP 1991–1993 data
0.464 ± 0.016 ± 0.017	2223	²⁰⁹ BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.37 ± 0.05 ± 0.02	458	²¹⁰ ALBRECHT	91D ARG	$E_{\text{cm}}^{ee} = 9.4$ – 10.6 GeV

208 BUSKULIC 96 quote the fraction of $\tau \rightarrow h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0) decays which originate in a $h^- \omega$ final state = 0.383 ± 0.029 . We divide this by the $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ branching fraction (0.888).

209 BALEST 95C quote the fraction of $\tau^- \rightarrow h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0) decays which originate in a $h^- \omega$ final state equals $0.412 \pm 0.014 \pm 0.015$. We divide this by the $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ branching fraction (0.888).

210 ALBRECHT 91D quote the fraction of $\tau^- \rightarrow h^- h^- h^+ \pi^0 \nu_\tau$ decays which originate in a $\pi^- \omega$ final state equals $0.33 \pm 0.04 \pm 0.02$. We divide this by the $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ branching fraction (0.888).

$\Gamma(h^- \omega \pi^0 \nu_\tau) / \Gamma_{\text{total}}$					Γ_{138} / Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.43 ± 0.05 OUR FIT					
0.43 ± 0.06 ± 0.05	7283	BUSKULIC	97C ALEP	1991–1994 LEP runs	

$\Gamma(h^- \omega 2\pi^0 \nu_\tau) / \Gamma_{\text{total}}$					Γ_{139} / Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.4 ± 0.4 ± 0.3	53	ANASTASSOV 01	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.89^{+0.74}_{-0.67} \pm 0.40$	19	ANDERSON 97	CLEO	Repl. by ANASTASSOV 01
---------------------------------	----	-------------	------	------------------------

$\Gamma(h^- \omega \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau)$					$\Gamma_{138} / \Gamma_{52}$
$\Gamma_{138} / \Gamma_{52} = \Gamma_{138} / (0.3431\Gamma_{33} + 0.3431\Gamma_{35} + 0.3431\Gamma_{38} + 0.3431\Gamma_{40} + 0.4307\Gamma_{45} + 0.6861\Gamma_{46} + \Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.9101\Gamma_{137} + 0.9101\Gamma_{138})$					

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0286 ± 0.0031 OUR FIT				
0.028 ± 0.003 ± 0.003	avg 430	211 BORTOLETTO93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

211 Not independent of BORTOLETTO 93 $\Gamma(\tau^- \rightarrow h^- \omega \pi^0 \nu_\tau) / \Gamma(\tau^- \rightarrow h^- h^- h^+ 2\pi^0 \nu_\tau \text{ (ex. } K^0))$ value.

$\Gamma(h^- \omega \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau \text{ (ex. } K^0))$					$\Gamma_{138} / \Gamma_{73}$
$\Gamma_{138} / \Gamma_{73} = \Gamma_{138} / (\Gamma_{74} + 0.236\Gamma_{117} + 0.888\Gamma_{138})$					

VALUE	DOCUMENT ID	TECN	COMMENT
0.81 ± 0.08 OUR FIT			
0.81 ± 0.06 ± 0.06	BORTOLETTO93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

$\Gamma(2h^- h^+ \omega \nu_\tau) / \Gamma_{\text{total}}$					Γ_{140} / Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.2 ± 0.2 ± 0.1	110	ANASTASSOV 01	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV	

$\Gamma(e^- \gamma)/\Gamma_{\text{total}}$ Γ_{141}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.7 \times 10^{-6}$	90	EDWARDS	97 CLEO	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 1.1 \times 10^{-4}$	90	ABREU	95U DLPH	1990–1993 LEP runs
$< 1.2 \times 10^{-4}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10$ GeV
$< 2.0 \times 10^{-4}$	90	KEH	88 CBAL	$E_{\text{cm}}^{ee} = 10$ GeV
$< 6.4 \times 10^{-4}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8$ GeV

$\Gamma(\mu^- \gamma)/\Gamma_{\text{total}}$ Γ_{142}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.1 \times 10^{-6}$	90	AHMED	00 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 3.0 \times 10^{-6}$	90	EDWARDS	97 CLEO	
$< 6.2 \times 10^{-5}$	90	ABREU	95U DLPH	1990–1993 LEP runs
$< 0.42 \times 10^{-5}$	90	BEAN	93 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
$< 3.4 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10$ GeV
$< 55 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8$ GeV

$\Gamma(e^- \pi^0)/\Gamma_{\text{total}}$ Γ_{143}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 3.7 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 17 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10$ GeV
$< 14 \times 10^{-5}$	90	KEH	88 CBAL	$E_{\text{cm}}^{ee} = 10$ GeV
$< 210 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8$ GeV

$\Gamma(\mu^- \pi^0)/\Gamma_{\text{total}}$ Γ_{144}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 4.0 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 4.4 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10$ GeV
$< 82 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8$ GeV

$\Gamma(e^- K_S^0)/\Gamma_{\text{total}}$ Γ_{145}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 9.1 \times 10^{-7}$	90	CHEN	02C CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 1.3 \times 10^{-3}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8$ GeV

$\Gamma(\mu^- K_S^0)/\Gamma_{\text{total}}$ Γ_{146}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.5 \times 10^{-7}$	90	CHEN	02C CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 1.0 \times 10^{-3}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

$\Gamma(e^- \eta)/\Gamma_{\text{total}}$ Γ_{147}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.2 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 6.3 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 24 \times 10^{-5}$	90	KEH	88 CBAL	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

$\Gamma(\mu^- \eta)/\Gamma_{\text{total}}$ Γ_{148}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.6 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 7.3 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

$\Gamma(e^- \rho^0)/\Gamma_{\text{total}}$ Γ_{149}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.0 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.42 \times 10^{-5}$	90	²¹² BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.9 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 37 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²¹² BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \rho^0)/\Gamma_{\text{total}}$ Γ_{150}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.57 \times 10^{-5}$	90	²¹³ BARTELT	94 CLEO	Repl. by BLISS 98
$< 2.9 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 44 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²¹³ BARTELT 94 assume phase space decays.

$\Gamma(e^- K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{151}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.1 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.63 × 10 ⁻⁵	90	²¹⁴ BARTELT	94 CLEO	Repl. by BLISS 98
<3.8 × 10 ⁻⁵	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

²¹⁴ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{152}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.5 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.94 × 10 ⁻⁵	90	²¹⁵ BARTELT	94 CLEO	Repl. by BLISS 98
<4.5 × 10 ⁻⁵	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

²¹⁵ BARTELT 94 assume phase space decays.

$\Gamma(e^- \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{153}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.4 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
••• We do not use the following data for averages, fits, limits, etc. •••				
<1.1 × 10 ⁻⁵	90	²¹⁶ BARTELT	94 CLEO	Repl. by BLISS 98

²¹⁶ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{154}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.5 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.87 × 10 ⁻⁵	90	²¹⁷ BARTELT	94 CLEO	Repl. by BLISS 98

²¹⁷ BARTELT 94 assume phase space decays.

$\Gamma(e^- \phi)/\Gamma_{\text{total}}$ Γ_{155}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.9 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \phi)/\Gamma_{\text{total}}$ Γ_{156}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.0 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{157}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.9 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 0.33 \times 10^{-5}$	90	²¹⁸ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.3 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$
$< 40 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²¹⁸ BARTELT 94 assume phase space decays.

$\Gamma(e^- \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{158}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.8 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 0.36 \times 10^{-5}$	90	²¹⁹ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.9 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$
$< 33 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²¹⁹ BARTELT 94 assume phase space decays.

$\Gamma(e^+ \mu^- \mu^-)/\Gamma_{\text{total}}$ **Γ_{159}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.5 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 0.35 \times 10^{-5}$	90	²²⁰ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.8 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 1.6 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²²⁰ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{160}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.7 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 0.34 \times 10^{-5}$	90	²²¹ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.4 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$
$< 44 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²²¹ BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ e^- e^-)/\Gamma_{\text{total}}$ Γ_{161}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.5 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.34 × 10 ⁻⁵	90	²²² BARTELT	94 CLEO	Repl. by BLISS 98
<1.4 × 10 ⁻⁵	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
<1.6 × 10 ⁻⁵	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²²² BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{162}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.9 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.43 × 10 ⁻⁵	90	²²³ BARTELT	94 CLEO	Repl. by BLISS 98
< 1.9 × 10 ⁻⁵	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
< 1.7 × 10 ⁻⁵	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$
<49 × 10 ⁻⁵	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²²³ BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{163}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.2 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.44 × 10 ⁻⁵	90	²²⁴ BARTELT	94 CLEO	Repl. by BLISS 98
<2.7 × 10 ⁻⁵	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
<6.0 × 10 ⁻⁵	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²²⁴ BARTELT 94 assume phase space decays.

$\Gamma(e^+ \pi^- \pi^-)/\Gamma_{\text{total}}$ Γ_{164}/Γ

Test of lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.9 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.44 × 10 ⁻⁵	90	²²⁵ BARTELT	94 CLEO	Repl. by BLISS 98
<1.8 × 10 ⁻⁵	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
<1.7 × 10 ⁻⁵	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²²⁵ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{165}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.2 × 10⁻⁶	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.74 \times 10^{-5}$	90	²²⁶ BARTELT	94	CLEO	Repl. by BLISS 98
$<3.6 \times 10^{-5}$	90	ALBRECHT	92k	ARG	$E_{cm}^{ee} = 10$ GeV
$<3.9 \times 10^{-5}$	90	BOWCOCK	90	CLEO	$E_{cm}^{ee} = 10.4-10.9$

²²⁶ BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ \pi^- \pi^-)/\Gamma_{total}$ Γ_{166}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-6}$	90	BLISS	98	CLEO $E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.69 \times 10^{-5}$	90	²²⁷ BARTELT	94	CLEO	Repl. by BLISS 98
$<6.3 \times 10^{-5}$	90	ALBRECHT	92k	ARG	$E_{cm}^{ee} = 10$ GeV
$<3.9 \times 10^{-5}$	90	BOWCOCK	90	CLEO	$E_{cm}^{ee} = 10.4-10.9$

²²⁷ BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^+ K^-)/\Gamma_{total}$ Γ_{167}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.4 \times 10^{-6}$	90	BLISS	98	CLEO $E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.77 \times 10^{-5}$	90	²²⁸ BARTELT	94	CLEO	Repl. by BLISS 98
$<2.9 \times 10^{-5}$	90	ALBRECHT	92k	ARG	$E_{cm}^{ee} = 10$ GeV
$<5.8 \times 10^{-5}$	90	BOWCOCK	90	CLEO	$E_{cm}^{ee} = 10.4-10.9$

²²⁸ BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^- K^+)/\Gamma_{total}$ Γ_{168}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.8 \times 10^{-6}$	90	BLISS	98	CLEO $E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.46 \times 10^{-5}$	90	²²⁹ BARTELT	94	CLEO	Repl. by BLISS 98
$<5.8 \times 10^{-5}$	90	BOWCOCK	90	CLEO	$E_{cm}^{ee} = 10.4-10.9$

²²⁹ BARTELT 94 assume phase space decays.

$\Gamma(e^+ \pi^- K^-)/\Gamma_{total}$ Γ_{169}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-6}$	90	BLISS	98	CLEO $E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.45 \times 10^{-5}$	90	²³⁰ BARTELT	94	CLEO	Repl. by BLISS 98
$<2.0 \times 10^{-5}$	90	ALBRECHT	92k	ARG	$E_{cm}^{ee} = 10$ GeV
$<4.9 \times 10^{-5}$	90	BOWCOCK	90	CLEO	$E_{cm}^{ee} = 10.4-10.9$

²³⁰ BARTELT 94 assume phase space decays.

$\Gamma(e^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{170}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-6}$	90	CHEN	02C CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{171}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.0 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^+ K^- K^-)/\Gamma_{\text{total}}$ Γ_{172}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.8 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \pi^+ K^-)/\Gamma_{\text{total}}$ Γ_{173}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.5 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.87 \times 10^{-5}$	90	²³¹ BARTELT	94 CLEO	Repl. by BLISS 98
$<11 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$<7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³¹ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \pi^- K^+)/\Gamma_{\text{total}}$ Γ_{174}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.4 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.5 \times 10^{-5}$	90	²³² BARTELT	94 CLEO	Repl. by BLISS 98
$<7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³² BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ \pi^- K^-)/\Gamma_{\text{total}}$ Γ_{175}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.0 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.0 \times 10^{-5}$	90	²³³ BARTELT	94 CLEO	Repl. by BLISS 98
$<5.8 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$<4.0 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³³ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{176}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-6}$	90	CHEN	02C CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{177}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<15 \times 10^{-6}$	90	BLISS	98	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^+ K^- K^-)/\Gamma_{\text{total}}$ Γ_{178}/Γ

Test of lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.0 \times 10^{-6}$	90	BLISS	98	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{179}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.5 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{180}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<14 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- \eta \eta)/\Gamma_{\text{total}}$ Γ_{181}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<35 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \eta \eta)/\Gamma_{\text{total}}$ Γ_{182}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<60 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- \pi^0 \eta)/\Gamma_{\text{total}}$ Γ_{183}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<24 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \pi^0 \eta)/\Gamma_{\text{total}}$ Γ_{184}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<22 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\bar{p} \gamma)/\Gamma_{\text{total}}$ Γ_{185}/Γ

Test of lepton number and baryon number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 3.5 \times 10^{-6}$	90	GODANG	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<29 \times 10^{-5}$	90	ALBRECHT	92K	ARG $E_{\text{cm}}^{ee} = 10 \text{ GeV}$
----------------------	----	----------	-----	---

$\Gamma(\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{186}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<15 \times 10^{-6}$	90	GODANG	99 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<66 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

$\Gamma(\bar{p}2\pi^0)/\Gamma_{\text{total}}$ Γ_{187}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<33 \times 10^{-6}$	90	GODANG	99 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{188}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.9 \times 10^{-6}$	90	GODANG	99 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<130 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

$\Gamma(\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{189}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<27 \times 10^{-6}$	90	GODANG	99 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- \text{ light boson})/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ Γ_{190}/Γ_5

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.015	95	²³⁴ ALBRECHT	95G ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.018	95	²³⁵ ALBRECHT	90E ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$
<0.040	95	²³⁶ BALTRUSAIT..85	MRK3	$E_{\text{cm}}^{ee} = 3.77 \text{ GeV}$

²³⁴ ALBRECHT 95G limit holds for bosons with mass $< 0.4 \text{ GeV}$. The limit rises to 0.036 for a mass of 1.0 GeV, then falls to 0.006 at the upper mass limit of 1.6 GeV.

²³⁵ ALBRECHT 90E limit applies for spinless boson with mass $< 100 \text{ MeV}$, and rises to 0.050 for mass = 500 MeV.

²³⁶ BALTRUSAITIS 85 limit applies for spinless boson with mass $< 100 \text{ MeV}$.

$\Gamma(\mu^- \text{ light boson})/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ Γ_{191}/Γ_5

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.026	95	²³⁷ ALBRECHT	95G ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.033	95	²³⁸ ALBRECHT	90E ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$
<0.125	95	²³⁹ BALTRUSAIT..85	MRK3	$E_{\text{cm}}^{ee} = 3.77 \text{ GeV}$

²³⁷ ALBRECHT 95G limit holds for bosons with mass $< 1.3 \text{ GeV}$. The limit rises to 0.034 for a mass of 1.4 GeV, then falls to 0.003 at the upper mass limit of 1.6 GeV.

²³⁸ ALBRECHT 90E limit applies for spinless boson with mass $< 100 \text{ MeV}$, and rises to 0.071 for mass = 500 MeV.

²³⁹ BALTRUSAITIS 85 limit applies for spinless boson with mass $< 100 \text{ MeV}$.

τ -DECAY PARAMETERS

A REVIEW GOES HERE – Check our WWW List of Reviews

$\rho^\tau(e \text{ or } \mu)$ PARAMETER

($V-A$) theory predicts $\rho = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.745±0.008 OUR FIT				
0.749±0.008 OUR AVERAGE				
0.742±0.014±0.006	81k	HEISTER	01E ALEP	1991–1995 LEP runs
0.775±0.023±0.020	36k	ABREU	00L DLPH	1992–1995 runs
0.781±0.028±0.018	46k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.762±0.035	54k	ACCIARRI	98R L3	1991–1995 LEP runs
0.731±0.031		²⁴⁰ ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.72 ±0.09 ±0.03		²⁴¹ ABE	97O SLD	1993–1995 SLC runs
0.747±0.010±0.006	55k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
0.79 ±0.10 ±0.10	3732	FORD	87B MAC	$E_{cm}^{ee} = 29$ GeV
0.71 ±0.09 ±0.03	1426	BEHREND	85 CLEO	e^+e^- near $\Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.735±0.013±0.008	31k	AMMAR	97B CLEO	Repl. by ALEXANDER 97F
0.794±0.039±0.031	18k	ACCIARRI	96H L3	Repl. by ACCIARRI 98R
0.732±0.034±0.020	8.2k	²⁴² ALBRECHT	95 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.738±0.038		²⁴³ ALBRECHT	95C ARG	Repl. by ALBRECHT 98
0.751±0.039±0.022		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
0.742±0.035±0.020	8000	ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV

²⁴⁰ Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

²⁴¹ ABE 97O assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a ρ^τ value of $0.69 \pm 0.13 \pm 0.05$.

²⁴² Value is from a simultaneous fit for the ρ^τ and η^τ decay parameters to the lepton energy spectrum. Not independent of ALBRECHT 90E $\rho^\tau(e \text{ or } \mu)$ value which assumes $\eta^\tau = 0$. Result is strongly correlated with ALBRECHT 95C.

²⁴³ Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E.

$\rho^\tau(e)$ PARAMETER

($V-A$) theory predicts $\rho = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.747±0.010 OUR FIT				
0.744±0.010 OUR AVERAGE				
0.747±0.019±0.014	44k	HEISTER	01E ALEP	1991–1995 LEP runs
0.744±0.036±0.037	17k	ABREU	00L DLPH	1992–1995 runs
0.779±0.047±0.029	25k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.68 ±0.04 ±0.07		²⁴⁴ ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.71 ±0.14 ±0.05		ABE	97O SLD	1993–1995 SLC runs
0.747±0.012±0.004	34k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV

0.735±0.036±0.020	4.7k	245	ALBRECHT	95	ARG	$E_{cm}^{ee} = 9.5-10.6$ GeV
0.79 ±0.08 ±0.06	3230	246	ALBRECHT	93G	ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.64 ±0.06 ±0.07	2753		JANSSEN	89	CBAL	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.62 ±0.17 ±0.14	1823		FORD	87B	MAC	$E_{cm}^{ee} = 29$ GeV
0.60 ±0.13	699		BEHREND	85	CLEO	e^+e^- near $\Upsilon(4S)$
0.72 ±0.10 ±0.11	594		BACINO	79B	DLCO	$E_{cm}^{ee} = 3.5-7.4$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.732±0.014±0.009	19k		AMMAR	97B	CLEO	Repl. by ALEXANDER 97F
0.793±0.050±0.025			BUSKULIC	95D	ALEP	Repl. by HEISTER 01E
0.747±0.045±0.028	5106		ALBRECHT	90E	ARG	Repl. by ALBRECHT 95
244 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.						
245 ALBRECHT 95 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(h^+ h^- h^+ (\pi^0) \bar{\nu}_\tau)$ and their charged conjugates.						
246 ALBRECHT 93G use tau pair events of the type $\tau^- \tau^+ \rightarrow (\mu^- \bar{\nu}_\mu \nu_\tau)(e^+ \nu_e \bar{\nu}_\tau)$ and their charged conjugates.						

$\rho^\tau(\mu)$ PARAMETER

(V-A) theory predicts $\rho = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.763±0.020 OUR FIT				
0.770±0.022 OUR AVERAGE				
0.776±0.045±0.019	46k	HEISTER	01E ALEP	1991-1995 LEP runs
0.999±0.098±0.045	22k	ABREU	00L DLPH	1992-1995 runs
0.777±0.044±0.016	27k	ACKERSTAFF	99D OPAL	1990-1995 LEP runs
0.69 ±0.06 ±0.06		247 ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5-10.6$ GeV
0.54 ±0.28 ±0.14		ABE	97O SLD	1993-1995 SLC runs
0.750±0.017±0.045	22k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
0.76 ±0.07 ±0.08	3230	ALBRECHT	93G ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.734±0.055±0.027	3041	ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.89 ±0.14 ±0.08	1909	FORD	87B MAC	$E_{cm}^{ee} = 29$ GeV
0.81 ±0.13	727	BEHREND	85 CLEO	e^+e^- near $\Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.747±0.048±0.044	13k	AMMAR	97B CLEO	Repl. by ALEXANDER 97F
0.693±0.057±0.028		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
247 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.				

$\xi^\tau(e \text{ or } \mu)$ PARAMETER

(V-A) theory predicts $\xi = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.985±0.030 OUR FIT				
0.981±0.031 OUR AVERAGE				
0.986±0.068±0.031	81k	HEISTER	01E ALEP	1991-1995 LEP runs
0.929±0.070±0.030	36k	ABREU	00L DLPH	1992-1995 runs
0.98 ±0.22 ±0.10	46k	ACKERSTAFF	99D OPAL	1990-1995 LEP runs
0.70 ±0.16	54k	ACCIARRI	98R L3	1991-1995 LEP runs
1.03 ±0.11		248 ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5-10.6$ GeV
1.05 ±0.35 ±0.04		249 ABE	97O SLD	1993-1995 SLC runs
1.007±0.040±0.015	55k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.94 ± 0.21 ± 0.07	18k	ACCIARRI	96H L3	Repl. by ACCIARRI 98R
0.97 ± 0.14		250 ALBRECHT	95C ARG	Repl. by ALBRECHT 98
1.18 ± 0.15 ± 0.16		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
0.90 ± 0.15 ± 0.10	3230	251 ALBRECHT	93G ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV

248 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

249 ABE 97O assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a ξ^τ value of $1.02 \pm 0.36 \pm 0.05$.

250 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 95C uses events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(h^+ h^- h^+ \bar{\nu}_\tau)$ and their charged conjugates.

251 ALBRECHT 93G measurement determines $|\xi^\tau|$ for the case $\xi^\tau(e) = \xi^\tau(\mu)$, but the authors point out that other LEP experiments determine the sign to be positive.

$\xi^\tau(e)$ PARAMETER

(V-A) theory predicts $\xi = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.994 ± 0.040 OUR FIT				
1.00 ± 0.04 OUR AVERAGE				
1.011 ± 0.094 ± 0.038	44k	HEISTER	01E ALEP	1991–1995 LEP runs
1.01 ± 0.12 ± 0.05	17k	ABREU	00L DLPH	1992–1995 runs
1.13 ± 0.39 ± 0.14	25k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
1.11 ± 0.20 ± 0.08		252 ALBRECHT	98 ARG	$E_{\text{cm}}^{ee} = 9.5\text{--}10.6$ GeV
1.16 ± 0.52 ± 0.06		ABE	97O SLD	1993–1995 SLC runs
0.979 ± 0.048 ± 0.016	34k	ALEXANDER	97F CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.03 ± 0.23 ± 0.09		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
--------------------	--	----------	----------	----------------------

252 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

$\xi^\tau(\mu)$ PARAMETER

(V-A) theory predicts $\xi = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.030 ± 0.059 OUR FIT				
1.06 ± 0.06 OUR AVERAGE				
1.030 ± 0.120 ± 0.050	46k	HEISTER	01E ALEP	1991–1995 LEP runs
1.16 ± 0.19 ± 0.06	22k	ABREU	00L DLPH	1992–1995 runs
0.79 ± 0.41 ± 0.09	27k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
1.26 ± 0.27 ± 0.14		253 ALBRECHT	98 ARG	$E_{\text{cm}}^{ee} = 9.5\text{--}10.6$ GeV
0.75 ± 0.50 ± 0.14		ABE	97O SLD	1993–1995 SLC runs
1.054 ± 0.069 ± 0.047	22k	ALEXANDER	97F CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.23 ± 0.22 ± 0.10		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
--------------------	--	----------	----------	----------------------

253 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

$\eta^\tau(e \text{ or } \mu)$ PARAMETER $(V-A)$ theory predicts $\eta = 0$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.013±0.020 OUR FIT				
0.015±0.021 OUR AVERAGE				
0.012±0.026±0.004	81k	HEISTER	01E ALEP	1991–1995 LEP runs
−0.005±0.036±0.037		ABREU	00L DLPH	1992–1995 runs
0.027±0.055±0.005	46k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.27 ±0.14	54k	ACCIARRI	98R L3	1991–1995 LEP runs
−0.13 ±0.47 ±0.15		ABE	97O SLD	1993–1995 SLC runs
−0.015±0.061±0.062	31k	AMMAR	97B CLEO	$E_{cm}^{ee} = 10.6$ GeV
0.03 ±0.18 ±0.12	8.2k	ALBRECHT	95 ARG	$E_{cm}^{ee} = 9.5$ –10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.25 ±0.17 ±0.11	18k	ACCIARRI	96H L3	Repl. by ACCIARRI 98R
−0.04 ±0.15 ±0.11		BUSKULIC	95D ALEP	Repl. by HEISTER 01E

 $\eta^\tau(\mu)$ PARAMETER $(V-A)$ theory predicts $\eta = 0$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.094±0.073 OUR FIT				
0.17 ±0.15 OUR AVERAGE Error includes scale factor of 1.2.				
0.160±0.150±0.060	46k	HEISTER	01E ALEP	1991–1995 LEP runs
0.72 ±0.32 ±0.15		ABREU	00L DLPH	1992–1995 runs
−0.59 ±0.82 ±0.45	254	ABE	97O SLD	1993–1995 SLC runs
0.010±0.149±0.171	13k	255 AMMAR	97B CLEO	$E_{cm}^{ee} = 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.010±0.065±0.001	27k	256 ACKERSTAFF	99D OPAL	1990–1995 LEP runs
−0.24 ±0.23 ±0.18		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
254 Highly correlated (corr. = 0.92) with ABE 97O $\rho^\tau(\mu)$ measurement.				
255 Highly correlated (corr. = 0.949) with AMMAR 97B $\rho^\tau(\mu)$ value.				
256 ACKERSTAFF 99D result is dominated by a constraint on η^τ from the OPAL measurements of the τ lifetime and $B(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)$ assuming lepton universality for the total coupling strength.				

 $(\delta\xi)^\tau(e \text{ or } \mu)$ PARAMETER $(V-A)$ theory predicts $(\delta\xi) = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.746±0.021 OUR FIT				
0.744±0.022 OUR AVERAGE				
0.776±0.045±0.024	81k	HEISTER	01E ALEP	1991–1995 LEP runs
0.779±0.070±0.028	36k	ABREU	00L DLPH	1992–1995 runs
0.65 ±0.14 ±0.07	46k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.70 ±0.11	54k	ACCIARRI	98R L3	1991–1995 LEP runs
0.63 ±0.09	257	ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5$ –10.6 GeV
0.88 ±0.27 ±0.04	258	ABE	97O SLD	1993–1995 SLC runs
0.745±0.026±0.009	55k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.81 ± 0.14 ± 0.06	18k	ACCIARRI	96H L3	Repl. by ACCIARRI 98R
0.65 ± 0.12		259 ALBRECHT	95C ARG	Repl. by ALBRECHT 98
0.88 ± 0.11 ± 0.07		BUSKULIC	95D ALEP	Repl. by HEISTER 01E

257 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

258 ABE 970 assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a $(\rho\xi)^\tau$ value of $0.87 \pm 0.27 \pm 0.04$.

259 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 95C uses events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(h^+ h^- h^+ \bar{\nu}_\tau)$ and their charged conjugates.

$(\delta\xi)^\tau(e)$ PARAMETER

(V-A) theory predicts $(\delta\xi) = 0.75$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.734 ± 0.028 OUR FIT

0.731 ± 0.029 OUR AVERAGE

0.778 ± 0.066 ± 0.024	44k	HEISTER	01E ALEP	1991–1995 LEP runs
0.85 ± 0.12 ± 0.04	17k	ABREU	00L DLPH	1992–1995 runs
0.72 ± 0.31 ± 0.14	25k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.56 ± 0.14 ± 0.06		260 ALBRECHT	98 ARG	$E_{\text{cm}}^{ee} = 9.5\text{--}10.6$ GeV
0.85 ± 0.43 ± 0.08		ABE	97O SLD	1993–1995 SLC runs
0.720 ± 0.032 ± 0.010	34k	ALEXANDER	97F CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.11 ± 0.17 ± 0.07		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
--------------------	--	----------	----------	----------------------

260 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

$(\delta\xi)^\tau(\mu)$ PARAMETER

(V-A) theory predicts $(\delta\xi) = 0.75$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.778 ± 0.037 OUR FIT

0.79 ± 0.04 OUR AVERAGE

0.786 ± 0.066 ± 0.028	46k	HEISTER	01E ALEP	1991–1995 LEP runs
0.86 ± 0.13 ± 0.04	22k	ABREU	00L DLPH	1992–1995 runs
0.63 ± 0.23 ± 0.05	27k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.73 ± 0.18 ± 0.10		261 ALBRECHT	98 ARG	$E_{\text{cm}}^{ee} = 9.5\text{--}10.6$ GeV
0.82 ± 0.32 ± 0.07		ABE	97O SLD	1993–1995 SLC runs
0.786 ± 0.041 ± 0.032	22k	ALEXANDER	97F CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.71 ± 0.14 ± 0.06		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
--------------------	--	----------	----------	----------------------

261 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

$\xi^T(\pi)$ PARAMETER

($V-A$) theory predicts $\xi^T(\pi) = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.993±0.022 OUR FIT

0.994±0.023 OUR AVERAGE

0.994±0.020±0.014	27k	HEISTER	01E ALEP	1991–1995 LEP runs
0.81 ±0.17 ±0.02		ABE	97O SLD	1993–1995 SLC runs
1.03 ±0.06 ±0.04	2.0k	COAN	97 CLEO	$E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.987±0.057±0.027		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
0.95 ±0.11 ±0.05		²⁶² BUSKULIC	94D ALEP	1990+1991 LEP run

²⁶² Superseded by BUSKULIC 95D.

$\xi^T(\rho)$ PARAMETER

($V-A$) theory predicts $\xi^T(\rho) = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.994±0.008 OUR FIT

0.994±0.009 OUR AVERAGE

0.987±0.012±0.011	59k	HEISTER	01E ALEP	1991–1995 LEP runs
0.99 ±0.12 ±0.04		ABE	97O SLD	1993–1995 SLC runs
0.995±0.010±0.003	66k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.022±0.028±0.030	1.7k	²⁶³ ALBRECHT	94E ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.045±0.058±0.032		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
1.03 ±0.11 ±0.05		²⁶⁴ BUSKULIC	94D ALEP	1990+1991 LEP run

²⁶³ ALBRECHT 94E measure the square of this quantity and use the sign determined by ALBRECHT 90I to obtain the quoted result.

²⁶⁴ Superseded by BUSKULIC 95D.

$\xi^T(a_1)$ PARAMETER

($V-A$) theory predicts $\xi^T(a_1) = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

1.001±0.027 OUR FIT

1.002±0.028 OUR AVERAGE

1.000±0.016±0.024	35k	²⁶⁵ HEISTER	01E ALEP	1991–1995 LEP runs
1.02 ±0.13 ±0.03	17.2k	ASNER	00 CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.29 ±0.26 ±0.11	7.4k	²⁶⁶ ACKERSTAFF	97R OPAL	1992–1994 LEP runs
0.85 ^{+0.15} _{-0.17} ±0.05		ALBRECHT	95C ARG	$E_{cm}^{ee} = 9.5-10.6$ GeV
1.25 ±0.23 ^{+0.15} _{-0.08}	7.5k	ALBRECHT	93C ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.08 ^{+0.46} _{-0.41} ^{+0.14} _{-0.25}	2.6k	²⁶⁷ AKERS	95P OPAL	Repl. by ACKER-STAFF 97R
0.937±0.116±0.064		BUSKULIC	95D ALEP	Repl. by HEISTER 01E

- 265 HEISTER 01E quote $1.000 \pm 0.016 \pm 0.013 \pm 0.020$ where the errors are statistical, systematic, and an uncertainty due to the final state model. We combine the systematic error and model uncertainty.
- 266 ACKERSTAFF 97R obtain this result with a model independent fit to the hadronic structure functions. Fitting with the model of Kuhn and Santamaria (ZPHY **C48**, 445 (1990)) gives $0.87 \pm 0.16 \pm 0.04$, and with the model of of Isgur *et al.* (PR **D39**,1357 (1989)) they obtain $1.20 \pm 0.21 \pm 0.14$.
- 267 AKERS 95P obtain this result with a model independent fit to the hadronic structure functions. Fitting with the model of Kuhn and Santamaria (ZPHY **C48**, 445 (1990)) gives $0.87 \pm 0.27^{+0.05}_{-0.06}$, and with the model of of Isgur *et al.* (PR **D39**,1357 (1989)) they obtain $1.10 \pm 0.31^{+0.13}_{-0.14}$.

ξ^T (all hadronic modes) PARAMETER

($V-A$) theory predicts $\xi^T = 1$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.995±0.007 OUR FIT				
0.997±0.007 OUR AVERAGE				
$0.992 \pm 0.007 \pm 0.008$	102k	268 HEISTER	01E ALEP	1991–1995 LEP runs
$0.997 \pm 0.027 \pm 0.011$	39k	269 ABREU	00L DLPH	1992–1995 runs
$1.02 \pm 0.13 \pm 0.03$	17.2k	270 ASNER	00 CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.032 ± 0.031	37k	271 ACCIARRI	98R L3	1991–1995 LEP runs
$0.93 \pm 0.10 \pm 0.04$		ABE	97O SLD	1993–1995 SLC runs
$1.29 \pm 0.26 \pm 0.11$	7.4k	272 ACKERSTAFF	97R OPAL	1992–1994 LEP runs
$0.995 \pm 0.010 \pm 0.003$	66k	273 ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
$1.03 \pm 0.06 \pm 0.04$	2.0k	274 COAN	97 CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.017 ± 0.039		275 ALBRECHT	95C ARG	$E_{cm}^{ee} = 9.5–10.6$ GeV
$1.25 \pm 0.23^{+0.15}_{-0.08}$	7.5k	276 ALBRECHT	93C ARG	$E_{cm}^{ee} = 9.4–10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.970 \pm 0.053 \pm 0.011$	14k	277 ACCIARRI	96H L3	Repl. by ACCIARRI 98R
$1.08^{+0.46}_{-0.41}^{+0.14}_{-0.25}$	2.6k	278 AKERS	95P OPAL	Repl. by ACKERSTAFF 97R
$1.006 \pm 0.032 \pm 0.019$		279 BUSKULIC	95D ALEP	Repl. by HEISTER 01E
$1.022 \pm 0.028 \pm 0.030$	1.7k	280 ALBRECHT	94E ARG	$E_{cm}^{ee} = 9.4–10.6$ GeV
$0.99 \pm 0.07 \pm 0.04$		281 BUSKULIC	94D ALEP	1990+1991 LEP run

- 268 HEISTER 01E quote $0.992 \pm 0.007 \pm 0.006 \pm 0.005$ where the errors are statistical, systematic, and an uncertainty due to the final state model. We combine the systematic error and model uncertainty. They use $\tau \rightarrow \pi \nu_\tau$, $\tau \rightarrow K \nu_\tau$, $\tau \rightarrow \rho \nu_\tau$, and $\tau \rightarrow a_1 \nu_\tau$ decays.
- 269 ABREU 00L use $\tau^- \rightarrow h^- \geq 0\pi^0 \nu_\tau$ decays.
- 270 ASNER 00 use $\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau$ decays.
- 271 ACCIARRI 98R use $\tau \rightarrow \pi \nu_\tau$, $\tau \rightarrow K \nu_\tau$, and $\tau \rightarrow \rho \nu_\tau$ decays.
- 272 ACKERSTAFF 97R use $\tau \rightarrow a_1 \nu_\tau$ decays.
- 273 ALEXANDER 97F use $\tau \rightarrow \rho \nu_\tau$ decays.
- 274 COAN 97 use $h^+ h^-$ energy correlations.
- 275 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E.
- 276 Uses $\tau \rightarrow a_1 \nu_\tau$ decays. Replaced by ALBRECHT 95C.
- 277 ACCIARRI 96H use $\tau \rightarrow \pi \nu_\tau$, $\tau \rightarrow K \nu_\tau$, and $\tau \rightarrow \rho \nu_\tau$ decays.
- 278 AKERS 95P use $\tau \rightarrow a_1 \nu_\tau$ decays.

- 279 BUSKULIC 95D use $\tau \rightarrow \pi\nu_\tau$, $\tau \rightarrow \rho\nu_\tau$, and $\tau \rightarrow a_1\nu_\tau$ decays.
 280 ALBRECHT 94E measure the square of this quantity and use the sign determined by ALBRECHT 90I to obtain the quoted result. Uses $\tau \rightarrow a_1\nu_\tau$ decays. Replaced by ALBRECHT 95C.
 281 BUSKULIC 94D use $\tau \rightarrow \pi\nu_\tau$ and $\tau \rightarrow \rho\nu_\tau$ decays. Superseded by BUSKULIC 95D.

τ REFERENCES

ABBIENDI	03	PL B551 35	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
INAMI	03	PL B551 16	K. Inami <i>et al.</i>	(BELLE Collab.)
CHEN	02C	PR D66 071101R	S. Chen <i>et al.</i>	(CLEO Collab.)
ABBIENDI	01J	EPJ C19 653	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	01M	EPJ C20 617	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	01F	PL B507 47	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACHARD	01D	PL B519 189	P. Achard <i>et al.</i>	(L3 Collab.)
ANASTASSOV	01	PRL 86 4467	A. Anastassov <i>et al.</i>	(CLEO Collab.)
HEISTER	01E	EPJ C22 217	A. Heister <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	00A	PL B492 23	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00C	EPJ C13 213	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00D	EPJ C13 197	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	00L	EPJ C16 229	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00B	PL B479 67	M. Acciarri <i>et al.</i>	(L3 Collab.)
AHMED	00	PR D61 071101R	S. Ahmed <i>et al.</i>	(CLEO Collab.)
ALBRECHT	00	PL B485 37	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BERGFELD	00	PRL 84 830	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BROWDER	00	PR D61 052004	T.E. Browder <i>et al.</i>	(CLEO Collab.)
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GONZALEZ-S...	00	NP B582 3	G.A. Gonzalez-Sprinberg <i>et al.</i>	
ABBIENDI	99H	PL B447 134	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	99X	EPJ C10 201	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	99D	EPJ C8 3	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	99E	EPJ C8 183	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	99K	EPJ C10 1	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)
BISHAI	99	PRL 82 281	M. Bishai <i>et al.</i>	(CLEO Collab.)
GODANG	99	PR D59 091303	R. Godang <i>et al.</i>	(CLEO Collab.)
RICHICHI	99	PR D60 112002	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ACCIARRI	98C	PL B426 207	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98E	PL B434 169	M. Aciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98R	PL B438 405	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	98M	EPJ C4 193	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98N	PL B431 188	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ALBRECHT	98	PL B431 179	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARATE	98	EPJ C1 65	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98E	EPJ C4 29	R. Barate <i>et al.</i>	(ALEPH Collab.)
BLISS	98	PR D57 5903	D.W. Bliss <i>et al.</i>	(CLEO Collab.)
ABE	97O	PRL 78 4691	K. Abe <i>et al.</i>	(SLD Collab.)
ACKERSTAFF	97J	PL B404 213	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97L	ZPHY C74 403	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97R	ZPHY C75 593	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ALEXANDER	97F	PR D56 5320	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	97B	PRL 78 4686	R. Ammar <i>et al.</i>	(CLEO Collab.)
ANASTASSOV	97	PR D55 2559	A. Anastassov <i>et al.</i>	(CLEO Collab.)
Also	98B	PR D58 119903 (erratum)	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ANDERSON	97	PRL 79 3814	S. Anderson <i>et al.</i>	(CLEO Collab.)
AVERY	97	PR D55 R1119	P. Avery <i>et al.</i>	(CLEO Collab.)
BARATE	97I	ZPHY C74 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	97R	PL B414 362	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERGFELD	97	PRL 79 2406	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BONVICINI	97	PRL 79 1221	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUSKULIC	97C	ZPHY C74 263	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
COAN	97	PR D55 7291	T.E. Coan <i>et al.</i>	(CLEO Collab.)
EDWARDS	97	PR D55 R3919	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
EDWARDS	97B	PR D56 R5297	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ESCRIBANO	97	PL B395 369	R. Escribano, E. Masso	(BARC, PARIT)
ABREU	96B	PL B365 448	P. Abreu <i>et al.</i>	(DELPHI Collab.)

ACCIARRI	96H	PL B377 313	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	96K	PL B389 187	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALAM	96	PRL 76 2637	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96E	PRPL 276 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	96D	PL B369 163	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	96E	PL B374 341	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	96S	PL B388 437	G. Alexander <i>et al.</i>	(OPAL Collab.)
BAI	96	PR D53 20	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALEST	96	PL B388 402	R. Balest <i>et al.</i>	(CLEO Collab.)
BARTELT	96	PRL 76 4119	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
BUSKULIC	96	ZPHY C70 579	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96C	ZPHY C70 561	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
COAN	96	PR D53 6037	T.E. Coan <i>et al.</i>	(CLEO Collab.)
ABE	95Y	PR D52 4828	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	95T	PL B357 715	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95U	PL B359 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	95	PL B345 93	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	95F	PL B352 487	M. Acciarri <i>et al.</i>	(L3 Collab.)
AKERS	95F	ZPHY C66 31	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95I	ZPHY C66 543	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95P	ZPHY C67 45	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95Y	ZPHY C68 555	R. Akers <i>et al.</i>	(OPAL Collab.)
ALBRECHT	95	PL B341 441	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	95C	PL B349 576	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	95G	ZPHY C68 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	95H	ZPHY C68 215	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	95C	PRL 75 3809	R. Balest <i>et al.</i>	(CLEO Collab.)
BERNABEU	95	NP B436 474	J. Bernabeu <i>et al.</i>	
BUSKULIC	95C	PL B346 371	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	95D	PL B346 379	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
Also	95P	PL B363 265 erratum	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	94K	PL B334 435	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	94E	PL B328 207	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	94G	PL B339 278	R. Akers <i>et al.</i>	(OPAL Collab.)
ALBRECHT	94E	PL B337 383	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ARTUSO	94	PRL 72 3762	M. Artuso <i>et al.</i>	(CLEO Collab.)
BARTELT	94	PRL 73 1890	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
BATTLE	94	PRL 73 1079	M. Battle <i>et al.</i>	(CLEO Collab.)
BAUER	94	PR D50 R13	D.A. Bauer <i>et al.</i>	(TPC/2gamma Collab.)
BUSKULIC	94D	PL B321 168	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	94E	PL B332 209	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	94F	PL B332 219	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBAUT	94B	PRL 73 934	D. Gibaut <i>et al.</i>	(CLEO Collab.)
ADRIANI	93M	PRPL 236 1	O. Adriani <i>et al.</i>	(L3 Collab.)
ALBRECHT	93C	ZPHY C58 61	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93G	PL B316 608	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	93	PR D47 R3671	R. Balest <i>et al.</i>	(CLEO Collab.)
BEAN	93	PRL 70 138	A. Bean <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	93	PRL 71 1791	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ESCRIBANO	93	PL B301 419	R. Escribano, E. Masso	(BARC)
PROCARIO	93	PRL 70 1207	M. Procaro <i>et al.</i>	(CLEO Collab.)
ABREU	92N	ZPHY C55 555	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	92F	PL B281 405	D.P. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	92H	PL B288 373	P.D. Acton <i>et al.</i>	(OPAL Collab.)
AKERIB	92	PRL 69 3610	D.S. Akerib <i>et al.</i>	(CLEO Collab.)
Also	93B	PRL 71 3395 (erratum)	D.S. Akerib <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92D	ZPHY C53 367	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92K	ZPHY C55 179	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92M	PL B292 221	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92Q	ZPHY C56 339	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMMAR	92	PR D45 3976	R. Ammar <i>et al.</i>	(CLEO Collab.)
ARTUSO	92	PRL 69 3278	M. Artuso <i>et al.</i>	(CLEO Collab.)
BAI	92	PRL 69 3021	J.Z. Bai <i>et al.</i>	(BES Collab.)
BATTLE	92	PL B291 488	M. Battle <i>et al.</i>	(CLEO Collab.)
BUSKULIC	92J	PL B297 459	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
DECAMP	92C	ZPHY C54 211	D. Decamp <i>et al.</i>	(ALEPH Collab.)
ADEVA	91F	PL B265 451	B. Adeva <i>et al.</i>	(L3 Collab.)
ALBRECHT	91D	PL B260 259	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	91D	PL B266 201	G. Alexander <i>et al.</i>	(OPAL Collab.)
ANTREASYAN	91	PL B259 216	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
GRIFOLS	91	PL B255 611	J.A. Grifols, A. Mendez	(BARC)

SAMUEL	91B	PRL 67 668	M.A. Samuel, G.W. Li, R. Mendel	(OKSU, WONT)
Also	92B	PRL 69 995	M.A. Samuel, G.W. Li, R. Mendel	(OKSU, WONT)
Erratum.				
ABACHI	90	PR D41 1414	S. Abachi <i>et al.</i>	(HRS Collab.)
ALBRECHT	90E	PL B246 278	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90I	PL B250 164	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BEHREND	90	ZPHY C46 537	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BOWCOCK	90	PR D41 805	T.J.V. Bowcock <i>et al.</i>	(CLEO Collab.)
DELAGUILA	90	PL B252 116	F. del Aguila, M. Sher	(BARC, WILL)
GOLDBERG	90	PL B251 223	M. Goldberg <i>et al.</i>	(CLEO Collab.)
WU	90	PR D41 2339	D.Y. Wu <i>et al.</i>	(Mark II Collab.)
ABACHI	89B	PR D40 902	S. Abachi <i>et al.</i>	(HRS Collab.)
BEHREND	89B	PL B222 163	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
JANSSEN	89	PL B228 273	H. Janssen <i>et al.</i>	(Crystal Ball Collab.)
KLEINWORT	89	ZPHY C42 7	C. Kleinwort <i>et al.</i>	(JADE Collab.)
ADEVA	88	PR D38 2665	B. Adeva <i>et al.</i>	(Mark-J Collab.)
ALBRECHT	88B	PL B202 149	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88L	ZPHY C41 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88M	ZPHY C41 405	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMIDEI	88	PR D37 1750	D. Amidei <i>et al.</i>	(Mark II Collab.)
BEHREND	88	PL B200 226	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BRAUNSCH...	88C	ZPHY C39 331	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
KEH	88	PL B212 123	S. Keh <i>et al.</i>	(Crystal Ball Collab.)
TSCHIRHART	88	PL B205 407	R. Tschirhart <i>et al.</i>	(HRS Collab.)
ABACHI	87B	PL B197 291	S. Abachi <i>et al.</i>	(HRS Collab.)
ABACHI	87C	PRL 59 2519	S. Abachi <i>et al.</i>	(HRS Collab.)
ADLER	87B	PRL 59 1527	J. Adler <i>et al.</i>	(Mark III Collab.)
AIHARA	87B	PR D35 1553	H. Aihara <i>et al.</i>	(TPC Collab.)
AIHARA	87C	PRL 59 751	H. Aihara <i>et al.</i>	(TPC Collab.)
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87P	PL B199 580	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BAND	87	PL B198 297	H.R. Band <i>et al.</i>	(MAC Collab.)
BAND	87B	PRL 59 415	H.R. Band <i>et al.</i>	(MAC Collab.)
BARINGER	87	PRL 59 1993	P. Baringer <i>et al.</i>	(CLEO Collab.)
BEBEK	87C	PR D36 690	C. Bebek <i>et al.</i>	(CLEO Collab.)
BURCHAT	87	PR D35 27	P.R. Burchat <i>et al.</i>	(Mark II Collab.)
BYLSMA	87	PR D35 2269	B.G. Bylsma <i>et al.</i>	(HRS Collab.)
COFFMAN	87	PR D36 2185	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
DERRICK	87	PL B189 260	M. Derrick <i>et al.</i>	(HRS Collab.)
FORD	87	PR D35 408	W.T. Ford <i>et al.</i>	(MAC Collab.)
FORD	87B	PR D36 1971	W.T. Ford <i>et al.</i>	(MAC Collab.)
GAN	87	PRL 59 411	K.K. Gan <i>et al.</i>	(Mark II Collab.)
GAN	87B	PL B197 561	K.K. Gan <i>et al.</i>	(Mark II Collab.)
AIHARA	86E	PRL 57 1836	H. Aihara <i>et al.</i>	(TPC Collab.)
BARTEL	86D	PL B182 216	W. Bartel <i>et al.</i>	(JADE Collab.)
PDG	86	PL 170B	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
RUCKSTUHL	86	PRL 56 2132	W. Ruckstuhl <i>et al.</i>	(DELCO Collab.)
SCHMIDKE	86	PRL 57 527	W.B. Schmidke <i>et al.</i>	(Mark II Collab.)
YELTON	86	PRL 56 812	J.M. Yelton <i>et al.</i>	(Mark II Collab.)
ALTHOFF	85	ZPHY C26 521	M. Althoff <i>et al.</i>	(TASSO Collab.)
ASH	85B	PRL 55 2118	W.W. Ash <i>et al.</i>	(MAC Collab.)
BALTRUSAIT...	85	PRL 55 1842	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BARTEL	85F	PL 161B 188	W. Bartel <i>et al.</i>	(JADE Collab.)
BEHREND	85	PR D32 2468	S. Behrends <i>et al.</i>	(CLEO Collab.)
BELTRAMI	85	PRL 54 1775	I. Beltrami <i>et al.</i>	(HRS Collab.)
BERGER	85	ZPHY C28 1	C. Berger <i>et al.</i>	(PLUTO Collab.)
BURCHAT	85	PRL 54 2489	P.R. Burchat <i>et al.</i>	(Mark II Collab.)
FERNANDEZ	85	PRL 54 1624	E. Fernandez <i>et al.</i>	(MAC Collab.)
MILLS	85	PRL 54 624	G.B. Mills <i>et al.</i>	(DELCO Collab.)
AIHARA	84C	PR D30 2436	H. Aihara <i>et al.</i>	(TPC Collab.)
BEHREND	84	ZPHY C23 103	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
MILLS	84	PRL 52 1944	G.B. Mills <i>et al.</i>	(DELCO Collab.)
BEHREND	83C	PL 127B 270	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
SILVERMAN	83	PR D27 1196	D.J. Silverman, G.L. Shaw	(UCI)
BEHREND	82	PL 114B 282	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BLOCKER	82B	PRL 48 1586	C.A. Blocker <i>et al.</i>	(Mark II Collab.)
BLOCKER	82D	PL 109B 119	C.A. Blocker <i>et al.</i>	(Mark II Collab.)
FELDMAN	82	PRL 48 66	G.J. Feldman <i>et al.</i>	(Mark II Collab.)
HAYES	82	PR D25 2869	K.G. Hayes <i>et al.</i>	(Mark II Collab.)
BERGER	81B	PL 99B 489	C. Berger <i>et al.</i>	(PLUTO Collab.)
DORFAN	81	PRL 46 215	J.M. Dorfan <i>et al.</i>	(Mark II Collab.)

BRANDELIK	80	PL 92B 199	R. Brandelik <i>et al.</i>	(TASSO Collab.)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also	81	SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34	1471.	
BACINO	79B	PRL 42 749	W.J. Bacino <i>et al.</i>	(DELCO Collab.)
KIRKBY	79	SLAC-PUB-2419	J. Kirkby	(SLAC) J
		Batavia Lepton Photon Conference.		
BACINO	78B	PRL 41 13	W.J. Bacino <i>et al.</i>	(DELCO Collab.) J
Also	78	Tokyo Conf. 249	J. Kirz	(STON)
Also	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
BRANDELIK	78	PL 73B 109	R. Brandelik <i>et al.</i>	(DASP Collab.) J
FELDMAN	78	Tokyo Conf. 777	G.J. Feldman	(SLAC) J
JAROS	78	PRL 40 1120	J. Jaros <i>et al.</i>	(SLAC, LBL, NWES, HAWA)
PERL	75	PRL 35 1489	M.L. Perl <i>et al.</i>	(LBL, SLAC)

————— **OTHER RELATED PAPERS** —————

RAHAL-CAL...	98	IJMP A13 695	G. Rahal-Callot	(ETH)
GENTILE	96	PRPL 274 287	S. Gentile, M. Pohl	(ROMAI, ETH)
WEINSTEIN	93	ARNPS 43 457	A.J. Weinstein, R. Stroynowski	(CIT, SMU)
PERL	92	RPP 55 653	M.L. Perl	(SLAC)
PICH	90	MPL A5 1995	A. Pich	(VALE)
BARISH	88	PRPL 157 1	B.C. Barish, R. Stroynowski	(CIT)
GAN	88	IJMP A3 531	K.K. Gan, M.L. Perl	(SLAC)
HAYES	88	PR D38 3351	K.G. Hayes, M.L. Perl	(SLAC)
PERL	80	ARNPS 30 299	M.L. Perl	(SLAC)