

# $\Sigma(1840) P_{13}$

$$I(J^P) = 1(\frac{3}{2}^+) \text{ Status: } *$$

**OMITTED FROM SUMMARY TABLE**

For the time being, we list together here all resonance claims in the  $P_{13}$  wave between 1700 and 1900 MeV.

### $\Sigma(1840)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>\approx 1840</math> OUR ESTIMATE</b>			
1798 or 1802	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel
$1720 \pm 30$	<sup>2</sup> BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
$1925 \pm 200$	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$
$1840 \pm 10$	LANGBEIN	72	IPWA $\bar{K}N$ multichannel

### $\Sigma(1840)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
93 or 93	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel
$120 \pm 30$	<sup>2</sup> BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
$65^{+50}_{-20}$	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$
$120 \pm 10$	LANGBEIN	72	IPWA $\bar{K}N$ multichannel

### $\Sigma(1840)$ DECAY MODES

Mode
$\Gamma_1 \quad N\bar{K}$
$\Gamma_2 \quad \Lambda\pi$
$\Gamma_3 \quad \Sigma\pi$

### $\Sigma(1840)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on  $\Lambda$  and  $\Sigma$  Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0 or 0	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel
$0.37 \pm 0.13$	LANGBEIN	72	IPWA $\bar{K}N$ multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1840) \rightarrow \Lambda\pi$	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$+0.03$ or $+0.03$	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel
$+0.11 \pm 0.02$	<sup>2</sup> BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
$+0.06 \pm 0.04$	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$
$+0.122 \pm 0.078$	DEVENISH	74B	Fixed- $t$ dispersion rel.
$0.20 \pm 0.04$	LANGBEIN	72	IPWA $\bar{K}N$ multichannel

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1840) \rightarrow \Sigma\pi$				$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
−0.04 or −0.04	<sup>1</sup> MARTIN	77	DPWA	$\bar{K}N$ multichannel
0.15 ± 0.04	LANGBEIN	72	IPWA	$\bar{K}N$ multichannel

### $\Sigma(1840)$ FOOTNOTES

<sup>1</sup> The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

<sup>2</sup> From solution 1 of BAILLON 75; not present in solution 2.

### $\Sigma(1840)$ REFERENCES

MARTIN	77	NP B127 349	B.R. Martin, M.K. Pidcock, R.G. Moorhouse	(LOUC+) IJP
Also	77B	NP B126 266	B.R. Martin, M.K. Pidcock	(LOUC)
Also	77C	NP B126 285	B.R. Martin, M.K. Pidcock	(LOUC) IJP
BAILLON	75	NP B94 39	P.H. Baillon, P.J. Litchfield	(CERN, RHEL) IJP
VANHORN	75	NP B87 145	A.J. van Horn	(LBL) IJP
Also	75B	NP B87 157	A.J. van Horn	(LBL) IJP
DEVENISH	74B	NP B81 330	R.C.E. Devenish, C.D. Froggatt, B.R. Martin	(DESY+)
LANGBEIN	72	NP B47 477	W. Langbein, F. Wagner	(MPIM) IJP