$f_0(1710)$

$I^G(J^{PC}) = 0^+(0^{++})$

THE $f_0(1710)$

Updated April 2002 by M. Doser (CERN).

The $f_0(1710)$ is seen in the radiative decay $J/\psi(1S) \to \gamma f_0(1710)$; therefore $C = +1$. It decays into 2η and $K^0_S K^0_S$, which implies $I^G J^{PC} = 0^+(even)^{++}$. The spin of the $f_0(1710)$ has been controversial, but evidence for spin 0 has accumulated recently in all production modes.

An analysis of radiative $J/\psi(1S)$ decays at BES into $\pi^+\pi^-\pi^+\pi^-$ (BAI 00) clearly favors spin 0. Combined amplitude analyses of the K^+K^-, K_SK_S, and $\pi^+\pi^-$ systems produced in $J/\psi(1S)$ radiative decay by MARK III (CHEN 91 and more recently DUNWOODIE 97) find a large spin-0 component, as well as reproducing known parameters of the $f_2(1270)$ and $f_2'(1525)$. In addition, a recent reanalysis (BUGG 95) of the 4π channel from MARK III, allowing both $\rho\rho$ and two $\pi\pi$ S waves, also finds a 0^{++} assignment for the $f_0(1710)$. Earlier analyses of this final state (BISELLO 89B, BALTRUSAITIS 86B) found only pseudoscalar activity in the $f_0(1710)$ region, but considered only the process $J/\psi(1S) \to \gamma\rho\rho$. Similarly, earlier analyses of the K^+K^- system based on less statistics (BALTRUSAITIS 87, BAI 96) found a spin of 2 for the $f_0(1710)$.

A similar situation is present in central production, with earlier analyses favoring spin 2 over spin 0 (ARMSTRONG 89D). More recent analyses with greater statistics [BARBERIS 99 (K^+K^-,K_SK_S), BARBERIS 99B ($\pi^+\pi^-$), and FRENCH 99 (K^+K^-)], however, clearly indicate spin 0, and exclude spin 2. Generally, analyses preferring spin 2 concentrate on angular distributions in the $f_J(1710)$ region, and do not include possible interferences or distortion due to the nearby $f_2'(1525)$.

The $f_0(1710)$ is also observed in $K\overline{K}$ (FALVARD 88) in $J/\psi(1S) \to \omega K\overline{K}$ and $J/\psi(1S) \to \phi K\overline{K}$, but with no spin-parity analysis, as well as in $\eta\eta$ in radiative $J/\psi(1S)$ decays (EDWARDS 82). It is also clearly seen in 300-GeV/c pp central production in both K^+K^- and $K^0_SK^0_S$ (ARMSTRONG 89D). Mass and width are determined via a fit to non-interfering Breit-Wigners over a polynomial background, which leads to large systematic errors for the width. ARMSTRONG 93C also sees a broad peak in $\eta\eta$ at 1747 MeV, which may be the $f_0(1710)$.

This resonance is not observed in the hypercharge-exchange reactions $K^-p \to K^0_SK^0_S\Lambda$ (ASTON 88D) and $K^-p \to K^0_SK^0_SY^*$ (BOLONKIN 86); these non-observations are explained by a spin of 0 (LINDENBAUM 92). It is not observed in $\overline{p}p$ interactions, neither via its $\pi\pi$ nor its $\eta\eta$ decay (AMSLER 02). A possible observation in $\gamma\gamma$ collisions leading to K_SK_S (BRACCINI 99, but no spin determination), and a non-observation in $\gamma\gamma \to \pi^+\pi^-$ (BARATE 00E), are consistent with a large $\overline{s}s$ component.

References

References may be found at the end of the $f_0(1710)$ Listing.

$f_0(1710)$ MASS

VALUE (MeV)		*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
1714± 5	**OUR AVERAGE**					
1740± 4	$^{+10}_{-25}$		[1] BAI	03G	BES	$J/\psi \to \gamma K\overline{K}$
1740^{+30}_{-25}			[1] BAI	00A	BES	$J/\psi \to \gamma(\pi^+\pi^-\pi^+\pi^-)$
1698±18			[2] BARBERIS	00E		450 $pp \to p_f\eta\eta p_s$
1710±12	±11		[3] BARBERIS	99D	OMEG	450 $pp \to K^+K^-$, $\pi^+\pi^-$
1710±25			[4] FRENCH	99		300 $pp \to p_f(K^+K^-)p_s$
1707±10			[5] AUGUSTIN	88	DM2	$J/\psi \to \gamma K^+K^-$, $K^0_SK^0_S$
1698±15			[5] AUGUSTIN	87	DM2	$J/\psi \to \gamma\pi^+\pi^-$
1720±10	±10		[6] BALTRUSAIT...	87	MRK3	$J/\psi \to \gamma K^+K^-$
1742±15			[5] WILLIAMS	84	MPSF	200 $\pi^-N \to 2K^0_S$X
1670±50			BLOOM	83	CBAL	$J/\psi \to \gamma 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1726± 7	74	[7]	CHEKANOV	04	ZEUS	$ep \to K^0_S K^0_S X$
1732+15		[8]	ANISOVICH	03	RVUE	
1682±16			TIKHOMIROV	03	SPEC	40.0 π^- C → $K^0_S K^0_S K^0_L$ X
1670±26	3651	[1,9]	NICHITIU	02	OBLX	
1767±14	221	[10]	ACCIARRI	01H	L3	$\gamma\gamma \to K^0_S K^0_S$, E^{ee}_{cm}= 91, 183–209 GeV
1770±12		[11,12]	ANISOVICH	99B	SPEC	0.6–1.2 $p\overline{p} \to \eta\eta\pi^0$
1730±15		[1]	BARBERIS	99	OMEG	450 $pp \to p_s p_f K^+ K^-$
1750±20		[1]	BARBERIS	99B	OMEG	450 $pp \to p_s p_f \pi^+ \pi^-$
1750±30		[13]	ANISOVICH	98B	RVUE	Compilation
1720±39			BAI	98H	BES	$J/\psi \to \gamma\pi^0\pi^0$
1775± 1.5	57	[14]	BARKOV	98		$\pi^- p \to K^0_S K^0_S n$
1690±11		[15]	ABREU	96C	DLPH	$Z^0 \to K^+K^-$ + X
1696± 5 $^{+9}_{-34}$		[6]	BAI	96C	BES	$J/\psi \to \gamma K^+K^-$
1781± 8 $^{+10}_{-31}$		[1]	BAI	96C	BES	$J/\psi \to \gamma K^+K^-$
1768±14			BALOSHIN	95	SPEC	40 π^- C → $K^0_S K^0_S$ X
1750±15		[16]	BUGG	95	MRK3	$J/\psi \to \gamma\pi^+\pi^-\pi^+\pi^-$
1620±16		[6]	BUGG	95	MRK3	$J/\psi \to \gamma\pi^+\pi^-\pi^+\pi^-$
1748±10		[5]	ARMSTRONG	93C	E760	$\overline{p}p \to \pi^0\eta\eta \to 6\gamma$
~ 1750			BREAKSTONE	93	SFM	$pp \to pp\pi^+\pi^-\pi^+\pi^-$
1744±15		[17]	ALDE	92D	GAM2	38 $\pi^- p \to \eta\eta n$
1713±10		[18]	ARMSTRONG	89D	OMEG	300 $pp \to pp K^+K^-$
1706±10		[18]	ARMSTRONG	89D	OMEG	300 $pp \to pp K^0_S K^0_S$
1700±15		[6]	BOLONKIN	88	SPEC	40 $\pi^- p \to K^0_S K^0_S n$
1720±60		[1]	BOLONKIN	88	SPEC	40 $\pi^- p \to K^0_S K^0_S n$
1638±10		[19]	FALVARD	88	DM2	$J/\psi \to \phi K^+K^-$, $K^0_S K^0_S$
1690± 4		[20]	FALVARD	88	DM2	$J/\psi \to \phi K^+K^-$, $K^0_S K^0_S$
1755± 8		[21]	ALDE	86C	GAM2	38 $\pi^- p \to n2\eta$
1730 $^{+2}_{-10}$		[22]	LONGACRE	86	RVUE	22 $\pi^- p \to n2K^0_S$
1650±50			BURKE	82	MRK2	$J/\psi \to \gamma 2\rho$
1640±50		[23,24]	EDWARDS	82D	CBAL	$J/\psi \to \gamma 2\eta$
1730±10 ±20		[25]	ETKIN	82C	MPS	23 $\pi^- p \to n2K^0_S$

[1] $J^P = 0^+$.
[2] T-matrix pole.
[3] Supersedes BARBERIS 99 and BARBERIS 99B.
[4] $J^P = 0^+$, supersedes by ARMSTRONG 89D.
[5] No J^{PC} determination.
[6] $J^P = 2^+$.
[7] Systematic errors not estimated.

[8] K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \to \pi^0\pi^0 n$, $\pi^- p \to K\overline{K} n$, $\pi^+\pi^- \to \pi^+\pi^-$, $\overline{p}p \to \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$, $\pi^+\pi^-\pi^0$, $K^+K^-\pi^0$, $K^0_S K^0_S \pi^0$, $K^+ K^0_S \pi^-$ at rest, $\overline{p}n \to \pi^-\pi^-\pi^+$, $K^0_S K^-\pi^0$, $K^0_S K^0_S \pi^-$ at rest.

[9] Decaying to $f_0(1370)\pi\pi$.

[10] Spin 2 dominant, isospin not determined, could also be I=1.

[11] $J^P = 0^+$.

[12] Not seen by AMSLER 02.

[13] T-matrix pole, assuming $J^P = 0^+$

[14] No J^{PC} determination.

[15] No J^{PC} determination, width not determined.

[16] From a fit to the 0^+ partial wave.

[17] ALDE 92D combines all the GAMS-2000 data.

[18] $J^P = 2^+$, superseded by FRENCH 99.

[19] From an analysis ignoring interference with $f_2'(1525)$.

[20] From an analysis including interference with $f_2'(1525)$.

[21] Superseded by ALDE 92D.

[22] Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

[23] $J^P = 2^+$ preferred.

[24] From fit neglecting nearby $f_2'(1525)$. Replaced by BLOOM 83.

[25] Superseded by LONGACRE 86.

$f_0(1710)$ WIDTH

VALUE (MeV)	*CL%*	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
140 ± 10 OUR AVERAGE			Error includes scale factor of 1.2.			
$166 \pm 5 \pm 15 \atop - 8 - 10$			[26] BAI	03G	BES	$J/\psi \to \gamma K\overline{K}$
$120 \pm 50 \atop - 40$			[26] BAI	00A	BES	$J/\psi \to \gamma(\pi^+\pi^-\pi^+\pi^-)$
120 ± 26			[27] BARBERIS	00E		450 $pp \to p_f \eta\eta p_s$
126 ± 16 ±18			[28] BARBERIS	99D	OMEG	450 $pp \to K^+K^-, \pi^+\pi^-$
105 ± 34			[29] FRENCH	99		300 $pp \to p_f(K^+K^-)p_s$
166.4± 33.2			[30] AUGUSTIN	88	DM2	$J/\psi \to \gamma K^+K^-, K^0_S K^0_S$
136 ± 28			[30] AUGUSTIN	87	DM2	$J/\psi \to \gamma\pi^+\pi^-$
130 ± 20			[31] BALTRUSAIT..	87	MRK3	$J/\psi \to \gamma K^+K^-$
57 ± 38			[5] WILLIAMS	84	MPSF	200 $\pi^- N \to 2K^0_S X$
160 ± 80			BLOOM	83	CBAL	$J/\psi \to \gamma 2\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$38 \pm 20 \atop - 14$		74	[32] CHEKANOV	04	ZEUS	$ep \to K^0_S K^0_S X$
144 ± 30			[33,34] ANISOVICH	03	RVUE	
$320 \pm 50 \atop - 20$			[34,35] ANISOVICH	03	RVUE	
102 ± 26			TIKHOMIROV	03	SPEC	40.0 $\pi^- C \to K^0_S K^0_S K^0_L X$

267	± 44		3651	[26,36] NICHITIU	02	OBLX	
187	± 60		221	[37] ACCIARRI	01H	L3	$\gamma\gamma \to K^0_S K^0_S$, E^{ee}_{cm}= 91, 183–209 GeV
220	± 40			[38,39] ANISOVICH	99B	SPEC	0.6–1.2 $p\overline{p} \to \eta\eta\pi^0$
100	± 25			[26] BARBERIS	99	OMEG	450 $pp \to p_s p_f K^+ K^-$
160	± 30			[26] BARBERIS	99B	OMEG	450 $pp \to p_s p_f \pi^+ \pi^-$
250	±140			[40] ANISOVICH	98B	RVUE	Compilation
30	± 7		57	[41] BARKOV	98		$\pi^- p \to K^0_S K^0_S n$
103	± 18	$^{+30}_{-11}$		[31] BAI	96C	BES	$J/\psi \to \gamma K^+ K^-$
85	± 24	$^{+22}_{-19}$		[26] BAI	96C	BES	$J/\psi \to \gamma K^+ K^-$
56	± 19			BALOSHIN	95	SPEC	40 π^- C $\to K^0_S K^0_S$ X
160	± 40			[42] BUGG	95	MRK3	$J/\psi \to \gamma\pi^+\pi^-\pi^+\pi^-$
160	$^{+60}_{-20}$			[31] BUGG	95	MRK3	$J/\psi \to \gamma\pi^+\pi^-\pi^+\pi^-$
264	± 25			[30] ARMSTRONG	93C	E760	$\overline{p}p \to \pi^0\eta\eta \to 6\gamma$
200	to 300			BREAKSTONE	93	SFM	$pp \to pp\pi^+\pi^-\pi^+\pi^-$
< 80		90		[43] ALDE	92D	GAM2	38 $\pi^- p \to \eta\eta N^*$
181	± 30			[44] ARMSTRONG	89D	OMEG	300 $pp \to pp K^+ K^-$
104	± 30			[44] ARMSTRONG	89D	OMEG	300 $pp \to pp K^0_S K^0_S$
30	± 20			[31] BOLONKIN	88	SPEC	40 $\pi^- p \to K^0_S K^0_S n$
350	±150			[26] BOLONKIN	88	SPEC	40 $\pi^- p \to K^0_S K^0_S n$
148	± 17			[45] FALVARD	88	DM2	$J/\psi \to \phi K^+ K^-$, $K^0_S K^0_S$
184	± 6			[46] FALVARD	88	DM2	$J/\psi \to \phi K^+ K^-$, $K^0_S K^0_S$
122	$^{+74}_{-15}$			[47] LONGACRE	86	RVUE	22 $\pi^- p \to n 2K^0_S$
200	±100			BURKE	82	MRK2	$J/\psi \to \gamma 2\rho$
220	$^{+100}_{-70}$			[48,49] EDWARDS	82D	CBAL	$J/\psi \to \gamma 2\eta$
200.0	$^{+156.0}_{-9.0}$			[50] ETKIN	82B	MPS	23 $\pi^- p \to n 2K^0_S$

[26] $J^P = 0^+$.
[27] T-matrix pole.
[28] Supersedes BARBERIS 99 and BARBERIS 99B.
[29] $J^P = 0^+$, supersedes by ARMSTRONG 89D.
[30] No J^{PC} determination.
[31] $J^P = 2^+$.
[32] Systematic errors not estimated.
[33] (Solution I)

[34] K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \to \pi^0\pi^0 n$, $\pi^- p \to K\overline{K} n$, $\pi^+\pi^- \to \pi^+\pi^-$, $\overline{p}p \to \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$, $\pi^+\pi^-\pi^0$, $K^+K^-\pi^0$, $K^0_S K^0_S \pi^0$, $K^+ K^0_S \pi^-$ at rest, $\overline{p}n \to \pi^-\pi^-\pi^+$, $K^0_S K^-\pi^0$, $K^0_S K^0_S \pi^-$ at rest.

[35] (Solution I)

[36] Decaying to $f_0(1370)\pi\pi$.

[37] Spin 2 dominant, isospin not determined, could also be I=1.

[38] $J^P = 0^+$.

[39] Not seen by AMSLER 02.

[40] T-matrix pole, assuming $J^P = 0^+$

[41] No J^{PC} determination.

[42] From a fit to the 0^+ partial wave.

[43] ALDE 92D combines all the GAMS-2000 data.

[44] $J^P = 2^+$, (0^+ excluded).

[45] From an analysis ignoring interference with $f'_2(1525)$.

[46] From an analysis including interference with $f'_2(1525)$.

[47] Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

[48] $J^P = 2^+$ preferred.

[49] From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.

[50] From an amplitude analysis of the $K^0_S K^0_S$ system, superseded by LONGACRE 86.

$f_0(1710)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)
Γ_1	$K\overline{K}$	seen
Γ_2	$\eta\eta$	seen
Γ_3	$\pi\pi$	seen
Γ_4	$\gamma\gamma$	

$f_0(1710)$ Γ(i)$\Gamma(\gamma\gamma)/\Gamma$(total)

$\Gamma(K\overline{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\quad\quad$ $\Gamma_1\Gamma_4/\Gamma$

VALUE (eV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<110	95	[52] BEHREND	89C	CELL	$\gamma\gamma \to K^0_S K^0_S$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$49 \pm 11 \pm 13$		[53] ACCIARRI	01H	L3	$\gamma\gamma \to K^0_S K^0_S$, E^{ee}_{cm}= 91, 183–209 GeV
<480	95	ALBRECHT	90G	ARG	$\gamma\gamma \to K^+K^-$
<280	95	[52] ALTHOFF	85B	TASS	$\gamma\gamma \to K\overline{K}\pi$

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\quad\quad$ $\Gamma_3\Gamma_4/\Gamma$

VALUE (keV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<0.82	95	[51] BARATE	00E	ALEP	$\gamma\gamma \to \pi^+\pi^-$

[51] Assuming spin 0.

[52] Assuming helicity 2.

[53] Spin 2 dominant, isospin not determined, could also be I=1.

$f_0(1710)$ BRANCHING RATIOS

$\Gamma(K\overline{K})/\Gamma_{total}$ Γ_1/Γ

VALUE		DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.38^{+0.09}_{-0.19}$	[54,55]	LONGACRE	86	MPS	22 $\pi^- p \rightarrow n 2K^0_S$

$\Gamma(\eta\eta)/\Gamma_{total}$ Γ_2/Γ

VALUE		DOCUMENT ID		TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.18^{+0.03}_{-0.13}$	[54,55]	LONGACRE	86	RVUE

$\Gamma(\pi\pi)/\Gamma_{total}$ Γ_3/Γ

VALUE		DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen		AMSLER	02	CBAR	0.9 $\overline{p}p \rightarrow \pi^0\eta\eta$, $\pi^0\pi^0\pi^0$
$0.039^{+0.002}_{-0.024}$	[54,55]	LONGACRE	86	RVUE	

$\Gamma(\pi\pi)/\Gamma(K\overline{K})$ Γ_3/Γ_1

VALUE		DOCUMENT ID		TECN	COMMENT
0.2 ±0.024±0.036		BARBERIS	99D	OMEG	450 $pp \rightarrow K^+K^-$, $\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$5.8\ ^{+9.1}_{-5.5}$	[56]	ANISOVICH	02D	SPEC	Combined fit
0.39±0.14		ARMSTRONG	91	OMEG	300 $pp \rightarrow pp\pi\pi$, $pp K\overline{K}$

$\Gamma(\eta\eta)/\Gamma(K\overline{K})$ Γ_2/Γ_1

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
0.48±0.15		BARBERIS	00E		450 $pp \rightarrow p_f\eta\eta p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.46^{+0.70}_{-0.38}$		[56] ANISOVICH	02D	SPEC	Combined fit
<0.02	90	[57] PROKOSHKIN	91	GA24	300 $\pi^- p \rightarrow \pi^- p\eta\eta$

[54] From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2.
[55] Fit with constrained inelasticity.
[56] From a combined K-matrix analysis of Crystal Barrel (0. $p\overline{p} \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\overline{K} n$) data.
[57] Combining results of GAM4 with those of ARMSTRONG 89D.

$f_0(1710)$ REFERENCES

CHEKANOV	04	PL B578 33	S. Chekanov *et al.*	(ZEUS Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich *et al.*	
BAI	03G	PR D68 052003	J.Z. Bai *et al.*	(BES Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov *et al.*	
		Translated from YAF 66 860.		
AMSLER	02	EPJ C23 29	C. Amsler *et al.*	
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich *et al.*	
		Translated from YAF 65 1583.		
NICHITIU	02	PL B545 261	F. Nichitiu *et al.*	(OBELIX Collab.)
ACCIARRI	01H	PL B501 173	M. Acciarri *et al.*	(L3 Collab.)
BAI	00A	PL B472 207	J.Z. Bai *et al.*	(BES Collab.)
BARATE	00E	PL B472 189	R. Barate *et al.*	(ALEPH Collab.)
BARBERIS	00E	PL B479 59	D. Barberis *et al.*	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich *et al.*	
BARBERIS	99	PL B453 305	D. Barberis *et al.*	(Omega Expt.)
BARBERIS	99B	PL B453 316	D. Barberis *et al.*	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis *et al.*	(Omega Expt.)
FRENCH	99	PL B460 213	B. French *et al.*	(WA76 Collab.)
ANISOVICH	98B	UFN 41 419	V.V. Anisovich *et al.*	
BAI	98H	PRL 81 1179	J.Z. Bai *et al.*	(BES Collab.)
BARKOV	98	JEPTL 68 764	B.P. Barkov *et al.*	
ABREU	96C	PL B379 309	P. Abreu *et al.*	(DELPHI Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai *et al.*	(BES Collab.)
BALOSHIN	95	PAN 58 46	O.N. Baloshin *et al.*	(ITEP)
		Translated from YAF 58 50.		
BUGG	95	PL B353 378	D.V. Bugg *et al.*	(LOQM, PNPI, WASH)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong *et al.*	(FNAL, FERR, GENO+)
BREAKSTONE	93	ZPHY C58 251	A.M. Breakstone *et al.*	(IOWA, CERN, DORT+)
ALDE	92D	PL B284 457	D.M. Alde *et al.*	(GAM2 Collab.)
Also	91	SJNP 54 451	D.M. Alde *et al.*	(GAM2 Collab.)
		Translated from YAF 54 745.		
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong *et al.*	(ATHU, BARI, BIRM+)
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2, GAM4 Collab.)
		Translated from DANS 316 900.		
ALBRECHT	90G	ZPHY C48 183	H. Albrecht *et al.*	(ARGUS Collab.)
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
BEHREND	89C	ZPHY C43 91	H.J. Behrend *et al.*	(CELLO Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin *et al.*	(DM2 Collab.)
BOLONKIN	88	NP B309 426	B.V. Bolonkin *et al.*	(ITEP, SERP)
FALVARD	88	PR D38 2706	A. Falvard *et al.*	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin *et al.*	(LALO, CLER, FRAS+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis *et al.*	(Mark III Collab.)
ALDE	86C	PL B182 105	D.M. Alde *et al.*	(SERP, BELG, LANL, LAPP)
LONGACRE	86	PL B177 223	R.S. Longacre *et al.*	(BNL, BRAN, CUNY+)
ALTHOFF	85B	ZPHY C29 189	M. Althoff *et al.*	(TASSO Collab.)
WILLIAMS	84	PR D30 877	E.G.H. Williams *et al.*	(VAND, NDAM, TUFTS+)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
BURKE	82	PRL 49 632	D.L. Burke *et al.*	(LBL, SLAC)
EDWARDS	82D	PRL 48 458	C. Edwards *et al.*	(CIT, HARV, PRIN+)
ETKIN	82B	PR D25 1786	A. Etkin *et al.*	(BNL, CUNY, TUFTS, VAND)
ETKIN	82C	PR D25 2446	A. Etkin *et al.*	(BNL, CUNY, TUFTS, VAND)

OTHER RELATED PAPERS

ANISOVICH	03B	PAN 66 741	V.V. Anisovich, V.A. Nikonov, A.V. Sarantsev
		Translated from YAF 66 772.	
AMSLER	02B	PL B541 22	C. Amsler
JIN	02	PR D66 057505	H. Jin, X. Zhang
KLEEFELD	02	PR D66 034007	F. Kleefeld *et al.*
RUPP	02	PR D65 078501	G. Rupp, E. vanBeveren, M.D. Scadron
SHAKIN	02	PR D65 078502	C.M. Shakin, H. Wang
TESHIMA	02	JPG 28 1391	T. Teshima, I. Kitamura, N. Morisita
VOLKOV	02	PAN 65 1657	M.K. Volkov, V.L. Yudichev
		Translated from YAF 65 1701.	
LI	01B	EPJ C19 529	D.-M. Li, H. Yu, Q.-X. Shen
VOLKOV	01	PAN 64 2006	M.K. Volkov, V.L. Yudichev
		Translated from YAF 64 2091.	
ANISOVICH	99H	PL B467 289	A.V. Anisovich, V.V. Anisovich
GRYGOREV	99	PAN 62 470	V.K. Grygorev *et al.*
		Translated from YAF 62 513.	

PROKOSHKIN	99	PAN 62 356 Translated from YAF 62 396.	Yu.D. Prokoshkin	
ANISOVICH	97	PL B395 123	A.V. Anisovich, A.V. Sarantsev	(PNPI)
LINDENBAUM	92	PL B274 492	S.J. Lindenbaum, R.S. Longacre	(BNL)
BISELLO	89B	PR D39 701	G. Busetto *et al.*	(DM2 Collab.)
ASTON	88D	NP B301 525	D. Aston *et al.*	(SLAC, NAGO, CINC, INUS)
AKESSON	86	NP B264 154	T. Akesson *et al.*	(Axial Field Spec. Collab.)
ARMSTRONG	86B	PL 167B 133	T.A. Armstrong *et al.*	(ATHU, BARI, BIRM+)
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis *et al.*	(Mark III Collab.)
ALTHOFF	83	PL 121B 216	M. Althoff *et al.*	(TASSO Collab.)
BARNETT	83B	PL 120B 455	B. Barnett *et al.*	(JHU)
BARNES	82B	NP B198 380	T. Barnes, F.E. Close, S. Monaghan	(RHEL, OXFTP)
TANIMOTO	82	PL 116B 198	M. Tanimoto	(BIEL)