

LIGHT UNFLAVORED MESONS

($S = C = B = 0$)

For $I = 1$ (π, ρ, ω): $u\bar{d}, (u\bar{u}-d\bar{d})/\sqrt{2}, d\bar{u}$;
 for $I = 0$ ($\eta, \eta', h, h', \omega, \phi, f, f'$): $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

π^\pm

$$I^G(J^P) = 1^-(0^-)$$

Mass $m = 139.57018 \pm 0.00035$ MeV ($S = 1.2$)

Mean life $\tau = (2.6033 \pm 0.0005) \times 10^{-8}$ s ($S = 1.2$)

$$c\tau = 7.8045 \text{ m}$$

$\pi^\pm \rightarrow \ell^\pm \nu \gamma$ form factors [a]

$$F_V = 0.017 \pm 0.008$$

$$F_A = 0.0116 \pm 0.0016 \quad (S = 1.3)$$

$$R = 0.059^{+0.009}_{-0.008}$$

π^- modes are charge conjugates of the modes below.

For decay limits to particles which are not established, see the appropriate Search sections (Massive Neutrino Peak Search Test, A^0 (axion), and Other Light Boson (X^0) Searches, etc.).

π^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
$\mu^+ \nu_\mu$	[b] (99.98770 ± 0.00004) %		30
$\mu^+ \nu_\mu \gamma$	[c] (2.00 ± 0.25) × 10 ⁻⁴		30
$e^+ \nu_e$	[b] (1.230 ± 0.004) × 10 ⁻⁴		70
$e^+ \nu_e \gamma$	[c] (1.61 ± 0.23) × 10 ⁻⁷		70
$e^+ \nu_e \pi^0$	(1.025 ± 0.034) × 10 ⁻⁸		4
$e^+ \nu_e e^+ e^-$	(3.2 ± 0.5) × 10 ⁻⁹		70
$e^+ \nu_e \nu \bar{\nu}$	< 5	× 10 ⁻⁶ 90%	70

Lepton Family number (LF) or Lepton number (L) violating modes

$\mu^+ \bar{\nu}_e$	L	[d] < 1.5	× 10 ⁻³ 90%	30
$\mu^+ \nu_e$	LF	[d] < 8.0	× 10 ⁻³ 90%	30
$\mu^- e^+ e^+ \nu$	LF	< 1.6	× 10 ⁻⁶ 90%	30



$$I^G(J^{PC}) = 1^-(0^{-+})$$

Mass $m = 134.9766 \pm 0.0006$ MeV (S = 1.1)
 $m_{\pi^\pm} - m_{\pi^0} = 4.5936 \pm 0.0005$ MeV
 Mean life $\tau = (8.4 \pm 0.6) \times 10^{-17}$ s (S = 3.0)
 $c\tau = 25.1$ nm

For decay limits to particles which are not established, see the appropriate Search sections (A^0 (axion), and Other Light Boson (X^0) Searches, etc.).

π^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
2γ	(98.798 ± 0.032) %	S=1.1	67
$e^+ e^- \gamma$	(1.198 ± 0.032) %	S=1.1	67
γ positronium	$(1.82 \pm 0.29) \times 10^{-9}$		67
$e^+ e^+ e^- e^-$	$(3.14 \pm 0.30) \times 10^{-5}$		67
$e^+ e^-$	$(6.2 \pm 0.5) \times 10^{-8}$		67
4γ	< 2	$\times 10^{-8}$ CL=90%	67
$\nu \bar{\nu}$	$[e] < 8.3$	$\times 10^{-7}$ CL=90%	67
$\nu_e \bar{\nu}_e$	< 1.7	$\times 10^{-6}$ CL=90%	67
$\nu_\mu \bar{\nu}_\mu$	< 3.1	$\times 10^{-6}$ CL=90%	67
$\nu_\tau \bar{\nu}_\tau$	< 2.1	$\times 10^{-6}$ CL=90%	67
$\gamma \nu \bar{\nu}$	< 6	$\times 10^{-4}$ CL=90%	67

Charge conjugation (C) or Lepton Family number (LF) violating modes

3γ	C	< 3.1	$\times 10^{-8}$ CL=90%	67
$\mu^+ e^-$	LF	< 3.8	$\times 10^{-10}$ CL=90%	26
$\mu^- e^+$	LF	< 3.4	$\times 10^{-9}$ CL=90%	26
$\mu^+ e^- + \mu^- e^+$	LF	< 1.72	$\times 10^{-8}$ CL=90%	26



$$I^G(J^{PC}) = 0^+(0^{-+})$$

Mass $m = 547.75 \pm 0.12$ MeV [f] (S = 2.6)
 Full width $\Gamma = 1.29 \pm 0.07$ keV [g]

C-nonconserving decay parameters

$\pi^+ \pi^- \pi^0$ Left-right asymmetry = $(0.09 \pm 0.17) \times 10^{-2}$
 $\pi^+ \pi^- \pi^0$ Sextant asymmetry = $(0.18 \pm 0.16) \times 10^{-2}$
 $\pi^+ \pi^- \pi^0$ Quadrant asymmetry = $(-0.17 \pm 0.17) \times 10^{-2}$
 $\pi^+ \pi^- \gamma$ Left-right asymmetry = $(0.9 \pm 0.4) \times 10^{-2}$
 $\pi^+ \pi^- \gamma$ β (D-wave) = -0.02 ± 0.07 (S = 1.3)

Dalitz plot parameter

$$\pi^0 \pi^0 \pi^0 \quad \alpha = -0.031 \pm 0.004 \quad (S = 1.1)$$

η DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Neutral modes			
neutral modes	(72.0 \pm 0.5) %	S=1.3	–
2γ	[<i>g</i>] (39.43 \pm 0.26) %	S=1.2	274
$3\pi^0$	(32.51 \pm 0.29) %	S=1.2	179
$\pi^0 2\gamma$	(7.2 \pm 1.4) $\times 10^{-4}$		257
other neutral modes	< 2.8 %	CL=90%	–
Charged modes			
charged modes	(28.0 \pm 0.5) %	S=1.3	–
$\pi^+ \pi^- \pi^0$	(22.6 \pm 0.4) %	S=1.3	174
$\pi^+ \pi^- \gamma$	(4.68 \pm 0.11) %	S=1.2	236
$e^+ e^- \gamma$	(6.0 \pm 0.8) $\times 10^{-3}$	S=1.4	274
$\mu^+ \mu^- \gamma$	(3.1 \pm 0.4) $\times 10^{-4}$		253
$e^+ e^-$	< 7.7 $\times 10^{-5}$	CL=90%	274
$\mu^+ \mu^-$	(5.8 \pm 0.8) $\times 10^{-6}$		253
$e^+ e^- e^+ e^-$	< 6.9 $\times 10^{-5}$	CL=90%	274
$\pi^+ \pi^- e^+ e^-$	(4.0 $^{+14.0}_{-2.7}$) $\times 10^{-4}$	S=5.8	235
$\pi^+ \pi^- 2\gamma$	< 2.0 $\times 10^{-3}$		236
$\pi^+ \pi^- \pi^0 \gamma$	< 5 $\times 10^{-4}$	CL=90%	174
$\pi^0 \mu^+ \mu^- \gamma$	< 3 $\times 10^{-6}$	CL=90%	210
Charge conjugation (C), Parity (P), Charge conjugation \times Parity (CP), or Lepton Family number (LF) violating modes			
$\pi^+ \pi^-$	<i>P, CP</i> < 3.3	$\times 10^{-4}$	CL=90% 236
$\pi^0 \pi^0$	<i>P, CP</i> < 4.3	$\times 10^{-4}$	CL=90% 238
3γ	<i>C</i> < 5	$\times 10^{-4}$	CL=95% 274
$4\pi^0$	<i>P, CP</i> < 6.9	$\times 10^{-7}$	CL=90% 40
$\pi^0 e^+ e^-$	<i>C</i> [<i>h</i>] < 4	$\times 10^{-5}$	CL=90% 257
$\pi^0 \mu^+ \mu^-$	<i>C</i> [<i>h</i>] < 5	$\times 10^{-6}$	CL=90% 210
$\mu^+ e^- + \mu^- e^+$	<i>LF</i> < 6	$\times 10^{-6}$	CL=90% 264

$f_0(600)$ [i]
or σ

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass $m = (400\text{--}1200)$ MeV

Full width $\Gamma = (600\text{--}1000)$ MeV

$f_0(600)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi$	dominant	—
$\gamma\gamma$	seen	—

$\rho(770)$ [j]

$$I^G(J^{PC}) = 1^+(1^{--})$$

Mass $m = 775.8 \pm 0.5$ MeV

Full width $\Gamma = 150.3 \pm 1.6$ MeV

$\Gamma_{ee} = 7.02 \pm 0.11$ keV

$\rho(770)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\pi\pi$	~ 100	%	364
$\rho(770)^\pm$ decays			
$\pi^\pm\gamma$	$(4.5 \pm 0.5) \times 10^{-4}$	S=2.2	375
$\pi^\pm\eta$	$< 6 \times 10^{-3}$	CL=84%	153
$\pi^\pm\pi^+\pi^-\pi^0$	$< 2.0 \times 10^{-3}$	CL=84%	254
$\rho(770)^0$ decays			
$\pi^+\pi^-\gamma$	$(9.9 \pm 1.6) \times 10^{-3}$		362
$\pi^0\gamma$	$(6.0 \pm 1.3) \times 10^{-4}$	S=1.1	376
$\eta\gamma$	$(3.0 \pm 0.4) \times 10^{-4}$	S=1.4	195
$\pi^0\pi^0\gamma$	$(4.5 \pm 0.8) \times 10^{-5}$		364
$\mu^+\mu^-$	[k] $(4.55 \pm 0.28) \times 10^{-5}$		373
e^+e^-	[k] $(4.67 \pm 0.09) \times 10^{-5}$		388
$\pi^+\pi^-\pi^0$	$(1.01^{+0.54}_{-0.36} \pm 0.34) \times 10^{-4}$		323
$\pi^+\pi^-\pi^+\pi^-$	$(1.8 \pm 0.9) \times 10^{-5}$		251
$\pi^+\pi^-\pi^0\pi^0$	$< 4 \times 10^{-5}$	CL=90%	257

$\omega(782)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 782.59 \pm 0.11$ MeV (S = 1.7)

Full width $\Gamma = 8.49 \pm 0.08$ MeV

$\Gamma_{ee} = 0.60 \pm 0.02$ keV

$\omega(782)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\pi^+\pi^-\pi^0$	(89.1 \pm 0.7) %	S=1.1	327
$\pi^0\gamma$	(8.92 ^{+0.28} _{-0.24}) %	S=1.1	380
$\pi^+\pi^-$	(1.70 \pm 0.27) %	S=1.4	366
neutrals (excluding $\pi^0\gamma$)	(1.4 ^{+7.0} _{-0.9}) $\times 10^{-3}$		—
$\eta\gamma$	(4.9 \pm 0.5) $\times 10^{-4}$		200
$\pi^0e^+e^-$	(5.9 \pm 1.9) $\times 10^{-4}$		380
$\pi^0\mu^+\mu^-$	(9.6 \pm 2.3) $\times 10^{-5}$		349
e^+e^-	(7.14 \pm 0.13) $\times 10^{-5}$	S=1.1	391
$\pi^+\pi^-\pi^0\pi^0$	< 2 %	CL=90%	262
$\pi^+\pi^-\gamma$	< 3.6 $\times 10^{-3}$	CL=95%	366
$\pi^+\pi^-\pi^+\pi^-$	< 1 $\times 10^{-3}$	CL=90%	256
$\pi^0\pi^0\gamma$	(6.7 \pm 1.1) $\times 10^{-5}$		367
$\eta\pi^0\gamma$	< 3.3 $\times 10^{-5}$	CL=90%	162
$\mu^+\mu^-$	(9.0 \pm 3.1) $\times 10^{-5}$		377
3γ	< 1.9 $\times 10^{-4}$	CL=95%	391
Charge conjugation (C) violating modes			
$\eta\pi^0$	C < 1 $\times 10^{-3}$	CL=90%	162
$3\pi^0$	C < 3 $\times 10^{-4}$	CL=90%	330

$\eta'(958)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Mass $m = 957.78 \pm 0.14$ MeV

Full width $\Gamma = 0.202 \pm 0.016$ MeV (S = 1.3)

$\eta'(958)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\pi^+\pi^-\eta$	(44.3 \pm 1.5) %	S=1.2	232
$\rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$)	(29.5 \pm 1.0) %	S=1.2	165
$\pi^0\pi^0\eta$	(20.9 \pm 1.2) %	S=1.2	239
$\omega\gamma$	(3.03 \pm 0.31) %		159
$\gamma\gamma$	(2.12 \pm 0.14) %	S=1.3	479
$3\pi^0$	(1.56 \pm 0.26) $\times 10^{-3}$		430
$\mu^+\mu^-\gamma$	(1.04 \pm 0.26) $\times 10^{-4}$		467
$\pi^+\pi^-\pi^0$	< 5 %	CL=90%	428

$\pi^0 \rho^0$	< 4	%	CL=90%	110
$\pi^+ \pi^+ \pi^- \pi^-$	< 1	%	CL=90%	372
$\pi^+ \pi^+ \pi^- \pi^-$ neutrals	< 1	%	CL=95%	–
$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	< 1	%	CL=90%	298
6π	< 1	%	CL=90%	211
$\pi^+ \pi^- e^+ e^-$	< 6	$\times 10^{-3}$	CL=90%	458
$\gamma e^+ e^-$	< 9	$\times 10^{-4}$	CL=90%	479
$\pi^0 \gamma \gamma$	< 8	$\times 10^{-4}$	CL=90%	469
$4\pi^0$	< 5	$\times 10^{-4}$	CL=90%	380
$e^+ e^-$	< 2.1	$\times 10^{-7}$	CL=90%	479

**Charge conjugation (C), Parity (P),
Lepton family number (LF) violating modes**

$\pi^+ \pi^-$	<i>P, CP</i>	< 2	%	CL=90%	458
$\pi^0 \pi^0$	<i>P, CP</i>	< 9	$\times 10^{-4}$	CL=90%	459
$\pi^0 e^+ e^-$	<i>C</i>	[<i>h</i>] < 1.4	$\times 10^{-3}$	CL=90%	469
$\eta e^+ e^-$	<i>C</i>	[<i>h</i>] < 2.4	$\times 10^{-3}$	CL=90%	322
3γ	<i>C</i>	< 1.0	$\times 10^{-4}$	CL=90%	479
$\mu^+ \mu^- \pi^0$	<i>C</i>	[<i>h</i>] < 6.0	$\times 10^{-5}$	CL=90%	445
$\mu^+ \mu^- \eta$	<i>C</i>	[<i>h</i>] < 1.5	$\times 10^{-5}$	CL=90%	273
$e\mu$	<i>LF</i>	< 4.7	$\times 10^{-4}$	CL=90%	473

$f_0(980)$ [¹]

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass $m = 980 \pm 10$ MeV

Full width $\Gamma = 40$ to 100 MeV

$f_0(980)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi$	dominant	471
$K\bar{K}$	seen	†
$\gamma\gamma$	seen	490

$a_0(980)$ [¹]

$$I^G(J^{PC}) = 1^-(0^{++})$$

Mass $m = 984.7 \pm 1.2$ MeV ($S = 1.5$)

Full width $\Gamma = 50$ to 100 MeV

$a_0(980)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\eta\pi$	dominant	322
$K\bar{K}$	seen	†
$\gamma\gamma$	seen	492

$\phi(1020)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 1019.456 \pm 0.020$ MeV (S = 1.1)

Full width $\Gamma = 4.26 \pm 0.05$ MeV (S = 1.7)

$\phi(1020)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$K^+ K^-$	(49.1 \pm 0.6) %	S=1.2	127
$K_L^0 K_S^0$	(34.0 \pm 0.5) %	S=1.1	110
$\rho\pi + \pi^+\pi^-\pi^0$	(15.4 \pm 0.5) %	S=1.3	—
$\eta\gamma$	(1.295 \pm 0.025) %	S=1.1	363
$\pi^0\gamma$	(1.23 \pm 0.10) $\times 10^{-3}$		501
e^+e^-	(2.98 \pm 0.04) $\times 10^{-4}$	S=1.1	510
$\mu^+\mu^-$	(2.85 \pm 0.19) $\times 10^{-4}$		499
ηe^+e^-	(1.15 \pm 0.10) $\times 10^{-4}$		363
$\pi^+\pi^-$	(7.3 \pm 1.3) $\times 10^{-5}$		490
$\omega\pi^0$	(5.2 $\begin{smallmatrix} +1.3 \\ -1.1 \end{smallmatrix}$) $\times 10^{-5}$		172
$\omega\gamma$	< 5 %	CL=84%	209
$\rho\gamma$	< 1.2 $\times 10^{-5}$	CL=90%	215
$\pi^+\pi^-\gamma$	(4.1 \pm 1.3) $\times 10^{-5}$		490
$f_0(980)\gamma$	(4.40 \pm 0.21) $\times 10^{-4}$		39
$\pi^0\pi^0\gamma$	(1.09 \pm 0.06) $\times 10^{-4}$		492
$\pi^+\pi^-\pi^+\pi^-$	(3.9 $\begin{smallmatrix} +2.8 \\ -2.2 \end{smallmatrix}$) $\times 10^{-6}$		410
$\pi^+\pi^+\pi^-\pi^-\pi^0$	< 4.6 $\times 10^{-6}$	CL=90%	342
$\pi^0 e^+ e^-$	(1.12 \pm 0.28) $\times 10^{-5}$		501
$\pi^0\eta\gamma$	(8.3 \pm 0.5) $\times 10^{-5}$		346
$a_0(980)\gamma$	(7.6 \pm 0.6) $\times 10^{-5}$		34
$\eta'(958)\gamma$	(6.2 \pm 0.7) $\times 10^{-5}$	S=1.1	60
$\eta\pi^0\pi^0\gamma$	< 2 $\times 10^{-5}$	CL=90%	293
$\mu^+\mu^-\gamma$	(1.4 \pm 0.5) $\times 10^{-5}$		499
$\rho\gamma\gamma$	< 5 $\times 10^{-4}$	CL=90%	215
$\eta\pi^+\pi^-$	< 1.8 $\times 10^{-5}$	CL=90%	288
$\eta\mu^+\mu^-$	< 9.4 $\times 10^{-6}$	CL=90%	321

$h_1(1170)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Mass $m = 1170 \pm 20$ MeV

Full width $\Gamma = 360 \pm 40$ MeV

$h_1(1170)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho\pi$	seen	307

$b_1(1235)$

$$I^G(J^{PC}) = 1^+(1^-)$$

Mass $m = 1229.5 \pm 3.2$ MeV (S = 1.6)

Full width $\Gamma = 142 \pm 9$ MeV (S = 1.2)

$b_1(1235)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\omega\pi$	dominant		348
[D/S amplitude ratio = 0.277 ± 0.027]			
$\pi^\pm\gamma$	$(1.6 \pm 0.4) \times 10^{-3}$		607
$\eta\rho$	seen		†
$\pi^+\pi^+\pi^-\pi^0$	< 50 %	84%	535
$(K\bar{K})^\pm\pi^0$	< 8 %	90%	248
$K_S^0 K_L^0 \pi^\pm$	< 6 %	90%	235
$K_S^0 K_S^0 \pi^\pm$	< 2 %	90%	235
$\phi\pi$	< 1.5 %	84%	147

$a_1(1260)$ ^[m]

$$I^G(J^{PC}) = 1^-(1^{++})$$

Mass $m = 1230 \pm 40$ MeV ^[n]

Full width $\Gamma = 250$ to 600 MeV

$a_1(1260)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$(\rho\pi)_{S\text{-wave}}$	seen	353
$(\rho\pi)_{D\text{-wave}}$	seen	353
$(\rho(1450)\pi)_{S\text{-wave}}$	seen	†
$(\rho(1450)\pi)_{D\text{-wave}}$	seen	†
$\sigma\pi$	seen	–
$f_0(980)\pi$	not seen	189
$f_0(1370)\pi$	seen	–
$f_2(1270)\pi$	seen	†
$K\bar{K}^*(892) + \text{c.c.}$	seen	†
$\pi\gamma$	seen	608

$f_2(1270)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 1275.4 \pm 1.2$ MeV

Full width $\Gamma = 185.1^{+3.5}_{-2.6}$ MeV (S = 1.5)

$f_2(1270)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\pi\pi$	$(84.8^{+2.5}_{-1.3})\%$	S=1.3	623
$\pi^+\pi^-\pi^0$	$(7.1^{+1.5}_{-2.7})\%$	S=1.3	563

$K\bar{K}$	(4.6 ± 0.4) %	S=2.7	404
$2\pi^+ 2\pi^-$	(2.8 ± 0.4) %	S=1.2	559
$\eta\eta$	(4.5 ± 1.0) × 10 ⁻³	S=2.4	327
$4\pi^0$	(3.0 ± 1.0) × 10 ⁻³		565
$\gamma\gamma$	(1.41 ± 0.13) × 10 ⁻⁵		638
$\eta\pi\pi$	< 8 × 10 ⁻³	CL=95%	478
$K^0 K^- \pi^+ + \text{c.c.}$	< 3.4 × 10 ⁻³	CL=95%	293
$e^+ e^-$	< 6 × 10 ⁻¹⁰	CL=90%	638

$f_1(1285)$

$$J^{PC} = 0^+(1^+)$$

Mass $m = 1281.8 \pm 0.6$ MeV (S = 1.6)

Full width $\Gamma = 24.1 \pm 1.1$ MeV (S = 1.3)

$f_1(1285)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
4π	(33.1 ^{+2.1} _{-1.8}) %	S=1.3	568
$\pi^0 \pi^0 \pi^+ \pi^-$	(22.0 ^{+1.4} _{-1.2}) %	S=1.3	566
$2\pi^+ 2\pi^-$	(11.0 ^{+0.7} _{-0.6}) %	S=1.3	563
$\rho^0 \pi^+ \pi^-$	(11.0 ^{+0.7} _{-0.6}) %	S=1.3	336
$\rho^0 \rho^0$	seen		†
$4\pi^0$	< 7 × 10 ⁻⁴	CL=90%	568
$\eta\pi\pi$	(52 ± 16) %		482
$a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$]	(36 ± 7) %		234
$\eta\pi\pi$ [excluding $a_0(980)\pi$]	(16 ± 7) %		482
$K\bar{K}\pi$	(9.0 ± 0.4) %	S=1.1	308
$K\bar{K}^*(892)$	not seen		†
$\gamma\rho^0$	(5.5 ± 1.3) %	S=2.8	406
$\phi\gamma$	(7.4 ± 2.6) × 10 ⁻⁴		236

$\eta(1295)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

Mass $m = 1294 \pm 4$ MeV (S = 1.6)

Full width $\Gamma = 55 \pm 5$ MeV

$\eta(1295)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\eta\pi^+\pi^-$	seen	487
$a_0(980)\pi$	seen	244
$\eta\pi^0\pi^0$	seen	490
$\eta(\pi\pi)_{S\text{-wave}}$	seen	—

$\pi(1300)$

$$I^G(J^{PC}) = 1^-(0^-+)$$

Mass $m = 1300 \pm 100$ MeV [η]

Full width $\Gamma = 200$ to 600 MeV

$\pi(1300)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho\pi$	seen	404
$\pi(\pi\pi)_{S\text{-wave}}$	seen	—

$a_2(1320)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

Mass $m = 1318.3 \pm 0.6$ MeV (S = 1.2)

Full width $\Gamma = 107 \pm 5$ MeV [η]

$a_2(1320)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\rho\pi$	(70.1 \pm 2.7) %	S=1.2	416
$\eta\pi$	(14.5 \pm 1.2) %		535
$\omega\pi\pi$	(10.6 \pm 3.2) %	S=1.3	366
$K\bar{K}$	(4.9 \pm 0.8) %		437
$\eta'(958)\pi$	(5.3 \pm 0.9) $\times 10^{-3}$		288
$\pi^\pm\gamma$	(2.68 \pm 0.31) $\times 10^{-3}$		652
$\gamma\gamma$	(9.4 \pm 0.7) $\times 10^{-6}$		659
$\pi^+\pi^-\pi^-$	< 8 %	CL=90%	621
e^+e^-	< 6 $\times 10^{-9}$	CL=90%	659

$f_0(1370)$ [1]

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass $m = 1200$ to 1500 MeV
Full width $\Gamma = 200$ to 500 MeV

$f_0(1370)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi$	seen	—
4π	seen	—
$4\pi^0$	seen	—
$2\pi^+2\pi^-$	seen	—
$\pi^+\pi^-2\pi^0$	seen	—
$\rho\rho$	dominant	—
$2(\pi\pi)_{S\text{-wave}}$	seen	—
$\pi(1300)\pi$	seen	—
$a_1(1260)\pi$	seen	—
$\eta\eta$	seen	—
$K\bar{K}$	seen	—
$\gamma\gamma$	seen	—
e^+e^-	not seen	—

$\pi_1(1400)$ [o]

$$I^G(J^{PC}) = 1^-(1^{-+})$$

Mass $m = 1376 \pm 17$ MeV
Full width $\Gamma = 300 \pm 40$ MeV

$\pi_1(1400)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\eta\pi^0$	seen	570
$\eta\pi^-$	seen	569

$\eta(1405)$ [p]
was $\eta(1440)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Mass $m = 1410.3 \pm 2.6$ MeV [n] ($S = 2.2$)
Full width $\Gamma = 51 \pm 4$ MeV [n] ($S = 2.2$)

$\eta(1405)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}\pi$	seen	425
$\eta\pi\pi$	seen	563
$a_0(980)\pi$	seen	342

$\eta(\pi\pi)$ S-wave	seen	—
$f_0(980)\eta$	seen	†
4π	seen	639
$K^*(892)K$	seen	127

$f_1(1420)$ [q]

$$I^G(J^{PC}) = 0^+(1^{++})$$

Mass $m = 1426.3 \pm 0.9$ MeV (S = 1.1)

Full width $\Gamma = 54.9 \pm 2.6$ MeV

$f_1(1420)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}\pi$	dominant	438
$K\bar{K}^*(892) + \text{c.c.}$	dominant	163
$\eta\pi\pi$	possibly seen	573
$\phi\gamma$	seen	349

$\omega(1420)$ [r]

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass m (1400–1450) MeV

Full width Γ (180–250) MeV

$\omega(1420)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho\pi$	dominant	488
$\omega\pi\pi$	seen	—
$b_1(1235)\pi$	seen	—
e^+e^-	seen	—

$a_0(1450)$ [l]

$$I^G(J^{PC}) = 1^-(0^{++})$$

Mass $m = 1474 \pm 19$ MeV

Full width $\Gamma = 265 \pm 13$ MeV

$a_0(1450)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\eta$	seen	627
$\pi\eta'(958)$	seen	410
$K\bar{K}$	seen	547
$\omega\pi\pi$	seen	484

$\rho(1450)$ [s]

$$J^{PC} = 1^{+}(1^{-}-)$$

Mass $m = 1465 \pm 25$ MeV [n]

Full width $\Gamma = 400 \pm 60$ MeV [n]

$\rho(1450)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\pi\pi$	seen		720
4π	seen		669
$\omega\pi$	<2.0 %	95%	512
e^+e^-	seen		732
$\eta\rho$	<4 %		310
$a_2(1320)\pi$	not seen		55
$\phi\pi$	<1 %		360
$K\bar{K}$	< 1.6×10^{-3}	95%	541
$\eta\gamma$	possibly seen		630

$\eta(1475)$ [ρ]
was $\eta(1440)$

$$J^{PC} = 0^{+}(0^{-}+)$$

Mass $m = 1476 \pm 4$ MeV (S = 1.4)

Full width $\Gamma = 87 \pm 9$ MeV (S = 1.6)

$\eta(1475)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}\pi$	dominant	477
$K\bar{K}^*(892)+$ c.c.	seen	245
$a_0(980)\pi$	seen	393
$\gamma\gamma$	seen	738

$f_0(1500)$ [o]

$$J^{PC} = 0^{+}(0^{++})$$

Mass $m = 1507 \pm 5$ MeV (S = 1.2)

Full width $\Gamma = 109 \pm 7$ MeV

$f_0(1500)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	p (MeV/c)
$\eta\eta'(958)$	(1.9 ± 0.8) %	1.7	34
$\eta\eta$	(5.1 ± 0.9) %	1.4	518
4π	(49.5 ± 3.3) %	1.2	692
$4\pi^0$	seen		692
$2\pi^+2\pi^-$	seen		688

$\pi\pi$	$(34.9 \pm 2.3) \%$	1.2	741
$\pi^+\pi^-$	seen		741
$2\pi^0$	seen		741
$K\bar{K}$	$(8.6 \pm 1.0) \%$	1.1	569
$\gamma\gamma$	not seen		754

$f'_2(1525)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 1525 \pm 5$ MeV [n]

Full width $\Gamma = 73^{+6}_{-5}$ MeV [n]

$f'_2(1525)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}$	$(88.8 \pm 3.1) \%$	581
$\eta\eta$	$(10.3 \pm 3.1) \%$	530
$\pi\pi$	$(8.2 \pm 1.5) \times 10^{-3}$	750
$\gamma\gamma$	$(1.11 \pm 0.14) \times 10^{-6}$	763

$\pi_1(1600)$ [o]

$$I^G(J^{PC}) = 1^-(1^{-+})$$

Mass $m = 1596^{+25}_{-14}$ MeV

Full width $\Gamma = 312^{+64}_{-24}$ MeV ($S = 1.1$)

$\pi_1(1600)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi\pi$	seen	769
$\rho^0\pi^-$	seen	600
$f_2(1270)\pi^-$	not seen	259
$\eta'(958)\pi^-$	seen	497

$\eta_2(1645)$

$$I^G(J^{PC}) = 0^+(2^{-+})$$

Mass $m = 1617 \pm 5$ MeV

Full width $\Gamma = 181 \pm 11$ MeV

$\eta_2(1645)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$a_2(1320)\pi$	seen	242
$K\bar{K}\pi$	seen	580
$K^*\bar{K}$	seen	404
$\eta\pi^+\pi^-$	seen	685
$a_0(980)\pi$	seen	496
$f_2(1270)\eta$	not seen	†

$\omega(1650)$ [^t]
was $\omega(1600)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 1670 \pm 30$ MeV

Full width $\Gamma = 315 \pm 35$ MeV

$\omega(1650)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho\pi$	seen	646
$\omega\pi\pi$	seen	617
$\omega\eta$	seen	500
e^+e^-	seen	835

$\omega_3(1670)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

Mass $m = 1667 \pm 4$ MeV

Full width $\Gamma = 168 \pm 10$ MeV [ⁿ]

$\omega_3(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho\pi$	seen	645
$\omega\pi\pi$	seen	615
$b_1(1235)\pi$	possibly seen	361

$\pi_2(1670)$

$$I^G(J^{PC}) = 1^-(2^{-+})$$

Mass $m = 1672.4 \pm 3.2$ MeV [ⁿ] (S = 1.4)

Full width $\Gamma = 259 \pm 9$ MeV [ⁿ] (S = 1.3)

$\pi_2(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
3π	(95.8±1.4) %		809
$f_2(1270)\pi$	(56.2±3.2) %		329
$\rho\pi$	(31 ±4) %		648
$\sigma\pi$	(10.9±3.4) %		—
$(\pi\pi)$ S-wave	(8.7±3.4) %		—
$K\bar{K}^*(892)+$ c.c.	(4.2±1.4) %		455
$\omega\rho$	(2.7±1.1) %		303
$\rho(1450)\pi$	< 3.6	$\times 10^{-3}$	97.7% 148
$b_1(1235)\pi$	< 1.9	$\times 10^{-3}$	97.7% 366

$\phi(1680)$

$$J^{PC} = 0^-(1^--)$$

Mass $m = 1680 \pm 20$ MeV [n]
 Full width $\Gamma = 150 \pm 50$ MeV [n]

$\phi(1680)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}^*(892) + \text{c.c.}$	dominant	462
$K_S^0 K \pi$	seen	621
$K\bar{K}$	seen	680
$e^+ e^-$	seen	840
$\omega \pi \pi$	not seen	623

$\rho_3(1690)$

$$J^{PC} = 1^+(3^--)$$

Mass $m = 1688.8 \pm 2.1$ MeV [n]
 Full width $\Gamma = 161 \pm 10$ MeV [n] (S = 1.5)

$\rho_3(1690)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	p (MeV/c)
4π	(71.1 \pm 1.9) %		790
$\pi^\pm \pi^+ \pi^- \pi^0$	(67 \pm 22) %		787
$\omega \pi$	(16 \pm 6) %		655
$\pi \pi$	(23.6 \pm 1.3) %		834
$K\bar{K} \pi$	(3.8 \pm 1.2) %		629
$K\bar{K}$	(1.58 \pm 0.26) %	1.2	685
$\eta \pi^+ \pi^-$	seen		727
$\rho(770)\eta$	seen		520
$\pi \pi \rho$	seen		633
Excluding 2ρ and $a_2(1320)\pi$.			
$a_2(1320)\pi$	seen		307
$\rho\rho$	seen		333

$\rho(1700)$ [s]

$$J^{PC} = 1^+(1^--)$$

Mass $m = 1720 \pm 20$ MeV [n] ($\eta\rho^0$ and $\pi^+\pi^-$ modes)
 Full width $\Gamma = 250 \pm 100$ MeV [n] ($\eta\rho^0$ and $\pi^+\pi^-$ modes)

$\rho(1700)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$2(\pi^+ \pi^-)$	large	803
$\rho \pi \pi$	dominant	653
$\rho^0 \pi^+ \pi^-$	large	650

$\rho^\pm \pi^\mp \pi^0$	large	651
$a_1(1260)\pi$	seen	404
$h_1(1170)\pi$	seen	447
$\pi(1300)\pi$	seen	349
$\rho\rho$	seen	371
$\pi^+\pi^-$	seen	849
$\pi\pi$	seen	849
$K\bar{K}^*(892) + \text{c.c.}$	seen	496
$\eta\rho$	seen	544
$a_2(1320)\pi$	not seen	334
$K\bar{K}$	seen	704
e^+e^-	seen	860
$\pi^0\omega$	seen	674

$f_0(1710)$ [*u*]

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass $m = 1714 \pm 5$ MeV

Full width $\Gamma = 140 \pm 10$ MeV ($S = 1.2$)

$f_0(1710)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}$	seen	701
$\eta\eta$	seen	659
$\pi\pi$	seen	846

$\pi(1800)$

$$I^G(J^{PC}) = 1^-(0^{-+})$$

Mass $m = 1812 \pm 14$ MeV ($S = 2.3$)

Full width $\Gamma = 207 \pm 13$ MeV

$\pi(1800)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi^+\pi^-\pi^-$	seen	879
$f_0(600)\pi^-$	seen	—
$f_0(980)\pi^-$	seen	631
$f_0(1370)\pi^-$	seen	—
$f_0(1500)\pi^-$	not seen	248
$\rho\pi^-$	not seen	732
$\eta\eta\pi^-$	seen	661
$a_0(980)\eta$	seen	469
$f_0(1500)\pi^-$	seen	248

$\eta\eta'(958)\pi^-$	seen	376
$K_0^*(1430)K^-$	seen	†
$K^*(892)K^-$	not seen	570

$\phi_3(1850)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

Mass $m = 1854 \pm 7$ MeV

Full width $\Gamma = 87^{+28}_{-23}$ MeV (S = 1.2)

$\phi_3(1850)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}$	seen	785
$K\bar{K}^*(892) + \text{c.c.}$	seen	602

$f_2(1950)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 1945 \pm 13$ MeV (S = 1.6)

Full width $\Gamma = 475 \pm 19$ MeV

$f_2(1950)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K^*(892)\bar{K}^*(892)$	seen	389
$\pi^+\pi^-$	seen	963
4π	seen	925
$\eta\eta$	seen	804
$K\bar{K}$	seen	838
$\gamma\gamma$	seen	973

$f_2(2010)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 2011^{+60}_{-80}$ MeV

Full width $\Gamma = 202 \pm 60$ MeV

$f_2(2010)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\phi\phi$	seen	†

$a_4(2040)$

$$I^G(J^{PC}) = 1^-(4^{++})$$

Mass $m = 2010 \pm 12$ MeV

Full width $\Gamma = 353 \pm 40$ MeV

$a_4(2040)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K \bar{K}$	seen	875
$\pi^+ \pi^- \pi^0$	seen	981
$\rho \pi$	seen	849
$f_2(1270) \pi$	seen	590
$\eta \pi^0$	seen	925
$\eta'(958) \pi$	seen	769

$f_4(2050)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

Mass $m = 2034 \pm 11$ MeV ($S = 1.6$)

Full width $\Gamma = 222 \pm 19$ MeV ($S = 1.8$)

$f_4(2050)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\omega \omega$	not seen	650
$\pi \pi$	$(17.0 \pm 1.5) \%$	1008
$K \bar{K}$	$(6.8^{+3.4}_{-1.8}) \times 10^{-3}$	889
$\eta \eta$	$(2.1 \pm 0.8) \times 10^{-3}$	857
$4\pi^0$	$< 1.2 \%$	972
$a_2(1320) \pi$	seen	579

$f_2(2300)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 2297 \pm 28$ MeV

Full width $\Gamma = 149 \pm 40$ MeV

$f_2(2300)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\phi \phi$	seen	529
$K \bar{K}$	seen	1037
$\gamma \gamma$	seen	1149

$f_2(2340)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 2339 \pm 60$ MeVFull width $\Gamma = 319^{+80}_{-70}$ MeV

$f_2(2340)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\phi\phi$	seen	573

NOTES

- [a] See the “Note on $\pi^\pm \rightarrow \ell^\pm \nu \gamma$ and $K^\pm \rightarrow \ell^\pm \nu \gamma$ Form Factors” in the π^\pm Particle Listings for definitions and details.
- [b] Measurements of $\Gamma(e^+ \nu_e)/\Gamma(\mu^+ \nu_\mu)$ always include decays with γ 's, and measurements of $\Gamma(e^+ \nu_e \gamma)$ and $\Gamma(\mu^+ \nu_\mu \gamma)$ never include low-energy γ 's. Therefore, since no clean separation is possible, we consider the modes with γ 's to be subreactions of the modes without them, and let $[\Gamma(e^+ \nu_e) + \Gamma(\mu^+ \nu_\mu)]/\Gamma_{\text{total}} = 100\%$.
- [c] See the π^\pm Particle Listings for the energy limits used in this measurement; low-energy γ 's are not included.
- [d] Derived from an analysis of neutrino-oscillation experiments.
- [e] Astrophysical and cosmological arguments give limits of order 10^{-13} ; see the π^0 Particle Listings.
- [f] Due to a new measurement in the average, this is 0.45 MeV larger than the mass we gave in our 2002 edition, 547.30 ± 0.12 MeV.
- [g] Due to removing an old measurement from the average, this is 0.11 keV larger than the width we gave in our 2002 edition, 1.18 ± 0.11 keV. See the $\Gamma(2\gamma)$ data block in the Data Listings.
- [h] C parity forbids this to occur as a single-photon process.
- [i] See the “Note on scalar mesons” in the $f_0(1370)$ Particle Listings . The interpretation of this entry as a particle is controversial.
- [j] See the “Note on $\rho(770)$ ” in the $\rho(770)$ Particle Listings .
- [k] The $\omega\rho$ interference is then due to $\omega\rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+ \mu^-) = \Gamma(\rho^0 \rightarrow e^+ e^-) \times 0.99785$.
- [l] See the “Note on scalar mesons” in the $f_0(1370)$ Particle Listings .
- [m] See the “Note on $a_1(1260)$ ” in the $a_1(1260)$ Particle Listings .
- [n] This is only an educated guess; the error given is larger than the error on the average of the published values. See the Particle Listings for details.

- [o] See the “Note on non- $q\bar{q}$ mesons” in the Particle Listings (see the index for the page number).
- [p] See the “Note on the $\eta(1405)$ ” in the $\eta(1405)$ Particle Listings.
- [q] See the “Note on the $f_1(1420)$ ” in the $\eta(1405)$ Particle Listings.
- [r] See also the $\omega(1650)$ Particle Listings.
- [s] See the “Note on the $\rho(1450)$ and the $\rho(1700)$ ” in the $\rho(1700)$ Particle Listings.
- [t] See also the $\omega(1420)$ Particle Listings.
- [u] See the “Note on $f_0(1710)$ ” in the $f_0(1710)$ Particle Listings .