

# $\Upsilon(1S)$

$$J^{PC} = 0^-(1^{--})$$

## $\Upsilon(1S)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9460.30 ± 0.26 OUR AVERAGE</b>	Error includes scale factor of 3.3.		
9460.51 ± 0.09 ± 0.05	<sup>1</sup> ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
9459.97 ± 0.11 ± 0.07	MACKAY	84 REDE	$e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9460.60 ± 0.09 ± 0.05	<sup>2,3</sup> BARU	92B REDE	$e^+ e^- \rightarrow$ hadrons
9460.59 ± 0.12	BARU	86 REDE	$e^+ e^- \rightarrow$ hadrons
9460.6 ± 0.4	<sup>3,4</sup> ARTAMONOV 84	REDE	$e^+ e^- \rightarrow$ hadrons
<sup>1</sup> Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).			
<sup>2</sup> Superseding BARU 86.			
<sup>3</sup> Superseded by ARTAMONOV 00.			
<sup>4</sup> Value includes data of ARTAMONOV 82.			

## $\Upsilon(1S)$ WIDTH

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
<b>53.0 ± 1.5 OUR EVALUATION</b>	See the Note on Width Determinations of the $\Upsilon$ states

## $\Upsilon(1S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $\tau^+ \tau^-$	(2.67 <sup>+0.14</sup> <sub>-0.16</sub> ) %	
$\Gamma_2$ $e^+ e^-$	(2.38 ± 0.11) %	
$\Gamma_3$ $\mu^+ \mu^-$	(2.48 ± 0.05) %	
<b>Hadronic decays</b>		
$\Gamma_4$ $\eta'(958)$ anything	(2.8 ± 0.4) %	
$\Gamma_5$ $J/\psi(1S)$ anything	(6.5 ± 0.7) × 10 <sup>-4</sup>	
$\Gamma_6$ $\chi_{c0}$ anything	< 5 × 10 <sup>-3</sup>	90%
$\Gamma_7$ $\chi_{c1}$ anything	(2.3 ± 0.7) × 10 <sup>-4</sup>	
$\Gamma_8$ $\chi_{c2}$ anything	(3.4 ± 1.0) × 10 <sup>-4</sup>	
$\Gamma_9$ $\psi(2S)$ anything	(2.7 ± 0.9) × 10 <sup>-4</sup>	
$\Gamma_{10}$ $\rho\pi$	< 2 × 10 <sup>-4</sup>	90%
$\Gamma_{11}$ $\pi^+ \pi^-$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_{12}$ $K^+ K^-$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_{13}$ $p\bar{p}$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_{14}$ $\pi^0 \pi^+ \pi^-$	< 1.84 × 10 <sup>-5</sup>	90%
$\Gamma_{15}$ $D^*(2010)^\pm$ anything		

### Radiative decays

$\Gamma_{16}$	$\gamma \pi^+ \pi^-$	$(6.3 \pm 1.8) \times 10^{-5}$	
$\Gamma_{17}$	$\gamma \pi^0 \pi^0$	$(1.7 \pm 0.7) \times 10^{-5}$	
$\Gamma_{18}$	$\gamma 2h^+ 2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$	
$\Gamma_{19}$	$\gamma 3h^+ 3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$	
$\Gamma_{20}$	$\gamma 4h^+ 4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$	
$\Gamma_{21}$	$\gamma \pi^+ \pi^- K^+ K^-$	$(2.9 \pm 0.9) \times 10^{-4}$	
$\Gamma_{22}$	$\gamma 2\pi^+ 2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$	
$\Gamma_{23}$	$\gamma 3\pi^+ 3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$	
$\Gamma_{24}$	$\gamma 2\pi^+ 2\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$	
$\Gamma_{25}$	$\gamma \pi^+ \pi^- p \bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$	
$\Gamma_{26}$	$\gamma 2\pi^+ 2\pi^- p \bar{p}$	$(4 \pm 6) \times 10^{-5}$	
$\Gamma_{27}$	$\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$	
$\Gamma_{28}$	$\gamma \eta'(958)$	$< 1.6$	$\times 10^{-5}$ 90%
$\Gamma_{29}$	$\gamma \eta$	$< 2.1$	$\times 10^{-5}$ 90%
$\Gamma_{30}$	$\gamma f'_2(1525)$	$< 1.4$	$\times 10^{-4}$ 90%
$\Gamma_{31}$	$\gamma f_2(1270)$	$(8 \pm 4)$	$\times 10^{-5}$
$\Gamma_{32}$	$\gamma \eta(1405)$	$< 8.2$	$\times 10^{-5}$ 90%
$\Gamma_{33}$	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$< 2.6$	$\times 10^{-4}$ 90%
$\Gamma_{34}$	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	$< 2$	$\times 10^{-4}$ 90%
$\Gamma_{35}$	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	$< 1.5$	$\times 10^{-5}$ 90%
$\Gamma_{36}$	$\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-$	$< 1.2$	$\times 10^{-5}$ 90%
$\Gamma_{37}$	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$< 1.6$	$\times 10^{-5}$ 90%
$\Gamma_{38}$	$\gamma \eta(2225) \rightarrow \gamma \phi \phi$	$< 3$	$\times 10^{-3}$ 90%
$\Gamma_{39}$	$\gamma X$ ( $X =$ pseudoscalar with $m < 7.2$ GeV)	$< 3$	$\times 10^{-5}$ 90%
$\Gamma_{40}$	$\gamma X \bar{X}$ ( $X \bar{X} =$ vectors with $m < 3.1$ GeV)	$< 1$	$\times 10^{-3}$ 90%

### $\mathcal{R}(1S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$

$\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-) / \Gamma_{\text{total}}$				$\Gamma_2 \Gamma_3 / \Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>31.2 \pm 1.6 \pm 1.7</math></b>	KOBEL	92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_0 \Gamma_2 / \Gamma$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>1.216 \pm 0.027</math> OUR AVERAGE</b>				
$1.187 \pm 0.023 \pm 0.031$	<sup>5</sup> BARU	92B	MD1	$e^+ e^- \rightarrow$ hadrons
$1.23 \pm 0.02 \pm 0.05$	<sup>5</sup> JAKUBOWSKI	88	CBAL	$e^+ e^- \rightarrow$ hadrons
$1.37 \pm 0.06 \pm 0.09$	<sup>6</sup> GILES	84B	CLEO	$e^+ e^- \rightarrow$ hadrons
$1.23 \pm 0.08 \pm 0.04$	<sup>6</sup> ALBRECHT	82	DASP	$e^+ e^- \rightarrow$ hadrons
$1.13 \pm 0.07 \pm 0.11$	<sup>6</sup> NICZYPORUK	82	LENA	$e^+ e^- \rightarrow$ hadrons
$1.09 \pm 0.25$	<sup>6</sup> BOCK	80	CNTR	$e^+ e^- \rightarrow$ hadrons
$1.35 \pm 0.14$	<sup>7</sup> BERGER	79	PLUT	$e^+ e^- \rightarrow$ hadrons

<sup>5</sup> Radiative corrections evaluated following KURAEV 85.

<sup>6</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

<sup>7</sup> Radiative corrections reevaluated by ALEXANDER 89 using  $B(\mu\mu) = 0.026$ .

## $\Upsilon(1S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

$\Gamma_2$

VALUE (keV)

DOCUMENT ID

**1.314 ± 0.029 OUR EVALUATION**

## $\Upsilon(1S)$ BRANCHING RATIOS

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma$

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

**0.0267<sup>+0.0014</sup><sub>-0.0016</sub> OUR AVERAGE**

0.0261 ± 0.0012<sup>+0.0009</sup><sub>-0.0013</sub>

25k

CINABRO

94B CLE2

$e^+e^- \rightarrow \tau^+\tau^-$

0.027 ± 0.004 ± 0.002

<sup>8</sup> ALBRECHT

85C ARG

$\Upsilon(2S) \rightarrow$

0.034 ± 0.004 ± 0.004

GILES

83 CLEO

$e^+e^- \rightarrow \tau^+\tau^-$   
 $\pi^+\pi^-\tau^+\tau^-$

<sup>8</sup> Using  $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$ ; not used for width evaluations.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

$\Gamma_3/\Gamma$

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

**0.0248 ± 0.0005 OUR AVERAGE**

0.0249 ± 0.0002 ± 0.0007

345k

ADAMS

05 CLEO

$e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.0249 ± 0.0008 ± 0.0013

ALEXANDER

98 CLE2

$\Upsilon(2S) \rightarrow$

0.0212 ± 0.0020 ± 0.0010

<sup>9</sup> BARU

92 MD1

$\pi^+\pi^-\mu^+\mu^-$   
 $e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.0231 ± 0.0012 ± 0.0010

<sup>9</sup> KOBEL

92 CBAL

$e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.0252 ± 0.0007 ± 0.0007

CHEN

89B CLEO

$e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.0261 ± 0.0009 ± 0.0011

KAARSBERG

89 CSB2

$e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.0230 ± 0.0025 ± 0.0013

86

ALBRECHT

87 ARG

$\Upsilon(2S) \rightarrow$

0.029 ± 0.003 ± 0.002

864

BESSON

84 CLEO

$\pi^+\pi^-\mu^+\mu^-$   
 $\Upsilon(2S) \rightarrow$

0.027 ± 0.003 ± 0.003

ANDREWS

83 CLEO

$\pi^+\pi^-\mu^+\mu^-$   
 $e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.032 ± 0.013 ± 0.003

ALBRECHT

82 DASP

$e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.038 ± 0.015 ± 0.002

NICZYPORUK

82 LENA

$e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.014<sup>+0.034</sup><sub>-0.014</sub>

BOCK

80 CNTR

$e^+e^- \rightarrow$   
 $\mu^+\mu^-$

0.022 ± 0.020

BERGER

79 PLUT

$e^+e^- \rightarrow$   
 $\mu^+\mu^-$

<sup>9</sup> Taking into account interference between the resonance and continuum.

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$			$\Gamma_2/\Gamma$		
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>0.0238±0.0011 OUR AVERAGE</b>					
0.0229±0.0008±0.0011		ALEXANDER	98 CLE2	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$	
0.0242±0.0014±0.0014	307	ALBRECHT	87 ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$	
0.028 ±0.003 ±0.002	826	BESSION	84 CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$	
0.051 ±0.030		BERGER	80C PLUT	$e^+e^- \rightarrow e^+e^-$	

$\Gamma(\eta'(958) \text{ anything})/\Gamma_{\text{total}}$			$\Gamma_4/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>0.028±0.004±0.002</b>					
	ARTUSO	03 CLE2	$\Upsilon(1S) \rightarrow \eta' \text{ anything}$		

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$			$\Gamma_5/\Gamma$		
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.65±0.07 OUR AVERAGE</b>					
0.64±0.04±0.06		730±40	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi X$
1.1 ±0.4 ±0.2		10	FULTON	89 CLEO	$e^+e^- \rightarrow \mu^+\mu^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.68	90		ALBRECHT	92J ARG	$e^+e^- \rightarrow e^+e^- X,$ $e^+e^- \rightarrow \mu^+\mu^- X$
<1.7	90		MASCHMANN	90 CBAL	$e^+e^- \rightarrow \text{hadrons}$
<20	90		NICZYPORUK	83 LENA	

<sup>10</sup> Using  $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$ .

$\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$			$\Gamma_6/\Gamma_5$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;7.4</b>					
	90	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi X$	

$\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$			$\Gamma_7/\Gamma_5$		
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>0.35±0.08±0.06</b>					
	52 ± 12	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi X$	

$\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$			$\Gamma_8/\Gamma_5$		
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>0.52±0.12±0.09</b>					
	47 ± 11	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi X$	

$\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$			$\Gamma_9/\Gamma_5$		
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>0.41±0.11±0.08</b>					
	42 ± 11	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi \pi^+ \pi^- X$	

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$			$\Gamma_{11}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;5</b>					
	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow \pi^+\pi^-$	

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow K^+ K^-$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	<sup>11</sup> BARU	96 MD1	$\Upsilon(1S) \rightarrow p\bar{p}$

<sup>11</sup>Supersedes BARU 92 in this node.

$\Gamma(\pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.84</b>	90	ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma X)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$

( $X$  = pseudoscalar with  $m < 7.2$  GeV)

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3</b>	90	<sup>12</sup> BALEST	95 CLEO	$e^+ e^- \rightarrow \gamma + X$

<sup>12</sup>For a noninteracting pseudoscalar  $X$  with mass  $< 7.2$  GeV.

$\Gamma(\gamma X \bar{X})/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$

( $X \bar{X}$  = vectors with  $m < 3.1$  GeV)

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1</b>	90	<sup>13</sup> BALEST	95 CLEO	$e^+ e^- \rightarrow \gamma + X \bar{X}$

<sup>13</sup>For a noninteracting vector  $X$  with mass  $< 3.1$  GeV.

$\Gamma(\gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>6.3 \pm 1.2 \pm 1.3</math></b>	<sup>14</sup> ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

<sup>14</sup>For  $m_{\pi\pi} > 1$  GeV.

$\Gamma(\gamma \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>1.7 \pm 0.6 \pm 0.3</math></b>	<sup>15</sup> ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

<sup>15</sup>For  $m_{\pi\pi} > 1$  GeV.

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.5 \pm 0.7 \pm 0.5</math></b>	$26 \pm 7$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.9 \pm 0.7 \pm 0.6</math></b>	$29 \pm 8$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma \pi^+ \pi^- \rho\bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.5 \pm 0.5 \pm 0.3</math></b>	$22 \pm 6$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

**$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.2 ± 0.2</b>	2 ± 2	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

**$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.5 ± 0.9 ± 0.8</b>	17 ± 5	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

**$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.4 ± 0.9 ± 0.8</b>	18 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

**$\Gamma(\gamma 2\pi^+ 2\pi^- \rho\bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.4 ± 0.4 ± 0.4</b>	7 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

**$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>7.0 ± 1.1 ± 1.0</b>	80 ± 12	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

**$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.4 ± 1.5 ± 1.3</b>	39 ± 11	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

**$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>7.4 ± 2.5 ± 2.5</b>	36 ± 12	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

**$\Gamma(\rho\pi)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 2</b>	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10	90	BLINOV	90 MD1	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
<21	90	NICZYPORUK	83 LENA	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$

**$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<19	90	<sup>16</sup> ALBRECHT	92J ARG	$e^+ e^- \rightarrow D^0 \pi^\pm X$

<sup>16</sup> For  $x_p > 0.2$ .

**$\Gamma(\gamma\eta(1405))/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;8.2</b>	90	<sup>17</sup> FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$

<sup>17</sup> Includes unknown branching ratio of  $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$ .

**$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$**   **$\Gamma_{28}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 1.6</b>	90	RICHICHI	01B CLE2	$\Upsilon(1S) \rightarrow \gamma\eta' \rightarrow \gamma\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<130	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$
------	----	---------	---------	-------------------------------------

**$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$**   **$\Gamma_{29}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 2.1</b>	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<28.2	90	MASEK	02 CLEO	$\eta \rightarrow \gamma\gamma$
< 6.7	90	MASEK	02 CLEO	$\eta \rightarrow \pi^0\pi^0\pi^0$
< 2.6	90	MASEK	02 CLEO	$\eta \rightarrow \pi^+\pi^-\pi^0$
<35	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

**$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$**   **$\Gamma_{30}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;14</b>	90	18 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19.4	90	18 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
-------	----	-------------	--------	---

<sup>18</sup> Assuming  $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$ .

**$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$**   **$\Gamma_{33}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 2.6</b>	90	19 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.3	90	19 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<19	90	19 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$
< 8	90	20 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$
<24	90	21 SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

<sup>19</sup> Assuming  $B(f_0(1710) \rightarrow K\bar{K}) = 0.38$ .

<sup>20</sup> Assuming  $B(f_0(1710) \rightarrow \pi\pi) = 0.04$ .

<sup>21</sup> Assuming  $B(f_0(1710) \rightarrow \eta\eta) = 0.18$ .

**$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$**   **$\Gamma_{31}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>8.1 \pm 2.3^{+2.9}_{-2.7}</math></b>		22 ANASTASSOV	99 CLE2	$e^+ e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	22 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$
<13	90	22 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$
<81	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

<sup>22</sup> Using  $B(f_2(1270) \rightarrow \pi\pi) = 0.84$ .

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$   $\Gamma_{35} / \Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 1.6	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<b>&lt; 1.5</b>	90	<sup>23</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 2.9	90	<sup>23</sup> ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 20	90	<sup>23</sup> BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

<sup>23</sup> Including unknown branching ratio of  $f_J(2220) \rightarrow K^+ K^-$ .

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{36} / \Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.2</b>	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p}) / \Gamma_{\text{total}}$   $\Gamma_{37} / \Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.6</b>	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma p \bar{p}$

$\Gamma(\gamma \eta(2225) \rightarrow \gamma \phi \phi) / \Gamma_{\text{total}}$   $\Gamma_{38} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.003</b>	90	<sup>24</sup> BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

<sup>24</sup> Assuming that the  $\eta(2225)$  decays only into  $\phi \phi$ .

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$   $\Gamma_{34} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.0002</b>	90	<sup>25</sup> BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

<sup>25</sup> Assuming that the  $f_0(2200)$  decays only into  $K^+ K^-$ .

### $\Upsilon(1S)$ REFERENCES

ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)



KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)

### OTHER RELATED PAPERS

KOENIGS...	86	DESY 86/136	K. Koenigsmann	(DESY)
ALBRECHT	84	PL 134B 137	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
BERGER	78	PL 76B 243	C. Berger <i>et al.</i>	(PLUTO Collab.)
BIELEIN	78	PL 78B 360	J.K. Bienlein <i>et al.</i>	(DESY, HAMB, HEIDP+)
DARDEN	78	PL 76B 246	C.W. Darden <i>et al.</i>	(DESY, DORT, HEIDH+)
GARELICK	78	PR D18 945	D.A. Garelick <i>et al.</i>	(NEAS, WASH, TUFTS)
KAPLAN	78	PRL 40 435	D.M. Kaplan <i>et al.</i>	(STON, FNAL, COLU)
YOH	78	PRL 41 684	J.K. Yoh <i>et al.</i>	(COLU, FNAL, STON)
COBB	77	PL 72B 273	J.H. Cobb <i>et al.</i>	(BNL, CERN, SYRA, YALE)
HERB	77	PRL 39 252	S.W. Herb <i>et al.</i>	(COLU, FNAL, STON)
INNES	77	PRL 39 1240	W.R. Innes <i>et al.</i>	(COLU, FNAL, STON)