



$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = \frac{2}{3} e \quad \text{Charm} = +1$$

c-QUARK MASS

The *c*-quark mass corresponds to the “running” mass $m_c(\mu = m_c)$ in the $\overline{\text{MS}}$ scheme. We have converted masses in other schemes to the $\overline{\text{MS}}$ scheme using two-loop QCD perturbation theory with $\alpha_s(\mu=m_c) = 0.39$. The range 1.0–1.4 GeV for the $\overline{\text{MS}}$ mass corresponds to 1.47–1.83 GeV for the pole mass (see the “Note on Quark Masses”).

<u>VALUE (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.15 to 1.35 OUR EVALUATION			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.29 ± 0.07	1 HOANG	04	THEO $\overline{\text{MS}}$ scheme
1.319 ± 0.028	2 DEDIVITIIS	03	LATT $\overline{\text{MS}}$ scheme
1.19 ± 0.11	3 EIDEMULLER	03	THEO $\overline{\text{MS}}$ scheme
1.289 ± 0.043	4 ERLER	03	THEO $\overline{\text{MS}}$ scheme
1.26 ± 0.02	5 ZYABLYUK	03	THEO $\overline{\text{MS}}$ scheme
1.26 ± 0.04 ± 0.12	6 BECIREVIC	02	LATT $\overline{\text{MS}}$ scheme
1.301 ± 0.034	7 ROLF	02	LATT $\overline{\text{MS}}$ scheme
1.23 ± 0.09	8 EIDEMULLER	01	THEO $\overline{\text{MS}}$ scheme
1.304 ± 0.027	9 KUHN	01	THEO $\overline{\text{MS}}$ scheme
1.04 ± 0.04	10 MARTIN	01	THEO $\overline{\text{MS}}$ scheme
1.1 ± 0.04	11 NARISON	01B	THEO $\overline{\text{MS}}$ scheme
1.37 ± 0.09	12 PENARROCHA	01	THEO $\overline{\text{MS}}$ scheme
1.210 ± 0.070 ± 0.080	13 PINEDA	01	THEO $\overline{\text{MS}}$ scheme
1.3 ± 0.3 ± 0.3	14 ASTIER	00D	NOMD
1.79 ± 0.38	15 VILAIN	99	THEO $\overline{\text{MS}}$ scheme

¹ HOANG 04 determines $\overline{m}_c(\overline{m}_c)$ from moments at order α_s^2 of the charm production cross-section in e^+e^- annihilation.

² DEDIVITIIS 03 use a quenched lattice computation of heavy-heavy and heavy-light meson masses.

³ EIDEMULLER 03 determines m_b and m_c using QCD sum rules.

⁴ ERLER 03 determines m_b and m_c using QCD sum rules. Includes recent BES data.

⁵ ZYABLYUK 03 determines m_c by using QCD sum rules in the pseudoscalar channel and comparing with the η_c mass.

⁶ BECIREVIC 02 uses Monte-Carlo calculations of lattice Ward identities and the D_s mass. The authors estimate an error of about 5% for use of the quenched approximation, not included in systematic error of 0.12.

⁷ ROLF 02 determines m_c from a quenched lattice calculation of the D_s mass. The error estimate is for all systematics except the quenched approximation, including lattice spacing effects, finite volume effects, excited states contamination, rounding errors, and the scale uncertainty. The authors estimate the uncertainty due to the quenched approximation may be about 3%.

⁸ EIDEMULLER 01 result is QCD sum rule analysis of charmonium using NRQCD at next-to-next-to-leading order.

⁹ KUHN 01 uses an analysis of the e^+e^- total cross section to hadrons.

¹⁰ MARTIN 01 obtain a pole mass of 1.33–1.4 GeV from an analysis of R , the rate for $e^+e^- \rightarrow$ hadrons. We have converted this to the $\overline{\text{MS}}$ scheme using the two-loop formula.

