



$$J = \frac{1}{2}$$

τ discovery paper was PERL 75. $e^+e^- \rightarrow \tau^+\tau^-$ cross-section threshold behavior and magnitude are consistent with pointlike spin-1/2 Dirac particle. BRANDELIK 78 ruled out pointlike spin-0 or spin-1 particle. FELDMAN 78 ruled out $J = 3/2$. KIRKBY 79 also ruled out $J=\text{integer}$, $J = 3/2$.

τ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
$1776.99^{+0.29}_{-0.26}$				OUR AVERAGE
$1775.1 \pm 1.6 \pm 1.0$	13.3k	¹ ABBIENDI	00A OPAL	1990–1995 LEP runs
$1778.2 \pm 0.8 \pm 1.2$		ANASTASSOV 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
$1776.96^{+0.18+0.25}_{-0.21-0.17}$	65	² BAI	96 BES	$E_{\text{cm}}^{ee} = 3.54\text{--}3.57$ GeV
$1776.3 \pm 2.4 \pm 1.4$	11k	³ ALBRECHT	92M ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
$1783 \begin{smallmatrix} +3 \\ -4 \end{smallmatrix}$	692	⁴ BACINO	78B DLCO	$E_{\text{cm}}^{ee} = 3.1\text{--}7.4$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1777.8 \pm 0.7 \pm 1.7$	35k	⁵ BALEST	93 CLEO	Repl. by ANASTASSOV 97
$1776.9 \begin{smallmatrix} +0.4 \\ -0.5 \end{smallmatrix} \pm 0.2$	14	⁶ BAI	92 BES	Repl. by BAI 96

¹ ABBIENDI 00A fit τ pseudomass spectrum in $\tau \rightarrow \pi^\pm \leq 2\pi^0 \nu_\tau$ and $\tau \rightarrow \pi^\pm \pi^+ \pi^- \leq 1\pi^0 \nu_\tau$ decays. Result assumes $m_{\nu_\tau} = 0$.

² BAI 96 fit $\sigma(e^+e^- \rightarrow \tau^+\tau^-)$ at different energies near threshold.

³ ALBRECHT 92M fit τ pseudomass spectrum in $\tau^- \rightarrow 2\pi^- \pi^+ \nu_\tau$ decays. Result assumes $m_{\nu_\tau} = 0$.

⁴ BACINO 78B value comes from $e^\pm X^\mp$ threshold. Published mass 1782 MeV increased by 1 MeV using the high precision $\psi(2S)$ mass measurement of ZHOLENTZ 80 to eliminate the absolute SPEAR energy calibration uncertainty.

⁵ BALEST 93 fit spectra of minimum kinematically allowed τ mass in events of the type $e^+e^- \rightarrow \tau^+\tau^- \rightarrow (\pi^+ n\pi^0 \nu_\tau)(\pi^- m\pi^0 \nu_\tau)$ $n \leq 2, m \leq 2, 1 \leq n+m \leq 3$. If $m_{\nu_\tau} \neq 0$, result increases by $(m_{\nu_\tau}^2/1100)$ MeV.

⁶ BAI 92 fit $\sigma(e^+e^- \rightarrow \tau^+\tau^-)$ near threshold using $e\mu$ events.

$$(m_{\tau^+} - m_{\tau^-})/m_{\text{average}}$$

A test of CPT invariance.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	ABBIENDI	00A OPAL	1990–1995 LEP runs

τ MEAN LIFE

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
290.6 ± 1.0	OUR AVERAGE			
290.9 ± 1.4 ± 1.0		ABDALLAH	04T DLPH	1991-1995 LEP runs
293.2 ± 2.0 ± 1.5		ACCIARRI	00B L3	1991-1995 LEP runs
290.1 ± 1.5 ± 1.1		BARATE	97R ALEP	1989-1994 LEP runs
289.2 ± 1.7 ± 1.2		ALEXANDER	96E OPAL	1990-1994 LEP runs
289.0 ± 2.8 ± 4.0	57.4k	BALEST	96 CLEO	$E_{cm}^{ee} = 10.6$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
291.2 ± 2.0 ± 1.2		BARATE	97I ALEP	Repl. by BARATE 97R
291.4 ± 3.0		ABREU	96B DLPH	Repl. by ABDALLAH 04T
290.1 ± 4.0	34k	ACCIARRI	96K L3	Repl. by ACCIARRI 00B
297 ± 9 ± 5	1671	ABE	95Y SLD	1992-1993 SLC runs
304 ± 14 ± 7	4100	BATTLE	92 CLEO	$E_{cm}^{ee} = 10.6$ GeV
301 ± 29	3780	KLEINWORT	89 JADE	$E_{cm}^{ee} = 35-46$ GeV
288 ± 16 ± 17	807	AMIDEI	88 MRK2	$E_{cm}^{ee} = 29$ GeV
306 ± 20 ± 14	695	BRAUNSCH...	88C TASS	$E_{cm}^{ee} = 36$ GeV
299 ± 15 ± 10	1311	ABACHI	87C HRS	$E_{cm}^{ee} = 29$ GeV
295 ± 14 ± 11	5696	ALBRECHT	87P ARG	$E_{cm}^{ee} = 9.3-10.6$ GeV
309 ± 17 ± 7	3788	BAND	87B MAC	$E_{cm}^{ee} = 29$ GeV
325 ± 14 ± 18	8470	BEBEK	87C CLEO	$E_{cm}^{ee} = 10.5$ GeV
460 ± 190	102	FELDMAN	82 MRK2	$E_{cm}^{ee} = 29$ GeV

τ MAGNETIC MOMENT ANOMALY

The q^2 dependence is expected to be small providing no thresholds are nearby.

$$\mu_{\tau}/(e\hbar/2m_{\tau})-1 = (g_{\tau}-2)/2$$

For a theoretical calculation [$(g_{\tau}-2)/2 = 11773(3) \times 10^{-7}$], see SAMUEL 91B.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
> -0.052 and < 0.013 (CL = 95%) OUR LIMIT				
> -0.052 and < 0.013	95	⁷ ABDALLAH	04K DLPH	$e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ at LEP2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 0.107	95	⁸ ACHARD	04G L3	$e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ at LEP2
> -0.007 and < 0.005	95	⁹ GONZALEZ-S..00	RVUE	$e^+e^- \rightarrow \tau^+\tau^-$ and $W \rightarrow \tau\nu_{\tau}$
> -0.052 and < 0.058	95	¹⁰ ACCIARRI	98E L3	1991-1995 LEP runs
> -0.068 and < 0.065	95	¹¹ ACKERSTAFF	98N OPAL	1990-1995 LEP runs
> -0.004 and < 0.006	95	¹² ESCRIBANO	97 RVUE	$Z \rightarrow \tau^+\tau^-$ at LEP
< 0.01	95	¹³ ESCRIBANO	93 RVUE	$Z \rightarrow \tau^+\tau^-$ at LEP
< 0.12	90	GRIFOLS	91 RVUE	$Z \rightarrow \tau\tau\gamma$ at LEP
< 0.023	95	¹⁴ SILVERMAN	83 RVUE	$e^+e^- \rightarrow \tau^+\tau^-$ at PETRA

- ⁷ ABDALLAH 04K limit is derived from $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ total cross-section measurements at \sqrt{s} between 183 and 208 GeV. In addition to the limits, the authors also quote a value of -0.018 ± 0.017 .
- ⁸ ACHARD 04G limit is derived from $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ total cross-section measurements at \sqrt{s} between 189 and 206 GeV, and is on the absolute value of the magnetic moment anomaly.
- ⁹ GONZALEZ-SPRINBERG 00 use data on tau lepton production at LEP1, SLC, and LEP2, and data from colliders and LEP2 to determine limits. Assume imaginary component is zero.
- ¹⁰ ACCIARRI 98E use $Z \rightarrow \tau^+\tau^-\gamma$ events. In addition to the limits, the authors also quote a value of $0.004 \pm 0.027 \pm 0.023$.
- ¹¹ ACKERSTAFF 98N use $Z \rightarrow \tau^+\tau^-\gamma$ events. The limit applies to an average of the form factor for off-shell τ 's having p^2 ranging from m_τ^2 to $(M_Z - m_\tau)^2$.
- ¹² ESCRIBANO 97 use preliminary experimental results.
- ¹³ ESCRIBANO 93 limit derived from $\Gamma(Z \rightarrow \tau^+\tau^-)$, and is on the absolute value of the magnetic moment anomaly.
- ¹⁴ SILVERMAN 83 limit is derived from $e^+e^- \rightarrow \tau^+\tau^-$ total cross-section measurements for q^2 up to $(37 \text{ GeV})^2$.

τ ELECTRIC DIPOLE MOMENT (d_τ)

A nonzero value is forbidden by both T invariance and P invariance.

The q^2 dependence is expected to be small providing no thresholds are nearby.

Re(d_τ)

VALUE ($10^{-16} e\text{cm}$)	CL%	DOCUMENT ID	TECN	COMMENT
– 0.22 to 0.45	95	¹⁵ INAMI	03 BELL	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 3.7	95	¹⁶ ABDALLAH	04K DLPH	$e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ at LEP2
< 11.4	95	¹⁷ ACHARD	04G L3	$e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ at LEP2
< 4.6	95	¹⁸ ALBRECHT	00 ARG	$E_{\text{cm}}^{ee} = 10.4 \text{ GeV}$
> –3.1 and < 3.1	95	ACCIARRI	98E L3	1991–1995 LEP runs
> –3.8 and < 3.6	95	¹⁹ ACKERSTAFF	98N OPAL	1990–1995 LEP runs
< 0.11	95	^{20,21} ESCRIBANO	97 RVUE	$Z \rightarrow \tau^+\tau^-$ at LEP
< 0.5	95	²² ESCRIBANO	93 RVUE	$Z \rightarrow \tau^+\tau^-$ at LEP
< 7	90	GRIFOLS	91 RVUE	$Z \rightarrow \tau\tau\gamma$ at LEP
< 1.6	90	DELAGUILA	90 RVUE	$e^+e^- \rightarrow \tau^+\tau^-$ $E_{\text{cm}}^{ee} = 35 \text{ GeV}$

¹⁵ INAMI 03 use $e^+e^- \rightarrow \tau^+\tau^-$ events.

¹⁶ ABDALLAH 04K limit is derived from $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ total cross-section measurements at \sqrt{s} between 183 and 208 GeV and is on the absolute value of d_τ .

¹⁷ ACHARD 04G limit is derived from $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ total cross-section measurements at \sqrt{s} between 189 and 206 GeV, and is on the absolute value of d_τ .

¹⁸ ALBRECHT 00 use $e^+e^- \rightarrow \tau^+\tau^-$ events. Limit is on the absolute value of Re(d_τ).

¹⁹ ACKERSTAFF 98N use $Z \rightarrow \tau^+\tau^-\gamma$ events. The limit applies to an average of the form factor for off-shell τ 's having p^2 ranging from m_τ^2 to $(M_Z - m_\tau)^2$.

²⁰ ESCRIBANO 97 derive the relationship $|d_\tau| = \cot \theta_W |d_\tau^W|$ using effective Lagrangian methods, and use a conference result $|d_\tau^W| < 5.8 \times 10^{-18} \text{ e cm}$ at 95% CL (L. Silvestris, ICHEP96) to obtain this result.

²¹ ESCRIBANO 97 use preliminary experimental results.

²² ESCRIBANO 93 limit derived from $\Gamma(Z \rightarrow \tau^+ \tau^-)$, and is on the absolute value of the electric dipole moment.

Im(d_τ)

VALUE (10^{-16} e cm)	CL%	DOCUMENT ID	TECN	COMMENT
-0.25 to 0.008	95	²³ INAMI	03 BELL	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.8	95	²⁴ ALBRECHT	00 ARG	$E_{\text{cm}}^{ee} = 10.4 \text{ GeV}$
-------	----	------------------------	--------	---

²³ INAMI 03 use $e^+ e^- \rightarrow \tau^+ \tau^-$ events.

²⁴ ALBRECHT 00 use $e^+ e^- \rightarrow \tau^+ \tau^-$ events. Limit is on the absolute value of Im(d_τ).

τ WEAK DIPOLE MOMENT (d_τ^W)

A nonzero value is forbidden by *CP* invariance.

The q^2 dependence is expected to be small providing no thresholds are nearby.

Re(d_τ^W)

VALUE (10^{-17} e cm)	CL%	DOCUMENT ID	TECN	COMMENT
<0.50	95	²⁵ HEISTER	03F ALEP	1990–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.0	90	²⁵ ACCIARRI	98C L3	1991–1995 LEP runs
<0.56	95	ACKERSTAFF	97L OPAL	1991–1995 LEP runs
<0.78	95	²⁶ AKERS	95F OPAL	Repl. by ACKER-STAFF 97L
<1.5	95	²⁶ BUSKULIC	95C ALEP	Repl. by HEISTER 03F
<7.0	95	²⁶ ACTON	92F OPAL	$Z \rightarrow \tau^+ \tau^-$ at LEP
<3.7	95	²⁶ BUSKULIC	92J ALEP	Repl. by BUSKULIC 95C

²⁵ Limit is on the absolute value of the real part of the weak dipole moment.

²⁶ Limit is on the absolute value of the real part of the weak dipole moment, and applies for $q^2 = m_Z^2$.

Im(d_τ^W)

VALUE (10^{-17} e cm)	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	95	²⁷ HEISTER	03F ALEP	1990–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	95	ACKERSTAFF	97L OPAL	1991–1995 LEP runs
<4.5	95	²⁸ AKERS	95F OPAL	Repl. by ACKER-STAFF 97L

²⁷ HEISTER 03F limit is on the absolute value of the imaginary part of the weak dipole moment.

²⁸ Limit is on the absolute value of the imaginary part of the weak dipole moment, and applies for $q^2 = m_Z^2$.

τ WEAK ANOMALOUS MAGNETIC DIPOLE MOMENT (α_τ^W)

Electroweak radiative corrections are expected to contribute at the 10^{-6} level. See BERNABEU 95.

The q^2 dependence is expected to be small providing no thresholds are nearby.

Re(α_τ^W)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-3}$	95	²⁹ HEISTER	03F ALEP	1990–1995 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> -0.0024 and < 0.0025	95	³⁰ GONZALEZ-S..00	RVUE	$e^+e^- \rightarrow \tau^+\tau^-$ and $W \rightarrow \tau\nu_\tau$
$<4.5 \times 10^{-3}$	90	²⁹ ACCIARRI	98C L3	1991–1995 LEP runs
²⁹ Limit is on the absolute value of the real part of the weak anomalous magnetic dipole moment.				
³⁰ GONZALEZ-SPRINBERG 00 use data on tau lepton production at LEP1, SLC, and LEP2, and data from colliders and LEP2 to determine limits. Assume imaginary component is zero.				

Im(α_τ^W)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.7 \times 10^{-3}$	95	³¹ HEISTER	03F ALEP	1990–1995 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<9.9 \times 10^{-3}$	90	³¹ ACCIARRI	98C L3	1991–1995 LEP runs
³¹ Limit is on the absolute value of the imaginary part of the weak anomalous magnetic dipole moment.				

τ^- DECAY MODES

τ^+ modes are charge conjugates of the modes below. “ h^\pm ” stands for π^\pm or K^\pm . “ ℓ ” stands for e or μ . “Neutrals” stands for γ 's and/or π^0 's.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Modes with one charged particle		
Γ_1 particle $^- \geq 0$ neutrals $\geq 0K^0\nu_\tau$ (“1-prong”)	(85.35±0.07) %	S=1.1
Γ_2 particle $^- \geq 0$ neutrals $\geq 0K_L^0\nu_\tau$	(84.72±0.07) %	S=1.1
Γ_3 $\mu^- \bar{\nu}_\mu \nu_\tau$	[a] (17.36±0.06) %	
Γ_4 $\mu^- \bar{\nu}_\mu \nu_\tau \gamma$	[b] (3.6 ±0.4) × 10 ⁻³	
Γ_5 $e^- \bar{\nu}_e \nu_\tau$	[a] (17.84±0.06) %	
Γ_6 $e^- \bar{\nu}_e \nu_\tau \gamma$	[b] (1.75±0.18) %	
Γ_7 $h^- \geq 0K_L^0 \nu_\tau$	(12.30±0.11) %	S=1.4
Γ_8 $h^- \nu_\tau$	(11.75±0.11) %	S=1.4
Γ_9 $\pi^- \nu_\tau$	[a] (11.06±0.11) %	S=1.4
Γ_{10} $K^- \nu_\tau$	[a] (6.86±0.23) × 10 ⁻³	

Γ_{11}	$h^- \geq 1$ neutrals ν_τ		$(36.92 \pm 0.14) \%$	S=1.1
Γ_{12}	$h^- \pi^0 \nu_\tau$		$(25.87 \pm 0.13) \%$	S=1.1
Γ_{13}	$\pi^- \pi^0 \nu_\tau$	[a]	$(25.41 \pm 0.13) \%$	S=1.1
Γ_{14}	$\pi^- \pi^0$ non- $\rho(770) \nu_\tau$		$(3.0 \pm 3.2) \times 10^{-3}$	
Γ_{15}	$K^- \pi^0 \nu_\tau$	[a]	$(4.54 \pm 0.27) \times 10^{-3}$	
Γ_{16}	$h^- \geq 2\pi^0 \nu_\tau$		$(10.76 \pm 0.15) \%$	S=1.1
Γ_{17}	$h^- 2\pi^0 \nu_\tau$		$(9.39 \pm 0.14) \%$	S=1.1
Γ_{18}	$h^- 2\pi^0 \nu_\tau$ (ex. K^0)		$(9.23 \pm 0.14) \%$	S=1.1
Γ_{19}	$\pi^- 2\pi^0 \nu_\tau$ (ex. K^0)	[a]	$(9.17 \pm 0.14) \%$	S=1.1
Γ_{20}	$\pi^- 2\pi^0 \nu_\tau$ (ex. K^0),		$< 9 \times 10^{-3}$	CL=95%
Γ_{21}	scalar $\pi^- 2\pi^0 \nu_\tau$ (ex. K^0),		$< 7 \times 10^{-3}$	CL=95%
Γ_{22}	vector $K^- 2\pi^0 \nu_\tau$ (ex. K^0)	[a]	$(5.8 \pm 2.3) \times 10^{-4}$	
Γ_{23}	$h^- \geq 3\pi^0 \nu_\tau$		$(1.37 \pm 0.11) \%$	S=1.1
Γ_{24}	$h^- 3\pi^0 \nu_\tau$		$(1.21 \pm 0.10) \%$	
Γ_{25}	$\pi^- 3\pi^0 \nu_\tau$ (ex. K^0)	[a]	$(1.08 \pm 0.10) \%$	
Γ_{26}	$K^- 3\pi^0 \nu_\tau$ (ex. K^0, η)	[a]	$(3.7^{+2.2}_{-1.9}) \times 10^{-4}$	
Γ_{27}	$h^- 4\pi^0 \nu_\tau$ (ex. K^0)		$(1.6 \pm 0.6) \times 10^{-3}$	
Γ_{28}	$h^- 4\pi^0 \nu_\tau$ (ex. K^0, η)	[a]	$(1.0^{+0.6}_{-0.5}) \times 10^{-3}$	
Γ_{29}	$K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma \nu_\tau$		$(1.56 \pm 0.04) \%$	
Γ_{30}	$K^- \geq 1 (\pi^0 \text{ or } K^0 \text{ or } \gamma) \nu_\tau$		$(8.76 \pm 0.34) \times 10^{-3}$	

Modes with K^0 's

Γ_{31}	K_S^0 (particles) $^- \nu_\tau$		$(9.2 \pm 0.4) \times 10^{-3}$	S=1.1
Γ_{32}	$h^- \bar{K}^0 \nu_\tau$		$(1.05 \pm 0.04) \%$	S=1.1
Γ_{33}	$\pi^- \bar{K}^0 \nu_\tau$	[a]	$(8.9 \pm 0.4) \times 10^{-3}$	S=1.1
Γ_{34}	$\pi^- \bar{K}^0$ (non- $K^*(892)^-$) ν_τ		$< 1.7 \times 10^{-3}$	CL=95%
Γ_{35}	$K^- K^0 \nu_\tau$	[a]	$(1.54 \pm 0.16) \times 10^{-3}$	
Γ_{36}	$K^- K^0 \geq 0\pi^0 \nu_\tau$		$(3.08 \pm 0.24) \times 10^{-3}$	
Γ_{37}	$h^- \bar{K}^0 \pi^0 \nu_\tau$		$(5.2 \pm 0.4) \times 10^{-3}$	
Γ_{38}	$\pi^- \bar{K}^0 \pi^0 \nu_\tau$	[a]	$(3.7 \pm 0.4) \times 10^{-3}$	
Γ_{39}	$\bar{K}^0 \rho^- \nu_\tau$		$(2.2 \pm 0.5) \times 10^{-3}$	
Γ_{40}	$K^- K^0 \pi^0 \nu_\tau$	[a]	$(1.54 \pm 0.20) \times 10^{-3}$	
Γ_{41}	$\pi^- \bar{K}^0 \geq 1\pi^0 \nu_\tau$		$(3.2 \pm 1.0) \times 10^{-3}$	
Γ_{42}	$\pi^- \bar{K}^0 \pi^0 \pi^0 \nu_\tau$		$(2.6 \pm 2.4) \times 10^{-4}$	
Γ_{43}	$K^- K^0 \pi^0 \pi^0 \nu_\tau$		$< 1.6 \times 10^{-4}$	CL=95%
Γ_{44}	$\pi^- K^0 \bar{K}^0 \nu_\tau$		$(1.59 \pm 0.29) \times 10^{-3}$	S=1.1
Γ_{45}	$\pi^- K_S^0 K_S^0 \nu_\tau$	[a]	$(2.4 \pm 0.5) \times 10^{-4}$	
Γ_{46}	$\pi^- K_S^0 K_L^0 \nu_\tau$	[a]	$(1.10 \pm 0.28) \times 10^{-3}$	S=1.1
Γ_{47}	$\pi^- K^0 \bar{K}^0 \pi^0 \nu_\tau$		$(3.1 \pm 2.3) \times 10^{-4}$	
Γ_{48}	$\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau$		$< 2.0 \times 10^{-4}$	CL=95%

Γ_{49}	$\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau$	$(3.1 \pm 1.2) \times 10^{-4}$	
Γ_{50}	$K^0 h^+ h^- h^- \geq 0$ neutrals ν_τ	$< 1.7 \times 10^{-3}$	CL=95%
Γ_{51}	$K^0 h^+ h^- h^- \nu_\tau$	$(2.3 \pm 2.0) \times 10^{-4}$	

Modes with three charged particles

Γ_{52}	$h^- h^- h^+ \geq 0$ neutrals $\geq 0 K_L^0 \nu_\tau$	$(15.20 \pm 0.07) \%$	S=1.1
Γ_{53}	$h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$) ("3-prong")	$(14.57 \pm 0.07) \%$	S=1.1
Γ_{54}	$h^- h^- h^+ \nu_\tau$	$(10.01 \pm 0.09) \%$	S=1.2
Γ_{55}	$h^- h^- h^+ \nu_\tau$ (ex. K^0)	$(9.65 \pm 0.09) \%$	S=1.2
Γ_{56}	$h^- h^- h^+ \nu_\tau$ (ex. K^0, ω)	$(9.61 \pm 0.09) \%$	S=1.2
Γ_{57}	$\pi^- \pi^+ \pi^- \nu_\tau$	$(9.46 \pm 0.10) \%$	S=1.2
Γ_{58}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0)	$(9.15 \pm 0.10) \%$	S=1.2
Γ_{59}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0), non-axial vector	$< 2.4 \%$	CL=95%
Γ_{60}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0, ω)	[a] $(9.11 \pm 0.10) \%$	S=1.2
Γ_{61}	$h^- h^- h^+ \geq 1$ neutrals ν_τ	$(5.19 \pm 0.10) \%$	S=1.2
Γ_{62}	$h^- h^- h^+ \geq 1$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$)	$(4.92 \pm 0.09) \%$	S=1.3
Γ_{63}	$h^- h^- h^+ \pi^0 \nu_\tau$	$(4.53 \pm 0.09) \%$	S=1.3
Γ_{64}	$h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0)	$(4.35 \pm 0.09) \%$	S=1.3
Γ_{65}	$h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0, ω)	$(2.62 \pm 0.09) \%$	S=1.2
Γ_{66}	$\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$	$(4.37 \pm 0.09) \%$	S=1.3
Γ_{67}	$\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0)	$(4.24 \pm 0.09) \%$	S=1.3
Γ_{68}	$\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0, ω)	[a] $(2.51 \pm 0.09) \%$	S=1.2
Γ_{69}	$h^- \rho \pi^0 \nu_\tau$		
Γ_{70}	$h^- \rho^+ h^- \nu_\tau$		
Γ_{71}	$h^- \rho^- h^+ \nu_\tau$		
Γ_{72}	$h^- h^- h^+ 2\pi^0 \nu_\tau$	$(5.5 \pm 0.4) \times 10^{-3}$	
Γ_{73}	$h^- h^- h^+ 2\pi^0 \nu_\tau$ (ex. K^0)	$(5.4 \pm 0.4) \times 10^{-3}$	
Γ_{74}	$h^- h^- h^+ 2\pi^0 \nu_\tau$ (ex. K^0, ω, η)	[a] $(1.1 \pm 0.4) \times 10^{-3}$	
Γ_{75}	$h^- h^- h^+ 3\pi^0 \nu_\tau$	[a] $(2.3 \pm 0.8) \times 10^{-4}$	S=1.5
Γ_{76}	$K^- h^+ h^- \geq 0$ neutrals ν_τ	$(7.1 \pm 0.4) \times 10^{-3}$	S=1.4
Γ_{77}	$K^- h^+ \pi^- \nu_\tau$ (ex. K^0)	$(4.9 \pm 0.4) \times 10^{-3}$	S=1.6
Γ_{78}	$K^- h^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0)	$(1.10 \pm 0.22) \times 10^{-3}$	
Γ_{79}	$K^- \pi^+ \pi^- \geq 0$ neutrals ν_τ	$(5.2 \pm 0.5) \times 10^{-3}$	S=1.4
Γ_{80}	$K^- \pi^+ \pi^- \geq 0 \pi^0 \nu_\tau$ (ex. K^0)	$(4.1 \pm 0.5) \times 10^{-3}$	S=1.4
Γ_{81}	$K^- \pi^+ \pi^- \nu_\tau$	$(3.9 \pm 0.4) \times 10^{-3}$	S=1.6
Γ_{82}	$K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0)	[a] $(3.4 \pm 0.4) \times 10^{-3}$	S=1.6
Γ_{83}	$K^- \rho^0 \nu_\tau \rightarrow$ $K^- \pi^+ \pi^- \nu_\tau$	$(1.6 \pm 0.6) \times 10^{-3}$	
Γ_{84}	$K^- \pi^+ \pi^- \pi^0 \nu_\tau$	$(1.22 \pm 0.26) \times 10^{-3}$	
Γ_{85}	$K^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0)	$(6.9 \pm 2.5) \times 10^{-4}$	

Γ_{86}	$K^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0, η)	[a]	$(6.3 \pm 2.5) \times 10^{-4}$	
Γ_{87}	$K^- \pi^+ K^- \geq 0$ neut. ν_τ		$< 9 \times 10^{-4}$	CL=95%
Γ_{88}	$K^- K^+ \pi^- \geq 0$ neut. ν_τ		$(1.95 \pm 0.18) \times 10^{-3}$	S=1.1
Γ_{89}	$K^- K^+ \pi^- \nu_\tau$	[a]	$(1.55 \pm 0.07) \times 10^{-3}$	
Γ_{90}	$K^- K^+ \pi^- \pi^0 \nu_\tau$	[a]	$(4.1 \pm 1.6) \times 10^{-4}$	S=1.1
Γ_{91}	$K^- K^+ K^- \geq 0$ neut. ν_τ		$< 2.1 \times 10^{-3}$	CL=95%
Γ_{92}	$K^- K^+ K^- \nu_\tau$		$< 3.7 \times 10^{-5}$	CL=90%
Γ_{93}	$\pi^- K^+ \pi^- \geq 0$ neut. ν_τ		$< 2.5 \times 10^{-3}$	CL=95%
Γ_{94}	$e^- e^- e^+ \bar{\nu}_e \nu_\tau$		$(2.8 \pm 1.5) \times 10^{-5}$	
Γ_{95}	$\mu^- e^- e^+ \bar{\nu}_\mu \nu_\tau$		$< 3.6 \times 10^{-5}$	CL=90%

Modes with five charged particles

Γ_{96}	$3h^- 2h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^- \pi^+$) ("5-prong")		$(1.00 \pm 0.06) \times 10^{-3}$	
Γ_{97}	$3h^- 2h^+ \nu_\tau$ (ex. K^0)	[a]	$(8.2 \pm 0.6) \times 10^{-4}$	
Γ_{98}	$3h^- 2h^+ \pi^0 \nu_\tau$ (ex. K^0)	[a]	$(1.81 \pm 0.27) \times 10^{-4}$	
Γ_{99}	$3h^- 2h^+ 2\pi^0 \nu_\tau$		$< 1.1 \times 10^{-4}$	CL=90%

Miscellaneous other allowed modes

Γ_{100}	$(5\pi)^- \nu_\tau$		$(8.0 \pm 0.7) \times 10^{-3}$	
Γ_{101}	$4h^- 3h^+ \geq 0$ neutrals ν_τ ("7-prong")		$< 2.4 \times 10^{-6}$	CL=90%
Γ_{102}	$X^-(S=-1) \nu_\tau$		$(2.93 \pm 0.08) \%$	S=1.1
Γ_{103}	$K^*(892)^- \geq 0$ neutrals \geq $0K_L^0 \nu_\tau$		$(1.42 \pm 0.18) \%$	S=1.4
Γ_{104}	$K^*(892)^- \nu_\tau$		$(1.29 \pm 0.05) \%$	
Γ_{105}	$K^*(892)^0 K^- \geq 0$ neutrals ν_τ		$(3.2 \pm 1.4) \times 10^{-3}$	
Γ_{106}	$K^*(892)^0 K^- \nu_\tau$		$(2.1 \pm 0.4) \times 10^{-3}$	
Γ_{107}	$\bar{K}^*(892)^0 \pi^- \geq 0$ neutrals ν_τ		$(3.8 \pm 1.7) \times 10^{-3}$	
Γ_{108}	$\bar{K}^*(892)^0 \pi^- \nu_\tau$		$(2.2 \pm 0.5) \times 10^{-3}$	
Γ_{109}	$(\bar{K}^*(892)\pi)^- \nu_\tau \rightarrow$ $\pi^- \bar{K}^0 \pi^0 \nu_\tau$		$(1.0 \pm 0.4) \times 10^{-3}$	
Γ_{110}	$K_1(1270)^- \nu_\tau$		$(4.7 \pm 1.1) \times 10^{-3}$	
Γ_{111}	$K_1(1400)^- \nu_\tau$		$(1.7 \pm 2.6) \times 10^{-3}$	S=1.7
Γ_{112}	$K^*(1410)^- \nu_\tau$		$(1.5 \pm_{-1.0}^{+1.4}) \times 10^{-3}$	
Γ_{113}	$K_0^*(1430)^- \nu_\tau$		$< 5 \times 10^{-4}$	CL=95%
Γ_{114}	$K_2^*(1430)^- \nu_\tau$		$< 3 \times 10^{-3}$	CL=95%
Γ_{115}	$a_0(980)^- \geq 0$ neutrals ν_τ			
Γ_{116}	$\eta \pi^- \nu_\tau$		$< 1.4 \times 10^{-4}$	CL=95%
Γ_{117}	$\eta \pi^- \pi^0 \nu_\tau$	[a]	$(1.74 \pm 0.24) \times 10^{-3}$	
Γ_{118}	$\eta \pi^- \pi^0 \pi^0 \nu_\tau$		$(1.5 \pm 0.5) \times 10^{-4}$	
Γ_{119}	$\eta K^- \nu_\tau$	[a]	$(2.7 \pm 0.6) \times 10^{-4}$	
Γ_{120}	$\eta K^*(892)^- \nu_\tau$		$(2.9 \pm 0.9) \times 10^{-4}$	

Γ_{121}	$\eta K^- \pi^0 \nu_\tau$	$(1.8 \pm 0.9) \times 10^{-4}$	
Γ_{122}	$\eta \bar{K}^0 \pi^- \nu_\tau$	$(2.2 \pm 0.7) \times 10^{-4}$	
Γ_{123}	$\eta \pi^+ \pi^- \pi^- \geq 0$ neutrals ν_τ	$< 3 \times 10^{-3}$	CL=90%
Γ_{124}	$\eta \pi^- \pi^+ \pi^- \nu_\tau$	$(2.3 \pm 0.5) \times 10^{-4}$	
Γ_{125}	$\eta a_1(1260)^- \nu_\tau \rightarrow \eta \pi^- \rho^0 \nu_\tau$	$< 3.9 \times 10^{-4}$	CL=90%
Γ_{126}	$\eta \eta \pi^- \nu_\tau$	$< 1.1 \times 10^{-4}$	CL=95%
Γ_{127}	$\eta \eta \pi^- \pi^0 \nu_\tau$	$< 2.0 \times 10^{-4}$	CL=95%
Γ_{128}	$\eta'(958) \pi^- \nu_\tau$	$< 7.4 \times 10^{-5}$	CL=90%
Γ_{129}	$\eta'(958) \pi^- \pi^0 \nu_\tau$	$< 8.0 \times 10^{-5}$	CL=90%
Γ_{130}	$\phi \pi^- \nu_\tau$	$< 2.0 \times 10^{-4}$	CL=90%
Γ_{131}	$\phi K^- \nu_\tau$	$< 6.7 \times 10^{-5}$	CL=90%
Γ_{132}	$f_1(1285) \pi^- \nu_\tau$	$(5.8 \pm 2.3) \times 10^{-4}$	
Γ_{133}	$f_1(1285) \pi^- \nu_\tau \rightarrow \eta \pi^- \pi^+ \pi^- \nu_\tau$	$(1.3 \pm 0.4) \times 10^{-4}$	
Γ_{134}	$\pi(1300)^- \nu_\tau \rightarrow (\rho \pi)^- \nu_\tau \rightarrow (3\pi)^- \nu_\tau$	$< 1.0 \times 10^{-4}$	CL=90%
Γ_{135}	$\pi(1300)^- \nu_\tau \rightarrow ((\pi \pi)_{S\text{-wave}} \pi)^- \nu_\tau \rightarrow (3\pi)^- \nu_\tau$	$< 1.9 \times 10^{-4}$	CL=90%
Γ_{136}	$h^- \omega \geq 0$ neutrals ν_τ	$(2.37 \pm 0.08) \%$	
Γ_{137}	$h^- \omega \nu_\tau$	[a] $(1.94 \pm 0.07) \%$	
Γ_{138}	$h^- \omega \pi^0 \nu_\tau$	[a] $(4.3 \pm 0.5) \times 10^{-3}$	
Γ_{139}	$h^- \omega 2\pi^0 \nu_\tau$	$(1.4 \pm 0.5) \times 10^{-4}$	
Γ_{140}	$2h^- h^+ \omega \nu_\tau$	$(1.20 \pm 0.22) \times 10^{-4}$	

Lepton Family number (*LF*), Lepton number (*L*), or Baryon number (*B*) violating modes

L means lepton number violation (e.g. $\tau^- \rightarrow e^+ \pi^- \pi^-$). Following common usage, *LF* means lepton family violation *and not* lepton number violation (e.g. $\tau^- \rightarrow e^- \pi^+ \pi^-$). *B* means baryon number violation.

Γ_{141}	$e^- \gamma$	<i>LF</i>	$< 3.9 \times 10^{-7}$	CL=90%
Γ_{142}	$\mu^- \gamma$	<i>LF</i>	$< 3.1 \times 10^{-7}$	CL=90%
Γ_{143}	$e^- \pi^0$	<i>LF</i>	$< 3.7 \times 10^{-6}$	CL=90%
Γ_{144}	$\mu^- \pi^0$	<i>LF</i>	$< 4.0 \times 10^{-6}$	CL=90%
Γ_{145}	$e^- K_S^0$	<i>LF</i>	$< 9.1 \times 10^{-7}$	CL=90%
Γ_{146}	$\mu^- K_S^0$	<i>LF</i>	$< 9.5 \times 10^{-7}$	CL=90%
Γ_{147}	$e^- \eta$	<i>LF</i>	$< 8.2 \times 10^{-6}$	CL=90%
Γ_{148}	$\mu^- \eta$	<i>LF</i>	$< 3.4 \times 10^{-7}$	CL=90%
Γ_{149}	$e^- \rho^0$	<i>LF</i>	$< 2.0 \times 10^{-6}$	CL=90%
Γ_{150}	$\mu^- \rho^0$	<i>LF</i>	$< 6.3 \times 10^{-6}$	CL=90%
Γ_{151}	$e^- K^*(892)^0$	<i>LF</i>	$< 5.1 \times 10^{-6}$	CL=90%
Γ_{152}	$\mu^- K^*(892)^0$	<i>LF</i>	$< 7.5 \times 10^{-6}$	CL=90%
Γ_{153}	$e^- \bar{K}^*(892)^0$	<i>LF</i>	$< 7.4 \times 10^{-6}$	CL=90%
Γ_{154}	$\mu^- \bar{K}^*(892)^0$	<i>LF</i>	$< 7.5 \times 10^{-6}$	CL=90%

Γ_{155}	$e^- \phi$	<i>LF</i>	< 6.9	$\times 10^{-6}$	CL=90%
Γ_{156}	$\mu^- \phi$	<i>LF</i>	< 7.0	$\times 10^{-6}$	CL=90%
Γ_{157}	$e^- e^+ e^-$	<i>LF</i>	< 2.0	$\times 10^{-7}$	CL=90%
Γ_{158}	$e^- \mu^+ \mu^-$	<i>LF</i>	< 2.0	$\times 10^{-7}$	CL=90%
Γ_{159}	$e^+ \mu^- \mu^-$	<i>LF</i>	< 1.3	$\times 10^{-7}$	CL=90%
Γ_{160}	$\mu^- e^+ e^-$	<i>LF</i>	< 1.9	$\times 10^{-7}$	CL=90%
Γ_{161}	$\mu^+ e^- e^-$	<i>LF</i>	< 1.1	$\times 10^{-7}$	CL=90%
Γ_{162}	$\mu^- \mu^+ \mu^-$	<i>LF</i>	< 1.9	$\times 10^{-7}$	CL=90%
Γ_{163}	$e^- \pi^+ \pi^-$	<i>LF</i>	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{164}	$e^+ \pi^- \pi^-$	<i>L</i>	< 1.9	$\times 10^{-6}$	CL=90%
Γ_{165}	$\mu^- \pi^+ \pi^-$	<i>LF</i>	< 8.2	$\times 10^{-6}$	CL=90%
Γ_{166}	$\mu^+ \pi^- \pi^-$	<i>L</i>	< 3.4	$\times 10^{-6}$	CL=90%
Γ_{167}	$e^- \pi^+ K^-$	<i>LF</i>	< 6.4	$\times 10^{-6}$	CL=90%
Γ_{168}	$e^- \pi^- K^+$	<i>LF</i>	< 3.8	$\times 10^{-6}$	CL=90%
Γ_{169}	$e^+ \pi^- K^-$	<i>L</i>	< 2.1	$\times 10^{-6}$	CL=90%
Γ_{170}	$e^- K_S^0 K_S^0$	<i>LF</i>	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{171}	$e^- K^+ K^-$	<i>LF</i>	< 6.0	$\times 10^{-6}$	CL=90%
Γ_{172}	$e^+ K^- K^-$	<i>L</i>	< 3.8	$\times 10^{-6}$	CL=90%
Γ_{173}	$\mu^- \pi^+ K^-$	<i>LF</i>	< 7.5	$\times 10^{-6}$	CL=90%
Γ_{174}	$\mu^- \pi^- K^+$	<i>LF</i>	< 7.4	$\times 10^{-6}$	CL=90%
Γ_{175}	$\mu^+ \pi^- K^-$	<i>L</i>	< 7.0	$\times 10^{-6}$	CL=90%
Γ_{176}	$\mu^- K_S^0 K_S^0$	<i>LF</i>	< 3.4	$\times 10^{-6}$	CL=90%
Γ_{177}	$\mu^- K^+ K^-$	<i>LF</i>	< 1.5	$\times 10^{-5}$	CL=90%
Γ_{178}	$\mu^+ K^- K^-$	<i>L</i>	< 6.0	$\times 10^{-6}$	CL=90%
Γ_{179}	$e^- \pi^0 \pi^0$	<i>LF</i>	< 6.5	$\times 10^{-6}$	CL=90%
Γ_{180}	$\mu^- \pi^0 \pi^0$	<i>LF</i>	< 1.4	$\times 10^{-5}$	CL=90%
Γ_{181}	$e^- \eta \eta$	<i>LF</i>	< 3.5	$\times 10^{-5}$	CL=90%
Γ_{182}	$\mu^- \eta \eta$	<i>LF</i>	< 6.0	$\times 10^{-5}$	CL=90%
Γ_{183}	$e^- \pi^0 \eta$	<i>LF</i>	< 2.4	$\times 10^{-5}$	CL=90%
Γ_{184}	$\mu^- \pi^0 \eta$	<i>LF</i>	< 2.2	$\times 10^{-5}$	CL=90%
Γ_{185}	$\bar{p} \gamma$	<i>L,B</i>	< 3.5	$\times 10^{-6}$	CL=90%
Γ_{186}	$\bar{p} \pi^0$	<i>L,B</i>	< 1.5	$\times 10^{-5}$	CL=90%
Γ_{187}	$\bar{p} 2\pi^0$	<i>L,B</i>	< 3.3	$\times 10^{-5}$	CL=90%
Γ_{188}	$\bar{p} \eta$	<i>L,B</i>	< 8.9	$\times 10^{-6}$	CL=90%
Γ_{189}	$\bar{p} \pi^0 \eta$	<i>L,B</i>	< 2.7	$\times 10^{-5}$	CL=90%
Γ_{190}	e^- light boson	<i>LF</i>	< 2.7	$\times 10^{-3}$	CL=95%
Γ_{191}	μ^- light boson	<i>LF</i>	< 5	$\times 10^{-3}$	CL=95%

[a] Basis mode for the τ .

[b] See the Particle Listings below for the energy limits used in this measurement.

CONSTRAINED FIT INFORMATION

An overall fit to 64 branching ratios uses 129 measurements and one constraint to determine 31 parameters. The overall fit has a $\chi^2 = 63.6$ for 99 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_5	3									
x_9	-5	-5								
x_{10}	0	0	-20							
x_{13}	-13	-14	-25	1						
x_{15}	0	0	1	-2	-20					
x_{19}	-13	-14	-25	0	-36	4				
x_{22}	0	0	1	-2	4	-19	-16			
x_{25}	-8	-8	-15	0	-18	4	-24	4		
x_{26}	0	0	1	-2	4	-18	3	-14	-22	
x_{28}	-4	-4	-7	0	-11	0	-12	0	-7	0
x_{33}	-1	-2	-13	0	-3	1	-8	1	-1	1
x_{35}	0	0	-4	-1	1	-11	-1	-9	2	-8
x_{38}	-2	-2	-3	0	-5	3	-5	2	-9	2
x_{40}	-1	-1	0	-1	1	-14	1	-11	-1	-11
x_{45}	0	0	-2	0	0	0	0	0	0	0
x_{46}	-2	-2	-2	0	-4	1	-4	1	-2	1
x_{60}	-1	-1	-2	0	-4	0	-3	0	-2	0
x_{68}	-1	-1	-3	0	-3	0	-4	0	-2	0
x_{74}	0	0	0	0	0	0	0	0	0	0
x_{75}	0	0	0	0	0	0	0	0	0	0
x_{82}	0	0	0	0	0	0	0	0	0	0
x_{86}	0	0	0	0	0	0	0	0	0	0
x_{89}	0	0	0	0	0	0	0	0	0	0
x_{90}	0	0	0	0	0	0	0	0	0	0
x_{97}	0	0	-1	0	-1	0	-1	0	0	0
x_{98}	0	0	0	0	0	0	0	0	0	0
x_{117}	-1	-1	-1	0	-2	0	-2	0	-1	0
x_{119}	0	0	0	0	0	-3	0	-3	-2	-2
x_{137}	-1	-1	-2	0	-3	0	-3	0	-1	0
x_{138}	-1	-1	-1	0	-2	0	-2	0	-1	0
	x_3	x_5	x_9	x_{10}	x_{13}	x_{15}	x_{19}	x_{22}	x_{25}	x_{26}

x33	-1									
x35	0	-5								
x38	-1	-7	0							
x40	0	-2	-16	-19						
x45	0	-2	-1	-2	0					
x46	-1	-12	-4	-10	-3	-3				
x60	-1	-7	-3	3	2	0	0			
x68	-1	4	2	-5	-2	0	0	-46		
x74	2	1	0	1	0	0	0	-8	-7	
x75	0	0	0	0	0	0	0	-3	-3	0
x82	0	-1	0	0	0	0	0	-32	-3	-1
x86	0	0	0	0	0	0	0	-3	-15	0
x89	0	0	0	0	0	0	0	-3	0	0
x90	0	0	0	0	0	0	0	-3	0	-1
x97	0	0	0	0	0	0	0	-1	-1	0
x98	0	0	0	0	0	0	0	0	0	0
x117	-14	0	0	0	0	0	0	-1	-1	-14
x119	0	0	-1	0	-2	0	0	0	0	0
x137	-1	1	1	-3	-1	0	0	-22	-29	-3
x138	-1	1	0	1	0	0	0	-9	-9	-44
	x28	x33	x35	x38	x40	x45	x46	x60	x68	x74

x82	0									
x86	0	0								
x89	0	-11	0							
x90	0	0	-49	0						
x97	0	0	0	0	0					
x98	0	0	0	0	0	-19				
x117	0	0	0	0	0	0	0			
x119	0	0	-6	0	0	0	0	0		
x137	-1	-2	1	0	2	0	0	0	0	
x138	-1	-1	0	0	-1	0	0	0	0	-4
	x75	x82	x86	x89	x90	x97	x98	x117	x119	x137

A REVIEW GOES HERE – Check our WWW List of Reviews

τ^- BRANCHING RATIOS

$$\Gamma(\text{particle}^- \geq 0 \text{ neutrals} \geq 0 K^0 \nu_\tau \text{ ("1-prong")}) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$


$$\Gamma_1 / \Gamma = (\Gamma_3 + \Gamma_5 + \Gamma_9 + \Gamma_{10} + \Gamma_{13} + \Gamma_{15} + \Gamma_{19} + \Gamma_{22} + \Gamma_{25} + \Gamma_{26} + \Gamma_{28} + \Gamma_{33} + \Gamma_{35} + \Gamma_{38} + \Gamma_{40} + 2\Gamma_{45} + \Gamma_{46} + 0.708\Gamma_{117} + 0.715\Gamma_{119} + 0.09\Gamma_{137} + 0.09\Gamma_{138}) / \Gamma$$

The charged particle here can be e , μ , or hadron. In many analyses, the sum of the topological branching fractions (1, 3, and 5 prongs) is constrained to be unity. Since the 5-prong fraction is very small, the measured 1-prong and 3-prong fractions are highly correlated and cannot be treated as independent quantities in our overall fit. We arbitrarily choose to use the 3-prong fraction in our fit, and leave the 1-prong fraction out. We do, however, use these 1-prong measurements in our average below. The measurements used only for the average are marked "avg," whereas "f&a" marks a result used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
85.35 ± 0.07	OUR FIT				Error includes scale factor of 1.1.
85.26 ± 0.13	OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
85.316 ± 0.093 ± 0.049	avg	78k	³² ABREU	01M DLPH	1992–1995 LEP runs
85.274 ± 0.105 ± 0.073	avg		³³ ACHARD	01D L3	1992–1995 LEP runs
84.48 ± 0.27 ± 0.23	avg		ACTON	92H OPAL	1990–1991 LEP runs
85.45 $\begin{smallmatrix} +0.69 \\ -0.73 \end{smallmatrix}$ ± 0.65	f&a		DECAMP	92C ALEP	1989–1990 LEP runs

³² The correlation coefficients between this measurement and the ABREU 01M measurements of $B(\tau \rightarrow 3\text{-prong})$ and $B(\tau \rightarrow 5\text{-prong})$ are -0.98 and -0.08 respectively.

³³ The correlation coefficients between this measurement and the ACHARD 01D measurements of $B(\tau \rightarrow "3\text{-prong}")$ and $B(\tau \rightarrow "5\text{-prong}")$ are -0.978 and -0.082 respectively.



$$\Gamma(\text{particle}^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_2 / \Gamma$$

$$\Gamma_2 / \Gamma = (\Gamma_3 + \Gamma_5 + \Gamma_9 + \Gamma_{10} + \Gamma_{13} + \Gamma_{15} + \Gamma_{19} + \Gamma_{22} + \Gamma_{25} + \Gamma_{26} + \Gamma_{28} + 0.6569\Gamma_{33} + 0.6569\Gamma_{35} + 0.6569\Gamma_{38} + 0.6569\Gamma_{40} + 1.0985\Gamma_{45} + 0.3139\Gamma_{46} + 0.708\Gamma_{117} + 0.715\Gamma_{119} + 0.09\Gamma_{137} + 0.09\Gamma_{138}) / \Gamma$$

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

84.72 ± 0.07 OUR FIT Error includes scale factor of 1.1.

85.1 ± 0.4 OUR AVERAGE

85.6 ± 0.6 ± 0.3	avg	3300	³⁴ ADEVA	91F L3	$E_{\text{cm}}^{ee} = 88.3\text{--}94.3 \text{ GeV}$
84.9 ± 0.4 ± 0.3	avg		BEHREND	89B CELL	$E_{\text{cm}}^{ee} = 14\text{--}47 \text{ GeV}$
84.7 ± 0.8 ± 0.6	avg		³⁵ AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
86.4 ± 0.3 ± 0.3			ABACHI	89B HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
87.1 ± 1.0 ± 0.7			³⁶ BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
87.2 ± 0.5 ± 0.8			SCHMIDKE	86 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
84.7 ± 1.1 $\begin{smallmatrix} +1.6 \\ -1.3 \end{smallmatrix}$		169	³⁷ ALTHOFF	85 TASS	$E_{\text{cm}}^{ee} = 34.5 \text{ GeV}$
86.1 ± 0.5 ± 0.9			BARTEL	85F JADE	$E_{\text{cm}}^{ee} = 34.6 \text{ GeV}$
87.8 ± 1.3 ± 3.9			³⁸ BERGER	85 PLUT	$E_{\text{cm}}^{ee} = 34.6 \text{ GeV}$
86.7 ± 0.3 ± 0.6			FERNANDEZ	85 MAC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

³⁴ Not independent of ADEVA 91F $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ value.

³⁵ Not independent of AIHARA 87B $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}$, and $\Gamma(h^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ values.

³⁶ Not independent of SCHMIDKE 86 value (also not independent of BURCHAT 87 value for $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$).

³⁷ Not independent of ALTHOFF 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}$, $\Gamma(h^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$, and $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ values.

³⁸ Not independent of (1-prong + $0\pi^0$) and (1-prong + $\geq 1\pi^0$) values.

$$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_3 / \Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

To minimize the effect of experiments with large systematic errors, we exclude experiments which together would contribute 5% of the weight in the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

17.36 ± 0.06 OUR FIT

17.33 ± 0.06 OUR AVERAGE

17.34 ± 0.09 ± 0.06	f&a	31.4k	ABBIENDI	03 OPAL	1990–1995 LEP runs
17.342 ± 0.110 ± 0.067	f&a	21.5k	³⁹ ACCIARRI	01F L3	1991–1995 LEP runs
17.325 ± 0.095 ± 0.077	f&a	27.7k	ABREU	99X DLPH	1991–1995 LEP runs
17.37 ± 0.08 ± 0.18	avg		⁴⁰ ANASTASSOV	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
17.31 ± 0.11 ± 0.05	f&a	20.7k	BUSKULIC	96C ALEP	1991–1993 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

17.02 ±0.19 ±0.24	6586	ABREU	95T DLPH	Repl. by ABREU 99X
17.36 ±0.27	7941	AKERS	95I OPAL	Repl. by ABBI-ENDI 03
17.6 ±0.4 ±0.4	2148	ADRIANI	93M L3	Repl. by ACCIARRI 01F
17.4 ±0.3 ±0.5		⁴¹ ALBRECHT	93G ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV
17.35 ±0.41 ±0.37	f&a	DECAMP	92C ALEP	1989–1990 LEP runs
17.7 ±0.8 ±0.4	568	BEHREND	90 CELL	$E_{cm}^{ee} = 35$ GeV
17.4 ±1.0	2197	ADEVA	88 MRKJ	$E_{cm}^{ee} = 14\text{--}16$ GeV
17.7 ±1.2 ±0.7		AIHARA	87B TPC	$E_{cm}^{ee} = 29$ GeV
18.3 ±0.9 ±0.8		BURCHAT	87 MRK2	$E_{cm}^{ee} = 29$ GeV
18.6 ±0.8 ±0.7	558	⁴² BARTEL	86D JADE	$E_{cm}^{ee} = 34.6$ GeV
12.9 ±1.7 $\begin{smallmatrix} +0.7 \\ -0.5 \end{smallmatrix}$		ALTHOFF	85 TASS	$E_{cm}^{ee} = 34.5$ GeV
18.0 ±0.9 ±0.5	473	⁴² ASH	85B MAC	$E_{cm}^{ee} = 29$ GeV
18.0 ±1.0 ±0.6		⁴³ BALTRUSAITIS	85 MRK3	$E_{cm}^{ee} = 3.77$ GeV
19.4 ±1.6 ±1.7	153	BERGER	85 PLUT	$E_{cm}^{ee} = 34.6$ GeV
17.6 ±2.6 ±2.1	47	BEHREND	83C CELL	$E_{cm}^{ee} = 34$ GeV
17.8 ±2.0 ±1.8		BERGER	81B PLUT	$E_{cm}^{ee} = 9\text{--}32$ GeV

³⁹ The correlation coefficient between this measurement and the ACCIARRI 01F measurement of $B(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)$ is 0.08.

⁴⁰ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(e\bar{\nu}_e \nu_\tau)$, $B(\mu\bar{\nu}_\mu \nu_\tau)/B(e\bar{\nu}_e \nu_\tau)$, $B(h^- \nu_\tau)$, and $B(h^- \nu_\tau)/B(e\bar{\nu}_e \nu_\tau)$ are 0.50, 0.58, 0.50, and 0.08 respectively.

⁴¹ Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ and ALBRECHT 93G $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{total}^2$ values.

⁴² Modified using $B(e^- \bar{\nu}_e \nu_\tau)/B(\text{"1 prong"})$ and $B(\text{"1 prong"}) = 0.855$.

⁴³ Error correlated with BALTRUSAITIS 85 $e\nu\bar{\nu}$ value.

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma_{total}$					Γ_4/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.361 ± 0.016 ± 0.035		44 BERGFELD	00 CLEO	$E_{cm}^{ee} = 10.6$ GeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.30 ±0.04 ±0.05	116	⁴⁵ ALEXANDER	96S OPAL	1991–1994 LEP runs
0.23 ±0.10	10	⁴⁶ WU	90 MRK2	$E_{cm}^{ee} = 29$ GeV

⁴⁴ BERGFELD 00 impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_\gamma^* > 10$ MeV. For $E_\gamma^* > 20$ MeV, they quote $(3.04 \pm 0.14 \pm 0.30) \times 10^{-3}$.

⁴⁵ ALEXANDER 96S impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_\gamma > 20$ MeV.

⁴⁶ WU 90 reports $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) = 0.013 \pm 0.006$, which is converted to $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma_{total}$ using $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma_{total} = 17.35\%$. Requirements on detected γ 's correspond to a τ rest frame energy cutoff $E_\gamma > 37$ MeV.

$\Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}$

Γ_5 / Γ

To minimize the effect of experiments with large systematic errors, we exclude experiments which together would contribute 5% of the weight in the average.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
17.84 ± 0.06				OUR FIT
17.81 ± 0.06				OUR AVERAGE
17.806 ± 0.104 ± 0.076	24.7k	⁴⁷ ACCIARRI	01F L3	1991–1995 LEP runs
17.81 ± 0.09 ± 0.06	33.1k	ABBIENDI	99H OPAL	1991–1995 LEP runs
17.877 ± 0.109 ± 0.110	23.3k	ABREU	99X DLPH	1991–1995 LEP runs
17.76 ± 0.06 ± 0.17		⁴⁸ ANASTASSOV	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
17.79 ± 0.12 ± 0.06	20.6k	BUSKULIC	96C ALEP	1991–1993 LEP runs
18.09 ± 0.45 ± 0.45		DECAMP	92C ALEP	1989–1990 LEP runs
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
17.78 ± 0.10 ± 0.09	25.3k	ALEXANDER	96D OPAL	Repl. by ABBI- ENDI 99H
17.51 ± 0.23 ± 0.31	5059	ABREU	95T DLPH	Repl. by ABREU 99X
17.9 ± 0.4 ± 0.4	2892	ADRIANI	93M L3	Repl. by ACCIA- RRI 01F
17.5 ± 0.3 ± 0.5		⁴⁹ ALBRECHT	93G ARG	$E_{\text{cm}}^{ee} = 9.4$ – 10.6 GeV
17.97 ± 0.14 ± 0.23	3970	AKERIB	92 CLEO	Repl. by ANAS- TASSOV 97
19.1 ± 0.4 ± 0.6	2960	⁵⁰ AMMAR	92 CLEO	$E_{\text{cm}}^{ee} = 10.5$ – 10.9 GeV
17.0 ± 0.5 ± 0.6	1.7k	ABACHI	90 HRS	$E_{\text{cm}}^{ee} = 29$ GeV
18.4 ± 0.8 ± 0.4	644	BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35$ GeV
16.3 ± 0.3 ± 3.2		JANSSEN	89 CBAL	$E_{\text{cm}}^{ee} = 9.4$ – 10.6 GeV
18.4 ± 1.2 ± 1.0		AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29$ GeV
19.1 ± 0.8 ± 1.1		BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
16.8 ± 0.7 ± 0.9	515	⁵⁰ BARTEL	86D JADE	$E_{\text{cm}}^{ee} = 34.6$ GeV
20.4 ± 3.0 +1.4 -0.9		ALTHOFF	85 TASS	$E_{\text{cm}}^{ee} = 34.5$ GeV
17.8 ± 0.9 ± 0.6	390	⁵⁰ ASH	85B MAC	$E_{\text{cm}}^{ee} = 29$ GeV
18.2 ± 0.7 ± 0.5		⁵¹ BALTRUSAITIS	85 MRK3	$E_{\text{cm}}^{ee} = 3.77$ GeV
13.0 ± 1.9 ± 2.9		BERGER	85 PLUT	$E_{\text{cm}}^{ee} = 34.6$ GeV
18.3 ± 2.4 ± 1.9	60	BEHREND	83C CELL	$E_{\text{cm}}^{ee} = 34$ GeV
16.0 ± 1.3	459	⁵² BACINO	78B DLCO	$E_{\text{cm}}^{ee} = 3.1$ – 7.4 GeV

⁴⁷ The correlation coefficient between this measurement and the ACCIARRI 01F measurement of $B(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)$ is 0.08.

⁴⁸ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu^- \bar{\nu}_\mu \nu_\tau)$, $B(\mu^- \bar{\nu}_\mu \nu_\tau) / B(e^- \bar{\nu}_e \nu_\tau)$, $B(h^- \nu_\tau)$, and $B(h^- \nu_\tau) / B(e^- \bar{\nu}_e \nu_\tau)$ are 0.50, -0.42, 0.48, and -0.39 respectively.

⁴⁹ Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma(e^- \bar{\nu}_e \nu_\tau)$ and ALBRECHT 93G $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}^2$ values.

⁵⁰ Modified using $B(e^- \bar{\nu}_e \nu_\tau) / B(\text{"1 prong"})$ and $B(\text{"1 prong"})$, = 0.855.

⁵¹ Error correlated with BALTRUSAITIS 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma_{\text{total}}$.

⁵² BACINO 78B value comes from fit to events with e^\pm and one other nonelectron charged prong.

$\Gamma(e^- \bar{\nu}_e \nu_\tau \gamma) / \Gamma_{\text{total}}$ Γ_6 / Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
1.75 ± 0.06 ± 0.17	⁵³ BERGFELD	00	CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

⁵³ BERGFELD 00 impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_\gamma^* > 10$ MeV.

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma(e^- \bar{\nu}_e \nu_\tau)$ Γ_3 / Γ_5

Standard Model prediction including mass effects is 0.9726.

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE	DOCUMENT ID	TECN	COMMENT
0.974 ± 0.004 OUR FIT			
0.978 ± 0.011 OUR AVERAGE			

0.9777 ± 0.0063 ± 0.0087 f&a ⁵⁴ ANASTASSOV 97 CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

0.997 ± 0.035 ± 0.040 f&a ALBRECHT 92D ARG $E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV

⁵⁴ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu^- \bar{\nu}_\mu \nu_\tau)$, $B(e^- \bar{\nu}_e \nu_\tau)$, $B(h^- \nu_\tau)$, and $B(h^- \nu_\tau) / B(e^- \bar{\nu}_e \nu_\tau)$ are 0.58, -0.42, 0.07, and 0.45 respectively.

$\Gamma(h^- \geq 0K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_7 / Γ

$$\Gamma_7 / \Gamma = (\Gamma_9 + \Gamma_{10} + \frac{1}{2}\Gamma_{33} + \frac{1}{2}\Gamma_{35} + \Gamma_{45}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
12.30 ± 0.11 OUR FIT				Error includes scale factor of 1.4.
12.44 ± 0.14 OUR AVERAGE				

12.44 ± 0.11 ± 0.11	f&a	15k	⁵⁵ BUSKULIC	96	ALEP	1991–1993 LEP run
12.47 ± 0.26 ± 0.43	f&a	2967	⁵⁶ ACCIARRI	95	L3	1992 LEP run
12.4 ± 0.7 ± 0.7	f&a	283	⁵⁷ ABREU	92N	DLPH	1990 LEP run
12.98 ± 0.44 ± 0.33	f&a		⁵⁸ DECAMP	92C	ALEP	1989–1990 LEP runs
12.1 ± 0.7 ± 0.5	f&a	309	ALEXANDER	91D	OPAL	1990 LEP run
11.3 ± 0.5 ± 0.8	avg	798	⁵⁹ FORD	87	MAC	$E_{\text{cm}}^{ee} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.7 ± 0.6 ± 0.8			⁶⁰ ALBRECHT	92D	ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
12.3 ± 0.9 ± 0.5	1338		BEHREND	90	CELL	$E_{\text{cm}}^{ee} = 35$ GeV
11.1 ± 1.1 ± 1.4			⁶¹ BURCHAT	87	MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
12.3 ± 0.6 ± 1.1	328		⁶² BARTEL	86D	JADE	$E_{\text{cm}}^{ee} = 34.6$ GeV
13.0 ± 2.0 ± 4.0			BERGER	85	PLUT	$E_{\text{cm}}^{ee} = 34.6$ GeV
11.2 ± 1.7 ± 1.2	34		⁶³ BEHREND	83C	CELL	$E_{\text{cm}}^{ee} = 34$ GeV

- ⁵⁵ BUSKULIC 96 quote $11.78 \pm 0.11 \pm 0.13$ We add 0.66 to undo their correction for unseen K_L^0 and modify the systematic error accordingly.
- ⁵⁶ ACCIARRI 95 with 0.65% added to remove their correction for $\pi^- K_L^0$ backgrounds.
- ⁵⁷ ABREU 92N with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.
- ⁵⁸ DECAMP 92C quote $B(h^- \geq 0 K_L^0 \geq 0 (K_S^0 \rightarrow \pi^+ \pi^-) \nu_\tau) = 13.32 \pm 0.44 \pm 0.33$.
We subtract 0.35 to correct for their inclusion of the K_S^0 decays.
- ⁵⁹ FORD 87 result for $B(\pi^- \nu_\tau)$ with 0.67% added to remove their K^- correction and adjusted for 1992 B("1 prong").
- ⁶⁰ Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$, $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)$, and $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ values.
- ⁶¹ BURCHAT 87 with 1.1% added to remove their correction for K^- and $K^*(892)^-$ backgrounds.
- ⁶² BARTEL 86D result for $B(\pi^- \nu_\tau)$ with 0.59% added to remove their K^- correction and adjusted for 1992 B("1 prong").
- ⁶³ BEHREND 83C quote $B(\pi^- \nu_\tau) = 9.9 \pm 1.7 \pm 1.3$ after subtracting 1.3 ± 0.5 to correct for $B(K^- \nu_\tau)$.

$\Gamma(h^- \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10})/\Gamma$
Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		DOCUMENT ID	TECN	COMMENT
11.75±0.11 OUR FIT		Error includes scale factor of 1.4.		
11.65±0.21 OUR AVERAGE		Error includes scale factor of 1.9.		
11.98±0.13±0.16	f&a	ACKERSTAFF 98M	OPAL	1991–1995 LEP runs
11.52±0.05±0.12	f&a	⁶⁴ ANASTASSOV 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

⁶⁴ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu^- \bar{\nu}_\mu \nu_\tau)$, $B(e^- \bar{\nu}_e \nu_\tau)$, $B(\mu^- \bar{\nu}_\mu \nu_\tau)/B(e^- \bar{\nu}_e \nu_\tau)$, and $B(h^- \nu_\tau)/B(e^- \bar{\nu}_e \nu_\tau)$ are 0.50, 0.48, 0.07, and 0.63 respectively.

$\Gamma(h^- \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ $\Gamma_8/\Gamma_5 = (\Gamma_9 + \Gamma_{10})/\Gamma_5$
Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE		DOCUMENT ID	TECN	COMMENT
0.659 ±0.007 OUR FIT		Error includes scale factor of 1.4.		
0.6484±0.0041±0.0060 avg		⁶⁵ ANASTASSOV 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

⁶⁵ The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu^- \bar{\nu}_\mu \nu_\tau)$, $B(e^- \bar{\nu}_e \nu_\tau)$, $B(\mu^- \bar{\nu}_\mu \nu_\tau)/B(e^- \bar{\nu}_e \nu_\tau)$, and $B(h^- \nu_\tau)$ are 0.08, -0.39, 0.45, and 0.63 respectively.

$\Gamma(\pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_9/Γ
Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
11.06±0.11 OUR FIT		Error includes scale factor of 1.4.			
11.07±0.18 OUR AVERAGE					
11.06±0.11±0.14	avg	⁶⁶ BUSKULIC	96	ALEP	LEP 1991–1993 data
11.7 ±0.4 ±1.8	f&a	1138	BLOCKER	82D MRK2	$E_{\text{cm}}^{ee} = 3.5\text{--}6.7$ GeV

⁶⁶ Not independent of BUSKULIC 96 $B(h^- \nu_\tau)$ and $B(K^- \nu_\tau)$ values.

$\Gamma(K^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.686±0.023 OUR FIT

0.685±0.023 OUR AVERAGE

0.658±0.027±0.029 ⁶⁷ ABBIENDI 01J OPAL 1990–1995 LEP runs

0.696±0.025±0.014 2032 BARATE 99K ALEP 1991–1995 LEP runs

0.85 ±0.18 27 ABREU 94K DLPH LEP 1992 Z data

0.66 ±0.07 ±0.09 99 BATTLE 94 CLEO $E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.72 ±0.04 ±0.04 728 BUSKULIC 96 ALEP Repl. by
BARATE 99K

0.59 ±0.18 16 MILLS 84 DLCO $E_{\text{cm}}^{ee} = 29$ GeV

1.3 ±0.5 15 BLOCKER 82B MRK2 $E_{\text{cm}}^{ee} = 3.9$ –6.7 GeV

⁶⁷ The correlation coefficient between this measurement and the ABBIENDI 01J $B(\tau^- \rightarrow K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma \nu_\tau)$ is 0.60.

$\Gamma(h^- \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{11}/Γ

$$\Gamma_{11}/\Gamma = (\Gamma_{13} + \Gamma_{15} + \Gamma_{19} + \Gamma_{22} + \Gamma_{25} + \Gamma_{26} + \Gamma_{28} + 0.157\Gamma_{33} + 0.157\Gamma_{35} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.0985\Gamma_{45} + 0.708\Gamma_{117} + 0.715\Gamma_{119} + 0.09\Gamma_{137} + 0.09\Gamma_{138})/\Gamma$$

VALUE (%) DOCUMENT ID TECN COMMENT

36.92±0.14 OUR FIT Error includes scale factor of 1.1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.14±0.33±0.58 ⁶⁸ AKERS 94E OPAL 1991–1992 LEP runs

38.4 ±1.2 ±1.0 ⁶⁹ BURCHAT 87 MRK2 $E_{\text{cm}}^{ee} = 29$ GeV

42.7 ±2.0 ±2.9 BERGER 85 PLUT $E_{\text{cm}}^{ee} = 34.6$ GeV

⁶⁸ Not independent of ACKERSTAFF 98M $B(h^- \pi^0 \nu_\tau)$ and $B(h^- \geq 2\pi^0 \nu_\tau)$ values.

⁶⁹ BURCHAT 87 quote for $B(\pi^\pm \geq 1 \text{ neutral } \nu_\tau) = 0.378 \pm 0.012 \pm 0.010$. We add 0.006 to account for contribution from $(K^{*-} \nu_\tau)$ which they fixed at BR = 0.013.

$\Gamma(h^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma = (\Gamma_{13} + \Gamma_{15})/\Gamma$

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

25.87±0.13 OUR FIT Error includes scale factor of 1.1.

25.76±0.15 OUR AVERAGE

25.89±0.17±0.29 ACKERSTAFF 98M OPAL 1991–1995 LEP runs

25.76±0.15±0.13 31k BUSKULIC 96 ALEP LEP 1991–1993 data

25.05±0.35±0.50 6613 ACCIARRI 95 L3 1992 LEP run

25.87±0.12±0.42 51k ⁷⁰ ARTUSO 94 CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

25.98±0.36±0.52 ⁷¹ AKERS 94E OPAL Repl. by ACKER-
STAFF 98M

22.9 ±0.8 ±1.3 283 ⁷² ABREU 92N DLPH $E_{\text{cm}}^{ee} = 88.2$ –94.2 GeV

23.1 ±0.4 ±0.9 1249 ⁷³ ALBRECHT 92Q ARG $E_{\text{cm}}^{ee} = 10$ GeV

25.02±0.64±0.88 1849 DECAMP 92C ALEP 1989–1990 LEP runs

22.0 ±0.8 ±1.9 779 ANTREASYAN 91 CBAL $E_{\text{cm}}^{ee} = 9.4$ –10.6 GeV

22.6 ±1.5 ±0.7 1101 BEHREND 90 CELL $E_{\text{cm}}^{ee} = 35$ GeV

23.1 ±1.9 ±1.6 BEHREND 84 CELL $E_{\text{cm}}^{ee} = 14,22$ GeV

⁷⁰ ARTUSO 94 reports the combined result from three independent methods, one of which (23% of the $\tau^- \rightarrow h^- \pi^0 \nu_\tau$) is normalized to the inclusive one-prong branching fraction, taken as 0.854 ± 0.004 . Renormalization to the present value causes negligible change.

⁷¹ AKERS 94E quote $(26.25 \pm 0.36 \pm 0.52) \times 10^{-2}$; we subtract 0.27% from their number to correct for $\tau^- \rightarrow h^- K_L^0 \nu_\tau$.

⁷² ABREU 92N with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.

⁷³ ALBRECHT 92Q with 0.5% added to remove their correction for $\tau^- \rightarrow K^*(892)^- \nu_\tau$ background.

$\Gamma(\pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{13}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

25.41 ± 0.13 OUR FIT Error includes scale factor of 1.1.

25.31 ± 0.18 OUR AVERAGE

25.30 ± 0.15 ± 0.13	avg	⁷⁴ BUSKULIC	96 ALEP	LEP 1991–1993
---------------------	-----	------------------------	---------	---------------

25.36 ± 0.44	avg	⁷⁵ ARTUSO	94 CLEO	data $E_{\text{cm}}^{ee} = 10.6$ GeV
--------------	-----	----------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

21.5 ± 0.4 ± 1.9	4400	^{76,77} ALBRECHT	88L ARG	$E_{\text{cm}}^{ee} = 10$ GeV
------------------	------	---------------------------	---------	-------------------------------

23.0 ± 1.3 ± 1.7	582	ADLER	87B MRK3	$E_{\text{cm}}^{ee} = 3.77$ GeV
------------------	-----	-------	----------	---------------------------------

25.8 ± 1.7 ± 2.5		⁷⁸ BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
------------------	--	-----------------------	---------	-------------------------------

22.3 ± 0.6 ± 1.4	629	⁷⁷ YELTON	86 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
------------------	-----	----------------------	---------	-------------------------------

⁷⁴ Not independent of BUSKULIC 96 B($h^- \pi^0 \nu_\tau$) and B($K^- \pi^0 \nu_\tau$) values.

⁷⁵ Not independent of ARTUSO 94 B($h^- \pi^0 \nu_\tau$) and BATTLE 94 B($K^- \pi^0 \nu_\tau$) values.

⁷⁶ The authors divide by $(\Gamma_3 + \Gamma_5 + \Gamma_9 + \Gamma_{10})/\Gamma = 0.467$ to obtain this result.

⁷⁷ Experiment had no hadron identification. Kaon corrections were made, but insufficient information is given to permit their removal.

⁷⁸ BURCHAT 87 value is not independent of YELTON 86 value. Nonresonant decays included.

$\Gamma(\pi^- \pi^0 \text{non-}\rho(770) \nu_\tau)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

0.3 ± 0.1 ± 0.3	⁷⁹ BEHREND	84 CELL	$E_{\text{cm}}^{ee} = 14,22$ GeV
------------------------	-----------------------	---------	----------------------------------

⁷⁹ BEHREND 84 assume a flat nonresonant mass distribution down to the $\rho(770)$ mass, using events with mass above 1300 to set the level.

$\Gamma(K^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.454 ± 0.027 OUR FIT

0.454 ± 0.030 OUR AVERAGE

0.471 ± 0.059 ± 0.023	360	ABBIENDI	04J OPAL	1991-1995 LEP runs
-----------------------	-----	----------	----------	--------------------

0.444 ± 0.026 ± 0.024	923	BARATE	99K ALEP	1991-1995 LEP runs
-----------------------	-----	--------	----------	--------------------

0.51 ± 0.10 ± 0.07	37	BATTLE	94 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
--------------------	----	--------	---------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.52 ± 0.04 ± 0.05	395	BUSKULIC	96 ALEP	Repl. by BARATE 99K
--------------------	-----	----------	---------	---------------------

$$\Gamma(h^- \geq 2\pi^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{16} / \Gamma$$

$$\Gamma_{16} / \Gamma = (\Gamma_{19} + \Gamma_{22} + \Gamma_{25} + \Gamma_{26} + \Gamma_{28} + 0.157\Gamma_{33} + 0.157\Gamma_{35} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.0985\Gamma_{45} + 0.319\Gamma_{117} + 0.322\Gamma_{119}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

10.76 ± 0.15 OUR FIT Error includes scale factor of 1.1.

10.0 ± 0.4 OUR AVERAGE

9.91 ± 0.31 ± 0.27	f&a	ACKERSTAFF	98M OPAL	1991–1995 LEP runs
--------------------	-----	------------	----------	--------------------

12.0 ± 1.4 ± 2.5	f&a	⁸⁰ BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
------------------	-----	-----------------------	---------	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.89 ± 0.34 ± 0.55		⁸¹ AKERS	94E OPAL	Repl. by ACKERSTAFF 98M
--------------------	--	---------------------	----------	-------------------------

14.0 ± 1.2 ± 0.6	938	⁸² BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35$ GeV
------------------	-----	-----------------------	---------	-------------------------------

13.9 ± 2.0 $\begin{smallmatrix} +1.9 \\ -2.2 \end{smallmatrix}$		⁸³ AIHARA	86E TPC	$E_{\text{cm}}^{ee} = 29$ GeV
---	--	----------------------	---------	-------------------------------

⁸⁰ Error correlated with BURCHAT 87 $\Gamma(\rho^- \nu_e) / \Gamma(\text{total})$ value.

⁸¹ AKERS 94E not independent of AKERS 94E $B(h^- \geq 1\pi^0 \nu_\tau)$ and $B(h^- \pi^0 \nu_\tau)$ measurements.

⁸² No independent of BEHREND 90 $\Gamma(h^- 2\pi^0 \nu_\tau (\text{exp. } K^0))$ and $\Gamma(h^- \geq 3\pi^0 \nu_\tau)$.

⁸³ AIHARA 86E (TPC) quote $B(2\pi^0 \pi^- \nu_\tau) + 1.6B(3\pi^0 \pi^- \nu_\tau) + 1.1B(\pi^0 \eta \pi^- \nu_\tau)$.

$$\Gamma(h^- 2\pi^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{17} / \Gamma$$

$$\Gamma_{17} / \Gamma = (\Gamma_{19} + \Gamma_{22} + 0.157\Gamma_{33} + 0.157\Gamma_{35}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

9.39 ± 0.14 OUR FIT Error includes scale factor of 1.1.

9.48 ± 0.13 ± 0.10	12k	⁸⁴ BUSKULIC	96 ALEP	LEP 1991–1993 data
---------------------------	-----	------------------------	---------	--------------------

⁸⁴ BUSKULIC 96 quote $9.29 \pm 0.13 \pm 0.10$. We add 0.19 to undo their correction for $\tau^- \rightarrow h^- K^0 \nu_\tau$.

$$\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \qquad \Gamma_{18} / \Gamma$$

$$\Gamma_{18} / \Gamma = (\Gamma_{19} + \Gamma_{22}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. f&a marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

9.23 ± 0.14 OUR FIT Error includes scale factor of 1.1.

9.08 ± 0.34 OUR AVERAGE

8.88 ± 0.37 ± 0.42	f&a	1060	ACCIARRI	95 L3	1992 LEP run
--------------------	-----	------	----------	-------	--------------

8.96 ± 0.16 ± 0.44	avg		⁸⁵ PROCARIO	93 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
--------------------	-----	--	------------------------	---------	---------------------------------------

10.38 ± 0.66 ± 0.82	f&a	809	⁸⁶ DECAMP	92C ALEP	1989–1990 LEP runs
---------------------	-----	-----	----------------------	----------	--------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.7 ± 0.5 $\begin{smallmatrix} +1.7 \\ -1.0 \end{smallmatrix}$	133	⁸⁷ ANTREASYAN	91 CBAL	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
--	-----	--------------------------	---------	---

10.0 ± 1.5 ± 1.1	333	⁸⁸ BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35$ GeV
------------------	-----	-----------------------	---------	-------------------------------

8.7 ± 0.4 ± 1.1	815	⁸⁹ BAND	87 MAC	$E_{\text{cm}}^{ee} = 29$ GeV
-----------------	-----	--------------------	--------	-------------------------------

6.2 ± 0.6 ± 1.2		⁹⁰ GAN	87 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
-----------------	--	-------------------	---------	-------------------------------

6.0 ± 3.0 ± 1.8		BEHREND	84 CELL	$E_{\text{cm}}^{ee} = 14,22$ GeV
-----------------	--	---------	---------	----------------------------------

⁸⁵ PROCARIO 93 entry is obtained from $B(h^- 2\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.

⁸⁶ We subtract 0.0015 to account for $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

⁸⁷ ANTREASYAN 91 subtract 0.001 to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

⁸⁸ BEHREND 90 subtract 0.002 to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

⁸⁹ BAND 87 assume $B(\pi^- 3\pi^0 \nu_\tau) = 0.01$ and $B(\pi^- \pi^0 \eta \nu_\tau) = 0.005$.

⁹⁰ GAN 87 analysis use photon multiplicity distribution.

$$\frac{\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))}{\Gamma(h^- \pi^0 \nu_\tau)} \quad \Gamma_{18}/\Gamma_{12}$$

$$\Gamma_{18}/\Gamma_{12} = (\Gamma_{19} + \Gamma_{22}) / (\Gamma_{13} + \Gamma_{15})$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.357 ± 0.006 OUR FIT			Error includes scale factor of 1.1.
0.342 ± 0.006 ± 0.016	⁹¹ PROCARIO 93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

⁹¹ PROCARIO 93 quote $0.345 \pm 0.006 \pm 0.016$ after correction for 2 kaon backgrounds assuming $B(K^{*-} \nu_\tau) = 1.42 \pm 0.18\%$ and $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$. We multiply by 0.990 ± 0.010 to remove these corrections to $B(h^- \pi^0 \nu_\tau)$.

$$\frac{\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))}{\Gamma_{\text{total}}} \quad \Gamma_{19}/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
9.17 ± 0.14 OUR FIT			Error includes scale factor of 1.1.
9.21 ± 0.13 ± 0.11	⁹² BUSKULIC 96	ALEP	LEP 1991–1993 data

⁹² Not independent of BUSKULIC 96 $B(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ and $B(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ values.

$$\frac{\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0), \text{ scalar})}{\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))} \quad \Gamma_{20}/\Gamma_{19}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.094	95	⁹³ BROWDER 00	CLEO	$4.7 \text{ fb}^{-1} E_{\text{cm}}^{ee} = 10.6$ GeV

⁹³ Model-independent limit from structure function analysis on contribution to $B(\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ from scalars.

$$\frac{\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0), \text{ vector})}{\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))} \quad \Gamma_{21}/\Gamma_{19}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.073	95	⁹⁴ BROWDER 00	CLEO	$4.7 \text{ fb}^{-1} E_{\text{cm}}^{ee} = 10.6$ GeV

⁹⁴ Model-independent limit from structure function analysis on contribution to $B(\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ from vectors.

$$\frac{\Gamma(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))}{\Gamma_{\text{total}}} \quad \Gamma_{22}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.058 ± 0.023 OUR FIT				
0.058 ± 0.024 OUR AVERAGE				
0.056 ± 0.020 ± 0.015	131	BARATE	99K ALEP	1991–1995 LEP runs
0.09 ± 0.10 ± 0.03	3	⁹⁵ BATTLE	94 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.08 ± 0.02 ± 0.02	59	BUSKULIC	96 ALEP	Repl. by BARATE 99K

⁹⁵ BATTLE 94 quote $0.14 \pm 0.10 \pm 0.03$ or $< 0.3\%$ at 90% CL. We subtract $(0.05 \pm 0.02)\%$ to account for $\tau^- \rightarrow K^- (K^0 \rightarrow \pi^0 \pi^0) \nu_\tau$ background.

$$\Gamma(h^- \geq 3\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{23} / \Gamma$$

$$\Gamma_{23} / \Gamma = (\Gamma_{25} + \Gamma_{26} + \Gamma_{28} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.0985\Gamma_{45} + 0.319\Gamma_{117} + 0.322\Gamma_{119}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.37 ± 0.11 OUR FIT	Error includes scale factor of 1.1.			
1.53 ± 0.40 ± 0.46	186	DECAMP	92C ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$3.2 \pm 1.0 \pm 1.0$		BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35 \text{ GeV}$

$$\Gamma(h^- 3\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{24} / \Gamma$$

$$\Gamma_{24} / \Gamma = (\Gamma_{25} + \Gamma_{26} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.322\Gamma_{119}) / \Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.21 ± 0.10 OUR FIT				
1.22 ± 0.10 OUR AVERAGE				
$1.24 \pm 0.09 \pm 0.11$	f&a 2.3k	⁹⁶ BUSKULIC	96 ALEP	LEP 1991–1993 data
$1.70 \pm 0.24 \pm 0.38$	f&a 293	ACCIARRI	95 L3	1992 LEP run
$1.15 \pm 0.08 \pm 0.13$	avg	⁹⁷ PROCARIO	93 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.0 \begin{smallmatrix} +1.4 & +1.1 \\ -0.1 & -0.1 \end{smallmatrix}$		⁹⁸ GAN	87 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

⁹⁶ BUSKULIC 96 quote $B(h^- 3\pi^0 \nu_\tau (\text{ex. } K^0)) = 1.17 \pm 0.09 \pm 0.11$. We add 0.07 to remove their correction for K^0 backgrounds.

⁹⁷ PROCARIO 93 entry is obtained from $B(h^- 3\pi^0 \nu_\tau) / B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.

⁹⁸ Highly correlated with GAN 87 $\Gamma(\eta \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ value. Authors quote $B(\pi^\pm 3\pi^0 \nu_\tau) + 0.67B(\pi^\pm \eta \pi^0 \nu_\tau) = 0.047 \pm 0.010 \pm 0.011$.

$$\Gamma(h^- 3\pi^0 \nu_\tau) / \Gamma(h^- \pi^0 \nu_\tau) \quad \Gamma_{24} / \Gamma_{12}$$

$$\Gamma_{24} / \Gamma_{12} = (\Gamma_{25} + \Gamma_{26} + 0.157\Gamma_{38} + 0.157\Gamma_{40} + 0.322\Gamma_{119}) / (\Gamma_{13} + \Gamma_{15})$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.047 ± 0.004 OUR FIT	Error includes scale factor of 1.1.		
0.044 ± 0.003 ± 0.005	⁹⁹ PROCARIO	93 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$

⁹⁹ PROCARIO 93 quote $0.041 \pm 0.003 \pm 0.005$ after correction for 2 kaon backgrounds assuming $B(K^{*-} \nu_\tau) = 1.42 \pm 0.18\%$ and $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$. We add 0.003 ± 0.003 and multiply the sum by 0.990 ± 0.010 to remove these corrections.

$$\Gamma(\pi^- 3\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{25} / \Gamma$$

VALUE (%)	DOCUMENT ID
1.08 ± 0.10 OUR FIT	

$\Gamma(K^- 3\pi^0 \nu_\tau (\text{ex. } K^0, \eta))/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.037^{+0.022}_{-0.019} OUR FIT

0.037 ± 0.021 ± 0.011 22 BARATE 99K ALEP 1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.05 ± 0.13 ¹⁰⁰ BUSKULIC 94E ALEP Repl. by BARATE 99K

¹⁰⁰ BUSKULIC 94E quote $B(K^- \geq 0\pi^0 \geq 0K^0 \nu_\tau) - [B(K^- \nu_\tau) + B(K^- \pi^0 \nu_\tau) + B(K^- K^0 \nu_\tau) + B(K^- \pi^0 \pi^0 \nu_\tau) + B(K^- \pi^0 K^0 \nu_\tau)] = 0.05 \pm 0.13\%$ accounting for common systematic errors in BUSKULIC 94E and BUSKULIC 94F measurements of these modes. We assume $B(K^- \geq 2K^0 \nu_\tau)$ and $B(K^- \geq 4\pi^0 \nu_\tau)$ are negligible.

$\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ Γ_{27}/Γ

$\Gamma_{27}/\Gamma = (\Gamma_{28} + 0.319\Gamma_{117})/\Gamma$

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.16 ± 0.06 OUR FIT

0.16 ± 0.06 OUR AVERAGE

0.16 ± 0.04 ± 0.09 232 ¹⁰¹ BUSKULIC 96 ALEP LEP 1991–1993 data

0.16 ± 0.05 ± 0.05 ¹⁰² PROCARIO 93 CLEO $E_{\text{cm}}^{ee} \approx 10.6$ GeV

¹⁰¹ BUSKULIC 96 quote result for $\tau^- \rightarrow h^- \geq 4\pi^0 \nu_\tau$. We assume $B(h^- \geq 5\pi^0 \nu_\tau)$ is negligible.

¹⁰² PROCARIO 93 quotes $B(h^- 4\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau) = 0.006 \pm 0.002 \pm 0.002$. We multiply by the ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$ to obtain $B(h^- 4\pi^0 \nu_\tau)$. PROCARIO 93 assume $B(h^- \geq 5\pi^0 \nu_\tau)$ is small and do not correct for it.

$\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex. } K^0, \eta))/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (%) DOCUMENT ID

0.10^{+0.06}_{-0.05} OUR FIT

$\Gamma(K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma \nu_\tau)/\Gamma_{\text{total}}$ Γ_{29}/Γ

$\Gamma_{29}/\Gamma = (\Gamma_{10} + \Gamma_{15} + \Gamma_{22} + \Gamma_{26} + \Gamma_{35} + \Gamma_{40} + 0.715\Gamma_{119})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

1.56 ± 0.04 OUR FIT

1.53 ± 0.04 OUR AVERAGE

1.528 ± 0.039 ± 0.040 f&a ¹⁰³ ABBIENDI 01J OPAL 1990–1995 LEP runs

1.520 ± 0.040 ± 0.041 avg 4006 ¹⁰⁴ BARATE 99K ALEP 1991–1995 LEP runs

1.54 ± 0.24 f&a ABREU 94K DLPH LEP 1992 Z data

1.70 ± 0.12 ± 0.19 f&a 202 ¹⁰⁵ BATTLE 94 CLEO $E_{\text{cm}}^{ee} \approx 10.6$ GeV

1.6 ± 0.4 ± 0.2 f&a 35 AIHARA 87B TPC $E_{\text{cm}}^{ee} = 29$ GeV

1.71 ± 0.29 f&a 53 MILLS 84 DLCO $E_{\text{cm}}^{ee} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.70 ± 0.05 ± 0.06 1610 ¹⁰⁶ BUSKULIC 96 ALEP Repl. by BARATE 99K

- 103 The correlation coefficient between this measurement and the ABBIENDI 01J $B(\tau^- \rightarrow K^- \nu_\tau)$ is 0.60.
- 104 Not independent of BARATE 99K $B(K^- \nu_\tau)$, $B(K^- \pi^0 \nu_\tau)$, $B(K^- 2\pi^0 \nu_\tau)$ (ex. K^0), $B(K^- 3\pi^0 \nu_\tau)$ (ex. K^0), $B(K^- K^0 \nu_\tau)$, and $B(K^- K^0 \pi^0 \nu_\tau)$ values.
- 105 BATTLE 94 quote $1.60 \pm 0.12 \pm 0.19$. We add 0.10 ± 0.02 to correct for their rejection of $K_S^0 \rightarrow \pi^+ \pi^-$ decays.
- 106 Not independent of BUSKULIC 96 $B(K^- \nu_\tau)$, $B(K^- \pi^0 \nu_\tau)$, $B(K^- 2\pi^0 \nu_\tau)$, $B(K^- K^0 \nu_\tau)$, and $B(K^- K^0 \pi^0 \nu_\tau)$ values.

$$\Gamma(K^- \geq 1(\pi^0 \text{ or } K^0 \text{ or } \gamma) \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{30} / \Gamma$$

$$\Gamma_{30} / \Gamma = (\Gamma_{15} + \Gamma_{22} + \Gamma_{26} + \Gamma_{35} + \Gamma_{40} + 0.715\Gamma_{119}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.876 ± 0.034 OUR FIT				
0.86 ± 0.05 OUR AVERAGE				
0.869 ± 0.031 ± 0.034	avg	107 ABBIENDI	01J OPAL	1990–1995 LEP runs
0.69 ± 0.25	avg	108 ABREU	94K DLPH	LEP 1992 Z data
1.2 ± 0.5 $\begin{smallmatrix} +0.2 \\ -0.4 \end{smallmatrix}$	f&a	9 AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29$ GeV

- 107 Not independent of ABBIENDI 01J $B(\tau^- \rightarrow K^- \nu_\tau)$ and $B(\tau^- \rightarrow K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma \nu_\tau)$ values.
- 108 Not independent of ABREU 94K $B(K^- \nu_\tau)$ and $B(K^- \geq 0 \text{ neutrals } \nu_\tau)$ measurements.

$$\Gamma(K_S^0(\text{particles})^- \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{31} / \Gamma$$

$$\Gamma_{31} / \Gamma = (\frac{1}{2}\Gamma_{33} + \frac{1}{2}\Gamma_{35} + \frac{1}{2}\Gamma_{38} + \frac{1}{2}\Gamma_{40} + \Gamma_{45} + \Gamma_{46}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.92 ± 0.04 OUR FIT Error includes scale factor of 1.1.				
0.97 ± 0.07 OUR AVERAGE				
0.970 ± 0.058 ± 0.062	929	BARATE	98E ALEP	1991–1995 LEP runs
0.97 ± 0.09 ± 0.06	141	AKERS	94G OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$$\Gamma(h^- \bar{K}^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{32} / \Gamma = (\Gamma_{33} + \Gamma_{35}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.05 ± 0.04 OUR FIT Error includes scale factor of 1.1.				
0.90 ± 0.07 OUR AVERAGE				
1.01 ± 0.11 ± 0.07	avg	555 109 BARATE	98E ALEP	1991–1995 LEP runs
0.855 ± 0.036 ± 0.073	f&a	1242 COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

- 109 Not independent of BARATE 98E $B(\tau^- \rightarrow \pi^- \bar{K}^0 \nu_\tau)$ and $B(\tau^- \rightarrow K^- K^0 \nu_\tau)$ values.

$\Gamma(\pi^- \bar{K}^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{33}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

0.89 ± 0.04 OUR FIT Error includes scale factor of 1.1.

0.88 ± 0.05 OUR AVERAGE Error includes scale factor of 1.2.

0.933 ± 0.068 ± 0.049	f&a	377	ABBIENDI	00C OPAL	1991–1995 LEP runs
0.928 ± 0.045 ± 0.034	f&a	937	110 BARATE	99K ALEP	1991–1995 LEP runs
0.855 ± 0.117 ± 0.066	avg	509	111 BARATE	98E ALEP	1991–1995 LEP runs
0.704 ± 0.041 ± 0.072	avg		112 COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
0.95 ± 0.15 ± 0.06	f&a		113 ACCIARRI	95F L3	1991–1993 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.79 ± 0.10 ± 0.09		98	114 BUSKULIC	96 ALEP	Repl. by BARATE 99K
--------------------	--	----	--------------	---------	---------------------

110 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

111 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays. Not independent of BARATE 98E $B(K^0 \text{ particles}^- \nu_\tau)$ value.

112 Not independent of COAN 96 $B(h^- K^0 \nu_\tau)$ and $B(K^- K^0 \nu_\tau)$ measurements.

113 ACCIARRI 95F do not identify π^-/K^- and assume $B(K^- K^0 \nu_\tau) = (0.29 \pm 0.12)\%$.

114 BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$\Gamma(\pi^- \bar{K}^0 (\text{non-}K^*(892)^-) \nu_\tau)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

<0.17	95	ACCIARRI	95F L3	1991–1993 LEP runs
-------	----	----------	--------	--------------------

$\Gamma(K^- K^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.154 ± 0.016 OUR FIT

0.158 ± 0.017 OUR AVERAGE

0.162 ± 0.021 ± 0.011	150	115 BARATE	99K ALEP	1991–1995 LEP runs
0.158 ± 0.042 ± 0.017	46	116 BARATE	98E ALEP	1991–1995 LEP runs
0.151 ± 0.021 ± 0.022	111	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ± 0.09 ± 0.02	13	117 BUSKULIC	96 ALEP	Repl. by BARATE 99K
--------------------	----	--------------	---------	---------------------

115 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

116 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

117 BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$\Gamma(K^- K^0 \geq 0\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{36}/\Gamma = (\Gamma_{35} + \Gamma_{40})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.308 ± 0.024 OUR FIT

0.330 ± 0.055 ± 0.039	124	ABBIENDI	00C OPAL	1991–1995 LEP runs
------------------------------	-----	----------	----------	--------------------

$\Gamma(h^- \bar{K}^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ $\Gamma_{37} / \Gamma = (\Gamma_{38} + \Gamma_{40}) / \Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.52 ± 0.04 OUR FIT

0.50 ± 0.06 OUR AVERAGE Error includes scale factor of 1.2.

0.446 ± 0.052 ± 0.046	avg	157	¹¹⁸ BARATE	98E ALEP	1991–1995 LEP runs
-----------------------	-----	-----	-----------------------	----------	--------------------

0.562 ± 0.050 ± 0.048	f&a	264	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
-----------------------	-----	-----	------	---------	---------------------------------------

¹¹⁸ Not independent of BARATE 98E $B(\tau^- \rightarrow \pi^- \bar{K}^0 \pi^0 \nu_\tau)$ and $B(\tau^- \rightarrow K^- K^0 \pi^0 \nu_\tau)$ values.

$\Gamma(\pi^- \bar{K}^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{38} / Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.37 ± 0.04 OUR FIT

0.36 ± 0.04 OUR AVERAGE

0.347 ± 0.053 ± 0.037	f&a	299	¹¹⁹ BARATE	99K ALEP	1991–1995 LEP runs
-----------------------	-----	-----	-----------------------	----------	--------------------

0.294 ± 0.073 ± 0.037	f&a	142	¹²⁰ BARATE	98E ALEP	1991–1995 LEP runs
-----------------------	-----	-----	-----------------------	----------	--------------------

0.417 ± 0.058 ± 0.044	avg	121	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
-----------------------	-----	-----	------	---------	---------------------------------------

0.41 ± 0.12 ± 0.03	f&a	122	ACCIARRI	95F L3	1991–1993 LEP runs
--------------------	-----	-----	----------	--------	--------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.32 ± 0.11 ± 0.05		23	¹²³ BUSKULIC	96 ALEP	Repl. by BARATE 99K
--------------------	--	----	-------------------------	---------	---------------------

¹¹⁹ BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

¹²⁰ BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

¹²¹ Not independent of COAN 96 $B(h^- K^0 \pi^0 \nu_\tau)$ and $B(K^- K^0 \pi^0 \nu_\tau)$ measurements.

¹²² ACCIARRI 95F do not identify π^- / K^- and assume $B(K^- K^0 \pi^0 \nu_\tau) = (0.05 \pm 0.05)\%$.

¹²³ BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$\Gamma(\bar{K}^0 \rho^- \nu_\tau) / \Gamma_{\text{total}}$ Γ_{39} / Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

0.22 ± 0.05 OUR AVERAGE

0.250 ± 0.057 ± 0.044	124	BARATE	99K ALEP	1991–1995 LEP runs
-----------------------	-----	--------	----------	--------------------

0.188 ± 0.054 ± 0.038	125	BARATE	98E ALEP	1991–1995 LEP runs
-----------------------	-----	--------	----------	--------------------

¹²⁴ BARATE 99K measure K^0 's by detecting K_L^0 's in hadron calorimeter. They determine the $\bar{K}^0 \rho^-$ fraction in $\tau^- \rightarrow \pi^- \bar{K}^0 \pi^0 \nu_\tau$ decays to be $(0.72 \pm 0.12 \pm 0.10)$ and multiply their $B(\pi^- \bar{K}^0 \pi^0 \nu_\tau)$ measurement by this fraction to obtain the quoted result.

¹²⁵ BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays. They determine the $\bar{K}^0 \rho^-$ fraction in $\tau^- \rightarrow \pi^- \bar{K}^0 \pi^0 \nu_\tau$ decays to be $(0.64 \pm 0.09 \pm 0.10)$ and multiply their $B(\pi^- \bar{K}^0 \pi^0 \nu_\tau)$ measurement by this fraction to obtain the quoted result.

$\Gamma(K^- K^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.154 ± 0.020 OUR FIT

0.144 ± 0.023 OUR AVERAGE

0.143 ± 0.025 ± 0.015	78	¹²⁶ BARATE	99K ALEP	1991–1995 LEP runs
-----------------------	----	-----------------------	----------	--------------------

0.152 ± 0.076 ± 0.021	15	¹²⁷ BARATE	98E ALEP	1991–1995 LEP runs
-----------------------	----	-----------------------	----------	--------------------

0.145 ± 0.036 ± 0.020	32	COAN	96 CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6$ GeV
-----------------------	----	------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.10 ± 0.05 ± 0.03	5	¹²⁸ BUSKULIC	96 ALEP	Repl. by BARATE 99K
--------------------	---	-------------------------	---------	---------------------

¹²⁶ BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

¹²⁷ BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

¹²⁸ BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$\Gamma(\pi^- \bar{K}^0 \geq 1\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{41}/\Gamma = (\Gamma_{38} + \Gamma_{42})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.324 ± 0.074 ± 0.066	148	ABBIENDI	00C OPAL	1991–1995 LEP runs
------------------------------	-----	----------	----------	--------------------

$\Gamma(\pi^- \bar{K}^0 \pi^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

0.26 ± 0.24			¹²⁹ BARATE	99R ALEP	1991–1995 LEP runs
--------------------	--	--	-----------------------	----------	--------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.66	95	17	¹³⁰ BARATE	99K ALEP	1991–1995 LEP runs
-------	----	----	-----------------------	----------	--------------------

0.58 ± 0.33 ± 0.14		5	¹³¹ BARATE	98E ALEP	1991–1995 LEP runs
--------------------	--	---	-----------------------	----------	--------------------

¹²⁹ BARATE 99R combine the BARATE 98E and BARATE 99K measurements to obtain this value.

¹³⁰ BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

¹³¹ BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

$\Gamma(K^- K^0 \pi^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.16 × 10⁻³	95	¹³² BARATE	99R ALEP	1991–1995 LEP runs
-----------------------------------	----	-----------------------	----------	--------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.18 × 10 ⁻³	95	¹³³ BARATE	99K ALEP	1991–1995 LEP runs
--------------------------	----	-----------------------	----------	--------------------

<0.39 × 10 ⁻³	95	¹³⁴ BARATE	98E ALEP	1991–1995 LEP runs
--------------------------	----	-----------------------	----------	--------------------

¹³² BARATE 99R combine the BARATE 98E and BARATE 99K bounds to obtain this value.

¹³³ BARATE 99K measure K^0 's by detecting K_L^0 's in hadron calorimeter.

¹³⁴ BARATE 98E reconstruct K^0 's by using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

$\Gamma(\pi^- K^0 \bar{K}^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{44}/\Gamma = (2\Gamma_{45} + \Gamma_{46})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.159 ± 0.029 OUR FIT				Error includes scale factor of 1.1.

0.153 ± 0.030 ± 0.016 avg	74	¹³⁵ BARATE	98E ALEP	1991–1995 LEP runs
----------------------------------	----	-----------------------	----------	--------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ± 0.12 ± 0.04		¹³⁶ ACCIARRI	95F L3	1991–1993 LEP runs
--------------------	--	-------------------------	--------	--------------------

¹³⁵ BARATE 98E obtain this value by adding twice their $B(\pi^- K_S^0 K_S^0 \nu_\tau)$ value to their $B(\pi^- K_S^0 K_L^0 \nu_\tau)$ value.

¹³⁶ ACCIARRI 95F assume $B(\pi^- K_S^0 K_S^0 \nu) = B(\pi^- K_S^0 K_L^0 \nu) = 1/2 B(\pi^- K_S^0 K_L^0 \nu)$.

$\Gamma(\pi^- K_S^0 K_S^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{45}/Γ

Bose-Einstein correlations might make the mixing fraction different than 1/4.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.024 ± 0.005 OUR FIT				

0.024 ± 0.005 OUR AVERAGE

0.026 ± 0.010 ± 0.005	6	BARATE	98E ALEP	1991–1995 LEP runs
-----------------------	---	--------	----------	--------------------

0.023 ± 0.005 ± 0.003	42	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
-----------------------	----	------	---------	---------------------------------------

$\Gamma(\pi^- K_S^0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.110 ± 0.028 OUR FIT				Error includes scale factor of 1.1.

0.101 ± 0.023 ± 0.013	68	BARATE	98E ALEP	1991–1995 LEP runs
------------------------------	----	--------	----------	--------------------

$\Gamma(\pi^- K^0 \bar{K}^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
(0.31 ± 0.23) × 10⁻³	¹³⁷ BARATE	99R ALEP	1991–1995 LEP runs

¹³⁷ BARATE 99R combine BARATE 98E $\Gamma(\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ measurements to obtain this value.

$\Gamma(\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.020	95	BARATE	98E ALEP	1991–1995 LEP runs

$\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.031 ± 0.011 ± 0.005	11	BARATE	98E ALEP	1991–1995 LEP runs

$\Gamma(K^0 h^+ h^- h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.17	95	TSCHIRHART	88 HRS	$E_{\text{cm}}^{ee} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.27	90	BELTRAMI	85 HRS	$E_{\text{cm}}^{ee} = 29$ GeV
--------	----	----------	--------	-------------------------------

$\Gamma(K^0 h^+ h^- h^- \nu_\tau)/\Gamma_{\text{total}}$					Γ_{51}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.023±0.019±0.007	6	138 BARATE	98E ALEP	1991–1995 LEP runs	
138 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.					

$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$					Γ_{52}/Γ
$\Gamma_{52}/\Gamma = (0.3431\Gamma_{33} + 0.3431\Gamma_{35} + 0.3431\Gamma_{38} + 0.3431\Gamma_{40} + 0.4307\Gamma_{45} + 0.6861\Gamma_{46} + \Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.9101\Gamma_{137} + 0.9101\Gamma_{138})/\Gamma$					

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
15.20± 0.07 OUR FIT	Error includes scale factor of 1.1.			
14.8 ± 0.4 OUR AVERAGE				
14.4 ± 0.6 ± 0.3		ADEVA	91F L3	$E_{\text{cm}}^{ee} = 88.3\text{--}94.3$ GeV
15.0 ± 0.4 ± 0.3		BEHREND	89B CELL	$E_{\text{cm}}^{ee} = 14\text{--}47$ GeV
15.1 ± 0.8 ± 0.6		AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
13.5 ± 0.3 ± 0.3		ABACHI	89B HRS	$E_{\text{cm}}^{ee} = 29$ GeV
12.8 ± 1.0 ± 0.7	139	BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
12.1 ± 0.5 ± 1.2		RUCKSTUHL	86 DLCO	$E_{\text{cm}}^{ee} = 29$ GeV
12.8 ± 0.5 ± 0.8	1420	SCHMIDKE	86 MRK2	$E_{\text{cm}}^{ee} = 29$ GeV
15.3 ± 1.1 $\begin{smallmatrix} +1.3 \\ -1.6 \end{smallmatrix}$	367	ALTHOFF	85 TASS	$E_{\text{cm}}^{ee} = 34.5$ GeV
13.6 ± 0.5 ± 0.8		BARTEL	85F JADE	$E_{\text{cm}}^{ee} = 34.6$ GeV
12.2 ± 1.3 ± 3.9	140	BERGER	85 PLUT	$E_{\text{cm}}^{ee} = 34.6$ GeV
13.3 ± 0.3 ± 0.6		FERNANDEZ	85 MAC	$E_{\text{cm}}^{ee} = 29$ GeV
24 ± 6	35	BRANDELIK	80 TASS	$E_{\text{cm}}^{ee} = 30$ GeV
32 ± 5	692	141 BACINO	78B DLCO	$E_{\text{cm}}^{ee} = 3.1\text{--}7.4$ GeV
35 ± 11		141 BRANDELIK	78 DASP	Assumes $V\text{--}A$ decay
18 ± 6.5	33	141 JAROS	78 MRK1	$E_{\text{cm}}^{ee} > 6$ GeV

139 BURCHAT 87 value is not independent of SCHMIDKE 86 value.
 140 Not independent of BERGER 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$, $\Gamma(h^- \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$, and $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$, and therefore not used in the fit.
 141 Low energy experiments are not in average or fit because the systematic errors in background subtraction are judged to be large.

$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^+ \pi^-) \text{ ("3-prong")})/\Gamma_{\text{total}}$					Γ_{53}/Γ
$\Gamma_{53}/\Gamma = (\Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.9101\Gamma_{137} + 0.9101\Gamma_{138})/\Gamma$					

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
14.57 ± 0.07 OUR FIT	Error includes scale factor of 1.1.			
14.59 ± 0.08 OUR AVERAGE	Error includes scale factor of 1.1.			
14.569±0.093±0.048	f&a	23k 142 ABREU	01M DLPH	1992–1995 LEP runs
14.556±0.105±0.076	f&a	143 ACHARD	01D L3	1992–1995 LEP runs

14.96 ±0.09 ±0.22	f&a	10.4k	AKERS	95Y OPAL	1991–1994 LEP runs
14.22 ±0.10 ±0.37	avg		¹⁴⁴ BALEST	95C CLEO	$E_{cm}^{ee} \approx 10.6$ GeV
• • •	We do not use the following data for averages, fits, limits, etc. • • •				
15.26 ±0.26 ±0.22			ACTON	92H OPAL	Repl. by AK-ERS 95Y
13.3 ±0.3 ±0.8			¹⁴⁵ ALBRECHT	92D ARG	$E_{cm}^{ee} = 9.4–10.6$ GeV
14.35 $\begin{smallmatrix} +0.40 \\ -0.45 \end{smallmatrix}$ ±0.24			DECAMP	92C ALEP	1989–1990 LEP runs

¹⁴²The correlation coefficients between this measurement and the ABREU 01M measurements of $B(\tau \rightarrow 1\text{-prong})$ and $B(\tau \rightarrow 5\text{-prong})$ are -0.98 and -0.08 respectively.

¹⁴³The correlation coefficients between this measurement and the ACHARD 01D measurements of $B(\tau \rightarrow \text{"1-prong"})$ and $B(\tau \rightarrow \text{"5-prong"})$ are -0.978 and -0.19 respectively.

¹⁴⁴Not independent of BALEST 95C $B(h^- h^- h^+ \nu_\tau)$ and $B(h^- h^- h^+ \pi^0 \nu_\tau)$ values, and BORTOLETTO 93 $B(h^- h^- h^+ 2\pi^0 \nu_\tau)/B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau)$ value.

¹⁴⁵This ALBRECHT 92D value is not independent of their $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}^2$ value.

$$\frac{\Gamma(h^- h^- h^+ \nu_\tau)}{\Gamma_{\text{total}}} / \frac{\Gamma_{54}}{\Gamma} = (0.3431\Gamma_{33} + 0.3431\Gamma_{35} + \Gamma_{60} + \Gamma_{82} + \Gamma_{89} + 0.0221\Gamma_{137}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

10.01 ± 0.09 OUR FIT Error includes scale factor of 1.2.

9.8 ± 0.6 OUR AVERAGE Error includes scale factor of 4.4. See the ideogram below.

7.6 ±0.1 ±0.5	avg	7.5k	¹⁴⁶ ALBRECHT	96E ARG	$E_{cm}^{ee} = 9.4–10.6$ GeV
9.92 ±0.10 ±0.09	f&a	11.2k	¹⁴⁷ BUSKULIC	96 ALEP	LEP 1991–1993 data
9.49 ±0.36 ±0.63	f&a		DECAMP	92C ALEP	1989–1990 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.7 ±0.7 ±0.3		694	¹⁴⁸ BEHREND	90 CELL	$E_{cm}^{ee} = 35$ GeV
7.0 ±0.3 ±0.7		1566	¹⁴⁹ BAND	87 MAC	$E_{cm}^{ee} = 29$ GeV
6.7 ±0.8 ±0.9			¹⁵⁰ BURCHAT	87 MRK2	$E_{cm}^{ee} = 29$ GeV
6.4 ±0.4 ±0.9			¹⁵¹ RUCKSTUHL	86 DLCO	$E_{cm}^{ee} = 29$ GeV
7.8 ±0.5 ±0.8		890	SCHMIDKE	86 MRK2	$E_{cm}^{ee} = 29$ GeV
8.4 ±0.4 ±0.7		1255	¹⁵¹ FERNANDEZ	85 MAC	$E_{cm}^{ee} = 29$ GeV
9.7 ±2.0 ±1.3			BEHREND	84 CELL	$E_{cm}^{ee} = 14,22$ GeV

¹⁴⁶ALBRECHT 96E not independent of ALBRECHT 93C $\Gamma(h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) \times \Gamma(\text{particle}^- \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}^2$ value.


¹⁴⁷BUSKULIC 96 quote $B(h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) = 9.50 \pm 0.10 \pm 0.11$. We add 0.42 to remove their K^0 correction and reduce the systematic error accordingly.

¹⁴⁸BEHREND 90 subtract 0.3% to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution to measured events.

¹⁴⁹BAND 87 subtract for charged kaon modes; not independent of FERNANDEZ 85 value.

¹⁵⁰BURCHAT 87 value is not independent of SCHMIDKE 86 value.

¹⁵¹ Value obtained by multiplying paper's $R = B(h^- h^- h^+ \nu_\tau)/B(3\text{-prong})$ by $B(3\text{-prong}) = 0.143$ and subtracting 0.3% for $K^*(892)$ background.



$$\Gamma(h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{55} / \Gamma$$

$$\Gamma_{55} / \Gamma = (\Gamma_{60} + \Gamma_{82} + \Gamma_{89} + 0.0221 \Gamma_{137}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
9.65±0.09 OUR FIT	Error includes scale factor of 1.2.				
9.57±0.11 OUR AVERAGE					
9.50±0.10±0.11	avg	11.2k	¹⁵² BUSKULIC	96 ALEP	LEP 1991–1993 data
9.87±0.10±0.24	avg		¹⁵³ AKERS	95Y OPAL	1991–1994 LEP runs
9.51±0.07±0.20	f&a	37.7k	BALEST	95C CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$

¹⁵² Not independent of BUSKULIC 96 $B(h^- h^- h^+ \nu_\tau)$ value.

¹⁵³ Not independent of AKERS 95Y $B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$ and $B(h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$ values.

$$\Gamma(h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-)) \quad \Gamma_{55} / \Gamma_{53}$$

$$(\text{"3-prong"})$$

$$\Gamma_{55} / \Gamma_{53} = (\Gamma_{60} + \Gamma_{82} + \Gamma_{89} + 0.0221 \Gamma_{137}) / (\Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285 \Gamma_{117} + 0.285 \Gamma_{119} + 0.9101 \Gamma_{137} + 0.9101 \Gamma_{138})$$

VALUE		DOCUMENT ID	TECN	COMMENT
0.662±0.006 OUR FIT	Error includes scale factor of 1.3.			
0.660±0.004±0.014		AKERS	95Y OPAL	1991–1994 LEP runs

$$\frac{\Gamma(h^- h^- h^+ \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}}}{\text{VALUE (\%)}} \quad \frac{\text{DOCUMENT ID}}{\text{9.61} \pm \text{0.09 OUR FIT}} \quad \text{Error includes scale factor of 1.2.} \quad \Gamma_{56} / \Gamma = (\Gamma_{60} + \Gamma_{82} + \Gamma_{89}) / \Gamma$$

$$\frac{\Gamma(\pi^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}}}{\text{VALUE (\%)}} \quad \frac{\text{DOCUMENT ID}}{\text{9.46} \pm \text{0.10 OUR FIT}} \quad \text{Error includes scale factor of 1.2.} \quad \Gamma_{57} / \Gamma = (0.3431 \Gamma_{33} + \Gamma_{60} + 0.0221 \Gamma_{137}) / \Gamma$$

$$\frac{\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}}{\text{VALUE (\%)}} \quad \frac{\text{EVTS}}{\text{9.15} \pm \text{0.10 OUR FIT}} \quad \frac{\text{DOCUMENT ID}}{\text{9.13} \pm \text{0.05} \pm \text{0.46}} \quad \frac{\text{TECN}}{\text{43k}} \quad \frac{\text{COMMENT}}{\text{154 BRIERE 03 CLE3 } E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}}$$

¹⁵⁴ 47% correlated with BRIERE 03 $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ and 71% correlated with $\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau$ because of a common 5% normalization error.

$$\frac{\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0), \text{non-axial vector}) / \Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))}{\text{VALUE}} \quad \frac{\text{CL\%}}{95} \quad \frac{\text{DOCUMENT ID}}{155} \quad \frac{\text{TECN}}{\text{ACKERSTAFF 97R OPAL}} \quad \frac{\text{COMMENT}}{\text{1992-1994 LEP runs}}$$

¹⁵⁵ Model-independent limit from structure function analysis on contribution to $B(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ from non-axial vectors.

$$\frac{\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}}}{\text{VALUE (\%)}} \quad \frac{\text{DOCUMENT ID}}{\text{9.11} \pm \text{0.10 OUR FIT}} \quad \text{Error includes scale factor of 1.2.} \quad \Gamma_{60} / \Gamma$$

$$\frac{\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}}}{\text{VALUE (\%)}} \quad \frac{\text{EVTS}}{\text{5.19} \pm \text{0.10 OUR FIT}} \quad \frac{\text{DOCUMENT ID}}{\text{5.19} \pm \text{0.10 OUR FIT}} \quad \frac{\text{TECN}}{\text{Error includes scale factor of 1.2.}} \quad \frac{\text{COMMENT}}{\text{Error includes scale factor of 1.2.}}$$

$$\Gamma_{61} / \Gamma = (0.3431 \Gamma_{38} + 0.3431 \Gamma_{40} + 0.4307 \Gamma_{45} + 0.6861 \Gamma_{46} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{86} + \Gamma_{90} + 0.285 \Gamma_{117} + 0.285 \Gamma_{119} + 0.888 \Gamma_{137} + 0.9101 \Gamma_{138}) / \Gamma$$

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
5.6 ± 0.7 ± 0.3	352	¹⁵⁶ BEHREND	90	CELL $E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$
4.2 ± 0.5 ± 0.9	203	¹⁵⁷ ALBRECHT	87L	ARG $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$
6.1 ± 0.8 ± 0.9		¹⁵⁸ BURCHAT	87	MRK2 $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
7.6 ± 0.4 ± 0.9	159,160	RUCKSTUHL	86	DLCO $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
4.7 ± 0.5 ± 0.8	530	¹⁶¹ SCHMIDKE	86	MRK2 $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
5.6 ± 0.4 ± 0.7	160	FERNANDEZ	85	MAC $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
6.2 ± 2.3 ± 1.7		BEHREND	84	CELL $E_{\text{cm}}^{\text{ee}} = 14, 22 \text{ GeV}$

¹⁵⁶ BEHREND 90 value is not independent of BEHREND 90 $B(3h \nu_\tau \geq 1 \text{ neutrals}) + B(5\text{-prong})$.

¹⁵⁷ ALBRECHT 87L measure the product of branching ratios $B(3\pi^\pm \pi^0 \nu_\tau) B((e\bar{\nu} \text{ or } \mu\bar{\nu} \text{ or } \pi \text{ or } K \text{ or } \rho) \nu_\tau) = 0.029$ and use the PDG 86 values for the second branching ratio which sum to 0.69 ± 0.03 to get the quoted value.

¹⁵⁸ BURCHAT 87 value is not independent of SCHMIDKE 86 value.

¹⁵⁹ Contributions from kaons and from $> 1\pi^0$ are subtracted. Not independent of $(3\text{-prong} + 0\pi^0)$ and $(3\text{-prong} + \geq 0\pi^0)$ values.

¹⁶⁰ Value obtained using paper's $R = B(h^- h^- h^+ \nu_\tau) / B(3\text{-prong})$ and current $B(3\text{-prong}) = 0.143$.

¹⁶¹ Not independent of SCHMIDKE 86 $h^- h^- h^+ \nu_\tau$ and $h^- h^- h^+ (\geq 0\pi^0) \nu_\tau$ values.

$$\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{62} / \Gamma$$

$$\Gamma_{62} / \Gamma = (\Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{86} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.888\Gamma_{137} + 0.9101\Gamma_{138}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.92±0.09 OUR FIT	Error includes scale factor of 1.3.			
5.07±0.24 OUR AVERAGE				
5.09±0.10±0.23	avg	¹⁶² AKERS	95Y OPAL	1991–1994 LEP runs
4.95±0.29±0.65	f&a	570 DECAMP	92C ALEP	1989–1990 LEP runs

¹⁶² Not independent of AKERS 95Y B($h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$)) and B($h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. K^0))/B($h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$)) values.

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{63} / \Gamma$$

$$\Gamma_{63} / \Gamma = (0.3431\Gamma_{38} + 0.3431\Gamma_{40} + \Gamma_{68} + \Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119} + 0.888\Gamma_{137} + 0.0221\Gamma_{138}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.53±0.09 OUR FIT	Error includes scale factor of 1.3.			
4.45±0.09±0.07	6.1k	¹⁶³ BUSKULIC	96 ALEP	LEP 1991–1993 data

¹⁶³ BUSKULIC 96 quote B($h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0)) = $4.30 \pm 0.09 \pm 0.09$. We add 0.15 to remove their K^0 correction and reduce the systematic error accordingly.

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{64} / \Gamma$$

$$\Gamma_{64} / \Gamma = (\Gamma_{68} + \Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119} + 0.888\Gamma_{137} + 0.0221\Gamma_{138}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.35±0.09 OUR FIT	Error includes scale factor of 1.3.			
4.23±0.06±0.22	7.2k	BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} \quad \Gamma_{65} / \Gamma = (\Gamma_{68} + \Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119}) / \Gamma$$

VALUE (%)	DOCUMENT ID
2.62±0.09 OUR FIT	Error includes scale factor of 1.2.

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{66} / \Gamma = (0.3431\Gamma_{38} + \Gamma_{68} + 0.888\Gamma_{137} + 0.0221\Gamma_{138}) / \Gamma$$

VALUE (%)	DOCUMENT ID
4.37±0.09 OUR FIT	Error includes scale factor of 1.3.

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{67} / \Gamma = (\Gamma_{68} + 0.888\Gamma_{137} + 0.0221\Gamma_{138}) / \Gamma$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
4.24±0.09 OUR FIT	Error includes scale factor of 1.3.		
4.19±0.10±0.21	¹⁶⁴ EDWARDS	00A CLEO	4.7 fb^{-1} $E_{\text{cm}}^{ee} = 10.6$ GeV

¹⁶⁴ EDWARDS 00A quote $(4.19 \pm 0.10) \times 10^{-2}$ with a 5% systematic error.

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} \quad \Gamma_{68} / \Gamma$$

VALUE (%)	DOCUMENT ID
2.51±0.09 OUR FIT	Error includes scale factor of 1.2.

$\Gamma(h^- \rho \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ $\Gamma_{69} / \Gamma_{63}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.30 \pm 0.04 \pm 0.02$	393	ALBRECHT	91D ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV

$\Gamma(h^- \rho^+ h^- \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ $\Gamma_{70} / \Gamma_{63}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.10 \pm 0.03 \pm 0.04$	142	ALBRECHT	91D ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV

$\Gamma(h^- \rho^- h^+ \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ $\Gamma_{71} / \Gamma_{63}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.26 \pm 0.05 \pm 0.01$	370	ALBRECHT	91D ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV

$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau) / \Gamma_{total}$ Γ_{72} / Γ
 $\Gamma_{72} / \Gamma = (0.4307\Gamma_{45} + \Gamma_{74} + 0.236\Gamma_{117} + 0.888\Gamma_{138}) / \Gamma$

VALUE (%)	DOCUMENT ID
0.55 ± 0.04 OUR FIT	

$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (ex.K^0)) / \Gamma_{total}$ Γ_{73} / Γ
 $\Gamma_{73} / \Gamma = (\Gamma_{74} + 0.236\Gamma_{117} + 0.888\Gamma_{138}) / \Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.54 ± 0.04 OUR FIT				
0.50 ± 0.07 ± 0.07	1.8k	BUSKULIC	96 ALEP	LEP 1991-1993 data

$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (ex.K^0)) / \Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau)$ $\Gamma_{73} / \Gamma_{52}$
 $\Gamma_{73} / \Gamma_{52} = (\Gamma_{74} + 0.236\Gamma_{117} + 0.888\Gamma_{138}) / (0.3431\Gamma_{33} + 0.3431\Gamma_{35} + 0.3431\Gamma_{38} + 0.3431\Gamma_{40} + 0.4307\Gamma_{45} + 0.6861\Gamma_{46} + \Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.9101\Gamma_{137} + 0.9101\Gamma_{138})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0354 ± 0.0028 OUR FIT				
0.034 ± 0.002 ± 0.003	668	BORTOLETTO93	CLEO	$E_{cm}^{ee} \approx 10.6$ GeV

$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (ex.K^0, \omega, \eta)) / \Gamma_{total}$ Γ_{74} / Γ

VALUE (%)	DOCUMENT ID
0.11 ± 0.04 OUR FIT	

$\Gamma(h^- h^- h^+ 3\pi^0 \nu_\tau) / \Gamma_{total}$ Γ_{75} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.023 ± 0.008 OUR FIT				Error includes scale factor of 1.5.
0.023 ± 0.005 OUR AVERAGE				

0.022 ± 0.003 ± 0.004	139	ANASTASSOV 01	CLEO	$E_{cm}^{ee} = 10.6$ GeV
0.11 ± 0.04 ± 0.05	440	¹⁶⁵ BUSKULIC	96 ALEP	LEP 1991-1993 data

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0285 ± 0.0056 ± 0.0051	57	ANDERSON 97	CLEO	Repl. by ANASTASSOV 01
--------------------------	----	-------------	------	------------------------

¹⁶⁵BUSKULIC 96 state their measurement is for $B(h^- h^- h^+ \geq 3\pi^0 \nu_\tau)$. We assume that $B(h^- h^- h^+ \geq 4\pi^0 \nu_\tau)$ is very small.

$$\Gamma(K^- h^+ h^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}}$$

$$\Gamma_{76} / \Gamma = (0.3431\Gamma_{35} + 0.3431\Gamma_{40} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{119}) / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
0.71 ± 0.04 OUR FIT				Error includes scale factor of 1.4.
<0.6	90	AIHARA	84C TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$$\Gamma(K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{77} / \Gamma = (\Gamma_{82} + \Gamma_{89}) / \Gamma$$

VALUE (%)	DOCUMENT ID
0.49 ± 0.04 OUR FIT	Error includes scale factor of 1.6.

$$\Gamma(K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$$

$$\Gamma_{77} / \Gamma_{58} = (\Gamma_{82} + \Gamma_{89}) / (\Gamma_{60} + 0.0221\Gamma_{137})$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
5.4 ± 0.4 OUR FIT				Error includes scale factor of 1.6.
5.44 ± 0.21 ± 0.53	7.9k	RICHICHI	99 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma(K^- h^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{78} / \Gamma = (\Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119}) / \Gamma$$

VALUE (%)	DOCUMENT ID
0.110 ± 0.022 OUR FIT	

$$\Gamma(K^- h^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0))$$

$$\Gamma_{78} / \Gamma_{67} = (\Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119}) / (\Gamma_{68} + 0.888\Gamma_{137} + 0.0221\Gamma_{138})$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.5 OUR FIT				
2.61 ± 0.45 ± 0.42	719	RICHICHI	99 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma(K^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}}$$

$$\Gamma_{79} / \Gamma = (0.3431\Gamma_{35} + 0.3431\Gamma_{40} + \Gamma_{82} + \Gamma_{86} + 0.285\Gamma_{119}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.52 ± 0.05 OUR FIT				Error includes scale factor of 1.4.
0.58^{+0.15}_{-0.13} ± 0.12	20	¹⁶⁶ BAUER	94 TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22 ^{+0.16} _{-0.13} ± 0.05	9	¹⁶⁷ MILLS	85 DLCO	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
---	---	----------------------	---------	---------------------------------------

¹⁶⁶We multiply 0.58% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

¹⁶⁷Error correlated with MILLS 85 ($K K \pi \nu$) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain the systematic error.

$$\Gamma(K^- \pi^+ \pi^- \geq 0 \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{80} / \Gamma = (\Gamma_{82} + \Gamma_{86} + 0.231\Gamma_{119}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.41 ± 0.05 OUR FIT			Error includes scale factor of 1.4.
0.30 ± 0.05 OUR AVERAGE			
0.343 ± 0.073 ± 0.031	avg	ABBIENDI	00D OPAL 1990–1995 LEP runs
0.275 ± 0.064	avg	¹⁶⁸ BARATE	98 ALEP 1991–1995 LEP runs

¹⁶⁸Not independent of BARATE 98 $\Gamma(\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}}$ and $\Gamma(\tau^- \rightarrow K^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ values.

$$\Gamma(K^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}}$$

$$\Gamma_{81} / \Gamma = (0.3431 \Gamma_{35} + \Gamma_{82}) / \Gamma$$

VALUE (%)

DOCUMENT ID

0.39 ± 0.04 OUR FIT Error includes scale factor of 1.6.

$$\Gamma(K^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$$

Γ_{82} / Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)

EVTS

DOCUMENT ID

TECN

COMMENT

0.34 ± 0.04 OUR FIT Error includes scale factor of 1.6.

0.33 ± 0.05 OUR AVERAGE Error includes scale factor of 1.8. See the ideogram below.

0.415 ± 0.053 ± 0.040 f&a 269 ABBIENDI 04J OPAL 1991-1995 LEP runs

0.384 ± 0.014 ± 0.038 f&a 3.5k ¹⁶⁹ BRIERE 03 CLE3 $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

0.346 ± 0.023 ± 0.056 avg 158 ¹⁷⁰ RICHICHI 99 CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

0.214 ± 0.037 ± 0.029 f&a BARATE 98 ALEP 1991-1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •


0.360 ± 0.082 ± 0.048 ABBIENDI 00D OPAL 1990-1995 LEP runs

¹⁶⁹ 47% correlated with BRIERE 03 $\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$ and 34% correlated with $\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau$ because of a common 5% normalization error.

¹⁷⁰ Not independent of RICHICHI 99

$\Gamma(\tau^- \rightarrow K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$, $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau) / \Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ values.

WEIGHTED AVERAGE
0.33 ± 0.05 (Error scaled by 1.8)



$\Gamma(K^- \rho^0 \nu_\tau \rightarrow K^- \pi^+ \pi^- \nu_\tau) / \Gamma(K^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ $\Gamma_{83} / \Gamma_{82}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.48 ± 0.14 ± 0.10	171 ASNER	00B CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.39 ± 0.14	172 BARATE	99R ALEP	1991–1995 LEP runs
-------------	------------	----------	--------------------

171 ASNER 00B assume $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0) decays proceed only through $K \rho$ and $K^* \pi$ intermediate states. They assume the resonance structure of $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0) decays is dominated by $K_1(1270)^-$ and $K_1(1400)^-$ resonances, and assume $B(K_1(1270) \rightarrow K^*(892)\pi) = (16 \pm 5)\%$, $B(K_1(1270) \rightarrow K\rho) = (42 \pm 6)\%$, and $B(K_1(1400) \rightarrow K\rho) = 0$.

172 BARATE 99R assume $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0) decays proceed only through $K \rho$ and $K^* \pi$ intermediate states. The quoted error is statistical only.

$\Gamma(K^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ $\Gamma_{84} / \Gamma = (0.3431\Gamma_{40} + \Gamma_{86} + 0.231\Gamma_{119}) / \Gamma$

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
12.2 ± 2.6 OUR FIT	

$\Gamma(K^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ $\Gamma_{85} / \Gamma = (\Gamma_{86} + 0.231\Gamma_{119}) / \Gamma$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.9 ± 2.5 OUR FIT				
7.0 ± 2.5 OUR AVERAGE				

7.5 ± 2.6 ± 1.8	avg	173 RICHICHI	99 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
6.1 ± 3.9 ± 1.8	f&a	BARATE	98 ALEP	1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17	95	ABBIENDI	00D OPAL	1990–1995 LEP runs
-----	----	----------	----------	--------------------

173 Not independent of RICHICHI 99

$\Gamma(\tau^- \rightarrow K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$, $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau) / \Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ values.

$\Gamma(K^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0, \eta)) / \Gamma_{\text{total}}$ Γ_{86} / Γ

Test of lepton family number conservation.

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
6.3 ± 2.5 OUR FIT	

$\Gamma(K^- \pi^+ K^- \geq 0 \text{ neut. } \nu_\tau) / \Gamma_{\text{total}}$ Γ_{87} / Γ

<u>VALUE (%)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.09	95	BAUER	94 TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$\Gamma(K^- K^+ \pi^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{88}/\Gamma = (\Gamma_{89} + \Gamma_{90})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.195±0.018 OUR FIT Error includes scale factor of 1.1.

0.203±0.031 OUR AVERAGE

0.159±0.053±0.020	f&a		ABBIENDI	00D	OPAL	1990–1995 LEP runs
0.238±0.042	avg	174	BARATE	98	ALEP	1991–1995 LEP runs
0.15 $\begin{smallmatrix} +0.09 \\ -0.07 \end{smallmatrix}$ ±0.03	f&a	4	175 BAUER	94	TPC	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

¹⁷⁴ Not independent of BARATE 98 $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ values.

¹⁷⁵ We multiply 0.15% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$\Gamma(K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{89}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.155±0.007 OUR FIT

0.154±0.009 OUR AVERAGE

0.155±0.006±0.009	f&a	932	176 BRIERE	03	CLE3	$E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$
0.087±0.056±0.040	avg		ABBIENDI	00D	OPAL	1990–1995 LEP runs
0.145±0.013±0.028	avg	2.3k	177 RICHICHI	99	CLEO	$E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$
0.163±0.021±0.017	f&a		BARATE	98	ALEP	1991–1995 LEP runs
0.22 $\begin{smallmatrix} +0.17 \\ -0.11 \end{smallmatrix}$ ±0.05	f&a	9	178 MILLS	85	DLCO	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

¹⁷⁶ 71% correlated with BRIERE 03 $\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$ and 34% correlated with $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ because of a common 5% normalization error.

¹⁷⁷ Not independent of RICHICHI 99 $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ values.

¹⁷⁸ Error correlated with MILLS 85 ($K \pi \pi \pi^0 \nu$) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain the systematic error.

$\Gamma(K^- K^+ \pi^- \nu_\tau)/\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ $\Gamma_{89}/\Gamma_{58} = \Gamma_{89}/(\Gamma_{60} + 0.0221\Gamma_{137})$

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

1.69±0.08 OUR FIT Error includes scale factor of 1.1.

1.60±0.15±0.30 2.3k RICHICHI 99 CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$

$\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ **Γ_{90}/Γ**

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.1±1.6 OUR FIT	Error includes scale factor of 1.1.				
4.4±1.8 OUR AVERAGE	Error includes scale factor of 1.1.				
3.3±1.8±0.7	avg	158	¹⁷⁹ RICHICHI	99	CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV
7.5±2.9±1.5	f&a		BARATE	98	ALEP 1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

<27	95	ABBIENDI	00D	OPAL	1990–1995 LEP runs
-----	----	----------	-----	------	--------------------

¹⁷⁹ Not independent of RICHICHI 99

$\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ values.

$\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0))$
 $\Gamma_{90}/\Gamma_{67} = \Gamma_{90}/(\Gamma_{68} + 0.888\Gamma_{137} + 0.0221\Gamma_{138})$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.0 ± 0.4 OUR FIT	Error includes scale factor of 1.1.			
0.79±0.44±0.16	158	¹⁸⁰ RICHICHI	99	CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

¹⁸⁰ RICHICHI 99 also quote a 95%CL upper limit of 0.0157 for this measurement.

$\Gamma(K^- K^+ K^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ **Γ_{91}/Γ**

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.21	95	BAUER	94	TPC $E_{\text{cm}}^{ee} = 29$ GeV

$\Gamma(K^- K^+ K^- \nu_\tau)/\Gamma_{\text{total}}$ **Γ_{92}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.7 × 10⁻⁵	90	BRIERE	03	CLE3 $E_{\text{cm}}^{ee} = 10.6$ GeV
<1.9 × 10 ⁻⁴	90	BARATE	98	ALEP 1991–1995 LEP runs

$\Gamma(\pi^- K^+ \pi^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ **Γ_{93}/Γ**

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.25	95	BAUER	94	TPC $E_{\text{cm}}^{ee} = 29$ GeV

$\Gamma(e^- e^- e^+ \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$ **Γ_{94}/Γ**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8±1.4±0.4	5	ALAM	96	CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\mu^- e^- e^+ \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$ **Γ_{95}/Γ**

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<3.6	90	ALAM	96	CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(3h^-2h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^- \pi^+) ("5\text{-prong"})) / \Gamma_{\text{total}} \quad \Gamma_{96} / \Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average. $\Gamma_{96} / \Gamma = (\Gamma_{97} + \Gamma_{98}) / \Gamma$

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.100 ± 0.006 OUR FIT

0.111 ± 0.008 OUR AVERAGE Error includes scale factor of 1.1.

0.115 ± 0.013 ± 0.006	f&a	112	¹⁸¹ ABREU	01M DLPH	1992–1995 LEP runs
0.170 ± 0.022 ± 0.026	f&a		¹⁸² ACHARD	01D L3	1992–1995 LEP runs
0.119 ± 0.013 ± 0.008	avg	119	¹⁸³ ACKERSTAFF	99E OPAL	1991–1995 LEP runs
0.097 ± 0.005 ± 0.011	f&a	419	GIBAUT	94B CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
0.102 ± 0.029	f&a	13	BYLSMA	87 HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.26 ± 0.06 ± 0.05			ACTON	92H OPAL	$E_{\text{cm}}^{ee} = 88.2\text{--}94.2 \text{ GeV}$
0.10 ^{+0.05} _{-0.04} ± 0.03			DECAMP	92C ALEP	1989–1990 LEP runs
0.16 ± 0.13 ± 0.04			BEHREND	89B CELL	$E_{\text{cm}}^{ee} = 14\text{--}47 \text{ GeV}$
0.3 ± 0.1 ± 0.2			BARTEL	85F JADE	$E_{\text{cm}}^{ee} = 34.6 \text{ GeV}$
0.13 ± 0.04		10	BELTRAMI	85 HRS	Repl. by BYLSMA 87
0.16 ± 0.08 ± 0.04		4	BURCHAT	85 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
1.0 ± 0.4		10	BEHREND	82 CELL	Repl. by BEHREND 89B

¹⁸¹ The correlation coefficients between this measurement and the ABREU 01M measurements of $B(\tau \rightarrow 1\text{-prong})$ and $B(\tau \rightarrow 3\text{-prong})$ are -0.08 and -0.08 respectively.

¹⁸² The correlation coefficients between this measurement and the ACHARD 01D measurements of $B(\tau \rightarrow "1\text{-prong}")$ and $B(\tau \rightarrow "3\text{-prong}")$ are -0.082 and -0.19 respectively.

¹⁸³ Not independent of ACKERSTAFF 99E $B(\tau^- \rightarrow 3h^-2h^+ \nu_\tau (\text{ex. } K^0))$ and $B(\tau^- \rightarrow 3h^-2h^+ \pi^0 \nu_\tau (\text{ex. } K^0))$ measurements.

$\Gamma(3h^-2h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{97} / \Gamma$

VALUE (%) EVTS DOCUMENT ID TECN COMMENT

0.082 ± 0.006 OUR FIT

0.076 ± 0.007 OUR AVERAGE

0.091 ± 0.014 ± 0.006		97	ACKERSTAFF	99E OPAL	1991–1995 LEP runs
0.080 ± 0.011 ± 0.013		58	BUSKULIC	96 ALEP	LEP 1991–1993 data
0.077 ± 0.005 ± 0.009		295	GIBAUT	94B CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
0.064 ± 0.023 ± 0.01		12	ALBRECHT	88B ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
0.051 ± 0.020		7	BYLSMA	87 HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.067 ± 0.030		5	¹⁸⁴ BELTRAMI	85 HRS	Repl. by BYLSMA 87
---------------	--	---	-------------------------	--------	--------------------

¹⁸⁴ The error quoted is statistical only.

$\Gamma(3h^- 2h^+ \pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.0181 ± 0.0027 OUR FIT
0.0172 ± 0.0027 OUR AVERAGE

0.017 ± 0.002 ± 0.002	231	ANASTASSOV 01	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
0.027 ± 0.018 ± 0.009	23	ACKERSTAFF 99E	OPAL	1991–1995 LEP runs
0.018 ± 0.007 ± 0.012	18	BUSKULIC 96	ALEP	LEP 1991–1993 data
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.019 ± 0.004 ± 0.004	31	GIBAUT	94B CLEO	Repl. by ANAS-TASSOV 01
0.051 ± 0.022	6	BYLSMA	87 HRS	$E_{\text{cm}}^{ee} = 29$ GeV
0.067 ± 0.030	5	¹⁸⁵ BELTRAMI	85 HRS	Repl. by BYLSMA 87

¹⁸⁵ The error quoted is statistical only.

$\Gamma(3h^- 2h^+ 2\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

< 0.011 90 GIBAUT 94B CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma((5\pi)^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{100}/Γ

$$\Gamma_{100}/\Gamma = (\Gamma_{28} + \Gamma_{45} + \Gamma_{74} + \Gamma_{97} + 0.553\Gamma_{117} + 0.888\Gamma_{138})/\Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

0.80 ± 0.07 OUR FIT

0.61 ± 0.06 ± 0.08 avg ¹⁸⁶ GIBAUT 94B CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

¹⁸⁶ Not independent of GIBAUT 94B $B(3h^- 2h^+ \nu_\tau)$, PROCARIO 93 $B(h^- 4\pi^0 \nu_\tau)$, and BORTOLETTO 93 $B(2h^- h^+ 2\pi^0 \nu_\tau)/B(\text{“3prong”})$ measurements. Result is corrected for η contributions.

$\Gamma(4h^- 3h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{“7-prong”}))/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 2.4 × 10⁻⁶ 90 EDWARDS 97B CLEO $E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.8 × 10⁻⁵ 95 ACKERSTAFF 97J OPAL 1990–1995 LEP runs

< 2.9 × 10⁻⁴ 90 BYLSMA 87 HRS $E_{\text{cm}}^{ee} = 29$ GeV

$\Gamma(X^- (S=-1) \nu_\tau)/\Gamma_{\text{total}}$ Γ_{102}/Γ

$$\Gamma_{102}/\Gamma = (\Gamma_{10} + \Gamma_{15} + \Gamma_{22} + \Gamma_{26} + \Gamma_{33} + \Gamma_{38} + \Gamma_{82} + \Gamma_{86} + \Gamma_{119})/\Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

2.93 ± 0.08 OUR FIT Error includes scale factor of 1.1.

2.87 ± 0.12 avg ¹⁸⁷ BARATE 99R ALEP 1991–1995 LEP runs

¹⁸⁷ BARATE 99R perform a combined analysis of all ALEPH LEP 1 data on τ branching fraction measurements for decay modes having total strangeness equal to -1 .


$\Gamma(K^*(892)^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ **Γ_{103} / Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.42 ± 0.18 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			

1.19 ± 0.15 ^{+0.13} _{-0.18}	104	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$
1.94 ± 0.27 ± 0.15	74	¹⁸⁸ AKERS	94G OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
1.43 ± 0.11 ± 0.13	475	¹⁸⁹ GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9 \text{ GeV}$

¹⁸⁸ AKERS 94G reject events in which a K_S^0 accompanies the $K^*(892)^-$. We do not correct for them.

¹⁸⁹ GOLDBERG 90 estimates that 10% of observed $K^*(892)^-$ are accompanied by a π^0 .



$\Gamma(K^*(892)^- \nu_\tau) / \Gamma_{\text{total}}$ **Γ_{104} / Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.29 ± 0.05 OUR AVERAGE				

1.326 ± 0.063		BARATE	99R ALEP	1991–1995 LEP runs
1.11 ± 0.12		¹⁹⁰ COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
1.42 ± 0.22 ± 0.09		¹⁹¹ ACCIARRI	95F L3	1991–1993 LEP runs
1.23 ± 0.21 ^{+0.11} _{-0.21}	54	¹⁹² ALBRECHT	88L ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
1.9 ± 0.3 ± 0.4	44	¹⁹³ TSCHIRHART	88 HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
1.5 ± 0.4 ± 0.4	15	¹⁹⁴ AIHARA	87C TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
1.3 ± 0.3 ± 0.3	31	YELTON	86 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.39 ± 0.09 ± 0.10		¹⁹⁵ BUSKULIC	96 ALEP	Repl. by BARATE 99R
1.45 ± 0.13 ± 0.11	273	¹⁹⁶ BUSKULIC	94F ALEP	Repl. by BUSKULIC 96
1.7 ± 0.7	11	DORFAN	81 MRK2	$E_{\text{cm}}^{ee} = 4.2\text{--}6.7 \text{ GeV}$

¹⁹⁰ Not independent of COAN 96 $B(\pi^- \bar{K}^0 \nu_\tau)$ and BATTLE 94 $B(K^- \pi^0 \nu_\tau)$ measurements. $K\pi$ final states are consistent with and assumed to originate from $K^*(892)^-$ production.

¹⁹¹ This result is obtained from their $B(\pi^- \bar{K}^0 \nu_\tau)$ assuming all those decays originate in $K^*(892)^-$ decays.

¹⁹² The authors divide by $\Gamma_2/\Gamma = 0.865$ to obtain this result.

¹⁹³ Not independent of TSCHIRHART 88 $\Gamma(\tau^- \rightarrow h^- \bar{K}^0 \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau)/\Gamma(\text{total})$.

¹⁹⁴ Decay π^- identified in this experiment, is assumed in the others.

¹⁹⁵ Not independent of BUSKULIC 96 $B(\pi^- \bar{K}^0 \nu_\tau)$ and $B(K^- \pi^0 \nu_\tau)$ measurements.

¹⁹⁶ BUSKULIC 94F obtain this result from BUSKULIC 94F $B(\bar{K}^0 \pi^- \nu_\tau)$ and BUSKULIC 94E $B(K^- \pi^0 \nu_\tau)$ assuming all of those decays originate in $K^*(892)^-$ decays.

$\Gamma(K^*(892)^- \nu_\tau)/\Gamma(\pi^- \pi^0 \nu_\tau)$ Γ_{104}/Γ_{13}

VALUE		DOCUMENT ID	TECN	COMMENT
0.075 ± 0.027		¹⁹⁷ ABREU	94K DLPH	LEP 1992 Z data

¹⁹⁷ ABREU 94K quote $B(\tau^- \rightarrow K^*(892)^- \nu_\tau)B(K^*(892)^- \rightarrow K^- \pi^0)/B(\tau^- \rightarrow \rho^- \nu_\tau) = 0.025 \pm 0.009$. We divide by $B(K^*(892)^- \rightarrow K^- \pi^0) = 0.333$ to obtain this result.

$\Gamma(K^*(892)^0 K^- \geq 0 \text{ neutrals} \nu_\tau)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.08 ± 0.12	119	GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9 \text{ GeV}$

$\Gamma(K^*(892)^0 K^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.21 ± 0.04 OUR AVERAGE				
0.213 ± 0.048		¹⁹⁸ BARATE	98 ALEP	1991–1995 LEP runs
0.20 ± 0.05 ± 0.04	47	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$

¹⁹⁸ BARATE 98 measure the $K^- (\rho^0 \rightarrow \pi^+ \pi^-)$ fraction in $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ decays to be $(35 \pm 11)\%$ and derive this result from their measurement of $\Gamma(\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ assuming the intermediate states are all $K^- \rho$ and $K^- K^*(892)^0$.

$\Gamma(\bar{K}^*(892)^0 \pi^- \geq 0 \text{ neutrals} \nu_\tau)/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.38 ± 0.11 ± 0.13	105	GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9 \text{ GeV}$

$\Gamma(\bar{K}^*(892)^0 \pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.22 ± 0.05 OUR AVERAGE				
0.209 ± 0.058		¹⁹⁹ BARATE	98 ALEP	1991–1995 LEP runs
0.25 ± 0.10 ± 0.05	27	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$

¹⁹⁹ BARATE 98 measure the $K^- K^*(892)^0$ fraction in $\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau$ decays to be $(87 \pm 13)\%$ and derive this result from their measurement of $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$.

$\Gamma((\bar{K}^*(892)\pi)^-\nu_\tau \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

0.10 ± 0.04 OUR AVERAGE

0.097 ± 0.044 ± 0.036	200 BARATE	99K ALEP	1991–1995 LEP runs
0.106 ± 0.037 ± 0.032	201 BARATE	98E ALEP	1991–1995 LEP runs

200 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter. They determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.72 \pm 0.12 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by one minus this fraction to obtain the quoted result.

201 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+\pi^-$ decays. They determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.64 \pm 0.09 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by one minus this fraction to obtain the quoted result.

$\Gamma(K_1(1270)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.47 ± 0.11 OUR AVERAGE

0.48 ± 0.11		BARATE	99R ALEP	1991–1995 LEP runs
0.41 ^{+0.41} _{-0.35} ± 0.10	5	202 BAUER	94 TPC	$E_{\text{cm}}^{ee} = 29$ GeV

202 We multiply 0.41% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$\Gamma(K_1(1400)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.17 ± 0.26 OUR AVERAGE Error includes scale factor of 1.7.

0.05 ± 0.17		BARATE	99R ALEP	1991–1995 LEP runs
0.76 ^{+0.40} _{-0.33} ± 0.20	11	203 BAUER	94 TPC	$E_{\text{cm}}^{ee} = 29$ GeV

203 We multiply 0.76% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$[\Gamma(K_1(1270)^-\nu_\tau) + \Gamma(K_1(1400)^-\nu_\tau)]/\Gamma_{\text{total}}$ $(\Gamma_{110} + \Gamma_{111})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.17 ^{+0.41} _{-0.37} ± 0.29	16	204 BAUER	94 TPC	1991–1995 LEP runs
---	----	-----------	--------	--------------------

204 We multiply 1.17% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error. Not independent of BAUER 94 $B(K_1(1270)^-\nu_\tau)$ and BAUER 94 $B(K_1(1400)^-\nu_\tau)$ measurements.

$\Gamma(K_1(1270)^-\nu_\tau)/[\Gamma(K_1(1270)^-\nu_\tau) + \Gamma(K_1(1400)^-\nu_\tau)]$ $\Gamma_{110}/(\Gamma_{110} + \Gamma_{111})$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.69 ± 0.15 OUR AVERAGE

0.71 ± 0.16 ± 0.11	205 ABBIENDI	00D OPAL	1990–1995 LEP runs
0.66 ± 0.19 ± 0.13	206 ASNER	00B CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

205 ABBIENDI 00D assume the resonance structure of $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ decays is dominated by the $K_1(1270)^-$ and $K_1(1400)^-$ resonances.

206 ASNER 00B assume the resonance structure of $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays is dominated by $K_1(1270)^-$ and $K_1(1400)^-$ resonances.

$\Gamma(K^*(1410)^- \nu_\tau) / \Gamma_{\text{total}}$ Γ_{112} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.5^{+1.4}_{-1.0}$	BARATE	99R ALEP	1991–1995 LEP runs

$\Gamma(K_0^*(1430)^- \nu_\tau) / \Gamma_{\text{total}}$ Γ_{113} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	95	BARATE	99R ALEP	1991–1995 LEP runs

$\Gamma(K_2^*(1430)^- \nu_\tau) / \Gamma_{\text{total}}$ Γ_{114} / Γ

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.3	95		TSCHIRHART 88	HRS	$E_{\text{cm}}^{ee} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.33	95	207	ACCIARRI	95F L3	1991–1993 LEP runs
<0.9	95	0	DORFAN	81 MRK2	$E_{\text{cm}}^{ee} = 4.2\text{--}6.7$ GeV

²⁰⁷ ACCIARRI 95F quote $B(\tau^- \rightarrow K^*(1430)^- \rightarrow \pi^- \bar{K}^0 \nu_\tau) < 0.11\%$. We divide by $B(K^*(1430)^- \rightarrow \pi^- \bar{K}^0) = 0.33$ to obtain the limit shown.

$\Gamma(a_0(980)^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}} \times B(a_0(980) \rightarrow K^0 K^-)$ $\Gamma_{115} / \Gamma \times B$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	GOLDBERG 90	CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9$ GeV

$\Gamma(\eta \pi^- \nu_\tau) / \Gamma_{\text{total}}$ Γ_{116} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 1.4	95	0	BARTELT	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.2	95		BUSKULIC	97C ALEP	1991–1994 LEP runs
< 3.4	95		ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
< 90	95		ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10$ GeV
<140	90		BEHREND	88 CELL	$E_{\text{cm}}^{ee} = 14\text{--}46.8$ GeV
<180	95		BARINGER	87 CLEO	$E_{\text{cm}}^{ee} = 10.5$ GeV
<250	90	0	COFFMAN	87 MRK3	$E_{\text{cm}}^{ee} = 3.77$ GeV
510 $\pm 100 \pm 120$		65	DERRICK	87 HRS	$E_{\text{cm}}^{ee} = 29$ GeV
<100	95		GAN	87B MRK2	$E_{\text{cm}}^{ee} = 29$ GeV

$\Gamma(\eta \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{117} / Γ

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.174 \pm 0.024 OUR FIT					
0.173 \pm 0.024 OUR AVERAGE					
0.18 $\pm 0.04 \pm 0.02$			BUSKULIC	97C ALEP	1991–1994 LEP runs
0.17 $\pm 0.02 \pm 0.02$		125	ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.10	95	ALBRECHT	88M ARG	$E_{cm}^{ee} \approx 10$ GeV
<2.10	95	BARINGER	87 CLEO	$E_{cm}^{ee} = 10.5$ GeV
$4.20^{+0.70}_{-1.20} \pm 1.60$		²⁰⁸ GAN	87 MRK2	$E_{cm}^{ee} = 29$ GeV

²⁰⁸ Highly correlated with GAN 87 $\Gamma(\pi^- 3\pi^0 \nu_\tau)/\Gamma(\text{total})$ value.

$\Gamma(\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{118}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.5±0.5		30	²⁰⁹ ANASTASSOV 01	CLEO	$E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.4 \pm 0.6 \pm 0.3$		15	²¹⁰ BERGFELD	97 CLEO	Repl. by ANAS-TASSOV 01
< 4.3	95		ARTUSO	92 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV
<120	95		ALBRECHT	88M ARG	$E_{cm}^{ee} \approx 10$ GeV

²⁰⁹ Weighted average of BERGFELD 97 and ANASTASSOV 01 value of $(1.5 \pm 0.6 \pm 0.3) \times 10^{-4}$ obtained using η 's reconstructed from $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.

²¹⁰ BERGFELD 97 reconstruct η 's using $\eta \rightarrow \gamma\gamma$ decays.

$\Gamma(\eta K^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.7±0.6 OUR FIT					
2.7±0.6 OUR AVERAGE					

$2.9^{+1.3}_{-1.2} \pm 0.7$			BUSKULIC	97c ALEP	1991–1994 LEP runs
$2.6 \pm 0.5 \pm 0.5$		85	BARTELT	96 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.7	95		ARTUSO	92 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV
------	----	--	--------	---------	--------------------------------

$\Gamma(\eta K^*(892)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.90±0.80±0.42	25	BISHAI	99 CLEO	$E_{cm}^{ee} = 10.6$ GeV

$\Gamma(\eta K^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.77±0.56±0.71	36	BISHAI	99 CLEO	$E_{cm}^{ee} = 10.6$ GeV

$\Gamma(\eta\bar{K}^0\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20±0.70±0.22	15	²¹¹ BISHAI	99 CLEO	$E_{cm}^{ee} = 10.6$ GeV

²¹¹ We multiply the BISHAI 99 measurement $B(\tau^- \rightarrow \eta K_S^0 \pi^- \nu_\tau) = (1.10 \pm 0.35 \pm 0.11) \times 10^{-4}$ by 2 to obtain the listed value.

$\Gamma(\eta\pi^+\pi^-\pi^-\geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.3	90	ABACHI	87B HRS	$E_{cm}^{ee} = 29$ GeV

$\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ **Γ_{124}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.3 ± 0.5	170	212 ANASTASSOV 01	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
---------------------------------	-----	-------------------	------	---------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.4^{+0.6}_{-0.5} \pm 0.6$	89	213 BERGFELD	97 CLEO	Repl. by ANASTASSOV 01
-----------------------------	----	--------------	---------	------------------------

212 Weighted average of BERGFELD 97 and ANASTASSOV 01 measurements using η 's reconstructed from $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow 3\pi^0$ decays.

213 BERGFELD 97 reconstruct η 's using $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi^0$ decays.

$\Gamma(\eta a_1(1260)^-\nu_\tau \rightarrow \eta\pi^-\rho^0\nu_\tau)/\Gamma_{\text{total}}$ **Γ_{125}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 3.9 \times 10^{-4}$	90	BERGFELD 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
---	----	-------------	------	---------------------------------

$\Gamma(\eta\eta\pi^-\nu_\tau)/\Gamma_{\text{total}}$ **Γ_{126}/Γ**

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 1.1	95	ARTUSO 92	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
------------------------------	----	-----------	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 83	95	ALBRECHT 88M ARG		$E_{\text{cm}}^{ee} \approx 10$ GeV
--------	----	------------------	--	-------------------------------------

$\Gamma(\eta\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ **Γ_{127}/Γ**

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 2.0	95	ARTUSO 92	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
------------------------------	----	-----------	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 90	95	ALBRECHT 88M ARG		$E_{\text{cm}}^{ee} \approx 10$ GeV
--------	----	------------------	--	-------------------------------------

$\Gamma(\eta'(958)\pi^-\nu_\tau)/\Gamma_{\text{total}}$ **Γ_{128}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 7.4 \times 10^{-5}$	90	BERGFELD 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
---	----	-------------	------	---------------------------------

$\Gamma(\eta'(958)\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ **Γ_{129}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 8.0 \times 10^{-5}$	90	BERGFELD 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
---	----	-------------	------	---------------------------------

$\Gamma(\phi\pi^-\nu_\tau)/\Gamma_{\text{total}}$ **Γ_{130}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 2.0 \times 10^{-4}$	90	214 AVERY 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
---	----	--------------	------	---------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.5 \times 10^{-4}$	90	ALBRECHT 95H ARG		$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
------------------------	----	------------------	--	---

214 AVERY 97 limit varies from $(1.2\text{--}2.0) \times 10^{-4}$ depending on decay model assumptions.

$\Gamma(\phi K^-\nu_\tau)/\Gamma_{\text{total}}$ **Γ_{131}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 6.7 \times 10^{-5}$	90	215 AVERY 97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
---	----	--------------	------	---------------------------------

215 AVERY 97 limit varies from $(5.4\text{--}6.7) \times 10^{-5}$ depending on decay model assumptions.

$\Gamma(f_1(1285)\pi^-\nu_\tau)/\Gamma_{\text{total}}$					Γ_{132}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$5.8^{+1.4}_{-1.3} \pm 1.8$	54	BERGFELD	97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(f_1(1285)\pi^-\nu_\tau \rightarrow \eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)$					$\Gamma_{133}/\Gamma_{124}$
VALUE		DOCUMENT ID	TECN	COMMENT	
0.55 ± 0.14		BERGFELD	97	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\pi(1300)^-\nu_\tau \rightarrow (\rho\pi)^-\nu_\tau \rightarrow (3\pi)^-\nu_\tau)/\Gamma_{\text{total}}$					Γ_{134}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.0 \times 10^{-4}$	90	ASNER	00	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(\pi(1300)^-\nu_\tau \rightarrow ((\pi\pi)_{S\text{-wave}}\pi)^-\nu_\tau \rightarrow (3\pi)^-\nu_\tau)/\Gamma_{\text{total}}$					Γ_{135}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.9 \times 10^{-4}$	90	ASNER	00	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(h^-\omega \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$					Γ_{136}/Γ
$\Gamma_{136}/\Gamma = (\Gamma_{137} + \Gamma_{138})/\Gamma$					

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
2.37 ± 0.08 OUR FIT					
1.65 ± 0.3 ± 0.2	avg	1513	ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10$ GeV

$\Gamma(h^-\omega\nu_\tau)/\Gamma_{\text{total}}$					Γ_{137}/Γ
Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.					

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
1.94 ± 0.07 OUR FIT					
1.92 ± 0.07 OUR AVERAGE					
1.91 ± 0.07 ± 0.06	f&a	5803	BUSKULIC	97C ALEP	1991–1994 LEP runs
1.95 ± 0.07 ± 0.11	avg	2223	²¹⁶ BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
1.60 ± 0.27 ± 0.41	f&a	139	BARINGER	87 CLEO	$E_{\text{cm}}^{ee} = 10.5$ GeV

²¹⁶ Not independent of BALEST 95C $B(\tau^- \rightarrow h^-\omega\nu_\tau)/B(\tau^- \rightarrow h^-h^+h^+\pi^0\nu_\tau)$ value.

$\Gamma(h^-\omega\nu_\tau)/\Gamma(h^-h^+h^+\pi^0\nu_\tau \text{ (ex. } K^0))$					Γ_{137}/Γ_{64}
$\Gamma_{137}/\Gamma_{64} = \Gamma_{137}/(\Gamma_{68} + \Gamma_{86} + \Gamma_{90} + 0.231\Gamma_{119} + 0.888\Gamma_{137} + 0.0221\Gamma_{138})$					

VALUE		EVTS	DOCUMENT ID	TECN	COMMENT
0.446 ± 0.015 OUR FIT					
0.453 ± 0.019 OUR AVERAGE					
0.431 ± 0.033		2350	²¹⁷ BUSKULIC	96 ALEP	LEP 1991–1993 data
0.464 ± 0.016 ± 0.017		2223	²¹⁸ BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.37 ± 0.05 ± 0.02		458	²¹⁹ ALBRECHT	91D ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV

217 BUSKULIC 96 quote the fraction of $\tau \rightarrow h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0) decays which originate in a $h^- \omega$ final state = 0.383 ± 0.029 . We divide this by the $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ branching fraction (0.888).

218 BALEST 95C quote the fraction of $\tau^- \rightarrow h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0) decays which originate in a $h^- \omega$ final state equals $0.412 \pm 0.014 \pm 0.015$. We divide this by the $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ branching fraction (0.888).

219 ALBRECHT 91D quote the fraction of $\tau^- \rightarrow h^- h^- h^+ \pi^0 \nu_\tau$ decays which originate in a $\pi^- \omega$ final state equals $0.33 \pm 0.04 \pm 0.02$. We divide this by the $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ branching fraction (0.888).

$\Gamma(h^- \omega \pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{138} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.43 ± 0.05 OUR FIT				
0.43 ± 0.06 ± 0.05	7283	BUSKULIC	97C ALEP	1991–1994 LEP runs

$\Gamma(h^- \omega 2\pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{139} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.4 ± 0.4 ± 0.3	53	ANASTASSOV 01	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.89^{+0.74}_{-0.67} \pm 0.40$	19	ANDERSON 97	CLEO	Repl. by ANASTASSOV 01
---------------------------------	----	-------------	------	------------------------

$\Gamma(h^- \omega \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau)$ $\Gamma_{138} / \Gamma_{52}$

$$\Gamma_{138} / \Gamma_{52} = \Gamma_{138} / (0.3431\Gamma_{33} + 0.3431\Gamma_{35} + 0.3431\Gamma_{38} + 0.3431\Gamma_{40} + 0.4307\Gamma_{45} + 0.6861\Gamma_{46} + \Gamma_{60} + \Gamma_{68} + \Gamma_{74} + \Gamma_{75} + \Gamma_{82} + \Gamma_{86} + \Gamma_{89} + \Gamma_{90} + 0.285\Gamma_{117} + 0.285\Gamma_{119} + 0.9101\Gamma_{137} + 0.9101\Gamma_{138})$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0286 ± 0.0031 OUR FIT				

0.028 ± 0.003 ± 0.003	avg	430 220 BORTOLETTO93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
------------------------------	-----	----------------------	------	---------------------------------------

220 Not independent of BORTOLETTO 93 $\Gamma(\tau^- \rightarrow h^- \omega \pi^0 \nu_\tau) / \Gamma(\tau^- \rightarrow h^- h^- h^+ 2\pi^0 \nu_\tau \text{ (ex. } K^0))$ value.

$\Gamma(h^- \omega \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau \text{ (ex. } K^0))$ $\Gamma_{138} / \Gamma_{73}$

$$\Gamma_{138} / \Gamma_{73} = \Gamma_{138} / (\Gamma_{74} + 0.236\Gamma_{117} + 0.888\Gamma_{138})$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.81 ± 0.08 OUR FIT			
0.81 ± 0.06 ± 0.06	BORTOLETTO93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

$\Gamma(2h^- h^+ \omega \nu_\tau) / \Gamma_{\text{total}}$ Γ_{140} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2 ± 0.2 ± 0.1	110	ANASTASSOV 01	CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$\Gamma(e^- \gamma)/\Gamma_{\text{total}}$ Γ_{141}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.9 \times 10^{-7}$	90	HAYASAKA 05	BELL	86.7 fb^{-1} , $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.7 \times 10^{-6}$	90	EDWARDS 97	CLEO	
$< 1.1 \times 10^{-4}$	90	ABREU 95U	DLPH	1990–1993 LEP runs
$< 1.2 \times 10^{-4}$	90	ALBRECHT 92k	ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 2.0 \times 10^{-4}$	90	KEH 88	CBAL	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 6.4 \times 10^{-4}$	90	HAYES 82	MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

$\Gamma(\mu^- \gamma)/\Gamma_{\text{total}}$ Γ_{142}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.1 \times 10^{-7}$	90	ABE 04B	BELL	86.3 fb^{-1} at $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.1 \times 10^{-6}$	90	AHMED 00	CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 3.0 \times 10^{-6}$	90	EDWARDS 97	CLEO	
$< 6.2 \times 10^{-5}$	90	ABREU 95U	DLPH	1990–1993 LEP runs
$< 0.42 \times 10^{-5}$	90	BEAN 93	CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 3.4 \times 10^{-5}$	90	ALBRECHT 92k	ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 55 \times 10^{-5}$	90	HAYES 82	MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

$\Gamma(e^- \pi^0)/\Gamma_{\text{total}}$ Γ_{143}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.7 \times 10^{-6}$	90	BONVICINI 97	CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 17 \times 10^{-5}$	90	ALBRECHT 92k	ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 14 \times 10^{-5}$	90	KEH 88	CBAL	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 210 \times 10^{-5}$	90	HAYES 82	MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

$\Gamma(\mu^- \pi^0)/\Gamma_{\text{total}}$ Γ_{144}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.0 \times 10^{-6}$	90	BONVICINI 97	CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 4.4 \times 10^{-5}$	90	ALBRECHT 92k	ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 82 \times 10^{-5}$	90	HAYES 82	MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

$\Gamma(e^- K_S^0)/\Gamma_{\text{total}}$ Γ_{145}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.1 \times 10^{-7}$	90	CHEN 02C	CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.3 \times 10^{-3}$	90	HAYES 82	MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$
------------------------	----	----------	------	--

$\Gamma(\mu^- K_S^0)/\Gamma_{\text{total}}$ Γ_{146}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.5 \times 10^{-7}$	90	CHEN	02C CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 1.0 \times 10^{-3}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

$\Gamma(e^- \eta)/\Gamma_{\text{total}}$ Γ_{147}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.2 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 6.3 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 24 \times 10^{-5}$	90	KEH	88 CBAL	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

$\Gamma(\mu^- \eta)/\Gamma_{\text{total}}$ Γ_{148}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.4 \times 10^{-7}$	90	ENARI	04 BELL	84.3 fb^{-1} at $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 9.6 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 7.3 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

$\Gamma(e^- \rho^0)/\Gamma_{\text{total}}$ Γ_{149}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.0 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 0.42 \times 10^{-5}$	90	²²¹ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.9 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 37 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²²¹ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \rho^0)/\Gamma_{\text{total}}$ Γ_{150}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 0.57 \times 10^{-5}$	90	²²² BARTELT	94 CLEO	Repl. by BLISS 98
$< 2.9 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 44 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²²² BARTELT 94 assume phase space decays.

$\Gamma(e^- K^*(892)^0)/\Gamma_{\text{total}}$ **Γ_{151}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.1 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.63 \times 10^{-5}$	90	²²³ BARTELT	94 CLEO	Repl. by BLISS 98
$<3.8 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

²²³ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- K^*(892)^0)/\Gamma_{\text{total}}$ **Γ_{152}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.5 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.94 \times 10^{-5}$	90	²²⁴ BARTELT	94 CLEO	Repl. by BLISS 98
$<4.5 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$

²²⁴ BARTELT 94 assume phase space decays.

$\Gamma(e^- \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ **Γ_{153}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.4 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-5}$	90	²²⁵ BARTELT	94 CLEO	Repl. by BLISS 98
-----------------------	----	------------------------	---------	-------------------

²²⁵ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ **Γ_{154}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.5 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.87 \times 10^{-5}$	90	²²⁶ BARTELT	94 CLEO	Repl. by BLISS 98
------------------------	----	------------------------	---------	-------------------

²²⁶ BARTELT 94 assume phase space decays.

$\Gamma(e^- \phi)/\Gamma_{\text{total}}$ **Γ_{155}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.9 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \phi)/\Gamma_{\text{total}}$ **Γ_{156}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.0 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{157}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.0 \times 10^{-7}$	90	AUBERT	04J BABR	$91.5 \text{ fb}^{-1} E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.5 \times 10^{-7}$	90	YUSA	04 BELL	87.1 fb^{-1} at $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 2.9 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 0.33 \times 10^{-5}$	90	²²⁷ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.3 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$
$< 40 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²²⁷ BARTELT 94 assume phase space decays.

$\Gamma(e^- \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{158}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.0 \times 10^{-7}$	90	YUSA	04 BELL	87.1 fb^{-1} at $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.3 \times 10^{-7}$	90	AUBERT	04J BABR	$91.5 \text{ fb}^{-1} E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 1.8 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 0.36 \times 10^{-5}$	90	²²⁸ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.9 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$
$< 33 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²²⁸ BARTELT 94 assume phase space decays.

$\Gamma(e^+ \mu^- \mu^-)/\Gamma_{\text{total}}$ **Γ_{159}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.3 \times 10^{-7}$	90	AUBERT	04J BABR	$91.5 \text{ fb}^{-1} E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.0 \times 10^{-7}$	90	YUSA	04 BELL	87.1 fb^{-1} at $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 1.5 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 0.35 \times 10^{-5}$	90	²²⁹ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.8 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 1.6 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²²⁹ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{160}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.9 \times 10^{-7}$	90	YUSA	04 BELL	87.1 fb^{-1} at $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.7 \times 10^{-7}$	90	AUBERT	04J BABR	91.5 fb^{-1} $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 1.7 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 0.34 \times 10^{-5}$	90	²³⁰ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.4 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$
$< 44 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²³⁰ BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ e^- e^-)/\Gamma_{\text{total}}$ **Γ_{161}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.1 \times 10^{-7}$	90	AUBERT	04J BABR	91.5 fb^{-1} $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.0 \times 10^{-7}$	90	YUSA	04 BELL	87.1 fb^{-1} at $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 1.5 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 0.34 \times 10^{-5}$	90	²³¹ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.4 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 1.6 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³¹ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{162}/Γ**

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.9 \times 10^{-7}$	90	AUBERT	04J BABR	91.5 fb^{-1} $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.0 \times 10^{-7}$	90	YUSA	04 BELL	87.1 fb^{-1} at $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 1.9 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
$< 0.43 \times 10^{-5}$	90	²³² BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.9 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 1.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$
$< 49 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{ee} = 3.8\text{--}6.8 \text{ GeV}$

²³² BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{163} / Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.2 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.44 \times 10^{-5}$	90	²³³ BARTELT	94 CLEO	Repl. by BLISS 98
$< 2.7 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 6.0 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³³ BARTELT 94 assume phase space decays.

$\Gamma(e^+ \pi^- \pi^-) / \Gamma_{\text{total}}$ Γ_{164} / Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.9 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.44 \times 10^{-5}$	90	²³⁴ BARTELT	94 CLEO	Repl. by BLISS 98
$< 1.8 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 1.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³⁴ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{165} / Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.2 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.74 \times 10^{-5}$	90	²³⁵ BARTELT	94 CLEO	Repl. by BLISS 98
$< 3.6 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 3.9 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³⁵ BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ \pi^- \pi^-) / \Gamma_{\text{total}}$ Γ_{166} / Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.4 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.69 \times 10^{-5}$	90	²³⁶ BARTELT	94 CLEO	Repl. by BLISS 98
$< 6.3 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 3.9 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³⁶ BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^+ K^-) / \Gamma_{\text{total}}$ Γ_{167} / Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.4 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.77 \times 10^{-5}$	90	²³⁷ BARTELT	94 CLEO	Repl. by BLISS 98
$< 2.9 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 5.8 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³⁷ BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^- K^+)/\Gamma_{\text{total}}$ Γ_{168}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.8 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<0.46 \times 10^{-5}$	90	²³⁸ BARTELT	94 CLEO	Repl. by BLISS 98
$<5.8 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³⁸ BARTELT 94 assume phase space decays.

$\Gamma(e^+ \pi^- K^-)/\Gamma_{\text{total}}$ Γ_{169}/Γ

Test of lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.1 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<0.45 \times 10^{-5}$	90	²³⁹ BARTELT	94 CLEO	Repl. by BLISS 98
$<2.0 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$<4.9 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²³⁹ BARTELT 94 assume phase space decays.

$\Gamma(e^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{170}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.2 \times 10^{-6}$	90	CHEN	02c CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{171}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.0 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^+ K^- K^-)/\Gamma_{\text{total}}$ Γ_{172}/Γ

Test of lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.8 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \pi^+ K^-)/\Gamma_{\text{total}}$ Γ_{173}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 7.5 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.87 \times 10^{-5}$	90	²⁴⁰ BARTELT	94 CLEO	Repl. by BLISS 98
$< 11 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$< 7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²⁴⁰ BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \pi^- K^+)/\Gamma_{\text{total}}$ Γ_{174}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.4 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.5 \times 10^{-5}$	90	²⁴¹ BARTELT	94 CLEO	Repl. by BLISS 98
$<7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²⁴¹ BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ \pi^- K^-)/\Gamma_{\text{total}}$ Γ_{175}/Γ

Test of lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.0 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<2.0 \times 10^{-5}$	90	²⁴² BARTELT	94 CLEO	Repl. by BLISS 98
$<5.8 \times 10^{-5}$	90	ALBRECHT	92k ARG	$E_{\text{cm}}^{ee} = 10 \text{ GeV}$
$<4.0 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{\text{cm}}^{ee} = 10.4\text{--}10.9$

²⁴² BARTELT 94 assume phase space decays.

$\Gamma(\mu^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{176}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.4 \times 10^{-6}$	90	CHEN	02c CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{177}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<15 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^+ K^- K^-)/\Gamma_{\text{total}}$ Γ_{178}/Γ

Test of lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.0 \times 10^{-6}$	90	BLISS	98 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{179}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.5 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{180}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<14 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- \eta \eta)/\Gamma_{\text{total}}$ Γ_{181}/Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<35 \times 10^{-6}$	90	BONVICINI	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \eta \eta) / \Gamma_{\text{total}}$ Γ_{182} / Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 60 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- \pi^0 \eta) / \Gamma_{\text{total}}$ Γ_{183} / Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 24 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\mu^- \pi^0 \eta) / \Gamma_{\text{total}}$ Γ_{184} / Γ

Test of lepton family number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 22 \times 10^{-6}$	90	BONVICINI	97	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\bar{p} \gamma) / \Gamma_{\text{total}}$ Γ_{185} / Γ

Test of lepton number and baryon number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 3.5 \times 10^{-6}$	90	GODANG	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 29 \times 10^{-5}$	90	ALBRECHT	92K	ARG $E_{\text{cm}}^{ee} = 10 \text{ GeV}$
-----------------------	----	----------	-----	---

$\Gamma(\bar{p} \pi^0) / \Gamma_{\text{total}}$ Γ_{186} / Γ

Test of lepton number and baryon number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 15 \times 10^{-6}$	90	GODANG	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 66 \times 10^{-5}$	90	ALBRECHT	92K	ARG $E_{\text{cm}}^{ee} = 10 \text{ GeV}$
-----------------------	----	----------	-----	---

$\Gamma(\bar{p} 2\pi^0) / \Gamma_{\text{total}}$ Γ_{187} / Γ

Test of lepton number and baryon number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 33 \times 10^{-6}$	90	GODANG	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(\bar{p} \eta) / \Gamma_{\text{total}}$ Γ_{188} / Γ

Test of lepton number and baryon number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 8.9 \times 10^{-6}$	90	GODANG	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 130 \times 10^{-5}$	90	ALBRECHT	92K	ARG $E_{\text{cm}}^{ee} = 10 \text{ GeV}$
------------------------	----	----------	-----	---

$\Gamma(\bar{p} \pi^0 \eta) / \Gamma_{\text{total}}$ Γ_{189} / Γ

Test of lepton number and baryon number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 27 \times 10^{-6}$	90	GODANG	99	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$\Gamma(e^- \text{ light boson})/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ **Γ_{190}/Γ_5**

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.015	95	243 ALBRECHT	95G ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.018	95	244 ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV
<0.040	95	245 BALTRUSAIT..85	MRK3	$E_{cm}^{ee} = 3.77$ GeV

243 ALBRECHT 95G limit holds for bosons with mass < 0.4 GeV. The limit rises to 0.036 for a mass of 1.0 GeV, then falls to 0.006 at the upper mass limit of 1.6 GeV.

244 ALBRECHT 90E limit applies for spinless boson with mass < 100 MeV, and rises to 0.050 for mass = 500 MeV.

245 BALTRUSAITIS 85 limit applies for spinless boson with mass < 100 MeV.

$\Gamma(\mu^- \text{ light boson})/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ **Γ_{191}/Γ_5**

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.026	95	246 ALBRECHT	95G ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.033	95	247 ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV
<0.125	95	248 BALTRUSAIT..85	MRK3	$E_{cm}^{ee} = 3.77$ GeV

246 ALBRECHT 95G limit holds for bosons with mass < 1.3 GeV. The limit rises to 0.034 for a mass of 1.4 GeV, then falls to 0.003 at the upper mass limit of 1.6 GeV.

247 ALBRECHT 90E limit applies for spinless boson with mass < 100 MeV, and rises to 0.071 for mass = 500 MeV.

248 BALTRUSAITIS 85 limit applies for spinless boson with mass < 100 MeV.

τ -DECAY PARAMETERS

A REVIEW GOES HERE – Check our WWW List of Reviews

$\rho^\tau(e \text{ or } \mu)$ PARAMETER

(V–A) theory predicts $\rho = 0.75$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.745±0.008 OUR FIT				
0.749±0.008 OUR AVERAGE				
0.742±0.014±0.006	81k	HEISTER	01E ALEP	1991–1995 LEP runs
0.775±0.023±0.020	36k	ABREU	00L DLPH	1992–1995 runs
0.781±0.028±0.018	46k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.762±0.035	54k	ACCIARRI	98R L3	1991–1995 LEP runs
0.731±0.031		249 ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.72 ±0.09 ±0.03		250 ABE	97O SLD	1993–1995 SLC runs
0.747±0.010±0.006	55k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
0.79 ±0.10 ±0.10	3732	FORD	87B MAC	$E_{cm}^{ee} = 29$ GeV
0.71 ±0.09 ±0.03	1426	BEHRENDIS	85 CLEO	$e^+ e^-$ near $\Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.735±0.013±0.008	31k	AMMAR	97B CLEO	Repl. by ALEXANDER 97F
0.794±0.039±0.031	18k	ACCIARRI	96H L3	Repl. by ACCIARRI 98R
0.732±0.034±0.020	8.2k	251 ALBRECHT	95 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.738±0.038		252 ALBRECHT	95C ARG	Repl. by ALBRECHT 98
0.751±0.039±0.022		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
0.742±0.035±0.020	8000	ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV

- 249 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.
- 250 ABE 97O assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a ρ^τ value of $0.69 \pm 0.13 \pm 0.05$.
- 251 Value is from a simultaneous fit for the ρ^τ and η^τ decay parameters to the lepton energy spectrum. Not independent of ALBRECHT 90E ρ^τ (e or μ) value which assumes $\eta^\tau = 0$. Result is strongly correlated with ALBRECHT 95C.
- 252 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E.

$\rho^\tau(e)$ PARAMETER

(V-A) theory predicts $\rho = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.747±0.010 OUR FIT				
0.744±0.010 OUR AVERAGE				
0.747±0.019±0.014	44k	HEISTER	01E ALEP	1991–1995 LEP runs
0.744±0.036±0.037	17k	ABREU	00L DLPH	1992–1995 runs
0.779±0.047±0.029	25k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.68 ±0.04 ±0.07		253 ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.71 ±0.14 ±0.05		ABE	97O SLD	1993–1995 SLC runs
0.747±0.012±0.004	34k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
0.735±0.036±0.020	4.7k	254 ALBRECHT	95 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.79 ±0.08 ±0.06	3230	255 ALBRECHT	93G ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV
0.64 ±0.06 ±0.07	2753	JANSSEN	89 CBAL	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV
0.62 ±0.17 ±0.14	1823	FORD	87B MAC	$E_{cm}^{ee} = 29$ GeV
0.60 ±0.13	699	BEHREND	85 CLEO	$e^+ e^-$ near $\Upsilon(4S)$
0.72 ±0.10 ±0.11	594	BACINO	79B DLCO	$E_{cm}^{ee} = 3.5\text{--}7.4$ GeV

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.732±0.014±0.009	19k	AMMAR	97B CLEO	Repl. by ALEXANDER 97F
0.793±0.050±0.025		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
0.747±0.045±0.028	5106	ALBRECHT	90E ARG	Repl. by ALBRECHT 95
253 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.				
254 ALBRECHT 95 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(h^+ h^- h^+ (\pi^0) \bar{\nu}_\tau)$ and their charged conjugates.				
255 ALBRECHT 93G use tau pair events of the type $\tau^- \tau^+ \rightarrow (\mu^- \bar{\nu}_\mu \nu_\tau)(e^+ \nu_e \bar{\nu}_\tau)$ and their charged conjugates.				

$\rho^\tau(\mu)$ PARAMETER

(V-A) theory predicts $\rho = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.763±0.020 OUR FIT				
0.770±0.022 OUR AVERAGE				
0.776±0.045±0.019	46k	HEISTER	01E ALEP	1991–1995 LEP runs
0.999±0.098±0.045	22k	ABREU	00L DLPH	1992–1995 runs
0.777±0.044±0.016	27k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.69 ±0.06 ±0.06		256 ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.54 ±0.28 ±0.14		ABE	97O SLD	1993–1995 SLC runs

$0.750 \pm 0.017 \pm 0.045$	22k	ALEXANDER	97F CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV
$0.76 \pm 0.07 \pm 0.08$	3230	ALBRECHT	93G ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
$0.734 \pm 0.055 \pm 0.027$	3041	ALBRECHT	90E ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
$0.89 \pm 0.14 \pm 0.08$	1909	FORD	87B MAC	$E_{\text{cm}}^{ee} = 29$ GeV
0.81 ± 0.13	727	BEHREND	85 CLEO	e^+e^- near $\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.747 \pm 0.048 \pm 0.044$	13k	AMMAR	97B CLEO	Repl. by ALEXANDER 97F
$0.693 \pm 0.057 \pm 0.028$		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
²⁵⁶ ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.				

$\xi^\tau(e \text{ or } \mu)$ PARAMETER

($V-A$) theory predicts $\xi = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.985 ± 0.030 OUR FIT

0.981 ± 0.031 OUR AVERAGE

$0.986 \pm 0.068 \pm 0.031$	81k	HEISTER	01E ALEP	1991–1995 LEP runs
$0.929 \pm 0.070 \pm 0.030$	36k	ABREU	00L DLPH	1992–1995 runs
$0.98 \pm 0.22 \pm 0.10$	46k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.70 ± 0.16	54k	ACCIARRI	98R L3	1991–1995 LEP runs
1.03 ± 0.11		²⁵⁷ ALBRECHT	98 ARG	$E_{\text{cm}}^{ee} = 9.5\text{--}10.6$ GeV
$1.05 \pm 0.35 \pm 0.04$		²⁵⁸ ABE	97O SLD	1993–1995 SLC runs
$1.007 \pm 0.040 \pm 0.015$	55k	ALEXANDER	97F CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.94 \pm 0.21 \pm 0.07$	18k	ACCIARRI	96H L3	Repl. by ACCIARRI 98R
0.97 ± 0.14		²⁵⁹ ALBRECHT	95C ARG	Repl. by ALBRECHT 98
$1.18 \pm 0.15 \pm 0.16$		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
$0.90 \pm 0.15 \pm 0.10$	3230	²⁶⁰ ALBRECHT	93G ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV

²⁵⁷ Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

²⁵⁸ ABE 97O assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a ξ^τ value of $1.02 \pm 0.36 \pm 0.05$.

²⁵⁹ Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 95C uses events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(h^+ h^- h^+ \bar{\nu}_\tau)$ and their charged conjugates.

²⁶⁰ ALBRECHT 93G measurement determines $|\xi^\tau|$ for the case $\xi^\tau(e) = \xi^\tau(\mu)$, but the authors point out that other LEP experiments determine the sign to be positive.

$\xi^\tau(e)$ PARAMETER

($V-A$) theory predicts $\xi = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.994 ± 0.040 OUR FIT

1.00 ± 0.04 OUR AVERAGE

$1.011 \pm 0.094 \pm 0.038$	44k	HEISTER	01E ALEP	1991–1995 LEP runs
$1.01 \pm 0.12 \pm 0.05$	17k	ABREU	00L DLPH	1992–1995 runs
$1.13 \pm 0.39 \pm 0.14$	25k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs

1.11 ±0.20 ±0.08		²⁶¹ ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
1.16 ±0.52 ±0.06		ABE	97O SLD	1993–1995 SLC runs
0.979±0.048±0.016	34k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.03 ±0.23 ±0.09		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
²⁶¹ ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.				

$\xi^\tau(\mu)$ PARAMETER

(V–A) theory predicts $\xi = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.030±0.059 OUR FIT				
1.06 ±0.06 OUR AVERAGE				
1.030±0.120±0.050	46k	HEISTER	01E ALEP	1991–1995 LEP runs
1.16 ±0.19 ±0.06	22k	ABREU	00L DLPH	1992–1995 runs
0.79 ±0.41 ±0.09	27k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
1.26 ±0.27 ±0.14		²⁶² ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.75 ±0.50 ±0.14		ABE	97O SLD	1993–1995 SLC runs
1.054±0.069±0.047	22k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.23 ±0.22 ±0.10		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
²⁶² ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.				

$\eta^\tau(e \text{ or } \mu)$ PARAMETER

(V–A) theory predicts $\eta = 0$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.013±0.020 OUR FIT				
0.015±0.021 OUR AVERAGE				
0.012±0.026±0.004	81k	HEISTER	01E ALEP	1991–1995 LEP runs
–0.005±0.036±0.037		ABREU	00L DLPH	1992–1995 runs
0.027±0.055±0.005	46k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.27 ±0.14	54k	ACCIARRI	98R L3	1991–1995 LEP runs
–0.13 ±0.47 ±0.15		ABE	97O SLD	1993–1995 SLC runs
–0.015±0.061±0.062	31k	AMMAR	97B CLEO	$E_{cm}^{ee} = 10.6$ GeV
0.03 ±0.18 ±0.12	8.2k	ALBRECHT	95 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.25 ±0.17 ±0.11	18k	ACCIARRI	96H L3	Repl. by ACCIARRI 98R
–0.04 ±0.15 ±0.11		BUSKULIC	95D ALEP	Repl. by HEISTER 01E

$\eta^\tau(\mu)$ PARAMETER

(V–A) theory predicts $\eta = 0$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.094±0.073 OUR FIT				
0.17 ±0.15 OUR AVERAGE Error includes scale factor of 1.2.				
0.160±0.150±0.060	46k	HEISTER	01E ALEP	1991–1995 LEP runs
0.72 ±0.32 ±0.15		ABREU	00L DLPH	1992–1995 runs
–0.59 ±0.82 ±0.45		²⁶³ ABE	97O SLD	1993–1995 SLC runs
0.010±0.149±0.171	13k	²⁶⁴ AMMAR	97B CLEO	$E_{cm}^{ee} = 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				

- 0.010±0.065±0.001 27k ²⁶⁵ACKERSTAFF 99D OPAL 1990–1995 LEP runs
 –0.24 ±0.23 ±0.18 BUSKULIC 95D ALEP Repl. by HEISTER 01E
- ²⁶³ Highly correlated (corr. = 0.92) with ABE 97O $\rho^T(\mu)$ measurement.
²⁶⁴ Highly correlated (corr. = 0.949) with AMMAR 97B $\rho^T(\mu)$ value.
²⁶⁵ ACKERSTAFF 99D result is dominated by a constraint on η^T from the OPAL measurements of the τ lifetime and $B(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)$ assuming lepton universality for the total coupling strength.

$(\delta\xi)^T(\text{e or } \mu)$ PARAMETER

(V–A) theory predicts $(\delta\xi) = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.746±0.021 OUR FIT				
0.744±0.022 OUR AVERAGE				
0.776±0.045±0.024	81k	HEISTER	01E ALEP	1991–1995 LEP runs
0.779±0.070±0.028	36k	ABREU	00L DLPH	1992–1995 runs
0.65 ±0.14 ±0.07	46k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.70 ±0.11	54k	ACCIARRI	98R L3	1991–1995 LEP runs
0.63 ±0.09		²⁶⁶ ALBRECHT	98 ARG	$E_{\text{cm}}^{ee} = 9.5\text{--}10.6$ GeV
0.88 ±0.27 ±0.04		²⁶⁷ ABE	97O SLD	1993–1995 SLC runs
0.745±0.026±0.009	55k	ALEXANDER	97F CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.81 ±0.14 ±0.06	18k	ACCIARRI	96H L3	Repl. by ACCIARRI 98R
0.65 ±0.12		²⁶⁸ ALBRECHT	95C ARG	Repl. by ALBRECHT 98
0.88 ±0.11 ±0.07		BUSKULIC	95D ALEP	Repl. by HEISTER 01E

- ²⁶⁶ Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.
²⁶⁷ ABE 97O assume $\eta^T = 0$ in their fit. Letting η^T vary in the fit gives a $(\rho\xi)^T$ value of $0.87 \pm 0.27 \pm 0.04$.
²⁶⁸ Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 95C uses events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(h^+ h^- h^+ \bar{\nu}_\tau)$ and their charged conjugates.

$(\delta\xi)^T(\text{e})$ PARAMETER

(V–A) theory predicts $(\delta\xi) = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.734±0.028 OUR FIT				
0.731±0.029 OUR AVERAGE				
0.778±0.066±0.024	44k	HEISTER	01E ALEP	1991–1995 LEP runs
0.85 ±0.12 ±0.04	17k	ABREU	00L DLPH	1992–1995 runs
0.72 ±0.31 ±0.14	25k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.56 ±0.14 ±0.06		²⁶⁹ ALBRECHT	98 ARG	$E_{\text{cm}}^{ee} = 9.5\text{--}10.6$ GeV
0.85 ±0.43 ±0.08		ABE	97O SLD	1993–1995 SLC runs
0.720±0.032±0.010	34k	ALEXANDER	97F CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.11 ±0.17 ±0.07		BUSKULIC	95D ALEP	Repl. by HEISTER 01E
------------------	--	----------	----------	----------------------

²⁶⁹ ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

$(\delta\xi)^T(\mu)$ PARAMETER

($V-A$) theory predicts $(\delta\xi) = 0.75$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.778±0.037 OUR FIT

0.79 ±0.04 OUR AVERAGE

0.786±0.066±0.028	46k	HEISTER	01E ALEP	1991–1995 LEP runs
0.86 ±0.13 ±0.04	22k	ABREU	00L DLPH	1992–1995 runs
0.63 ±0.23 ±0.05	27k	ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.73 ±0.18 ±0.10		²⁷⁰ ALBRECHT	98 ARG	$E_{cm}^{ee} = 9.5\text{--}10.6$ GeV
0.82 ±0.32 ±0.07		ABE	97O SLD	1993–1995 SLC runs
0.786±0.041±0.032	22k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.71 ±0.14 ±0.06 BUSKULIC 95D ALEP Repl. by HEISTER 01E
²⁷⁰ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

$\xi^T(\pi)$ PARAMETER

($V-A$) theory predicts $\xi^T(\pi) = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.993±0.022 OUR FIT

0.994±0.023 OUR AVERAGE

0.994±0.020±0.014	27k	HEISTER	01E ALEP	1991–1995 LEP runs
0.81 ±0.17 ±0.02		ABE	97O SLD	1993–1995 SLC runs
1.03 ±0.06 ±0.04	2.0k	COAN	97 CLEO	$E_{cm}^{ee} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.987±0.057±0.027 BUSKULIC 95D ALEP Repl. by HEISTER 01E
 0.95 ±0.11 ±0.05 ²⁷¹BUSKULIC 94D ALEP 1990+1991 LEP run
²⁷¹Superseded by BUSKULIC 95D.

$\xi^T(\rho)$ PARAMETER

($V-A$) theory predicts $\xi^T(\rho) = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.994±0.008 OUR FIT

0.994±0.009 OUR AVERAGE

0.987±0.012±0.011	59k	HEISTER	01E ALEP	1991–1995 LEP runs
0.99 ±0.12 ±0.04		ABE	97O SLD	1993–1995 SLC runs
0.995±0.010±0.003	66k	ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.022±0.028±0.030	1.7k	²⁷² ALBRECHT	94E ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.045±0.058±0.032 BUSKULIC 95D ALEP Repl. by HEISTER 01E
 1.03 ±0.11 ±0.05 ²⁷³BUSKULIC 94D ALEP 1990+1991 LEP run
²⁷²ALBRECHT 94E measure the square of this quantity and use the sign determined by ALBRECHT 90I to obtain the quoted result.
²⁷³Superseded by BUSKULIC 95D.

$\xi^T(a_1)$ PARAMETER(V-A) theory predicts $\xi^T(a_1) = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

1.001±0.027 OUR FIT**1.002±0.028 OUR AVERAGE**

1.000±0.016±0.024	35k	274 HEISTER	01E ALEP	1991–1995 LEP runs
1.02 ±0.13 ±0.03	17.2k	ASNER	00 CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.29 ±0.26 ±0.11	7.4k	275 ACKERSTAFF	97R OPAL	1992–1994 LEP runs
0.85 $\begin{smallmatrix} +0.15 \\ -0.17 \end{smallmatrix}$ ±0.05		ALBRECHT	95C ARG	$E_{cm}^{ee} = 9.5$ –10.6 GeV
1.25 ±0.23 $\begin{smallmatrix} +0.15 \\ -0.08 \end{smallmatrix}$	7.5k	ALBRECHT	93C ARG	$E_{cm}^{ee} = 9.4$ –10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.08 $\begin{smallmatrix} +0.46 \\ -0.41 \end{smallmatrix}$ $\begin{smallmatrix} +0.14 \\ -0.25 \end{smallmatrix}$	2.6k	276 AKERS	95P OPAL	Repl. by ACKER-STAFF 97R
0.937±0.116±0.064		BUSKULIC	95D ALEP	Repl. by HEISTER 01E

274 HEISTER 01E quote $1.000 \pm 0.016 \pm 0.013 \pm 0.020$ where the errors are statistical, systematic, and an uncertainty due to the final state model. We combine the systematic error and model uncertainty.

275 ACKERSTAFF 97R obtain this result with a model independent fit to the hadronic structure functions. Fitting with the model of Kuhn and Santamaria (ZPHY **C48**, 445 (1990)) gives $0.87 \pm 0.16 \pm 0.04$, and with the model of of Isgur *et al.* (PR **D39**,1357 (1989)) they obtain $1.20 \pm 0.21 \pm 0.14$.

276 AKERS 95P obtain this result with a model independent fit to the hadronic structure functions. Fitting with the model of Kuhn and Santamaria (ZPHY **C48**, 445 (1990)) gives $0.87 \pm 0.27 \begin{smallmatrix} +0.05 \\ -0.06 \end{smallmatrix}$, and with the model of of Isgur *et al.* (PR **D39**,1357 (1989)) they obtain $1.10 \pm 0.31 \begin{smallmatrix} +0.13 \\ -0.14 \end{smallmatrix}$.

 ξ^T (all hadronic modes) PARAMETER(V-A) theory predicts $\xi^T = 1$.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.995±0.007 OUR FIT**0.997±0.007 OUR AVERAGE**

0.992±0.007±0.008	102k	277 HEISTER	01E ALEP	1991–1995 LEP runs
0.997±0.027±0.011	39k	278 ABREU	00L DLPH	1992–1995 runs
1.02 ±0.13 ±0.03	17.2k	279 ASNER	00 CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.032±0.031	37k	280 ACCIARRI	98R L3	1991–1995 LEP runs
0.93 ±0.10 ±0.04		ABE	97O SLD	1993–1995 SLC runs
1.29 ±0.26 ±0.11	7.4k	281 ACKERSTAFF	97R OPAL	1992–1994 LEP runs
0.995±0.010±0.003	66k	282 ALEXANDER	97F CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.03 ±0.06 ±0.04	2.0k	283 COAN	97 CLEO	$E_{cm}^{ee} = 10.6$ GeV
1.017±0.039		284 ALBRECHT	95C ARG	$E_{cm}^{ee} = 9.5$ –10.6 GeV
1.25 ±0.23 $\begin{smallmatrix} +0.15 \\ -0.08 \end{smallmatrix}$	7.5k	285 ALBRECHT	93C ARG	$E_{cm}^{ee} = 9.4$ –10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.970±0.053±0.011	14k	286 ACCIARRI	96H L3	Repl. by ACCIARRI 98R
1.08 $\begin{smallmatrix} +0.46 \\ -0.41 \end{smallmatrix}$ $\begin{smallmatrix} +0.14 \\ -0.25 \end{smallmatrix}$	2.6k	287 AKERS	95P OPAL	Repl. by ACKER-STAFF 97R
1.006±0.032±0.019		288 BUSKULIC	95D ALEP	Repl. by HEISTER 01E
1.022±0.028±0.030	1.7k	289 ALBRECHT	94E ARG	$E_{cm}^{ee} = 9.4$ –10.6 GeV
0.99 ±0.07 ±0.04		290 BUSKULIC	94D ALEP	1990+1991 LEP run

- 277 HEISTER 01E quote $0.992 \pm 0.007 \pm 0.006 \pm 0.005$ where the errors are statistical, systematic, and an uncertainty due to the final state model. We combine the systematic error and model uncertainty. They use $\tau \rightarrow \pi \nu_\tau$, $\tau \rightarrow K \nu_\tau$, $\tau \rightarrow \rho \nu_\tau$, and $\tau \rightarrow a_1 \nu_\tau$ decays.
- 278 ABREU 00L use $\tau^- \rightarrow h^- \geq 0\pi^0 \nu_\tau$ decays.
- 279 ASNER 00 use $\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau$ decays.
- 280 ACCIARRI 98R use $\tau \rightarrow \pi \nu_\tau$, $\tau \rightarrow K \nu_\tau$, and $\tau \rightarrow \rho \nu_\tau$ decays.
- 281 ACKERSTAFF 97R use $\tau \rightarrow a_1 \nu_\tau$ decays.
- 282 ALEXANDER 97F use $\tau \rightarrow \rho \nu_\tau$ decays.
- 283 COAN 97 use $h^+ h^-$ energy correlations.
- 284 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E.
- 285 Uses $\tau \rightarrow a_1 \nu_\tau$ decays. Replaced by ALBRECHT 95C.
- 286 ACCIARRI 96H use $\tau \rightarrow \pi \nu_\tau$, $\tau \rightarrow K \nu_\tau$, and $\tau \rightarrow \rho \nu_\tau$ decays.
- 287 AKERS 95P use $\tau \rightarrow a_1 \nu_\tau$ decays.
- 288 BUSKULIC 95D use $\tau \rightarrow \pi \nu_\tau$, $\tau \rightarrow \rho \nu_\tau$, and $\tau \rightarrow a_1 \nu_\tau$ decays.
- 289 ALBRECHT 94E measure the square of this quantity and use the sign determined by ALBRECHT 90I to obtain the quoted result. Uses $\tau \rightarrow a_1 \nu_\tau$ decays. Replaced by ALBRECHT 95C.
- 290 BUSKULIC 94D use $\tau \rightarrow \pi \nu_\tau$ and $\tau \rightarrow \rho \nu_\tau$ decays. Superseded by BUSKULIC 95D.

τ REFERENCES

HAYASAKA	05	PL B613 20	K. Hayasaka <i>et al.</i>	(BELLE Collab.)
ABBIENDI	04J	EPJ C35 437	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	04K	EPJ C35 159	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	04T	EPJ C36 283	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABE	04B	PRL 92 171802	K. Abe <i>et al.</i>	(BELLE Collab.)
ACHARD	04G	PL B585 53	P. Achard <i>et al.</i>	(L3 Collab.)
AUBERT	04J	PRL 92 121801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ENARI	04	PRL 93 081803	Y. Enari <i>et al.</i>	(BELLE Collab.)
YUSA	04	PL B589 103	Y. Yusa <i>et al.</i>	(BELLE Collab.)
ABBIENDI	03	PL B551 35	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
BRIERE	03	PRL 90 181802	R. A. Briere <i>et al.</i>	(CLEO Collab.)
HEISTER	03F	EPJ C30 291	A. Heister <i>et al.</i>	(ALEPH Collab.)
INAMI	03	PL B551 16	K. Inami <i>et al.</i>	(BELLE Collab.)
CHEN	02C	PR D66 071101R	S. Chen <i>et al.</i>	(CLEO Collab.)
ABBIENDI	01J	EPJ C19 653	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	01M	EPJ C20 617	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	01F	PL B507 47	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACHARD	01D	PL B519 189	P. Achard <i>et al.</i>	(L3 Collab.)
ANASTASSOV	01	PRL 86 4467	A. Anastassov <i>et al.</i>	(CLEO Collab.)
HEISTER	01E	EPJ C22 217	A. Heister <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	00A	PL B492 23	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00C	EPJ C13 213	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00D	EPJ C13 197	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	00L	EPJ C16 229	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00B	PL B479 67	M. Acciarri <i>et al.</i>	(L3 Collab.)
AHMED	00	PR D61 071101R	S. Ahmed <i>et al.</i>	(CLEO Collab.)
ALBRECHT	00	PL B485 37	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BERGFELD	00	PRL 84 830	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BROWDER	00	PR D61 052004	T.E. Browder <i>et al.</i>	(CLEO Collab.)
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GONZALEZ-S...	00	NP B582 3	G.A. Gonzalez-Sprinberg <i>et al.</i>	
ABBIENDI	99H	PL B447 134	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	99X	EPJ C10 201	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	99D	EPJ C8 3	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	99E	EPJ C8 183	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	99K	EPJ C10 1	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)

BISHAI	99	PRL 82 281	M. Bishai <i>et al.</i>	(CLEO Collab.)
GODANG	99	PR D59 091303	R. Godang <i>et al.</i>	(CLEO Collab.)
RICHICHI	99	PR D60 112002	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ACCIARRI	98C	PL B426 207	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98E	PL B434 169	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98R	PL B438 405	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	98M	EPJ C4 193	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98N	PL B431 188	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ALBRECHT	98	PL B431 179	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARATE	98	EPJ C1 65	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98E	EPJ C4 29	R. Barate <i>et al.</i>	(ALEPH Collab.)
BLISS	98	PR D57 5903	D.W. Bliss <i>et al.</i>	(CLEO Collab.)
ABE	97O	PRL 78 4691	K. Abe <i>et al.</i>	(SLD Collab.)
ACKERSTAFF	97J	PL B404 213	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97L	ZPHY C74 403	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97R	ZPHY C75 593	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ALEXANDER	97F	PR D56 5320	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	97B	PRL 78 4686	R. Ammar <i>et al.</i>	(CLEO Collab.)
ANASTASSOV	97	PR D55 2559	A. Anastassov <i>et al.</i>	(CLEO Collab.)
Also	98B	PR D58 119903 (erratum)	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ANDERSON	97	PRL 79 3814	S. Anderson <i>et al.</i>	(CLEO Collab.)
VERY	97	PR D55 R1119	P. Avery <i>et al.</i>	(CLEO Collab.)
BARATE	97I	ZPHY C74 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	97R	PL B414 362	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERGFELD	97	PRL 79 2406	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BONVICINI	97	PRL 79 1221	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUSKULIC	97C	ZPHY C74 263	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
COAN	97	PR D55 7291	T.E. Coan <i>et al.</i>	(CLEO Collab.)
EDWARDS	97	PR D55 R3919	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
EDWARDS	97B	PR D56 R5297	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ESCRIBANO	97	PL B395 369	R. Escribano, E. Masso	(BARC, PARIT)
ABREU	96B	PL B365 448	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	96H	PL B377 313	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	96K	PL B389 187	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALAM	96	PRL 76 2637	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96E	PRPL 276 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	96D	PL B369 163	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	96E	PL B374 341	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	96S	PL B388 437	G. Alexander <i>et al.</i>	(OPAL Collab.)
BAI	96	PR D53 20	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALEST	96	PL B388 402	R. Balest <i>et al.</i>	(CLEO Collab.)
BARTELT	96	PRL 76 4119	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
BUSKULIC	96	ZPHY C70 579	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96C	ZPHY C70 561	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
COAN	96	PR D53 6037	T.E. Coan <i>et al.</i>	(CLEO Collab.)
ABE	95Y	PR D52 4828	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	95T	PL B357 715	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95U	PL B359 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	95	PL B345 93	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	95F	PL B352 487	M. Acciarri <i>et al.</i>	(L3 Collab.)
AKERS	95F	ZPHY C66 31	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95I	ZPHY C66 543	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95P	ZPHY C67 45	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95Y	ZPHY C68 555	R. Akers <i>et al.</i>	(OPAL Collab.)
ALBRECHT	95	PL B341 441	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	95C	PL B349 576	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	95G	ZPHY C68 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	95H	ZPHY C68 215	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	95C	PRL 75 3809	R. Balest <i>et al.</i>	(CLEO Collab.)
BERNABEU	95	NP B436 474	J. Bernabeu <i>et al.</i>	
BUSKULIC	95C	PL B346 371	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	95D	PL B346 379	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
Also	95P	PL B363 265 (erratum)	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	94K	PL B334 435	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	94E	PL B328 207	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	94G	PL B339 278	R. Akers <i>et al.</i>	(OPAL Collab.)
ALBRECHT	94E	PL B337 383	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ARTUSO	94	PRL 72 3762	M. Artuso <i>et al.</i>	(CLEO Collab.)
BARTELT	94	PRL 73 1890	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
BATTLE	94	PRL 73 1079	M. Battle <i>et al.</i>	(CLEO Collab.)
BAUER	94	PR D50 R13	D.A. Bauer <i>et al.</i>	(TPC/2gamma Collab.)

BUSKULIC	94D	PL B321 168	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	94E	PL B332 209	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	94F	PL B332 219	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBAUT	94B	PRL 73 934	D. Gibaut <i>et al.</i>	(CLEO Collab.)
ADRIANI	93M	PRPL 236 1	O. Adriani <i>et al.</i>	(L3 Collab.)
ALBRECHT	93C	ZPHY C58 61	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93G	PL B316 608	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	93	PR D47 R3671	R. Balest <i>et al.</i>	(CLEO Collab.)
BEAN	93	PRL 70 138	A. Bean <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	93	PRL 71 1791	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ESCRIBANO	93	PL B301 419	R. Escribano, E. Masso	(BARC)
PROCARIO	93	PRL 70 1207	M. Procaro <i>et al.</i>	(CLEO Collab.)
ABREU	92N	ZPHY C55 555	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	92F	PL B281 405	D.P. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	92H	PL B288 373	P.D. Acton <i>et al.</i>	(OPAL Collab.)
AKERIB	92	PRL 69 3610	D.S. Akerib <i>et al.</i>	(CLEO Collab.)
Also	93B	PRL 71 3395 (erratum)	D.S. Akerib <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92D	ZPHY C53 367	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92K	ZPHY C55 179	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92M	PL B292 221	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92Q	ZPHY C56 339	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMMAR	92	PR D45 3976	R. Ammar <i>et al.</i>	(CLEO Collab.)
ARTUSO	92	PRL 69 3278	M. Artuso <i>et al.</i>	(CLEO Collab.)
BAI	92	PRL 69 3021	J.Z. Bai <i>et al.</i>	(BES Collab.)
BATTLE	92	PL B291 488	M. Battle <i>et al.</i>	(CLEO Collab.)
BUSKULIC	92J	PL B297 459	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
DECAMP	92C	ZPHY C54 211	D. Decamp <i>et al.</i>	(ALEPH Collab.)
ADEVA	91F	PL B265 451	B. Adeva <i>et al.</i>	(L3 Collab.)
ALBRECHT	91D	PL B260 259	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	91D	PL B266 201	G. Alexander <i>et al.</i>	(OPAL Collab.)
ANTREASYAN	91	PL B259 216	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
GRIFOLS	91	PL B255 611	J.A. Grifols, A. Mendez	(BARC)
SAMUEL	91B	PRL 67 668	M.A. Samuel, G.W. Li, R. Mendel	(OKSU, WONT)
Also	92B	PRL 69 995	M.A. Samuel, G.W. Li, R. Mendel	(OKSU, WONT)
Erratum.				
ABACHI	90	PR D41 1414	S. Abachi <i>et al.</i>	(HRS Collab.)
ALBRECHT	90E	PL B246 278	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90I	PL B250 164	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BEHREND	90	ZPHY C46 537	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BOWCOCK	90	PR D41 805	T.J.V. Bowcock <i>et al.</i>	(CLEO Collab.)
DELAGUILA	90	PL B252 116	F. del Aguila, M. Sher	(BARC, WILL)
GOLDBERG	90	PL B251 223	M. Goldberg <i>et al.</i>	(CLEO Collab.)
WU	90	PR D41 2339	D.Y. Wu <i>et al.</i>	(Mark II Collab.)
ABACHI	89B	PR D40 902	S. Abachi <i>et al.</i>	(HRS Collab.)
BEHREND	89B	PL B222 163	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
JANSEN	89	PL B228 273	H. Janssen <i>et al.</i>	(Crystal Ball Collab.)
KLEINWORT	89	ZPHY C42 7	C. Kleinwort <i>et al.</i>	(JADE Collab.)
ADEVA	88	PR D38 2665	B. Adeva <i>et al.</i>	(Mark-J Collab.)
ALBRECHT	88B	PL B202 149	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88L	ZPHY C41 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88M	ZPHY C41 405	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMIDEI	88	PR D37 1750	D. Amidei <i>et al.</i>	(Mark II Collab.)
BEHREND	88	PL B200 226	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BRAUNSCH...	88C	ZPHY C39 331	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
KEH	88	PL B212 123	S. Keh <i>et al.</i>	(Crystal Ball Collab.)
TSCHIRHART	88	PL B205 407	R. Tschirhart <i>et al.</i>	(HRS Collab.)
ABACHI	87B	PL B197 291	S. Abachi <i>et al.</i>	(HRS Collab.)
ABACHI	87C	PRL 59 2519	S. Abachi <i>et al.</i>	(HRS Collab.)
ADLER	87B	PRL 59 1527	J. Adler <i>et al.</i>	(Mark III Collab.)
AIHARA	87B	PR D35 1553	H. Aihara <i>et al.</i>	(TPC Collab.)
AIHARA	87C	PRL 59 751	H. Aihara <i>et al.</i>	(TPC Collab.)
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87P	PL B199 580	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BAND	87	PL B198 297	H.R. Band <i>et al.</i>	(MAC Collab.)
BAND	87B	PRL 59 415	H.R. Band <i>et al.</i>	(MAC Collab.)
BARINGER	87	PRL 59 1993	P. Baringer <i>et al.</i>	(CLEO Collab.)
BEBEK	87C	PR D36 690	C. Bebek <i>et al.</i>	(CLEO Collab.)
BURCHAT	87	PR D35 27	P.R. Burchat <i>et al.</i>	(Mark II Collab.)
BYLSMA	87	PR D35 2269	B.G. Bylsma <i>et al.</i>	(HRS Collab.)
COFFMAN	87	PR D36 2185	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
DERRICK	87	PL B189 260	M. Derrick <i>et al.</i>	(HRS Collab.)

FORD	87	PR D35 408	W.T. Ford <i>et al.</i>	(MAC Collab.)
FORD	87B	PR D36 1971	W.T. Ford <i>et al.</i>	(MAC Collab.)
GAN	87	PRL 59 411	K.K. Gan <i>et al.</i>	(Mark II Collab.)
GAN	87B	PL B197 561	K.K. Gan <i>et al.</i>	(Mark II Collab.)
AIHARA	86E	PRL 57 1836	H. Aihara <i>et al.</i>	(TPC Collab.)
BARTEL	86D	PL B182 216	W. Bartel <i>et al.</i>	(JADE Collab.)
PDG	86	PL 170B	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
RÜCKSTUHL	86	PRL 56 2132	W. Ruckstuhl <i>et al.</i>	(DELCO Collab.)
SCHMIDKE	86	PRL 57 527	W.B. Schmidke <i>et al.</i>	(Mark II Collab.)
YELTON	86	PRL 56 812	J.M. Yelton <i>et al.</i>	(Mark II Collab.)
ALTHOFF	85	ZPHY C26 521	M. Althoff <i>et al.</i>	(TASSO Collab.)
ASH	85B	PRL 55 2118	W.W. Ash <i>et al.</i>	(MAC Collab.)
BALTRUSAITIS...	85	PRL 55 1842	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BARTEL	85F	PL 161B 188	W. Bartel <i>et al.</i>	(JADE Collab.)
BEHREND	85	PR D32 2468	S. Behrend <i>et al.</i>	(CLEO Collab.)
BELTRAMI	85	PRL 54 1775	I. Beltrami <i>et al.</i>	(HRS Collab.)
BERGER	85	ZPHY C28 1	C. Berger <i>et al.</i>	(PLUTO Collab.)
BURCHAT	85	PRL 54 2489	P.R. Burchat <i>et al.</i>	(Mark II Collab.)
FERNANDEZ	85	PRL 54 1624	E. Fernandez <i>et al.</i>	(MAC Collab.)
MILLS	85	PRL 54 624	G.B. Mills <i>et al.</i>	(DELCO Collab.)
AIHARA	84C	PR D30 2436	H. Aihara <i>et al.</i>	(TPC Collab.)
BEHREND	84	ZPHY C23 103	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
MILLS	84	PRL 52 1944	G.B. Mills <i>et al.</i>	(DELCO Collab.)
BEHREND	83C	PL 127B 270	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
SILVERMAN	83	PR D27 1196	D.J. Silverman, G.L. Shaw	(UCI)
BEHREND	82	PL 114B 282	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BLOCKER	82B	PRL 48 1586	C.A. Blocker <i>et al.</i>	(Mark II Collab.)
BLOCKER	82D	PL 109B 119	C.A. Blocker <i>et al.</i>	(Mark II Collab.) J
FELDMAN	82	PRL 48 66	G.A. Feldman <i>et al.</i>	(Mark II Collab.)
HAYES	82	PR D25 2869	K.G. Hayes <i>et al.</i>	(Mark II Collab.)
BERGER	81B	PL 99B 489	C. Berger <i>et al.</i>	(PLUTO Collab.)
DORFAN	81	PRL 46 215	J.M. Dorfan <i>et al.</i>	(Mark II Collab.)
BRANDELIK	80	PL 92B 199	R. Brandelik <i>et al.</i>	(TASSO Collab.)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also	81	SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34	1471.	
BACINO	79B	PRL 42 749	W.J. Bacino <i>et al.</i>	(DELCO Collab.)
KIRKBY	79	SLAC-PUB-2419	J. Kirkby	(SLAC) J
		Batavia Lepton Photon Conference.		
BACINO	78B	PRL 41 13	W.J. Bacino <i>et al.</i>	(DELCO Collab.) J
Also	78	Tokyo Conf. 249	J. Kirz	(STON)
Also	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
BRANDELIK	78	PL 73B 109	R. Brandelik <i>et al.</i>	(DASP Collab.) J
FELDMAN	78	Tokyo Conf. 777	G.J. Feldman	(SLAC) J
JAROS	78	PRL 40 1120	J. Jaros <i>et al.</i>	(SLAC, LBL, NWES, HAWA)
PERL	75	PRL 35 1489	M.L. Perl <i>et al.</i>	(LBL, SLAC)

OTHER RELATED PAPERS

RAHAL-CAL...	98	IJMP A13 695	G. Rahal-Callot	(ETH)
GENTILE	96	PRPL 274 287	S. Gentile, M. Pohl	(ROMAI, ETH)
WEINSTEIN	93	ARNPS 43 457	A.J. Weinstein, R. Stroynowski	(CIT, SMU)
PERL	92	RPP 55 653	M.L. Perl	(SLAC)
PICH	90	MPL A5 1995	A. Pich	(VALE)
BARISH	88	PRPL 157 1	B.C. Barish, R. Stroynowski	(CIT)
GAN	88	IJMP A3 531	K.K. Gan, M.L. Perl	(SLAC)
HAYES	88	PR D38 3351	K.G. Hayes, M.L. Perl	(SLAC)
PERL	80	ARNPS 30 299	M.L. Perl	(SLAC)