

# SEARCHES FOR MONOPOLES, SUPERSYMMETRY, TECHNICOLOR, COMPOSITENESS, EXTRA DIMENSIONS, etc.

## Magnetic Monopole Searches

Isolated supermassive monopole candidate events have not been confirmed. The most sensitive experiments obtain negative results.

Best cosmic-ray supermassive monopole flux limit:

$$\lt 1.0 \times 10^{-15} \text{ cm}^{-2}\text{sr}^{-1}\text{s}^{-1} \quad \text{for } 1.1 \times 10^{-4} < \beta < 0.1$$

## Supersymmetric Particle Searches

Limits are based on the Minimal Supersymmetric Standard Model.

Assumptions include: 1)  $\tilde{\chi}_1^0$  (or  $\tilde{\gamma}$ ) is lightest supersymmetric particle; 2)  $R$ -parity is conserved; 3) With the exception of  $\tilde{t}$  and  $\tilde{b}$ , all scalar quarks are assumed to be degenerate in mass and  $m_{\tilde{q}_R} = m_{\tilde{q}_L}$ . 4) Limits for sleptons refer to the  $\tilde{\ell}_R$  states.

See the Particle Listings for a Note giving details of supersymmetry.

$\tilde{\chi}_i^0$  — neutralinos (mixtures of  $\tilde{\gamma}$ ,  $\tilde{Z}^0$ , and  $\tilde{H}_i^0$ )

Mass  $m_{\tilde{\chi}_1^0} > 46 \text{ GeV}$ , CL = 95% [all  $\tan\beta$ , all  $\Delta m_0$ , all  $m_0$ ]

Mass  $m_{\tilde{\chi}_2^0} > 62.4 \text{ GeV}$ , CL = 95%

[ $1 < \tan\beta < 40$ , all  $m_0$ , all  $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$ ]

Mass  $m_{\tilde{\chi}_3^0} > 99.9 \text{ GeV}$ , CL = 95%

[ $1 < \tan\beta < 40$ , all  $m_0$ , all  $m_{\tilde{\chi}_3^0} - m_{\tilde{\chi}_1^0}$ ]

$\tilde{\chi}_i^\pm$  — charginos (mixtures of  $\tilde{W}^\pm$  and  $\tilde{H}_i^\pm$ )

Mass  $m_{\tilde{\chi}_1^\pm} > 94 \text{ GeV}$ , CL = 95%

[ $\tan\beta < 40$ ,  $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} > 3 \text{ GeV}$ , all  $m_0$ ]

$\tilde{e}$  — scalar electron (selectron)

$$\text{Mass } m > 73 \text{ GeV, CL} = 95\% \quad [\text{all } m_{\tilde{e}_R} - m_{\tilde{\chi}_1^0}]$$

$\tilde{\mu}$  — scalar muon (smuon)

$$\text{Mass } m > 94 \text{ GeV, CL} = 95\% \\ [1 \leq \tan\beta \leq 40, m_{\tilde{\mu}_R} - m_{\tilde{\chi}_1^0} > 10 \text{ GeV}]$$

$\tilde{\tau}$  — scalar tau (stau)

$$\text{Mass } m > 81.9 \text{ GeV, CL} = 95\% \\ [m_{\tilde{\tau}_R} - m_{\tilde{\chi}_1^0} > 15 \text{ GeV, all } \theta_\tau]$$

$\tilde{q}$  — scalar quark (squark)

These limits include the effects of cascade decays, evaluated assuming a fixed value of the parameters  $\mu$  and  $\tan\beta$ . The limits are weakly sensitive to these parameters over much of parameter space. Limits assume GUT relations between gaugino masses and the gauge coupling.

$$\text{Mass } m > 250 \text{ GeV, CL} = 95\% \quad [\tan\beta = 2, \mu < 0, A = 0]$$

$\tilde{b}$  — scalar bottom (sbottom)

$$\text{Mass } m > 89 \text{ GeV, CL} = 95\% \quad [m_{\tilde{b}_1} - m_{\tilde{\chi}_1^0} > 8 \text{ GeV, all } \theta_b]$$

$\tilde{t}$  — scalar top (stop)

$$\text{Mass } m > 95.7 \text{ GeV, CL} = 95\% \\ [\tilde{t} \rightarrow c\tilde{\chi}_1^0, \text{ all } \theta_t, m_{\tilde{t}} - m_{\tilde{\chi}_1^0} > 10 \text{ GeV}]$$

$\tilde{g}$  — gluino

The limits summarised here refer to the high-mass region ( $m_{\tilde{g}} \gtrsim 5 \text{ GeV}$ ), and include the effects of cascade decays, evaluated assuming a fixed value of the parameters  $\mu$  and  $\tan\beta$ . The limits are weakly sensitive to these parameters over much of parameter space. Limits assume GUT relations between gaugino masses and the gauge coupling,

$$\text{Mass } m > 195 \text{ GeV, CL} = 95\% \quad [\text{any } m_{\tilde{q}}]$$

$$\text{Mass } m > 300 \text{ GeV, CL} = 95\% \quad [m_{\tilde{q}} = m_{\tilde{g}}]$$

## Technicolor

Searches for a color-octet techni- $\rho$  constrain its mass to be greater than 260 to 480 GeV, depending on allowed decay channels. Similar bounds exist on the color-octet techni- $\omega$ .

## Quark and Lepton Compositeness, Searches for

### Scale Limits $\Lambda$ for Contact Interactions (the lowest dimensional interactions with four fermions)

If the Lagrangian has the form  $\pm \frac{g^2}{2\Lambda^2} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_L \gamma^\mu \psi_L$  (with  $g^2/4\pi$  set equal to 1), then we define  $\Lambda \equiv \Lambda_{LL}^\pm$ . For the full definitions and for other forms, see the Note in the Listings on Searches for Quark and Lepton Compositeness in the full *Review* and the original literature.

$\Lambda_{LL}^+(eeee)$	> 8.3 TeV, CL = 95%
$\Lambda_{LL}^-(eeee)$	> 10.3 TeV, CL = 95%
$\Lambda_{LL}^+(ee\mu\mu)$	> 8.5 TeV, CL = 95%
$\Lambda_{LL}^-(ee\mu\mu)$	> 6.3 TeV, CL = 95%
$\Lambda_{LL}^+(ee\tau\tau)$	> 5.4 TeV, CL = 95%
$\Lambda_{LL}^-(ee\tau\tau)$	> 6.5 TeV, CL = 95%
$\Lambda_{LL}^+(\ell\ell\ell\ell)$	> 9.0 TeV, CL = 95%
$\Lambda_{LL}^-(\ell\ell\ell\ell)$	> 7.8 TeV, CL = 95%
$\Lambda_{LL}^+(eeuu)$	> 23.3 TeV, CL = 95%
$\Lambda_{LL}^-(eeuu)$	> 12.5 TeV, CL = 95%
$\Lambda_{LL}^+(eedd)$	> 11.1 TeV, CL = 95%
$\Lambda_{LL}^-(eedd)$	> 26.4 TeV, CL = 95%
$\Lambda_{LL}^+(eccc)$	> 1.0 TeV, CL = 95%
$\Lambda_{LL}^-(eccc)$	> 2.1 TeV, CL = 95%
$\Lambda_{LL}^+(eebb)$	> 5.6 TeV, CL = 95%
$\Lambda_{LL}^-(eebb)$	> 4.9 TeV, CL = 95%
$\Lambda_{LL}^+(\mu\mu qq)$	> 2.9 TeV, CL = 95%
$\Lambda_{LL}^-(\mu\mu qq)$	> 4.2 TeV, CL = 95%
$\Lambda(\ell\nu\ell\nu)$	> 3.10 TeV, CL = 90%
$\Lambda(e\nu qq)$	> 2.81 TeV, CL = 95%
$\Lambda_{LL}^+(qqqq)$	> 2.7 TeV, CL = 95%
$\Lambda_{LL}^-(qqqq)$	> 2.4 TeV, CL = 95%
$\Lambda_{LL}^+(\nu\nu qq)$	> 5.0 TeV, CL = 95%
$\Lambda_{LL}^-(\nu\nu qq)$	> 5.4 TeV, CL = 95%

## Excited Leptons

The limits from  $\ell^{*+} \ell^{*-}$  do not depend on  $\lambda$  (where  $\lambda$  is the  $\ell \ell^*$  transition coupling). The  $\lambda$ -dependent limits assume chiral coupling.

$e^{*\pm}$  — excited electron

Mass  $m > 103.2$  GeV, CL = 95% (from  $e^* e^*$ )

Mass  $m > 255$  GeV, CL = 95% (from  $e e^*$ )

Mass  $m > 310$  GeV, CL = 95% (if  $\lambda_\gamma = 1$ )

$\mu^{*\pm}$  — excited muon

Mass  $m > 103.2$  GeV, CL = 95% (from  $\mu^* \mu^*$ )

Mass  $m > 190$  GeV, CL = 95% (from  $\mu \mu^*$ )

$\tau^{*\pm}$  — excited tau

Mass  $m > 103.2$  GeV, CL = 95% (from  $\tau^* \tau^*$ )

Mass  $m > 185$  GeV, CL = 95% (from  $\tau \tau^*$ )

$\nu^*$  — excited neutrino

Mass  $m > 102.6$  GeV, CL = 95% (from  $\nu^* \nu^*$ )

Mass  $m > 190$  GeV, CL = 95% (from  $\nu \nu^*$ )

$q^*$  — excited quark

Mass  $m > 45.6$  GeV, CL = 95% (from  $q^* q^*$ )

Mass  $m > 570$ , none 580–760 GeV, CL = 95% (from  $q^* X$ )

## Color Sextet and Octet Particles

Color Sextet Quarks ( $q_6$ )

Mass  $m > 84$  GeV, CL = 95% (Stable  $q_6$ )

Color Octet Charged Leptons ( $\ell_8$ )

Mass  $m > 86$  GeV, CL = 95% (Stable  $\ell_8$ )

Color Octet Neutrinos ( $\nu_8$ )

Mass  $m > 110$  GeV, CL = 90% ( $\nu_8 \rightarrow \nu g$ )

---

## Extra Dimensions

Please refer to the Extra Dimensions section of the full *Review* for a discussion of the model-dependence of these bounds, and further constraints.

### Constraints on the fundamental gravity scale

$$M_H > 1.1 \text{ TeV, CL} = 95\% \quad (\text{dim-8 operators; } p\bar{p} \rightarrow e^+e^-, \gamma\gamma)$$

$$M_D > 1.1 \text{ TeV, CL} = 95\% \quad (e^+e^- \rightarrow G\gamma; \text{2-flat dimensions})$$

$$M_D > 3\text{--}1000 \text{ TeV} \quad (\text{astrophys. and cosmology; 2-flat dimensions; limits depend on technique and assumptions})$$

### Constraints on the radius of the extra dimensions, for the case of two-flat dimensions of equal radii

$$r < 90\text{--}660 \text{ nm} \quad (\text{astrophysics; limits depend on technique and assumptions})$$

$$r < 0.22 \text{ mm, CL} = 95\% \quad (\text{direct tests of Newton's law; cited in Extra Dimensions review})$$

---