

$\psi(4040)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 $\psi(4040)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

4039± 1 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

4037± 2	¹ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4040± 1	² SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4040±10	BRANDELIK	78C DASP	$e^+ e^-$

¹ From a fit to Crystal Ball (OSTERHELD 86) data.

² From a fit to BES (BAI 02C) data.

 $\psi(4040)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

80±10 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

85±10	³ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
89± 6	⁴ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
52±10	BRANDELIK	78C DASP	$e^+ e^-$

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

 $\psi(4040)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $e^+ e^-$	$(1.07 \pm 0.16) \times 10^{-5}$	
Γ_2 $D^0 \bar{D}^0$	seen	
Γ_3 $D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen	
Γ_4 $D^*(2007)^0 \bar{D}^*(2007)^0$	seen	
Γ_5 $J/\psi(1S)$ hadrons		
Γ_6 $J/\psi \pi^+ \pi^-$	< 4	$\times 10^{-3}$ 90%
Γ_7 $J/\psi \pi^0 \pi^0$	< 2	$\times 10^{-3}$ 90%
Γ_8 $J/\psi \eta$	< 7	$\times 10^{-3}$ 90%
Γ_9 $J/\psi \pi^0$	< 2	$\times 10^{-3}$ 90%
Γ_{10} $J/\psi \pi^+ \pi^- \pi^0$	< 2	$\times 10^{-3}$ 90%
Γ_{11} $\chi_{c1} \gamma$	< 1.1	% 90%
Γ_{12} $\chi_{c2} \gamma$	< 1.7	% 90%
Γ_{13} $\chi_{c1} \pi^+ \pi^- \pi^0$	< 1.1	% 90%
Γ_{14} $\chi_{c2} \pi^+ \pi^- \pi^0$	< 3.2	% 90%
Γ_{15} $\phi \pi^+ \pi^-$	< 3	$\times 10^{-3}$ 90%
Γ_{16} $\mu^+ \mu^-$		

$\psi(4040)$ PARTIAL WIDTHS $\Gamma(e^+e^-)$ Γ_1 VALUE (keV) DOCUMENT ID TECN COMMENT**0.86±0.07 OUR ESTIMATE**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88±0.11 ⁵ SETH 05A RVUE $e^+e^- \rightarrow$ hadrons0.91±0.13 ⁶ SETH 05A RVUE $e^+e^- \rightarrow$ hadrons0.75±0.15 BRANDELIK 78C DASP e^+e^- ⁵ From a fit to Crystal Ball (OSTERHELD 86) data.⁶ From a fit to BES (BAI 02C) data. $\psi(4040)$ BRANCHING RATIOS $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_1/Γ VALUE (units 10⁻⁵) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1.0 FELDMAN 77 MRK1 e^+e^- $\Gamma(D^0\bar{D}^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$ Γ_2/Γ_3 VALUE DOCUMENT ID TECN COMMENT**0.05±0.03** ⁷ GOLDHABER 77 MRK1 e^+e^- $\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$ Γ_4/Γ_3 VALUE DOCUMENT ID TECN COMMENT**32.0±12.0** ⁷ GOLDHABER 77 MRK1 e^+e^- $\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ VALUE (units 10⁻³) CL% DOCUMENT ID TECN COMMENT<4 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons $\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ VALUE (units 10⁻³) CL% DOCUMENT ID TECN COMMENT<2 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons $\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$ Γ_8/Γ VALUE (units 10⁻³) CL% DOCUMENT ID TECN COMMENT<7 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons $\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ VALUE (units 10⁻³) CL% DOCUMENT ID TECN COMMENT<2 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons $\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ VALUE (units 10⁻³) CL% DOCUMENT ID TECN COMMENT<2 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<11	90	COAN	06	CLEO	3.97–4.06 $e^+e^- \rightarrow$ hadrons
$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<17	90	COAN	06	CLEO	3.97–4.06 $e^+e^- \rightarrow$ hadrons
$\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{13}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<11	90	COAN	06	CLEO	3.97–4.06 $e^+e^- \rightarrow$ hadrons
$\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{14}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<32	90	COAN	06	CLEO	3.97–4.06 $e^+e^- \rightarrow$ hadrons
$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{15}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<3	90	COAN	06	CLEO	3.97–4.06 $e^+e^- \rightarrow$ hadrons

⁷ Phase-space factor (p^3) explicitly removed.

$\psi(4040)$ REFERENCES

COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
Also		ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)

OTHER RELATED PAPERS

HEIKKILA	84	PR D29 110	K. Heikkila, N.A. Tornqvist, S. Ono	(HELS, AACHT)
ONO	84	ZPHY C26 307	S. Ono	(ORSAY)
SIEGRIST	82	PR D26 969	J.L. Siegrist <i>et al.</i>	(SLAC, LBL)
AUGUSTIN	75	PRL 34 764	J.E. Augustin <i>et al.</i>	(SLAC, LBL)
BACCI	75	PL 58B 481	C. Bacci <i>et al.</i>	(ROMA, FRAS)
BOYARSKI	75B	PRL 34 762	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)
ESPOSITO	75	PL 58B 478	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)