

Quark and Lepton Compositeness, Searches for

SEARCHES FOR QUARK AND LEPTON COMPOSITENESS

Revised 2001 by K. Hagiwara (KEK), and K. Hikasa and M. Tanabashi (Tohoku University).

If quarks and leptons are made of constituents, then at the scale of constituent binding energies, there should appear new interactions among quarks and leptons. At energies much below the compositeness scale (Λ), these interactions are suppressed by inverse powers of Λ . The dominant effect should come from the lowest dimensional interactions with four fermions (contact terms), whose most general chirally invariant form reads [1]

$$L = \frac{g^2}{2\Lambda^2} \left[\eta_{LL} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_L \gamma^\mu \psi_L + \eta_{RR} \bar{\psi}_R \gamma_\mu \psi_R \bar{\psi}_R \gamma^\mu \psi_R + 2\eta_{LR} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_R \gamma^\mu \psi_R \right]. \quad (1)$$

Chiral invariance provides a natural explanation why quark and lepton masses are much smaller than their inverse size Λ . We may determine the scale Λ unambiguously by using the above form of the effective interactions; the conventional method [1] is to fix its scale by setting $g^2/4\pi = g^2(\Lambda)/4\pi = 1$ for the new strong interaction coupling and by setting the largest magnitude of the coefficients $\eta_{\alpha\beta}$ to be unity. In the following, we denote

$$\begin{aligned} \Lambda &= \Lambda_{LL}^\pm \quad \text{for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, 0, 0), \\ \Lambda &= \Lambda_{RR}^\pm \quad \text{for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (0, \pm 1, 0), \\ \Lambda &= \Lambda_{VV}^\pm \quad \text{for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, \pm 1, \pm 1), \\ \Lambda &= \Lambda_{AA}^\pm \quad \text{for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, \pm 1, \mp 1), \end{aligned} \quad (2)$$

as typical examples. Such interactions can arise by constituent interchange (when the fermions have common constituents, *e.g.*, for $ee \rightarrow ee$) and/or by exchange of the binding quanta (whenever binding quanta couple to constituents of both particles).

Another typical consequence of compositeness is the appearance of excited leptons and quarks (ℓ^* and q^*). Phenomenologically, an excited lepton is defined to be a heavy lepton which shares leptonic quantum number with one of the existing leptons (an excited quark is defined similarly). For example, an excited electron e^* is characterized by a nonzero transition-magnetic coupling with electrons. Smallness of the lepton mass and the success of QED prediction for $g-2$ suggest chirality conservation, *i.e.*, an excited lepton should not couple to both left- and right-handed components of the corresponding lepton.

Excited leptons may be classified by $SU(2) \times U(1)$ quantum numbers. Typical examples are:

1. Sequential type

$$\begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_L, \quad [\nu_R^*], \quad \ell_R^*.$$

ν_R^* is necessary unless ν^* has a Majorana mass.

2. Mirror type

$$[\nu_L^*], \quad \ell_L^*, \quad \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_R.$$

3. Homodoublet type

$$\begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_L, \quad \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_R.$$

Similar classification can be made for excited quarks.

Excited fermions can be pair produced via their gauge couplings. The couplings of excited leptons with Z are listed

| | Sequential type | Mirror type | Homodoublet type |
|---------------|------------------------------------|------------------------------------|--------------------------|
| V^{ℓ^*} | $-\frac{1}{2} + 2 \sin^2 \theta_W$ | $-\frac{1}{2} + 2 \sin^2 \theta_W$ | $-1 + 2 \sin^2 \theta_W$ |
| A^{ℓ^*} | $-\frac{1}{2}$ | $+\frac{1}{2}$ | 0 |
| $V^{\nu_D^*}$ | $+\frac{1}{2}$ | $+\frac{1}{2}$ | +1 |
| $A^{\nu_D^*}$ | $+\frac{1}{2}$ | $-\frac{1}{2}$ | 0 |
| $V^{\nu_M^*}$ | 0 | 0 | — |
| $A^{\nu_M^*}$ | +1 | -1 | — |

in the following table (for notation see Eq. (1) in “Standard Model of Electroweak Interactions”):

Here ν_D^* (ν_M^*) stands for Dirac (Majorana) excited neutrino. The corresponding couplings of excited quarks can be easily obtained. Although form factor effects can be present for the gauge couplings at $q^2 \neq 0$, they are usually neglected.

In addition, transition magnetic type couplings with a gauge boson are expected. These couplings can be generally parameterized as follows:

$$\begin{aligned}
 \mathcal{L} = & \frac{\lambda_{\gamma}^{(f^*)} e}{2m_{f^*}} \bar{f}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) f F_{\mu\nu} \\
 & + \frac{\lambda_Z^{(f^*)} e}{2m_{f^*}} \bar{f}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) f Z_{\mu\nu} \\
 & + \frac{\lambda_W^{(\ell^*)} g}{2m_{\ell^*}} \bar{\ell}^* \sigma^{\mu\nu} \frac{1-\gamma_5}{2} \nu W_{\mu\nu} \\
 & + \frac{\lambda_W^{(\nu^*)} g}{2m_{\nu^*}} \bar{\nu}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) \ell W_{\mu\nu}^\dagger \\
 & + \text{h.c.} , \tag{3}
 \end{aligned}$$

where $g = e/\sin\theta_W$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the photon field strength, $Z_{\mu\nu} = \partial_\mu Z_\nu - \partial_\nu Z_\mu$, *etc.* The normalization of the coupling is chosen such that

$$\max(|\eta_L|, |\eta_R|) = 1 .$$

Chirality conservation requires

$$\eta_L \eta_R = 0 . \quad (4)$$

Some experimental analyses assume the relation $\eta_L = \eta_R = 1$, which violates chiral symmetry. We encode the results of such analyses if the crucial part of the cross section is proportional to the factor $\eta_L^2 + \eta_R^2$ and the limits can be reinterpreted as those for chirality conserving cases $(\eta_L, \eta_R) = (1, 0)$ or $(0, 1)$ after rescaling λ .

These couplings in Eq. (3) can arise from $SU(2) \times U(1)$ -invariant higher-dimensional interactions. A well-studied model is the interaction of homodoublet type ℓ^* with the Lagrangian [2,3]

$$\mathcal{L} = \frac{1}{2\Lambda} \bar{L}^* \sigma^{\mu\nu} (gf \frac{\tau^a}{2} W_{\mu\nu}^a + g' f' Y B_{\mu\nu}) \frac{1-\gamma_5}{2} L + \text{h.c.} , \quad (5)$$

where L denotes the lepton doublet (ν, ℓ) , Λ is the compositeness scale, g, g' are $SU(2)$ and $U(1)_Y$ gauge couplings, and $W_{\mu\nu}^a$ and $B_{\mu\nu}$ are the field strengths for $SU(2)$ and $U(1)_Y$ gauge fields. The same interaction occurs for mirror-type excited leptons. For sequential-type excited leptons, the ℓ^* and ν^* couplings become unrelated, and the couplings receive the extra suppression of $(250 \text{ GeV})/\Lambda$ or m_{L^*}/Λ . In any case, these couplings satisfy the relation

$$\lambda_W = -\sqrt{2} \sin^2 \theta_W (\lambda_Z \cot \theta_W + \lambda_\gamma) . \quad (6)$$

Additional coupling with gluons is possible for excited quarks:

$$\mathcal{L} = \frac{1}{2\Lambda} \bar{Q}^* \sigma^{\mu\nu} \left(g_s f_s \frac{\lambda^a}{2} G_{\mu\nu}^a + g f \frac{\tau^a}{2} W_{\mu\nu}^a + g' f' Y B_{\mu\nu} \right) \times \frac{1-\gamma_5}{2} Q + \text{h.c.}, \quad (7)$$

where Q denotes a quark doublet, g_s is the QCD gauge coupling, and $G_{\mu\nu}^a$ the gluon field strength.

It should be noted that the electromagnetic radiative decay of $\ell^*(\nu^*)$ is forbidden if $f = -f'$ ($f = f'$). These two possibilities ($f = f'$ and $f = -f'$) are investigated in many analyses of the LEP experiments above the Z pole.

Several different conventions are used by LEP experiments on Z pole to express the transition magnetic couplings. To facilitate comparison, we re-express these in terms of λ_Z and λ_γ using the following relations and taking $\sin^2\theta_W = 0.23$. We assume chiral couplings, *i.e.*, $|c| = |d|$ in the notation of Ref. 2.

1. ALEPH (charged lepton and neutrino)

$$\lambda_Z^{\text{ALEPH}} = \frac{1}{2} \lambda_Z \quad (1990 \text{ papers}) \quad (8a)$$

$$\frac{2c}{\Lambda} = \frac{\lambda_Z}{m_{\ell^*} [\text{or } m_{\nu^*}]} \quad (\text{for } |c| = |d|) \quad (8b)$$

2. ALEPH (quark)

$$\lambda_u^{\text{ALEPH}} = \frac{\sin\theta_W \cos\theta_W}{\sqrt{\frac{1}{4} - \frac{2}{3}\sin^2\theta_W + \frac{8}{9}\sin^4\theta_W}} \lambda_Z = 1.11\lambda_Z \quad (9)$$

3. L3 and DELPHI (charged lepton)

$$\lambda^{\text{L3}} = \lambda_Z^{\text{DELPHI}} = -\frac{\sqrt{2}}{\cot\theta_W - \tan\theta_W} \lambda_Z = -1.10\lambda_Z \quad (10)$$

4. L3 (neutrino)

$$f_Z^{L3} = \sqrt{2}\lambda_Z \quad (11)$$

5. OPAL (charged lepton)

$$\frac{f^{\text{OPAL}}}{\Lambda} = -\frac{2}{\cot\theta_W - \tan\theta_W} \frac{\lambda_Z}{m_{\ell^*}} = -1.56 \frac{\lambda_Z}{m_{\ell^*}} \quad (12)$$

6. OPAL (quark)

$$\frac{f^{\text{OPAL}}_c}{\Lambda} = \frac{\lambda_Z}{2m_{q^*}} \quad (\text{for } |c| = |d|) \quad (13)$$

7. DELPHI (charged lepton)

$$\lambda_\gamma^{\text{DELPHI}} = -\frac{1}{\sqrt{2}} \lambda_\gamma \quad (14)$$

If leptons are made of color triplet and antitriplet constituents, we may expect their color-octet partners. Transitions between the octet leptons (ℓ_8) and the ordinary lepton (ℓ) may take place via the dimension-five interactions

$$\mathcal{L} = \frac{1}{2\Lambda} \sum_{\ell} \left\{ \bar{\ell}_8^\alpha g_S F_{\mu\nu}^\alpha \sigma^{\mu\nu} (\eta_L \ell_L + \eta_R \ell_R) + h.c. \right\} \quad (15)$$

where the summation is over charged leptons and neutrinos. The leptonic chiral invariance implies $\eta_L \eta_R = 0$ as before.

References

1. E.J. Eichten, K.D. Lane, and M.E. Peskin, Phys. Rev. Lett. **50**, 811 (1983).
2. K. Hagiwara, S. Komamiya, and D. Zeppenfeld, Z. Phys. **C29**, 115 (1985).
3. N. Cabibbo, L. Maiani, and Y. Srivastava, Phys. Lett. **139B**, 459 (1984).

SCALE LIMITS for Contact Interactions: $\Lambda(\mathbf{e e e e})$

Limits are for Λ_{LL}^\pm only. For other cases, see each reference.

| $\Lambda_{LL}^+(\text{TeV})$ | $\Lambda_{LL}^-(\text{TeV})$ | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------------------------------|-----|------------------------|---------|---|
| >8.3 | >10.3 | 95 | ¹ BOURILKOV | 01 RVUE | $E_{\text{cm}} = 192\text{--}208 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|-------|----|--------------------------|----------|---|
| >4.7 | >6.1 | 95 | ² ABBIENDI | 04G | $E_{\text{cm}} = 130\text{--}207$ GeV |
| >3.8 | >5.6 | 95 | ABBIENDI | 00R OPAL | $E_{\text{cm}} = 189$ GeV |
| >4.4 | >5.4 | 95 | ABREU | 00S DLPH | $E_{\text{cm}} = 183\text{--}189$ GeV |
| >4.3 | >4.9 | 95 | ACCIARRI | 00P L3 | $E_{\text{cm}} = 130\text{--}189$ GeV |
| >3.5 | >3.2 | 95 | BARATE | 00I ALEP | $E_{\text{cm}} = 130\text{--}183$ GeV |
| >6.0 | >7.7 | 95 | ³ BOURILKOV | 00 RVUE | $E_{\text{cm}} = 183\text{--}189$ GeV |
| >3.1 | >3.8 | 95 | ABBIENDI | 99 OPAL | $E_{\text{cm}} = 130\text{--}136, 161\text{--}172, 183$ GeV |
| >2.2 | >2.8 | 95 | ABREU | 99A DLPH | $E_{\text{cm}} = 130\text{--}172$ GeV |
| >2.7 | >2.4 | 95 | ACCIARRI | 98J L3 | $E_{\text{cm}} = 130\text{--}172$ GeV |
| >3.0 | >2.5 | 95 | ACKERSTAFF | 98V OPAL | $E_{\text{cm}} = 130\text{--}172$ GeV |
| >2.4 | >2.2 | 95 | ACKERSTAFF | 97C OPAL | $E_{\text{cm}} = 130\text{--}136, 161$ GeV |
| >1.7 | >2.3 | 95 | ARIMA | 97 VNS | $E_{\text{cm}} = 57.77$ GeV |
| >1.6 | >2.0 | 95 | ⁴ BUSKULIC | 93Q ALEP | $E_{\text{cm}} = 88.25\text{--}94.25$ GeV |
| >1.6 | | 95 | ^{4,5} BUSKULIC | 93Q RVUE | |
| | >2.2 | 95 | BUSKULIC | 93Q RVUE | |
| | >3.6 | 95 | ⁶ KROHA | 92 RVUE | |
| >1.3 | | 95 | ⁶ KROHA | 92 RVUE | |
| >0.7 | >2.8 | 95 | BEHREND | 91C CELL | $E_{\text{cm}} = 35$ GeV |
| >1.3 | >1.3 | 95 | KIM | 89 AMY | $E_{\text{cm}} = 50\text{--}57$ GeV |
| >1.4 | >3.3 | 95 | ⁷ BRAUNSCH... | 88 TASS | $E_{\text{cm}} = 12\text{--}46.8$ GeV |
| >1.0 | >0.7 | 95 | ⁸ FERNANDEZ | 87B MAC | $E_{\text{cm}} = 29$ GeV |
| >1.1 | >1.4 | 95 | ⁹ BARTEL | 86C JADE | $E_{\text{cm}} = 12\text{--}46.8$ GeV |
| >1.17 | >0.87 | 95 | ¹⁰ DERRICK | 86 HRS | $E_{\text{cm}} = 29$ GeV |
| >1.1 | >0.76 | 95 | ¹¹ BERGER | 85B PLUT | $E_{\text{cm}} = 34.7$ GeV |

¹ A combined analysis of the data from ALEPH, DELPHI, L3, and OPAL.

² ABBIENDI 04G limits are from $e^+e^- \rightarrow e^+e^-$ cross section at $\sqrt{s} = 130\text{--}207$ GeV.

³ A combined analysis of the data from ALEPH, L3, and OPAL.

⁴ BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

⁵ This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

⁶ KROHA 92 limit is from fit to BERGER 85B, BARTEL 86C, DERRICK 86B, FERNANDEZ 87B, BRAUNSCHWEIG 88, BEHREND 91B, and BEHREND 91C. The fit gives $\eta/\Lambda_{LL}^2 = +0.230 \pm 0.206 \text{ TeV}^{-2}$.

⁷ BRAUNSCHWEIG 88 assumed $m_Z = 92$ GeV and $\sin^2\theta_W = 0.23$.

⁸ FERNANDEZ 87B assumed $\sin^2\theta_W = 0.22$.

⁹ BARTEL 86C assumed $m_Z = 93$ GeV and $\sin^2\theta_W = 0.217$.

¹⁰ DERRICK 86 assumed $m_Z = 93$ GeV and $g_V^2 = (-1/2 + 2\sin^2\theta_W)^2 = 0.004$.

¹¹ BERGER 85B assumed $m_Z = 93$ GeV and $\sin^2\theta_W = 0.217$.

SCALE LIMITS for Contact Interactions: $\Lambda(ee\mu\mu)$

Limits are for Λ_{LL}^\pm only. For other cases, see each reference.

| Λ_{LL}^+ (TeV) | Λ_{LL}^- (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|------------------------|-----|------------------------|--------|---------------------------------------|
| >8.1 | > 7.3 | 95 | ¹² ABBIENDI | 04G | $E_{\text{cm}} = 130\text{--}207$ GeV |
| > 8.5 | >3.8 | 95 | ACCIARRI | 00P L3 | $E_{\text{cm}} = 130\text{--}189$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|------|-------|----|---------------------------|-----|------|---|
| >7.3 | >4.6 | 95 | ABBIENDI | 00R | OPAL | $E_{\text{cm}} = 189$ GeV |
| >6.6 | >6.3 | 95 | ABREU | 00S | DLPH | $E_{\text{cm}} = 183\text{--}189$ GeV |
| >4.0 | >4.7 | 95 | BARATE | 00i | ALEP | $E_{\text{cm}} = 130\text{--}183$ GeV |
| >4.5 | >4.3 | 95 | ABBIENDI | 99 | OPAL | $E_{\text{cm}} = 130\text{--}136, 161\text{--}172, 183$ GeV |
| >3.4 | >2.7 | 95 | ABREU | 99A | DLPH | $E_{\text{cm}} = 130\text{--}172$ GeV |
| >3.6 | >2.4 | 95 | ACCIARRI | 98J | L3 | $E_{\text{cm}} = 130\text{--}172$ GeV |
| >2.9 | >3.4 | 95 | ACKERSTAFF | 98V | OPAL | $E_{\text{cm}} = 130\text{--}172$ GeV |
| >3.1 | >2.0 | 95 | MIURA | 98 | VNS | $E_{\text{cm}} = 57.77$ GeV |
| >2.4 | >2.9 | 95 | ACKERSTAFF | 97C | OPAL | $E_{\text{cm}} = 130\text{--}136, 161$ GeV |
| >1.7 | >2.2 | 95 | ¹³ VELISSARIS | 94 | AMY | $E_{\text{cm}} = 57.8$ GeV |
| >1.3 | >1.5 | 95 | ¹³ BUSKULIC | 93Q | ALEP | $E_{\text{cm}} = 88.25\text{--}94.25$ GeV |
| >2.6 | >1.9 | 95 | ^{13,14} BUSKULIC | 93Q | RVUE | |
| >2.3 | >2.0 | 95 | HOWELL | 92 | TOPZ | $E_{\text{cm}} = 52\text{--}61.4$ GeV |
| | >1.7 | 95 | ¹⁵ KROHA | 92 | RVUE | |
| >2.5 | >1.5 | 95 | BEHREND | 91C | CELL | $E_{\text{cm}} = 35\text{--}43$ GeV |
| >1.6 | >2.0 | 95 | ¹⁶ ABE | 90i | VNS | $E_{\text{cm}} = 50\text{--}60.8$ GeV |
| >1.9 | >1.0 | 95 | KIM | 89 | AMY | $E_{\text{cm}} = 50\text{--}57$ GeV |
| >2.3 | >1.3 | 95 | BRAUNSCH... | 88D | TASS | $E_{\text{cm}} = 30\text{--}46.8$ GeV |
| >4.4 | >2.1 | 95 | ¹⁷ BARTEL | 86C | JADE | $E_{\text{cm}} = 12\text{--}46.8$ GeV |
| >2.9 | >0.86 | 95 | ¹⁸ BERGER | 85 | PLUT | $E_{\text{cm}} = 34.7$ GeV |

¹² ABBIENDI 04G limits are from $e^+e^- \rightarrow \mu\mu$ cross section at $\sqrt{s} = 130\text{--}207$ GeV.

¹³ BUSKULIC 93Q and VELISSARIS 94 use the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

¹⁴ This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

¹⁵ KROHA 92 limit is from fit to BARTEL 86C, BEHREND 87C, BRAUNSCHWEIG 88D, BRAUNSCHWEIG 89C, ABE 90i, and BEHREND 91C. The fit gives $\eta/\Lambda_{LL}^2 = -0.155 \pm 0.095 \text{ TeV}^{-2}$.

¹⁶ ABE 90i assumed $m_Z = 91.163$ GeV and $\sin^2\theta_W = 0.231$.

¹⁷ BARTEL 86C assumed $m_Z = 93$ GeV and $\sin^2\theta_W = 0.217$.

¹⁸ BERGER 85 assumed $m_Z = 93$ GeV and $\sin^2\theta_W = 0.217$.

SCALE LIMITS for Contact Interactions: $\Lambda(ee\tau\tau)$

Limits are for Λ_{LL}^\pm only. For other cases, see each reference.

| Λ_{LL}^+ (TeV) | Λ_{LL}^- (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|-----|------------------------|--------|--|
| >4.9 | >7.2 | 95 | ¹⁹ ABBIENDI | 04G | $E_{\text{cm}} = 130\text{--}207$ GeV |
| >5.4 | >4.7 | 95 | ACCIARRI | 00P L3 | $E_{\text{cm}} = 130\text{--}189$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| >3.9 | >6.5 | 95 | ABBIENDI | 00R | OPAL $E_{\text{cm}} = 189$ GeV |
| >5.2 | >5.4 | 95 | ABREU | 00S | DLPH $E_{\text{cm}} = 183\text{--}189$ GeV |
| >3.9 | >3.7 | 95 | BARATE | 00i | ALEP $E_{\text{cm}} = 130\text{--}183$ GeV |
| >3.8 | >4.0 | 95 | ABBIENDI | 99 | OPAL $E_{\text{cm}} = 130\text{--}136, 161\text{--}172, 183$ GeV |
| >2.8 | >2.6 | 95 | ABREU | 99A | DLPH $E_{\text{cm}} = 130\text{--}172$ GeV |
| >2.4 | >2.8 | 95 | ACCIARRI | 98J | L3 $E_{\text{cm}} = 130\text{--}172$ GeV |
| >2.3 | >3.7 | 95 | ACKERSTAFF | 98V | OPAL $E_{\text{cm}} = 130\text{--}172$ GeV |

| | | | | | | |
|------|------|----|----------------|-----|------|--|
| >1.9 | >3.0 | 95 | ACKERSTAFF | 97C | OPAL | $E_{\text{cm}} = 130\text{--}136, 161$ GeV |
| >1.4 | >2.0 | 95 | 20 VELISSARIS | 94 | AMY | $E_{\text{cm}} = 57.8$ GeV |
| >1.0 | >1.5 | 95 | 20 BUSKULIC | 93Q | ALEP | $E_{\text{cm}} = 88.25\text{--}94.25$ GeV |
| >1.8 | >2.3 | 95 | 20,21 BUSKULIC | 93Q | RVUE | |
| >1.9 | >1.7 | 95 | HOWELL | 92 | TOPZ | $E_{\text{cm}} = 52\text{--}61.4$ GeV |
| >1.9 | >2.9 | 95 | 22 KROHA | 92 | RVUE | |
| >1.6 | >2.3 | 95 | BEHREND | 91C | CELL | $E_{\text{cm}} = 35\text{--}43$ GeV |
| >1.8 | >1.3 | 95 | 23 ABE | 90I | VNS | $E_{\text{cm}} = 50\text{--}60.8$ GeV |
| >2.2 | >3.2 | 95 | 24 BARTEL | 86 | JADE | $E_{\text{cm}} = 12\text{--}46.8$ GeV |

¹⁹ ABBIENDI 04G limits are from $e^+e^- \rightarrow \tau\tau$ cross section at $\sqrt{s} = 130\text{--}207$ GeV.

²⁰ BUSKULIC 93Q and VELISSARIS 94 use the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

²¹ This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

²² KROHA 92 limit is from fit to BARTEL 86C BEHREND 89B, BRAUNSCHWEIG 89C, ABE 90I, and BEHREND 91C. The fit gives $\eta/\Lambda_{LL}^2 = +0.095 \pm 0.120 \text{ TeV}^{-2}$.

²³ ABE 90I assumed $m_Z = 91.163$ GeV and $\sin^2\theta_W = 0.231$.

²⁴ BARTEL 86 assumed $m_Z = 93$ GeV and $\sin^2\theta_W = 0.217$.

SCALE LIMITS for Contact Interactions: $\Lambda(\ell\ell\ell)$

Lepton universality assumed. Limits are for Λ_{LL}^\pm only. For other cases, see each reference.

| Λ_{LL}^+ (TeV) | Λ_{LL}^- (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|-----|----------------|--------|--|
| >7.7 | >9.5 | 95 | 25 ABBIENDI | 04G | $E_{\text{cm}} = 130\text{--}207$ GeV |
| >9.0 | >5.2 | 95 | ACCIARRI | 00P L3 | $E_{\text{cm}} = 130\text{--}189$ GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| | | | 26 BABICH | 03 | RVUE |
| >6.4 | >7.2 | 95 | ABBIENDI | 00R | OPAL $E_{\text{cm}} = 189$ GeV |
| >7.3 | >7.8 | 95 | ABREU | 00S | DLPH $E_{\text{cm}} = 183\text{--}189$ GeV |
| >5.3 | >5.5 | 95 | BARATE | 00I | ALEP $E_{\text{cm}} = 130\text{--}183$ GeV |
| >5.2 | >5.3 | 95 | ABBIENDI | 99 | OPAL $E_{\text{cm}} = 130\text{--}136, 161\text{--}172, 183$ GeV |
| >4.4 | >4.2 | 95 | ABREU | 99A | DLPH $E_{\text{cm}} = 130\text{--}172$ GeV |
| >4.0 | >3.1 | 95 | 27 ACCIARRI | 98J | L3 $E_{\text{cm}} = 130\text{--}172$ GeV |
| >3.4 | >4.4 | 95 | ACKERSTAFF | 98V | OPAL $E_{\text{cm}} = 130\text{--}172$ GeV |
| >2.7 | >3.8 | 95 | ACKERSTAFF | 97C | OPAL $E_{\text{cm}} = 130\text{--}136, 161$ GeV |
| >3.0 | >2.3 | 95 | 27,28 BUSKULIC | 93Q | ALEP $E_{\text{cm}} = 88.25\text{--}94.25$ GeV |
| >3.5 | >2.8 | 95 | 28,29 BUSKULIC | 93Q | RVUE |
| >2.5 | >2.2 | 95 | 30 HOWELL | 92 | TOPZ $E_{\text{cm}} = 52\text{--}61.4$ GeV |
| >3.4 | >2.7 | 95 | 31 KROHA | 92 | RVUE |

²⁵ ABBIENDI 04G limits are from $e^+e^- \rightarrow \ell^+\ell^-$ cross section at $\sqrt{s} = 130\text{--}207$ GeV.

²⁶ BABICH 03 obtain a bound $-0.175 \text{ TeV}^{-2} < 1/\Lambda_{LL}^2 < 0.095 \text{ TeV}^{-2}$ (95%CL) in a model independent analysis allowing all of $\Lambda_{LL}, \Lambda_{LR}, \Lambda_{RL}, \Lambda_{RR}$ to coexist.

²⁷ From $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-,$ and $\tau^+\tau^-$.

²⁸ BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

²⁹This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

³⁰HOWELL 92 limit is from $e^+e^- \rightarrow \mu^+\mu^-$ and $\tau^+\tau^-$.

³¹KROHA 92 limit is from fit to most PEP/PETRA/TRISTAN data. The fit gives $\eta/\Lambda_{LL}^2 = -0.0200 \pm 0.0666 \text{ TeV}^{-2}$.

SCALE LIMITS for Contact Interactions: $\Lambda(eeqq)$

Limits are for Λ_{LL}^\pm only. For other cases, see each reference.

| Λ_{LL}^+ (TeV) | Λ_{LL}^- (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|-----|---------------|----------|---|
| >23.3 | >12.5 | 95 | 32 CHEUNG | 01B RVUE | (<i>eeuu</i>) |
| >11.1 | >26.4 | 95 | 32 CHEUNG | 01B RVUE | (<i>eedd</i>) |
| > 5.6 | >4.9 | 95 | 33 BARATE | 00I ALEP | (<i>eebb</i>) |
| > 1.0 | >2.1 | 95 | 34 ABREU | 99A DLPH | (<i>eecc</i>) |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| > 8.2 | >3.7 | 95 | 35 ABBIENDI | 04G | (<i>eeqq</i>) |
| > 5.9 | >9.1 | 95 | 35 ABBIENDI | 04G | (<i>eeuu</i>) |
| > 8.6 | >5.5 | 95 | 35 ABBIENDI | 04G | (<i>eedd</i>) |
| > 2.7 | >1.7 | 95 | CHEKANOV | 04B ZEUS | (<i>eeqq</i>) |
| > 2.8 | >1.6 | 95 | 36 ADLOFF | 03 H1 | (<i>eeqq</i>) |
| > 2.7 | >2.7 | 95 | 37 ACHARD | 02J L3 | (<i>eetc</i>) |
| > 5.5 | >3.1 | 95 | 38 ABBIENDI | 00R OPAL | (<i>eeqq</i>) |
| > 4.9 | >6.1 | 95 | 38 ABBIENDI | 00R OPAL | (<i>eeuu</i>) |
| > 5.7 | >4.5 | 95 | 38 ABBIENDI | 00R OPAL | (<i>eedd</i>) |
| > 4.2 | >2.8 | 95 | 39 ACCIARRI | 00P L3 | (<i>eeqq</i>) |
| > 2.4 | >1.3 | 95 | 40 ADLOFF | 00 H1 | (<i>eeqq</i>) |
| > 5.4 | >6.2 | 95 | 41 BARATE | 00I ALEP | (<i>eeqq</i>) |
| | | | 42 BREITWEG | 00B ZEUS | |
| > 4.4 | >2.8 | 95 | 43 ABBIENDI | 99 OPAL | (<i>eeqq</i>) |
| > 4.0 | >4.8 | 95 | 44 ABBIENDI | 99 OPAL | (<i>eebb</i>) |
| > 3.3 | >4.2 | 95 | 45 ABBOTT | 99D D0 | (<i>eeqq</i>) |
| > 2.4 | >2.8 | 95 | 34 ABREU | 99A DLPH | (<i>eeqq</i>) (<i>d</i> or <i>s</i> quark) |
| > 4.4 | >3.9 | 95 | 34 ABREU | 99A DLPH | (<i>eebb</i>) |
| > 1.0 | >2.4 | 95 | 34 ABREU | 99A DLPH | (<i>eeuu</i>) |
| > 4.0 | >3.4 | 95 | 46 ZARNECKI | 99 RVUE | (<i>eedd</i>) |
| > 4.3 | >5.6 | 95 | 46 ZARNECKI | 99 RVUE | (<i>eeuu</i>) |
| > 3.0 | >2.1 | 95 | 47 ACCIARRI | 98J L3 | (<i>eeqq</i>) |
| > 3.4 | >2.2 | 95 | 48 ACKERSTAFF | 98V OPAL | (<i>eeqq</i>) |
| > 4.0 | >2.8 | 95 | 49 ACKERSTAFF | 98V OPAL | (<i>eebb</i>) |
| > 9.3 | >12.0 | 95 | 50 BARGER | 98E RVUE | (<i>eeuu</i>) |
| > 8.8 | >11.9 | 95 | 50 BARGER | 98E RVUE | (<i>eedd</i>) |
| > 2.5 | >3.7 | 95 | 51 ABE | 97T CDF | (<i>eeqq</i>) (isosinglet) |
| > 2.5 | >2.1 | 95 | 52 ACKERSTAFF | 97C OPAL | (<i>eeqq</i>) |
| > 3.1 | >2.9 | 95 | 53 ACKERSTAFF | 97C OPAL | (<i>eebb</i>) |
| > 7.4 | >11.7 | 95 | 54 DEANDREA | 97 RVUE | <i>eeuu</i> , atomic parity violation |
| > 2.3 | >1.0 | 95 | 55 AID | 95 H1 | (<i>eeqq</i>) (<i>u, d</i> quarks) |
| 1.7 | >2.2 | 95 | 56 ABE | 91D CDF | (<i>eeqq</i>) (<i>u, d</i> quarks) |
| > 1.2 | | 95 | 57 ADACHI | 91 TOPZ | (<i>eeqq</i>) (flavor-universal) |

| | | | | | |
|--------|-------|----|-------------|----------|---------------------------------------|
| | >1.6 | 95 | 57 ADACHI | 91 TOPZ | (<i>eeqq</i>) (flavor-universal) |
| > 0.6 | >1.7 | 95 | 58 BEHREND | 91C CELL | (<i>eecc</i>) |
| > 1.1 | >1.0 | 95 | 58 BEHREND | 91C CELL | (<i>eebb</i>) |
| > 0.9 | | 95 | 59 ABE | 89L VNS | (<i>eeqq</i>) (flavor-universal) |
| | >1.7 | 95 | 59 ABE | 89L VNS | (<i>eeqq</i>) (flavor-universal) |
| > 1.05 | >1.61 | 95 | 60 HAGIWARA | 89 RVUE | (<i>eecc</i>) |
| > 1.21 | >0.53 | 95 | 61 HAGIWARA | 89 RVUE | (<i>eebb</i>) |

³² CHEUNG 01B is an update of BARGER 98E.

³³ BARATE 00I limits are from R_b and jet-charge asymmetry at 130–183 GeV.

³⁴ ABREU 99A limits are from flavor-tagged $e^+e^- \rightarrow q\bar{q}$ cross section at 130–172 GeV.

³⁵ ABBIENDI 04G limits are from $e^+e^- \rightarrow q\bar{q}$ cross section at $\sqrt{s} = 130$ –207 GeV.

³⁶ ADLOFF 03 limits are from the $d\sigma/dQ^2$ measurement of $e^\pm p \rightarrow e^\pm X$.

³⁷ ACHARD 02J limit is from the bound on the $e^+e^- \rightarrow t\bar{c}$ cross section. $\Lambda_{LL} = \Lambda_{LR} = \Lambda_{RL} = \Lambda_{RR}$ and $m_t = 175$ GeV are assumed.

³⁸ ABBIENDI 00R limits are from $e^+e^- \rightarrow q\bar{q}$ cross section at $\sqrt{s} = 130$ –189 GeV.

³⁹ ACCIARRI 00P limit is from $e^+e^- \rightarrow qq$ cross section at $\sqrt{s} = 130$ –189 GeV.

⁴⁰ ADLOFF 00 limits are from the Q^2 spectrum measurement of $e^+p \rightarrow e^+X$.

⁴¹ BARATE 00I limits are from $e^+e^- \rightarrow q\bar{q}$ cross section and jet-charge asymmetry at 130–183 GeV.

⁴² BREITWEG 00B limits are from Q^2 spectrum measurement of e^+p collisions. See their Table 3 for the limits of various models.

⁴³ ABBIENDI 99 limits are from $e^+e^- \rightarrow q\bar{q}$ cross section at 130–136, 161–172, 183 GeV.

⁴⁴ ABBIENDI 99 limits are from R_b at 130–136, 161–172, 183 GeV.

⁴⁵ ABBOTT 99D limits are from e^+e^- mass distribution in $p\bar{p} \rightarrow e^+e^-X$ at $E_{\text{cm}} = 1.8$ TeV.

⁴⁶ ZARNECKI 99 use data from HERA, LEP, Tevatron, and various low-energy experiments.

⁴⁷ ACCIARRI 98J limits are from $e^+e^- \rightarrow q\bar{q}$ cross section at $E_{\text{cm}} = 130$ –172 GeV.

⁴⁸ ACKERSTAFF 98V limits are from $e^+e^- \rightarrow q\bar{q}$ at $E_{\text{cm}} = 130$ –172 GeV.

⁴⁹ ACKERSTAFF 98V limits are from R_b measurements at $E_{\text{cm}} = 130$ –172 GeV.

⁵⁰ BARGER 98E use data from HERA, LEP, Tevatron, and various low-energy experiments.

⁵¹ ABE 97T limits are from e^+e^- mass distribution in $p\bar{p} \rightarrow e^+e^-X$ at $E_{\text{cm}} = 1.8$ TeV.

⁵² ACKERSTAFF 97C limits are from $e^+e^- \rightarrow q\bar{q}$ cross section at $E_{\text{cm}} = 130$ –136 GeV and 161 GeV.

⁵³ ACKERSTAFF 97C limits are R_b measurements at $E_{\text{cm}} = 133$ GeV and 161 GeV.

⁵⁴ DEANDREA 97 limit is from atomic parity violation of cesium. The limit is excluded if the contact interactions are parity conserving.

⁵⁵ AID 95 limits are from the Q^2 spectrum measurement of $ep \rightarrow eX$.

⁵⁶ ABE 91D limits are from e^+e^- mass distribution in $p\bar{p} \rightarrow e^+e^-X$ at $E_{\text{cm}} = 1.8$ TeV.

⁵⁷ ADACHI 91 limits are from differential jet cross section. Universality of $\Lambda(eeqq)$ for five flavors is assumed.

⁵⁸ BEHREND 91C is from data at $E_{\text{cm}} = 35$ –43 GeV.

⁵⁹ ABE 89L limits are from jet charge asymmetry. Universality of $\Lambda(eeqq)$ for five flavors is assumed.

⁶⁰ The HAGIWARA 89 limit is derived from forward-backward asymmetry measurements of D/D^* mesons by ALTHOFF 83C, BARTEL 84E, and BARINGER 88.

⁶¹ The HAGIWARA 89 limit is derived from forward-backward asymmetry measurement of b hadrons by BARTEL 84D.

SCALE LIMITS for Contact Interactions: $\Lambda(\mu\mu qq)$

| Λ_{LL}^+ (TeV) | Λ_{LL}^- (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|------------------------|-----|-------------------|---------|----------------------------|
| > 2.9 | > 4.2 | 95 | ⁶² ABE | 97T CDF | $(\mu\mu qq)$ (isosinglet) |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| >1.4 | >1.6 | 95 | ABE | 92B CDF | $(\mu\mu qq)$ (isosinglet) |
| ⁶² ABE 97T limits are from $\mu^+ \mu^-$ mass distribution in $\bar{p}p \rightarrow \mu^+ \mu^- X$ at $E_{cm}=1.8$ TeV. | | | | | |

SCALE LIMITS for Contact Interactions: $\Lambda(\ell\nu\ell\nu)$

| VALUE (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------------|---------|---|
| >3.10 | 90 | ⁶³ JODIDIO | 86 SPEC | $\Lambda_{LR}^\pm(\nu_\mu \nu_e \mu e)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| >3.8 | | ⁶⁴ DIAZCRUZ | 94 RVUE | $\Lambda_{LL}^+(\tau \nu_\tau e \nu_e)$ |
| >8.1 | | ⁶⁴ DIAZCRUZ | 94 RVUE | $\Lambda_{LL}^-(\tau \nu_\tau e \nu_e)$ |
| >4.1 | | ⁶⁵ DIAZCRUZ | 94 RVUE | $\Lambda_{LL}^+(\tau \nu_\tau \mu \nu_\mu)$ |
| >6.5 | | ⁶⁵ DIAZCRUZ | 94 RVUE | $\Lambda_{LL}^-(\tau \nu_\tau \mu \nu_\mu)$ |
| ⁶³ JODIDIO 86 limit is from $\mu^+ \rightarrow \bar{\nu}_\mu e^+ \nu_e$. Chirality invariant interactions $L = (g^2/\Lambda^2) [\eta_{LL} (\bar{\nu}_\mu L \gamma^\alpha \mu_L) (\bar{e} L \gamma_\alpha \nu_e L) + \eta_{LR} (\bar{\nu}_\mu L \gamma^\alpha \nu_e L) (\bar{e} R \gamma_\alpha \mu_R)]$ with $g^2/4\pi = 1$ and $(\eta_{LL}, \eta_{LR}) = (0, \pm 1)$ are taken. No limits are given for Λ_{LL}^\pm with $(\eta_{LL}, \eta_{LR}) = (\pm 1, 0)$. For more general constraints with right-handed neutrinos and chirality nonconserving contact interactions, see their text. | | | | |
| ⁶⁴ DIAZCRUZ 94 limits are from $\Gamma(\tau \rightarrow e \nu \nu)$ and assume flavor-dependent contact interactions with $\Lambda(\tau \nu_\tau e \nu_e) \ll \Lambda(\mu \nu_\mu e \nu_e)$. | | | | |
| ⁶⁵ DIAZCRUZ 94 limits are from $\Gamma(\tau \rightarrow \mu \nu \nu)$ and assume flavor-dependent contact interactions with $\Lambda(\tau \nu_\tau \mu \nu_\mu) \ll \Lambda(\mu \nu_\mu e \nu_e)$. | | | | |

SCALE LIMITS for Contact Interactions: $\Lambda(e\nu qq)$

| VALUE (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|------------------------|---------|---------|
| >2.81 | 95 | ⁶⁶ AFFOLDER | 01i CDF | |
| ⁶⁶ AFFOLDER 00i bound is for a scalar interaction $\bar{q}_R q_L \bar{\nu} e_L$. | | | | |

SCALE LIMITS for Contact Interactions: $\Lambda(qqqq)$

Limits are for Λ_{LL}^\pm with color-singlet isoscalar exchanges among u_L 's and d_L 's only, unless otherwise noted. See EICHTEN 84 for details.

| VALUE (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------|-----|----------------------|--------|---|
| >2.7 | 95 | ⁶⁷ ABBOTT | 99C D0 | $p\bar{p} \rightarrow$ dijet mass. Λ_{LL}^+ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------|----|----|---------|---------|--|
| >2.0 | 95 | 68 | ABBOTT | 00E D0 | H_T distribution; Λ_{LL}^+ |
| >2.1 | 95 | 69 | ABBOTT | 98G D0 | $p\bar{p} \rightarrow$ dijet angl. Λ_{LL}^+ |
| | | 70 | BERTRAM | 98 RVUE | $p\bar{p} \rightarrow$ dijet mass |
| | | 71 | ABE | 96 CDF | $p\bar{p} \rightarrow$ jets inclusive |
| >1.6 | 95 | 72 | ABE | 96S CDF | $p\bar{p} \rightarrow$ dijet angl.; Λ_{LL}^+ |
| >1.3 | 95 | 73 | ABE | 93G CDF | $p\bar{p} \rightarrow$ dijet mass |
| >1.4 | 95 | 74 | ABE | 92D CDF | $p\bar{p} \rightarrow$ jets inclusive |
| >1.0 | 99 | 75 | ABE | 92M CDF | $p\bar{p} \rightarrow$ dijet angl. |
| >0.825 | 95 | 76 | ALITTI | 91B UA2 | $p\bar{p} \rightarrow$ jets inclusive |
| >0.700 | 95 | 74 | ABE | 89 CDF | $p\bar{p} \rightarrow$ jets inclusive |
| >0.330 | 95 | 77 | ABE | 89H CDF | $p\bar{p} \rightarrow$ dijet angl. |
| >0.400 | 95 | 78 | ARNISON | 86C UA1 | $p\bar{p} \rightarrow$ jets inclusive |
| >0.415 | 95 | 79 | ARNISON | 86D UA1 | $p\bar{p} \rightarrow$ dijet angl. |
| >0.370 | 95 | 80 | APPEL | 85 UA2 | $p\bar{p} \rightarrow$ jets inclusive |
| >0.275 | 95 | 81 | BAGNAIA | 84C UA2 | Repl. by APPEL 85 |

⁶⁷ The quoted limit is from inclusive dijet mass spectrum in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV.

ABBOTT 99C also obtain $\Lambda_{LL}^- > 2.4$ TeV. All quarks are assumed composite.

⁶⁸ The quoted limit for ABBOTT 00E is from H_T distribution in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. CTEQ4M PDF and $\mu=E_T^{\text{max}}$ are assumed. For limits with different assumptions, see their Tables 2 and 3. All quarks are assumed composite.

⁶⁹ ABBOTT 98G limit is from dijet angular distribution in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. All quarks are assumed composite.

⁷⁰ BERTRAM 98 obtain limit on the scale of color-octet axial-vector flavor-universal contact interactions: $\Lambda_{A8} > 2.1$ TeV. They also obtain a limit $\Lambda_{V8} > 2.4$ TeV on a color-octet flavor-universal vectorial contact interaction.

⁷¹ ABE 96 finds that the inclusive jet cross section for $E_T > 200$ GeV is significantly higher than the $\mathcal{O}(\alpha_s^3)$ perturbative QCD prediction. This could be interpreted as the effect of a contact interaction with $\Lambda_{LL} \sim 1.6$ TeV. However, ABE 96 state that uncertainty in the parton distribution functions, higher-order QCD corrections, and the detector calibration may possibly account for the effect.

⁷² ABE 96S limit is from dijet angular distribution in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. The limit for Λ_{LL}^- is > 1.4 TeV. ABE 96S also obtain limits for flavor symmetric contact interactions among all quark flavors: $\Lambda_{LL}^+ > 1.8$ TeV and $\Lambda_{LL}^- > 1.6$ TeV.

⁷³ ABE 93G limit is from dijet mass distribution in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. The limit is the weakest from several choices of structure functions and renormalization scale.

⁷⁴ Limit is from inclusive jet cross-section data in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. The limit takes into account uncertainties in choice of structure functions and in choice of process scale.

⁷⁵ ABE 92M limit is from dijet angular distribution for $m_{\text{dijet}} > 550$ GeV in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV.

⁷⁶ ALITTI 91B limit is from inclusive jet cross section in $p\bar{p}$ collisions at $E_{\text{cm}}=630$ GeV. The limit takes into account uncertainties in choice of structure functions and in choice of process scale.

⁷⁷ ABE 89H limit is from dijet angular distribution for $m_{\text{dijet}} > 200$ GeV at the Fermilab Tevatron Collider with $E_{\text{cm}}=1.8$ TeV. The QCD prediction is quite insensitive to choice of structure functions and choice of process scale.

- ⁷⁸ ARNISON 86C limit is from the study of inclusive high- p_T jet distributions at the CERN $\bar{p}p$ collider ($E_{\text{cm}} = 546$ and 630 GeV). The QCD prediction renormalized to the low- p_T region gives a good fit to the data.
- ⁷⁹ ARNISON 86D limit is from the study of dijet angular distribution in the range $240 < m(\text{dijet}) < 300$ GeV at the CERN $\bar{p}p$ collider ($E_{\text{cm}} = 630$ GeV). QCD prediction using EHLQ structure function (EICHTEN 84) with $\Lambda_{\text{QCD}} = 0.2$ GeV for the choice of $Q^2 = p_T^2$ gives the best fit to the data.
- ⁸⁰ APPEL 85 limit is from the study of inclusive high- p_T jet distributions at the CERN $\bar{p}p$ collider ($E_{\text{cm}} = 630$ GeV). The QCD prediction renormalized to the low- p_T region gives a good description of the data.
- ⁸¹ BAGNAIA 84C limit is from the study of jet p_T and dijet mass distributions at the CERN $\bar{p}p$ collider ($E_{\text{cm}} = 540$ GeV). The limit suffers from the uncertainties in comparing the data with the QCD prediction.

SCALE LIMITS for Contact Interactions: $\Lambda(\nu\nu qq)$

Limits are for Λ_{LL}^{\pm} only. For other cases, see each reference.

| Λ_{LL}^+ (TeV) | Λ_{LL}^- (TeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|------------------------|-----|----------------------------|------|--------------------|
| >5.0 | >5.4 | 95 | ⁸² MCFARLAND 98 | CCFR | νN scattering |

⁸² MCFARLAND 98 assumed a flavor universal interaction. Neutrinos were mostly of muon type.

MASS LIMITS for Excited e (e^*)

Most e^+e^- experiments assume one-photon or Z exchange. The limits from some e^+e^- experiments which depend on λ have assumed transition couplings which are chirality violating ($\eta_L = \eta_R$). However they can be interpreted as limits for chirality-conserving interactions after multiplying the coupling value λ by $\sqrt{2}$; see Note.

Excited leptons have the same quantum numbers as other ortholeptons. See also the searches for ortholeptons in the "Searches for Heavy Leptons" section.

Limits for Excited e (e^*) from Pair Production

These limits are obtained from $e^+e^- \rightarrow e^{*+}e^{*-}$ and thus rely only on the (electroweak) charge of e^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the e^* coupling is assumed to be of sequential type. Possible t channel contribution from transition magnetic coupling is neglected. All limits assume a dominant $e^* \rightarrow e\gamma$ decay except the limits from $\Gamma(Z)$.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|---------------------------------|------|--|
| >103.2 | 95 | ⁸³ ABBIENDI 02G OPAL | | $e^+e^- \rightarrow e^*e^*$ Homodoublet type |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|--------|----|-------|--------------|----------|---------------------------------------|------------------|
| >102.8 | 95 | 84 | ACHARD | 03B L3 | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| >100.0 | 95 | 85 | ACCIARRI | 01D L3 | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| > 91.3 | 95 | 86 | ABBIENDI | 00I OPAL | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| > 94.2 | 95 | 87 | ACCIARRI | 00E L3 | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| > 90.7 | 95 | 88 | ABREU | 990 DLPH | | Homodoublet type |
| > 85.0 | 95 | 89 | ACKERSTAFF | 98C OPAL | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| | | 90 | BARATE | 98U ALEP | $Z \rightarrow e^* e^*$ | |
| > 79.6 | 95 | 91,92 | ABREU | 97B DLPH | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| > 77.9 | 95 | 91,93 | ABREU | 97B DLPH | $e^+ e^- \rightarrow e^* e^*$ | Sequential type |
| > 79.7 | 95 | 91 | ACCIARRI | 97G L3 | $e^+ e^- \rightarrow e^* e^*$ | Sequential type |
| > 79.9 | 95 | 91,94 | ACKERSTAFF | 97 OPAL | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| > 62.5 | 95 | 95 | ABREU | 96K DLPH | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| > 64.7 | 95 | 96 | ACCIARRI | 96D L3 | $e^+ e^- \rightarrow e^* e^*$ | Sequential type |
| > 66.5 | 95 | 96 | ALEXANDER | 96Q OPAL | $e^+ e^- \rightarrow e^* e^*$ | Homodoublet type |
| > 65.2 | 95 | 96 | BUSKULIC | 96W ALEP | $e^+ e^- \rightarrow e^* e^*$ | Sequential type |
| > 45.6 | 95 | | ADRIANI | 93M L3 | $Z \rightarrow e^* e^*$ | |
| > 45.6 | 95 | | ABREU | 92C DLPH | $Z \rightarrow e^* e^*$ | |
| > 29.8 | 95 | 97 | BARDADIN-... | 92 RVUE | $\Gamma(Z)$ | |
| > 26.1 | 95 | 98 | DECAMP | 92 ALEP | $Z \rightarrow e^* e^*$; $\Gamma(Z)$ | |
| > 46.1 | 95 | | DECAMP | 92 ALEP | $Z \rightarrow e^* e^*$ | |
| > 33 | 95 | 98 | ABREU | 91F DLPH | $Z \rightarrow e^* e^*$; $\Gamma(Z)$ | |
| > 45.0 | 95 | 99 | ADEVA | 90F L3 | $Z \rightarrow e^* e^*$ | |
| > 44.9 | 95 | | AKRAWY | 90I OPAL | $Z \rightarrow e^* e^*$ | |
| > 44.6 | 95 | 100 | DECAMP | 90G ALEP | $e^+ e^- \rightarrow e^* e^*$ | |
| > 30.2 | 95 | | ADACHI | 89B TOPZ | $e^+ e^- \rightarrow e^* e^*$ | |
| > 28.3 | 95 | | KIM | 89 AMY | $e^+ e^- \rightarrow e^* e^*$ | |
| > 27.9 | 95 | 101 | ABE | 88B VNS | $e^+ e^- \rightarrow e^* e^*$ | |

⁸³ From $e^+ e^-$ collisions at $\sqrt{s} = 183\text{--}209$ GeV. $f = f'$ is assumed.

⁸⁴ From $e^+ e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = f'$ is assumed. ACHARD 03B also obtain limit for $f = -f'$: $m_{e^*} > 96.6$ GeV.

⁸⁵ From $e^+ e^-$ collisions at $\sqrt{s} = 192\text{--}202$ GeV. $f=f'$ is assumed. ACCIARRI 01D also obtain limit for $f=-f'$: $m_{e^*} > 93.4$ GeV.

⁸⁶ From $e^+ e^-$ collisions at $\sqrt{s}=161\text{--}183$ GeV. $f=f'$ is assumed. ABBIENDI 00I also obtain limit for $f=-f'$ ($e^* \rightarrow \nu W$): $m_{e^*} > 86.0$ GeV.

⁸⁷ From $e^+ e^-$ collisions at $\sqrt{s}=189$ GeV. $f=f'$ is assumed. ACCIARRI 00E also obtain limit for $f=-f'$ ($e^* \rightarrow \nu W$): $m_{e^*} > 92.6$ GeV.

⁸⁸ From $e^+ e^-$ collisions at $\sqrt{s}= 183$ GeV. $f=f'$ is assumed. ABREU 990 also obtain limit for $f=-f'$ ($e^* \rightarrow \nu W$): $m_{e^*} > 81.3$ GeV.

⁸⁹ From $e^+ e^-$ collisions at $\sqrt{s}=170\text{--}172$ GeV. ACKERSTAFF 98C also obtain limit from $e^* \rightarrow \nu W$ decay mode: $m_{e^*} > 81.3$ GeV.

⁹⁰ BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.

⁹¹ From $e^+ e^-$ collisions at $\sqrt{s}= 161$ GeV.

⁹² ABREU 97B also obtain limit from charged current decay mode $e^* \rightarrow \nu W$, $m_{e^*} > 70.9$ GeV.

⁹³ ABREU 97B also obtain limit from charged current decay mode $e^* \rightarrow \nu W$, $m_{e^*} > 44.6$ GeV.

- 94 ACKERSTAFF 97 also obtain limit from charged current decay mode $e^* \rightarrow \nu W$, $m_{\nu_e^*} > 77.1$ GeV.
 95 From e^+e^- collisions at $\sqrt{s}=130\text{--}136$ GeV.
 96 From e^+e^- collisions at $\sqrt{s}=130\text{--}140$ GeV.
 97 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on $\Delta\Gamma(Z) < 36$ MeV.
 98 Limit is independent of e^* decay mode.
 99 ADEVA 90F is superseded by ADRIANI 93M.
 100 Superseded by DECAMP 92.
 101 ABE 88B limits assume $e^+e^- \rightarrow e^*e^{*-}$ with one photon exchange only and $e^* \rightarrow e\gamma$ giving $e e \gamma \gamma$.

Limits for Excited e (e^*) from Single Production

These limits are from $e^+e^- \rightarrow e^*e$, $W \rightarrow e^*\nu$, or $ep \rightarrow e^*X$ and depend on transition magnetic coupling between e and e^* . All limits assume $e^* \rightarrow e\gamma$ decay except as noted. Limits from LEP, UA2, and H1 are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda\text{--}m_{e^*}$ plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------|----------|---|
| >255 | 95 | 102 ADLOFF | 02B H1 | $ep \rightarrow e^*X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| >209 | 95 | 103 ACOSTA | 05B CDF | $p\bar{p} \rightarrow e^*X$ |
| >206 | 95 | 104 ACHARD | 03B L3 | $e^+e^- \rightarrow ee^*$ |
| >208 | 95 | 105 ABBIENDI | 02G OPAL | $e^+e^- \rightarrow ee^*$ |
| >228 | 95 | 106 CHEKANOV | 02D ZEUS | $ep \rightarrow e^*X$ |
| >202 | | 107 ACCIARRI | 01D L3 | $e^+e^- \rightarrow ee^*$ |
| | | 108 ABBIENDI | 00I OPAL | $e^+e^- \rightarrow ee^*$ |
| | | 109 ACCIARRI | 00E L3 | $e^+e^- \rightarrow ee^*$ |
| >223 | 95 | 110 ADLOFF | 00E H1 | $ep \rightarrow e^*X$ |
| | | 111 ABREU | 99O DLPH | $e^+e^- \rightarrow ee^*$ |
| none 20–170 | 95 | 112 ACCIARRI | 98T L3 | $e\gamma \rightarrow e^* \rightarrow e\gamma$ |
| | | 113 ACKERSTAFF | 98C OPAL | $e^+e^- \rightarrow ee^*$ |
| | | 114 BARATE | 98U ALEP | $e^+e^- \rightarrow ee^*$ |
| | | 115,116 ABREU | 97B DLPH | $e^+e^- \rightarrow ee^*$ |
| | | 115,117 ACCIARRI | 97G L3 | $e^+e^- \rightarrow ee^*$ |
| | | 118 ACKERSTAFF | 97 OPAL | $e^+e^- \rightarrow ee^*$ |
| | | 119 ADLOFF | 97 H1 | Lepton-flavor violation |
| none 30–200 | 95 | 120 BREITWEG | 97C ZEUS | $ep \rightarrow e^*X$ |
| | | 121 ABREU | 96K DLPH | $e^+e^- \rightarrow ee^*$ |
| | | 122 ACCIARRI | 96D L3 | $e^+e^- \rightarrow ee^*$ |
| | | 123 ALEXANDER | 96Q OPAL | $e^+e^- \rightarrow ee^*$ |
| | | 124 BUSKULIC | 96W ALEP | $e^+e^- \rightarrow ee^*$ |
| | | 125 DERRICK | 95B ZEUS | $ep \rightarrow e^*X$ |
| | | 126 ABT | 93 H1 | $ep \rightarrow e^*X$ |
| > 86 | 95 | ADRIANI | 93M L3 | $\lambda_\gamma > 0.04$ |

- | | | | | |
|------------|----|-------------|----------|---|
| > 89 | 95 | ADRIANI | 93M L3 | $Z \rightarrow ee^*$, $\lambda_Z > 0.5$ |
| | | 127 DERRICK | 93B ZEUS | Superseded by DERRICK 95B |
| > 88 | 95 | ABREU | 92C DLPH | $Z \rightarrow ee^*$, $\lambda_Z > 0.5$ |
| > 86 | 95 | ABREU | 92C DLPH | $e^+e^- \rightarrow ee^*$, $\lambda_\gamma > 0.1$ |
| > 91 | 95 | DECAMP | 92 ALEP | $Z \rightarrow ee^*$, $\lambda_Z > 1$ |
| > 88 | 95 | 128 ADEVA | 90F L3 | $Z \rightarrow ee^*$, $\lambda_Z > 0.5$ |
| > 86 | 95 | 128 ADEVA | 90F L3 | $Z \rightarrow ee^*$, $\lambda_Z > 0.04$ |
| > 87 | 95 | AKRAWY | 90I OPAL | $Z \rightarrow ee^*$, $\lambda_Z > 0.5$ |
| > 81 | 95 | 129 DECAMP | 90G ALEP | $Z \rightarrow ee^*$, $\lambda_Z > 1$ |
| > 50 | 95 | ADACHI | 89B TOPZ | $e^+e^- \rightarrow ee^*$, $\lambda_\gamma > 0.04$ |
| > 56 | 95 | KIM | 89 AMY | $e^+e^- \rightarrow ee^*$, $\lambda_\gamma > 0.03$ |
| none 23–54 | 95 | 130 ABE | 88B VNS | $e^+e^- \rightarrow ee^*$, $\lambda_\gamma > 0.04$ |
| > 75 | 95 | 131 ANSARI | 87D UA2 | $W \rightarrow e^*\nu$; $\lambda_W > 0.7$ |
| > 63 | 95 | 131 ANSARI | 87D UA2 | $W \rightarrow e^*\nu$; $\lambda_W > 0.2$ |
| > 40 | 95 | 131 ANSARI | 87D UA2 | $W \rightarrow e^*\nu$; $\lambda_W > 0.09$ |
- 102 ADLOFF 02B search for single e^* production in ep collisions with the decays $e^* \rightarrow e\gamma$, eZ , νW . $f = f' = \Lambda/m_{e^*}$ is assumed for the e^* coupling. See their Fig. 3 for the exclusion plot in the mass-coupling plane.
- 103 ACOSTA 05B search for single e^* production in $p\bar{p}$ collisions with the decays $e^* \rightarrow e\gamma$. $f = f' = \Lambda/m_{e^*}$ is assumed for the e^* coupling. See their Fig.3 for the exclusion limit in the mass-coupling plane.
- 104 ACHARD 03B result is from e^+e^- collisions at $\sqrt{s} = 189\text{--}209$ GeV. See their Fig. 4 for the exclusion plot in the mass-coupling plane.
- 105 ABBIENDI 02G result is from e^+e^- collisions at $\sqrt{s} = 183\text{--}209$ GeV. $f = f' = \Lambda/m_{e^*}$ is assumed for e^* coupling. See their Fig. 4c for the exclusion limit in the mass-coupling plane.
- 106 CHEKANOV 02D search for single e^* production in ep collisions with the decays $e^* \rightarrow e\gamma$, eZ , νW . $f = f' = \Lambda/m_{e^*}$ is assumed for the e^* coupling. See their Fig. 5a for the exclusion plot in the mass-coupling plane.
- 107 ACCIARRI 01D result is from e^+e^- collisions at $\sqrt{s} = 192\text{--}202$ GeV. $f=f'=\Lambda/m_{e^*}$ is assumed for the e^* coupling. See their Fig. 4 for limits in the mass-coupling plane.
- 108 ABBIENDI 00I result is from e^+e^- collisions at $\sqrt{s}=161\text{--}183$ GeV. See their Fig. 7 for limits in mass-coupling plane.
- 109 ACCIARRI 00E result is from e^+e^- collisions at $\sqrt{s}=189$ GeV. See their Fig. 3 for limits in mass-coupling plane.
- 110 ADLOFF 00E search for single e^* production in ep collisions with the decays $e^* \rightarrow e\gamma$, eZ , νW . $f=f'=\Lambda/m_{e^*}$ is assumed for the e^* coupling. See their Fig. 9 for the exclusion plot in the mass-coupling plane.
- 111 ABREU 99O result is from e^+e^- collisions at $\sqrt{s}= 183$ GeV. See their Figs. 4 and 5 for the exclusion limit in the mass-coupling plane.
- 112 ACCIARRI 98T search for single e^* production in quasi-real Compton scattering. The limit is for $|\lambda| > 1.0 \times 10^{-1}$ and non-chiral coupling of e^* . See their Fig. 7 for the exclusion plot in the mass-coupling plane.
- 113 ACKERSTAFF 98C from e^+e^- collisions at $\sqrt{s}=170\text{--}172$ GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.
- 114 BARATE 98U is from e^+e^- collision at $\sqrt{s}=M_Z$. See their Fig. 12 for limits in mass-coupling plane
- 115 From e^+e^- collisions at $\sqrt{s}= 161$ GeV.
- 116 See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane.

- 117 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.
- 118 ACKERSTAFF 97 result is from e^+e^- collisions at $\sqrt{s}=161$ GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 119 ADLOFF 97 search for single e^* production in ep collisions with the decays $e^* \rightarrow e\gamma, eZ, \nu W$. See their Fig. 4 for the rejection limits on the product of the production cross section and the branching ratio into a specific decay channel.
- 120 BREITWEG 97C search for single e^* production in ep collisions with the decays $e^* \rightarrow e\gamma, eZ, \nu W$. $f=f'=2\Lambda/m_{e^*}$ is assumed for the e^* coupling. See their Fig. 9 for the exclusion plot in the mass-coupling plane.
- 121 ABREU 96K result is from e^+e^- collisions at $\sqrt{s}=130-136$ GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.
- 122 ACCIARRI 96D result is from e^+e^- collisions at $\sqrt{s}=130-140$ GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.
- 123 ALEXANDER 96Q result is from e^+e^- collisions at $\sqrt{s}=130-140$ GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane.
- 124 BUSKULIC 96W result is from e^+e^- collisions at $\sqrt{s}=130-140$ GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 125 DERRICK 95B search for single e^* production via $e^*e\gamma$ coupling in ep collisions with the decays $e^* \rightarrow e\gamma, eZ, \nu W$. See their Fig. 13 for the exclusion plot in the $m_{e^*}-\lambda\gamma$ plane.
- 126 ABT 93 search for single e^* production via $e^*e\gamma$ coupling in ep collisions with the decays $e^* \rightarrow e\gamma, eZ, \nu W$. See their Fig. 4 for exclusion plot in the $m_{e^*}-\lambda\gamma$ plane.
- 127 DERRICK 93B search for single e^* production via $e^*e\gamma$ coupling in ep collisions with the decays $e^* \rightarrow e\gamma, eZ, \nu W$. See their Fig. 3 for exclusion plot in the $m_{e^*}-\lambda\gamma$ plane.
- 128 Superseded by ADRIANI 93M.
- 129 Superseded by DECAMP 92.
- 130 ABE 88B limits use $e^+e^- \rightarrow ee^*$ where t-channel photon exchange dominates giving $e\gamma(e)$ (quasi-real compton scattering).
- 131 ANSARI 87D is at $E_{cm} = 546-630$ GeV.

Limits for Excited e (e^*) from $e^+e^- \rightarrow \gamma\gamma$

These limits are derived from indirect effects due to e^* exchange in the t channel and depend on transition magnetic coupling between e and e^* . All limits are for $\lambda_\gamma = 1$. All limits except ABE 89J and ACHARD 02D are for nonchiral coupling with $\eta_L = \eta_R = 1$. We choose the chiral coupling limit as the best limit and list it in the Summary Table.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------|----------|-------------------------|
| >310 | 95 | ACHARD | 02D L3 | $\sqrt{s}=192-209$ GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| >356 | 95 | ¹³² ABDALLAH | 04N DLPH | $\sqrt{s}=161-208$ GeV |
| >311 | 95 | ABREU | 00A DLPH | $\sqrt{s}=189-202$ GeV |
| >283 | 95 | ¹³³ ACCIARRI | 00G L3 | $\sqrt{s}=183-189$ GeV |
| >306 | 95 | ABBIENDI | 99P OPAL | $\sqrt{s}=189$ GeV |
| >231 | 95 | ABREU | 98J DLPH | $\sqrt{s}=130-183$ GeV |
| >194 | 95 | ACKERSTAFF | 98 OPAL | $\sqrt{s}=130-172$ GeV |
| >227 | 95 | ACKER...,K... | 98B OPAL | $\sqrt{s}=183$ GeV |
| >250 | 95 | BARATE | 98J ALEP | $\sqrt{s}=183$ GeV |
| >160 | 95 | ¹³⁴ BARATE | 98U ALEP | |
| >210 | 95 | ¹³⁵ ACCIARRI | 97W L3 | $\sqrt{s}=161, 172$ GeV |

| | | | | |
|--------|----|------------------|----------|-------------------------|
| >129 | 95 | ACCIARRI | 96L L3 | $\sqrt{s}=133$ GeV |
| >147 | 95 | ALEXANDER | 96K OPAL | |
| >136 | 95 | BUSKULIC | 96Z ALEP | $\sqrt{s}=130, 136$ GeV |
| >146 | 95 | ACCIARRI | 95G L3 | |
| | | 136 BUSKULIC | 93Q ALEP | |
| >127 | 95 | 137 ADRIANI | 92B L3 | |
| >114 | 95 | 138 BARDADIN-... | 92 RVUE | |
| > 99 | 95 | DECAMP | 92 ALEP | |
| | | 139 SHIMOZAWA | 92 TOPZ | |
| >100 | 95 | ABREU | 91E DLPH | |
| >116 | 95 | AKRAWY | 91F OPAL | |
| > 83 | 95 | ADEVA | 90K L3 | |
| > 82 | 95 | AKRAWY | 90F OPAL | |
| > 68 | 95 | 140 ABE | 89J VNS | $\eta_L=1, \eta_R=0$ |
| > 90.2 | 95 | ADACHI | 89B TOPZ | |
| > 65 | 95 | KIM | 89 AMY | |

132 ABDALLAH 04N also obtain a limit on the excited electron mass with $e e^*$ chiral coupling, $m_{e^*} > 295$ GeV at 95% CL.

133 ACCIARRI 00G also obtain a limit on e^* with chiral coupling, $m_{e^*} > 213$ GeV.

134 BARATE 98U is from $e^+ e^-$ collision at $\sqrt{s}=M_Z$. See their Fig. 5 for limits in mass-coupling plane

135 ACCIARRI 97W also obtain a limit on e^* with chiral coupling, $m_{e^*} > 157$ GeV (95%CL).

136 BUSKULIC 93Q obtain $\Lambda^+ > 121$ GeV (95%CL) from ALEPH experiment and $\Lambda^+ > 135$ GeV from combined TRISTAN and ALEPH data. These limits roughly correspond to limits on m_{e^*} .

137 ADRIANI 92B superseded by ACCIARRI 95G.

138 BARDADIN-OTWINOWSKA 92 limit from fit to the combined data of DECAMP 92, ABREU 91E, ADEVA 90K, AKRAWY 91F.

139 SHIMOZAWA 92 fit the data to the limiting form of the cross section with $m_{e^*} \gg E_{cm}$ and obtain $m_{e^*} > 168$ GeV at 95%CL. Use of the full form would reduce this limit by a few GeV. The statistically unexpected large value is due to fluctuation in the data.

140 The ABE 89J limit assumes chiral coupling. This corresponds to $\lambda_\gamma = 0.7$ for nonchiral coupling.

Indirect Limits for Excited e (e^*)

These limits make use of loop effects involving e^* and are therefore subject to theoretical uncertainty.

| <u>VALUE (GeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|----------------|
|--------------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|-------------|----|------|--|
| 141 | DORENBOS... | 89 | CHRM | $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ and $\nu_\mu e \rightarrow \nu_\mu e$ |
| 142 | GRIFOLS | 86 | THEO | $\nu_\mu e \rightarrow \nu_\mu e$ |
| 143 | RENARD | 82 | THEO | $g-2$ of electron |

141 DORENBOSCH 89 obtain the limit $\lambda_\gamma^2 \Lambda_{cut}^2 / m_{e^*}^2 < 2.6$ (95% CL), where Λ_{cut} is the cutoff scale, based on the one-loop calculation by GRIFOLS 86. If one assumes that $\Lambda_{cut} = 1$ TeV and $\lambda_\gamma = 1$, one obtains $m_{e^*} > 620$ GeV. However, one generally expects $\lambda_\gamma \approx m_{e^*} / \Lambda_{cut}$ in composite models.

142 GRIFOLS 86 uses $\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ data from CHARM Collaboration to derive mass limits which depend on the scale of compositeness.

¹⁴³RENARD 82 derived from $g-2$ data limits on mass and couplings of e^* and μ^* . See figures 2 and 3 of the paper.

MASS LIMITS for Excited μ (μ^*)

Limits for Excited μ (μ^*) from Pair Production

These limits are obtained from $e^+e^- \rightarrow \mu^{*+}\mu^{*-}$ and thus rely only on the (electroweak) charge of μ^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the μ^* coupling is assumed to be of sequential type. All limits assume a dominant $\mu^* \rightarrow \mu\gamma$ decay except the limits from $\Gamma(Z)$.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------------|----------|--|
| >103.2 | 95 | ¹⁴⁴ ABBIENDI | 02G OPAL | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| >102.8 | 95 | ¹⁴⁵ ACHARD | 03B L3 | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| >100.2 | 95 | ¹⁴⁶ ACCIARRI | 01D L3 | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| > 91.3 | 95 | ¹⁴⁷ ABBIENDI | 00I OPAL | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| > 94.2 | 95 | ¹⁴⁸ ACCIARRI | 00E L3 | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| > 90.7 | 95 | ¹⁴⁹ ABREU | 99O DLPH | Homodoublet type |
| > 85.3 | 95 | ¹⁵⁰ ACKERSTAFF | 98C OPAL | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| | | ¹⁵¹ BARATE | 98U ALEP | $Z \rightarrow \mu^*\mu^*$ |
| > 79.6 | 95 | ^{152,153} ABREU | 97B DLPH | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| > 78.4 | 95 | ^{152,154} ABREU | 97B DLPH | $e^+e^- \rightarrow \mu^*\mu^*$ Sequential type |
| > 79.9 | 95 | ¹⁵² ACCIARRI | 97G L3 | $e^+e^- \rightarrow \mu^*\mu^*$ Sequential type |
| > 80.0 | 95 | ^{152,155} ACKERSTAFF | 97 OPAL | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| > 62.6 | 95 | ¹⁵⁶ ABREU | 96K DLPH | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| > 64.9 | 95 | ¹⁵⁷ ACCIARRI | 96D L3 | $e^+e^- \rightarrow \mu^*\mu^*$ Sequential type |
| > 66.8 | 95 | ¹⁵⁷ ALEXANDER | 96Q OPAL | $e^+e^- \rightarrow \mu^*\mu^*$ Homodoublet type |
| > 65.4 | 95 | ¹⁵⁷ BUSKULIC | 96W ALEP | $e^+e^- \rightarrow \mu^*\mu^*$ Sequential type |
| > 45.6 | 95 | ADRIANI | 93M L3 | $Z \rightarrow \mu^*\mu^*$ |
| > 45.6 | 95 | ABREU | 92C DLPH | $Z \rightarrow \mu^*\mu^*$ |
| > 29.8 | 95 | ¹⁵⁸ BARDADIN-... | 92 RVUE | $\Gamma(Z)$ |
| > 26.1 | 95 | ¹⁵⁹ DECAMP | 92 ALEP | $Z \rightarrow \mu^*\mu^*; \Gamma(Z)$ |
| > 46.1 | 95 | DECAMP | 92 ALEP | $Z \rightarrow \mu^*\mu^*$ |
| > 33 | 95 | ¹⁵⁹ ABREU | 91F DLPH | $Z \rightarrow \mu^*\mu^*; \Gamma(Z)$ |
| > 45.3 | 95 | ¹⁶⁰ ADEVA | 90F L3 | $Z \rightarrow \mu^*\mu^*$ |
| > 44.9 | 95 | AKRAWY | 90I OPAL | $Z \rightarrow \mu^*\mu^*$ |
| > 44.6 | 95 | ¹⁶¹ DECAMP | 90G ALEP | $e^+e^- \rightarrow \mu^*\mu^*$ |
| > 29.9 | 95 | ADACHI | 89B TOPZ | $e^+e^- \rightarrow \mu^*\mu^*$ |
| > 28.3 | 95 | KIM | 89 AMY | $e^+e^- \rightarrow \mu^*\mu^*$ |

¹⁴⁴ From e^+e^- collisions at $\sqrt{s} = 183-209$ GeV. $f = f'$ is assumed.

¹⁴⁵ From e^+e^- collisions at $\sqrt{s} = 189-209$ GeV. $f = f'$ is assumed. ACHARD 03B also obtain limit for $f = -f'$: $m_{\mu^*} > 96.6$ GeV.

¹⁴⁶ From e^+e^- collisions at $\sqrt{s} = 192-202$ GeV. $f=f'$ is assumed. ACCIARRI 01D also obtain limit for $f=-f'$: $m_{\mu^*} > 93.4$ GeV.

- 147 From e^+e^- collisions at $\sqrt{s}=161\text{--}183$ GeV. $f=f'$ is assumed. ABBIENDI 00I also obtain limit for $f=-f'$ ($\mu^* \rightarrow \nu W$): $m_{\mu^*} > 86.0$ GeV.
- 148 From e^+e^- collisions at $\sqrt{s}=189$ GeV. $f=f'$ is assumed. ACCIARRI 00E also obtain limit for $f=-f'$ ($\mu^* \rightarrow \nu W$): $m_{\mu^*} > 92.6$ GeV.
- 149 From e^+e^- collisions at $\sqrt{s}=183$ GeV. $f=f'$ is assumed. ABREU 990 also obtain limit for $f=-f'$ ($\mu^* \rightarrow \nu W$): $m_{\mu^*} > 81.3$ GeV.
- 150 From e^+e^- collisions at $\sqrt{s}=170\text{--}172$ GeV. ACKERSTAFF 98C also obtain limit from $\mu^* \rightarrow \nu W$ decay mode: $m_{\mu^*} > 81.3$ GeV.
- 151 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.
- 152 From e^+e^- collisions at $\sqrt{s}=161$ GeV.
- 153 ABREU 97B also obtain limit from charged current decay mode $\mu^* \rightarrow \nu W$, $m_{\mu^*} > 70.9$ GeV.
- 154 ABREU 97B also obtain limit from charged current decay mode $\mu^* \rightarrow \nu W$, $m_{\mu^*} > 44.6$ GeV.
- 155 ACKERSTAFF 97 also obtain limit from charged current decay mode $\mu^* \rightarrow \nu W$, $m_{\nu\mu} > 77.1$ GeV.
- 156 From e^+e^- collisions at $\sqrt{s}=130\text{--}136$ GeV.
- 157 From e^+e^- collisions at $\sqrt{s}=130\text{--}140$ GeV.
- 158 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on $\Delta\Gamma(Z) < 36$ MeV.
- 159 Limit is independent of μ^* decay mode.
- 160 Superseded by ADRIANI 93M.
- 161 Superseded by DECAMP 92.

Limits for Excited μ (μ^*) from Single Production

These limits are from $e^+e^- \rightarrow \mu^*\mu$ and depend on transition magnetic coupling between μ and μ^* . All limits assume $\mu^* \rightarrow \mu\gamma$ decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda\text{--}m_{\mu^*}$ plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------|----------|-------------------------------|
| >190 | 95 | 162 ABBIENDI | 02G OPAL | $e^+e^- \rightarrow \mu\mu^*$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| >180 | 95 | 163 ACHARD | 03B L3 | $e^+e^- \rightarrow \mu\mu^*$ |
| >178 | 95 | 164 ACCIARRI | 01D L3 | $e^+e^- \rightarrow \mu\mu^*$ |
| | | 165 ABBIENDI | 00I OPAL | $e^+e^- \rightarrow \mu\mu^*$ |
| | | 166 ACCIARRI | 00E L3 | $e^+e^- \rightarrow \mu\mu^*$ |
| | | 167 ABREU | 990 DLPH | $e^+e^- \rightarrow \mu\mu^*$ |
| | | 168 ACKERSTAFF | 98C OPAL | $e^+e^- \rightarrow \mu\mu^*$ |
| | | 169 BARATE | 98U ALEP | $Z \rightarrow \mu\mu^*$ |
| 170,171 | | ABREU | 97B DLPH | $e^+e^- \rightarrow \mu\mu^*$ |
| 170,172 | | ACCIARRI | 97G L3 | $e^+e^- \rightarrow \mu\mu^*$ |
| | | 173 ACKERSTAFF | 97 OPAL | $e^+e^- \rightarrow \mu\mu^*$ |
| | | 174 ABREU | 96K DLPH | $e^+e^- \rightarrow \mu\mu^*$ |

| | | | | |
|------|-----|------------|----------|--|
| | 175 | ACCIARRI | 96D L3 | $e^+ e^- \rightarrow \mu\mu^*$ |
| | 176 | ALEXANDER | 96Q OPAL | $e^+ e^- \rightarrow \mu\mu^*$ |
| | 177 | BUSKULIC | 96W ALEP | $e^+ e^- \rightarrow \mu\mu^*$ |
| > 89 | 95 | ADRIANI | 93M L3 | $Z \rightarrow \mu\mu^*, \lambda_Z > 0.5$ |
| > 88 | 95 | ABREU | 92C DLPH | $Z \rightarrow \mu\mu^*, \lambda_Z > 0.5$ |
| > 91 | 95 | DECAMP | 92 ALEP | $Z \rightarrow \mu\mu^*, \lambda_Z > 1$ |
| > 85 | 95 | 178 ADEVA | 90F L3 | $Z \rightarrow \mu\mu^*, \lambda_Z > 1$ |
| > 75 | 95 | 178 ADEVA | 90F L3 | $Z \rightarrow \mu\mu^*, \lambda_Z > 0.1$ |
| > 87 | 95 | AKRAWY | 90I OPAL | $Z \rightarrow \mu\mu^*, \lambda_Z > 1$ |
| > 80 | 95 | 179 DECAMP | 90G ALEP | $e^+ e^- \rightarrow \mu\mu^*, \lambda_Z=1$ |
| > 50 | 95 | ADACHI | 89B TOPZ | $e^+ e^- \rightarrow \mu\mu^*, \lambda_\gamma=0.7$ |
| > 46 | 95 | KIM | 89 AMY | $e^+ e^- \rightarrow \mu\mu^*, \lambda_\gamma=0.2$ |

162 ABBIENDI 02G result is from $e^+ e^-$ collisions at $\sqrt{s} = 183\text{--}209$ GeV. $f = f' = \Lambda/m_{\mu^*}$ is assumed for μ^* coupling. See their Fig. 4c for the exclusion limit in the mass-coupling plane.

163 ACHARD 03B result is from $e^+ e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = f' = \Lambda/m_{\mu^*}$ is assumed. See their Fig. 4 for the exclusion plot in the mass-coupling plane.

164 ACCIARRI 01D result is from $e^+ e^-$ collisions at $\sqrt{s} = 192\text{--}202$ GeV. $f=f'=\Lambda/m_{\mu^*}$ is assumed for the μ^* coupling. See their Fig. 4 for limits in the mass-coupling plane.

165 ABBIENDI 00I result is from $e^+ e^-$ collisions at $\sqrt{s}=161\text{--}183$ GeV. See their Fig. 7 for limits in mass-coupling plane.

166 ACCIARRI 00E result is from $e^+ e^-$ collisions at $\sqrt{s}=189$ GeV. See their Fig. 3 for limits in mass-coupling plane.

167 ABREU 99O result is from $e^+ e^-$ collisions at $\sqrt{s}= 183$ GeV. See their Figs. 4 and 5 for the exclusion limit in the mass-coupling plane.

168 ACKERSTAFF 98C from $e^+ e^-$ collisions at $\sqrt{s}=170\text{--}172$ GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.

169 BARATE 98U obtain limits on the $Z\mu\mu^*$ coupling. See their Fig. 12 for limits in mass-coupling plane

170 From $e^+ e^-$ collisions at $\sqrt{s}= 161$ GeV.

171 See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane.

172 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.

173 ACKERSTAFF 97 result is from $e^+ e^-$ collisions at $\sqrt{s}= 161$ GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

174 ABREU 96K result is from $e^+ e^-$ collisions at $\sqrt{s}= 130\text{--}136$ GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.

175 ACCIARRI 96D result is from $e^+ e^-$ collisions at $\sqrt{s}= 130\text{--}140$ GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.

176 ALEXANDER 96Q result is from $e^+ e^-$ collisions at $\sqrt{s}= 130\text{--}140$ GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane.

177 BUSKULIC 96W result is from $e^+ e^-$ collisions at $\sqrt{s}= 130\text{--}140$ GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

178 Superseded by ADRIANI 93M.

179 Superseded by DECAMP 92.

Indirect Limits for Excited μ (μ^*)

These limits make use of loop effects involving μ^* and are therefore subject to theoretical uncertainty.

| <u>VALUE (GeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|----------------|
|--------------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

180 RENARD 82 THEO $g-2$ of muon

180 RENARD 82 derived from $g-2$ data limits on mass and couplings of e^* and μ^* . See figures 2 and 3 of the paper.

MASS LIMITS for Excited τ (τ^*)

Limits for Excited τ (τ^*) from Pair Production

These limits are obtained from $e^+e^- \rightarrow \tau^{*+}\tau^{*-}$ and thus rely only on the (electroweak) charge of τ^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the τ^* coupling is assumed to be of sequential type. All limits assume a dominant $\tau^* \rightarrow \tau\gamma$ decay except the limits from $\Gamma(Z)$.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

| <u>VALUE (GeV)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| >103.2 | 95 | 181 ABBIENDI | 02G OPAL | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| >102.8 | 95 | 182 ACHARD | 03B L3 | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| > 99.8 | 95 | 183 ACCIARRI | 01D L3 | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| > 91.2 | 95 | 184 ABBIENDI | 00I OPAL | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| > 94.2 | 95 | 185 ACCIARRI | 00E L3 | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| > 89.7 | 95 | 186 ABREU | 99O DLPH | Homodoublet type |
| > 84.6 | 95 | 187 ACKERSTAFF | 98C OPAL | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| | | 188 BARATE | 98U ALEP | $Z \rightarrow \tau^*\tau^*$ |
| > 79.4 | 95 | 189,190 ABREU | 97B DLPH | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| > 77.4 | 95 | 189,191 ABREU | 97B DLPH | $e^+e^- \rightarrow \tau^*\tau^*$ Sequential type |
| > 79.3 | 95 | 189 ACCIARRI | 97G L3 | $e^+e^- \rightarrow \tau^*\tau^*$ Sequential type |
| > 79.1 | 95 | 189,192 ACKERSTAFF | 97 OPAL | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| > 62.2 | 95 | 193 ABREU | 96K DLPH | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| > 64.2 | 95 | 194 ACCIARRI | 96D L3 | $e^+e^- \rightarrow \tau^*\tau^*$ Sequential type |
| > 65.3 | 95 | 194 ALEXANDER | 96Q OPAL | $e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type |
| > 64.8 | 95 | 194 BUSKULIC | 96W ALEP | $e^+e^- \rightarrow \tau^*\tau^*$ Sequential type |
| > 45.6 | 95 | ADRIANI | 93M L3 | $Z \rightarrow \tau^*\tau^*$ |
| > 45.3 | 95 | ABREU | 92C DLPH | $Z \rightarrow \tau^*\tau^*$ |
| > 29.8 | 95 | 195 BARDADIN-... | 92 RVUE | $\Gamma(Z)$ |
| > 26.1 | 95 | 196 DECAMP | 92 ALEP | $Z \rightarrow \tau^*\tau^*$; $\Gamma(Z)$ |
| > 46.0 | 95 | DECAMP | 92 ALEP | $Z \rightarrow \tau^*\tau^*$ |
| > 33 | 95 | 196 ABREU | 91F DLPH | $Z \rightarrow \tau^*\tau^*$; $\Gamma(Z)$ |
| > 45.5 | 95 | 197 ADEVA | 90L L3 | $Z \rightarrow \tau^*\tau^*$ |
| > 44.9 | 95 | AKRAWY | 90I OPAL | $Z \rightarrow \tau^*\tau^*$ |
| > 41.2 | 95 | 198 DECAMP | 90G ALEP | $e^+e^- \rightarrow \tau^*\tau^*$ |
| > 29.0 | 95 | ADACHI | 89B TOPZ | $e^+e^- \rightarrow \tau^*\tau^*$ |

- 181 From e^+e^- collisions at $\sqrt{s} = 183\text{--}209$ GeV. $f = f'$ is assumed.
- 182 From e^+e^- collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = f'$ is assumed. ACHARD 03B also obtain limit for $f = -f'$: $m_{\tau^*} > 96.6$ GeV.
- 183 From e^+e^- collisions at $\sqrt{s} = 192\text{--}202$ GeV. $f=f'$ is assumed. ACCIARRI 01D also obtain limit for $f=-f'$: $m_{\tau^*} > 93.4$ GeV.
- 184 From e^+e^- collisions at $\sqrt{s}=161\text{--}183$ GeV. $f=f'$ is assumed. ABBIENDI 00I also obtain limit for $f=-f'$ ($\tau^* \rightarrow \nu W$): $m_{\tau^*} > 86.0$ GeV.
- 185 From e^+e^- collisions at $\sqrt{s}=189$ GeV. $f=f'$ is assumed. ACCIARRI 00E also obtain limit for $f=-f'$ ($\tau^* \rightarrow \nu W$): $m_{\tau^*} > 92.6$ GeV.
- 186 From e^+e^- collisions at $\sqrt{s}= 183$ GeV. $f=f'$ is assumed. ABREU 990 also obtain limit for $f=-f'$ ($\tau^* \rightarrow \nu W$): $m_{\tau^*} > 81.3$ GeV.
- 187 From e^+e^- collisions at $\sqrt{s}=170\text{--}172$ GeV. ACKERSTAFF 98C also obtain limit from $\tau^* \rightarrow \nu W$ decay mode: $m_{\tau^*} > 81.3$ GeV.
- 188 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.
- 189 From e^+e^- collisions at $\sqrt{s}= 161$ GeV.
- 190 ABREU 97B also obtain limit from charged current decay mode $\tau^* \rightarrow \nu W$, $m_{\tau^*} > 70.9$ GeV.
- 191 ABREU 97B also obtain limit from charged current decay mode $\tau^* \rightarrow \nu W$, $m_{\tau^*} > 44.6$ GeV.
- 192 ACKERSTAFF 97 also obtain limit from charged current decay mode $\tau^* \rightarrow \nu W$, $m_{\nu\tau^*} > 77.1$ GeV.
- 193 From e^+e^- collisions at $\sqrt{s}= 130\text{--}136$ GeV.
- 194 From e^+e^- collisions at $\sqrt{s}= 130\text{--}140$ GeV.
- 195 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on $\Delta\Gamma(Z)<36$ MeV.
- 196 Limit is independent of τ^* decay mode.
- 197 Superseded by ADRIANI 93M.
- 198 Superseded by DECAMP 92.

Limits for Excited τ (τ^*) from Single Production

These limits are from $e^+e^- \rightarrow \tau^*\tau$ and depend on transition magnetic coupling between τ and τ^* . All limits assume $\tau^* \rightarrow \tau\gamma$ decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda\text{--}m_{\tau^*}$ plane. See the original papers.

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------|----------|---------------------------------|
| >185 | 95 | 199 ABBIENDI | 02G OPAL | $e^+e^- \rightarrow \tau\tau^*$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| >180 | 95 | 200 ACHARD | 03B L3 | $e^+e^- \rightarrow \tau\tau^*$ |
| >173 | 95 | 201 ACCIARRI | 01D L3 | $e^+e^- \rightarrow \tau\tau^*$ |
| | | 202 ABBIENDI | 00I OPAL | $e^+e^- \rightarrow \tau\tau^*$ |
| | | 203 ACCIARRI | 00E L3 | $e^+e^- \rightarrow \tau\tau^*$ |
| | | 204 ABREU | 99O DLPH | $e^+e^- \rightarrow \tau\tau^*$ |
| | | 205 ACKERSTAFF | 98C OPAL | $e^+e^- \rightarrow \tau\tau^*$ |
| | | 206 BARATE | 98U ALEP | $Z \rightarrow \tau\tau^*$ |
| | | 207,208 ABREU | 97B DLPH | $e^+e^- \rightarrow \tau\tau^*$ |
| | | 207,209 ACCIARRI | 97G L3 | $e^+e^- \rightarrow \tau\tau^*$ |
| | | 210 ACKERSTAFF | 97 OPAL | $e^+e^- \rightarrow \tau\tau^*$ |

| | | | | | | |
|--------|--|-----------|---------|------|--|---|
| | 211 | ABREU | 96K | DLPH | $e^+e^- \rightarrow \tau\tau^*$ | |
| | 212 | ACCIARRI | 96D | L3 | $e^+e^- \rightarrow \tau\tau^*$ | |
| | 213 | ALEXANDER | 96Q | OPAL | $e^+e^- \rightarrow \tau\tau^*$ | |
| | 214 | BUSKULIC | 96W | ALEP | $e^+e^- \rightarrow \tau\tau^*$ | |
| > 88 | 95 | ADRIANI | 93M | L3 | $Z \rightarrow \tau\tau^*, \lambda_Z > 0.5$ | |
| > 87 | 95 | ABREU | 92C | DLPH | $Z \rightarrow \tau\tau^*, \lambda_Z > 0.5$ | |
| > 90 | 95 | DECAMP | 92 | ALEP | $Z \rightarrow \tau\tau^*, \lambda_Z > 0.18$ | |
| > 88 | 95 | 215 | ADEVA | 90L | L3 | $Z \rightarrow \tau\tau^*, \lambda_Z > 1$ |
| > 86.5 | 95 | AKRAWY | 90I | OPAL | $Z \rightarrow \tau\tau^*, \lambda_Z > 1$ | |
| > 59 | 95 | 216 | DECAMP | 90G | ALEP | $Z \rightarrow \tau\tau^*, \lambda_Z = 1$ |
| > 40 | 95 | 217 | BARTEL | 86 | JADE | $e^+e^- \rightarrow \tau\tau^*, \lambda_\gamma = 1$ |
| > 41.4 | 95 | 218 | BEHREND | 86 | CELL | $e^+e^- \rightarrow \tau\tau^*, \lambda_\gamma = 1$ |
| > 40.8 | 95 | 218 | BEHREND | 86 | CELL | $e^+e^- \rightarrow \tau\tau^*, \lambda_\gamma = 0.7$ |
| 199 | ABBIENDI 02G result is from e^+e^- collisions at $\sqrt{s} = 183\text{--}209$ GeV. $f = f' = \Lambda/m_{\tau^*}$ is assumed for τ^* coupling. See their Fig. 4c for the exclusion limit in the mass-coupling plane. | | | | | |
| 200 | ACHARD 03B result is from e^+e^- collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = f' = \Lambda/m_{\tau^*}$ is assumed. See their Fig. 4 for the exclusion plot in the mass-coupling plane. | | | | | |
| 201 | ACCIARRI 01D result is from e^+e^- collisions at $\sqrt{s} = 192\text{--}202$ GeV. $f=f'=\Lambda/m_{\tau^*}$ is assumed for the τ^* coupling. See their Fig. 4 for limits in the mass-coupling plane. | | | | | |
| 202 | ABBIENDI 00I result is from e^+e^- collisions at $\sqrt{s}=161\text{--}183$ GeV. See their Fig. 7 for limits in mass-coupling plane. | | | | | |
| 203 | ACCIARRI 00E result is from e^+e^- collisions at $\sqrt{s}=189$ GeV. See their Fig. 3 for limits in mass-coupling plane. | | | | | |
| 204 | ABREU 99O result is from e^+e^- collisions at $\sqrt{s}= 183$ GeV. See their Figs. 4 and 5 for the exclusion limit in the mass-coupling plane. | | | | | |
| 205 | ACKERSTAFF 98C from e^+e^- collisions at $\sqrt{s}=170\text{--}172$ GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane. | | | | | |
| 206 | BARATE 98U obtain limits on the $Z\tau\tau^*$ coupling. See their Fig. 12 for limits in mass-coupling plane | | | | | |
| 207 | From e^+e^- collisions at $\sqrt{s}= 161$ GeV. | | | | | |
| 208 | See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane. | | | | | |
| 209 | See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane. | | | | | |
| 210 | ACKERSTAFF 97 result is from e^+e^- collisions at $\sqrt{s}= 161$ GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane. | | | | | |
| 211 | ABREU 96K result is from e^+e^- collisions at $\sqrt{s}= 130\text{--}136$ GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane. | | | | | |
| 212 | ACCIARRI 96D result is from e^+e^- collisions at $\sqrt{s}= 130\text{--}140$ GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane. | | | | | |
| 213 | ALEXANDER 96Q result is from e^+e^- collisions at $\sqrt{s}= 130\text{--}140$ GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane. | | | | | |
| 214 | BUSKULIC 96W result is from e^+e^- collisions at $\sqrt{s}= 130\text{--}140$ GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane. | | | | | |
| 215 | Superseded by ADRIANI 93M. | | | | | |
| 216 | Superseded by DECAMP 92. | | | | | |
| 217 | BARTEL 86 is at $E_{\text{cm}} = 30\text{--}46.78$ GeV. | | | | | |
| 218 | BEHREND 86 limit is at $E_{\text{cm}} = 33\text{--}46.8$ GeV. | | | | | |

MASS LIMITS for Excited Neutrino (ν^*)

Limits for Excited ν (ν^*) from Pair Production

These limits are obtained from $e^+ e^- \rightarrow \nu^* \nu^*$ and thus rely only on the (electroweak) charge of ν^* . Form factor effects are ignored unless noted. The ν^* coupling is assumed to be of sequential type unless otherwise noted. All limits assume a dominant $\nu^* \rightarrow \nu \gamma$ decay except the limits from $\Gamma(Z)$.

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|--------------------|----------|--|
| >102.6 | 95 | 219 ACHARD | 03B L3 | $e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| | | 220 ABBIENDI | 04N OPAL | |
| > 99.4 | 95 | 221 ACCIARRI | 01D L3 | $e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type |
| > 91.2 | 95 | 222 ABBIENDI | 00I OPAL | $e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type |
| | | 223 ABBIENDI,G | 00D OPAL | |
| > 94.1 | 95 | 224 ACCIARRI | 00E L3 | $e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type |
| | | 225 ABBIENDI | 99F OPAL | |
| > 90.0 | 95 | 226 ABREU | 99O DLPH | Homodoublet type |
| > 84.9 | 95 | 227 ACKERSTAFF | 98C OPAL | $e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type |
| | | 228 BARATE | 98U ALEP | $Z \rightarrow \nu^* \nu^*$ |
| > 77.6 | 95 | 229,230 ABREU | 97B DLPH | $e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type |
| > 64.4 | 95 | 229,231 ABREU | 97B DLPH | $e^+ e^- \rightarrow \nu^* \nu^*$ Sequential type |
| > 71.2 | 95 | 229,232 ACCIARRI | 97G L3 | $e^+ e^- \rightarrow \nu^* \nu^*$ Sequential type |
| > 77.8 | 95 | 229,233 ACKERSTAFF | 97 OPAL | $e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type |
| > 61.4 | 95 | 234,235 ACCIARRI | 96D L3 | $e^+ e^- \rightarrow \nu^* \nu^*$ Sequential type |
| > 65.0 | 95 | 236,237 ALEXANDER | 96Q OPAL | $e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type |
| > 63.6 | 95 | 234 BUSKULIC | 96W ALEP | $e^+ e^- \rightarrow \nu^* \nu^*$ Sequential type |
| > 43.7 | 95 | 238 BARDADIN-... | 92 RVUE | $\Gamma(Z)$ |
| > 47 | 95 | 239 DECAMP | 92 ALEP | |
| > 42.6 | 95 | 240 DECAMP | 92 ALEP | $\Gamma(Z)$ |
| > 35.4 | 95 | 241,242 DECAMP | 90O ALEP | $\Gamma(Z)$ |
| > 46 | 95 | 242,243 DECAMP | 90O ALEP | |

219 From $e^+ e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = -f'$ is assumed. ACHARD 03B also obtain limit for $f = f'$: $m_{\nu_e^*} > 101.7$ GeV, $m_{\nu_\mu^*} > 101.8$ GeV, and $m_{\nu_\tau^*} > 92.9$ GeV.

See their Fig. 4 for the exclusion plot in the mass-coupling plane.

220 From $e^+ e^-$ collisions at $\sqrt{s} = 192\text{--}209$ GeV, ABBIENDI 04N obtain limit on $\sigma(e^+ e^- \rightarrow \nu^* \nu^*) B^2(\nu^* \rightarrow \nu \gamma)$. See their Fig.2. The limit ranges from 20 to 45fb for $m_{\nu^*} > 45$ GeV.

221 From $e^+ e^-$ collisions at $\sqrt{s} = 192\text{--}202$ GeV. $f=f'$ is assumed. ACCIARRI 01D also obtain limit for $f=-f'$: $m_{\nu_e^*} > 99.1$ GeV, $m_{\nu_\mu^*} > 99.3$ GeV, $m_{\nu_\tau^*} > 90.5$ GeV.

222 From $e^+ e^-$ collisions at $\sqrt{s}=161\text{--}183$ GeV. $f=-f'$ (photonic decay) is assumed. ABBIENDI 00I also obtain limit for $f=f'$ ($\nu^* \rightarrow \ell W$): $m_{\nu_e^*} > 91.1$ GeV, $m_{\nu_\mu^*} > 91.1$ GeV, $m_{\nu_\tau^*} > 83.1$ GeV.

223 From $e^+ e^-$ collisions at $\sqrt{s}= 189$ GeV. ABBIENDI,G 00D obtain limit on $\sigma(e^+ e^- \rightarrow \nu^* \nu^*) B(\nu^* \rightarrow \nu \gamma)^2$. See their Fig. 14. The limit ranges from 50 to 80 fb for $\sqrt{s}/2=95$ GeV $> m_{\nu^*} > 45$ GeV.

- 224 From e^+e^- collisions at $\sqrt{s}=189$ GeV. $f=-f'$ (photonic decay) is assumed. ACCIA-RRI 00E also obtain limit for $f=f'$ ($\nu^* \rightarrow \ell W$): $m_{\nu_e^*} > 93.9$ GeV, $m_{\nu_\mu^*} > 94.0$ GeV, $m_{\nu_\tau^*} > 91.5$ GeV.
- 225 From e^+e^- collisions at $\sqrt{s}=130-183$ GeV, ABBIENDI 99F obtain limit on $\sigma(e^+e^- \rightarrow \nu^*\nu^*) B(\nu^* \rightarrow \nu\gamma)^2$. See their Fig. 13. The limit ranges from 0.094 to 0.14 pb for $\sqrt{s}/2 > m_{\nu^*} > 45$ GeV.
- 226 From e^+e^- collisions at $\sqrt{s}=183$ GeV. $f=-f'$ is assumed. ABREU 99O also obtain limit for $f=f'$: $m_{\nu_{e^*}} > 87.3$ GeV, $m_{\nu_{\mu^*}} > 88.0$ GeV, $m_{\nu_{\tau^*}} > 81.0$ GeV.
- 227 From e^+e^- collisions at $\sqrt{s}=170-172$ GeV. ACKERSTAFF 98C also obtain limit from charged decay modes: $m_{\nu_e^*} > 84.1$ GeV, $m_{\nu_\mu^*} > 83.9$ GeV, and $m_{\nu_\tau^*} > 79.4$ GeV.
- 228 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.
- 229 From e^+e^- collisions at $\sqrt{s}=161$ GeV.
- 230 ABREU 97B also obtain limits from charged current decay modes, $m_{\nu^*} > 56.4$ GeV.
- 231 ABREU 97B also obtain limits from charged current decay modes, $m_{\nu^*} > 44.9$ GeV.
- 232 ACCIARRI 97G also obtain limits from charged current decay mode $\nu_e^* \rightarrow eW$, $m_{\nu^*} > 64.5$ GeV.
- 233 ACKERSTAFF 97 also obtain limits from charged current decay modes $m_{\nu_e^*} > 78.3$ GeV, $m_{\nu_\mu^*} > 78.9$ GeV, $m_{\nu_\tau^*} > 76.2$ GeV.
- 234 From e^+e^- collisions at $\sqrt{s}=130-140$ GeV.
- 235 ACCIARRI 96D also obtain limit from $\nu^* \rightarrow eW$ decay mode: $m_{\nu^*} > 57.3$ GeV.
- 236 From e^+e^- collisions at $\sqrt{s}=130-136$ GeV.
- 237 ALEXANDER 96Q also obtain limits from charged current decay modes: $m_{\nu_e^*} > 66.2$ GeV, $m_{\nu_\mu^*} > 66.5$ GeV, $m_{\nu_\tau^*} > 64.7$ GeV.
- 238 BARDADIN-OTWINOWSKA 92 limit is for Dirac ν^* . Based on $\Delta\Gamma(Z) < 36$ MeV. The limit is 36.4 GeV for Majorana ν^* , 45.4 GeV for homodoublet ν^* .
- 239 Limit is based on $B(Z \rightarrow \nu^*\bar{\nu}^*) \times B(\nu^* \rightarrow \nu\gamma)^2 < 5 \times 10^{-5}$ (95%CL) assuming Dirac ν^* , $B(\nu^* \rightarrow \nu\gamma) = 1$.
- 240 Limit is for Dirac ν^* . The limit is 34.6 GeV for Majorana ν^* , 45.4 GeV for homodoublet ν^* .
- 241 DECAMP 900 limit is from excess $\Delta\Gamma(Z) < 89$ MeV. The above value is for Dirac ν^* ; 26.6 GeV for Majorana ν^* ; 44.8 GeV for homodoublet ν^* .
- 242 Superseded by DECAMP 92.
- 243 DECAMP 900 limit based on $B(Z \rightarrow \nu^*\nu^*) \cdot B(\nu^* \rightarrow \nu\gamma)^2 < 7 \times 10^{-5}$ (95%CL), assuming Dirac ν^* , $B(\nu^* \rightarrow \nu\gamma) = 1$.

Limits for Excited ν (ν^*) from Single Production

These limits are from $e^+e^- \rightarrow \nu\nu^*$, $Z \rightarrow \nu\nu^*$, or $ep \rightarrow \nu^*X$ and depend on transition magnetic coupling between ν/e and ν^* . Assumptions about ν^* decay mode are given in footnotes.

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------|-----|-------------|--------|-------------------------------|
| >190 | 95 | 244 ACHARD | 03B L3 | $e^+e^- \rightarrow \nu\nu^*$ |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

| | | | | | | |
|-------------|---------|----------|--------------|------|---------------------------------|---|
| none 50–150 | 95 | 245 | ADLOFF | 02 | H1 | $ep \rightarrow \nu^* X$ |
| >158 | 95 | 246 | CHEKANOV | 02D | ZEUS | $ep \rightarrow \nu^* X$ |
| >171 | 95 | 247 | ACCIARRI | 01D | L3 | $e^+ e^- \rightarrow \nu \nu^*$ |
| | | 248 | ABBIENDI | 00I | OPAL | $e^+ e^- \rightarrow \nu \nu^*$ |
| | | 249 | ABBIENDI,G | 00D | OPAL | |
| | | 250 | ACCIARRI | 00E | L3 | $e^+ e^- \rightarrow \nu \nu^*$ |
| >114 | 95 | 251 | ADLOFF | 00E | H1 | $ep \rightarrow \nu^* X$ |
| | | 252 | ABBIENDI | 99F | OPAL | |
| | | 253 | ABREU | 99O | DLPH | $e^+ e^- \rightarrow \nu \nu^*$ |
| | | 254 | ACKERSTAFF | 98C | OPAL | $e^+ e^- \rightarrow \nu^* \nu^*$ Ho- modoublet type |
| | | 255 | BARATE | 98U | ALEP | $Z \rightarrow \nu \nu^*$ |
| | 256,257 | ABREU | 97B | DLPH | $e^+ e^- \rightarrow \nu \nu^*$ | |
| | | 258 | ABREU | 97I | DLPH | $\nu^* \rightarrow \ell W, \nu Z$ |
| | | 259 | ABREU | 97J | DLPH | $\nu^* \rightarrow \nu \gamma$ |
| | 256,260 | ACCIARRI | 97G | L3 | $e^+ e^- \rightarrow \nu \nu^*$ | |
| | | 261 | ACKERSTAFF | 97 | OPAL | $e^+ e^- \rightarrow \nu \nu^*$ |
| | | 262 | ADLOFF | 97 | H1 | Lepton-flavor violation |
| none 40–96 | 95 | 263 | BREITWEG | 97C | ZEUS | $ep \rightarrow \nu^* X$ |
| | | 264 | ACCIARRI | 96D | L3 | $e^+ e^- \rightarrow \nu \nu^*$ |
| | | 265 | ALEXANDER | 96Q | OPAL | $e^+ e^- \rightarrow \nu \nu^*$ |
| | | 266 | BUSKULIC | 96W | ALEP | $e^+ e^- \rightarrow \nu \nu^*$ |
| | | 267 | DERRICK | 95B | ZEUS | $ep \rightarrow \nu^* X$ |
| | | 268 | ABT | 93 | H1 | $ep \rightarrow \nu^* X$ |
| > 91 | 95 | | ADRIANI | 93M | L3 | $\lambda_Z > 1, \nu^* \rightarrow \nu \gamma$ |
| > 89 | 95 | | ADRIANI | 93M | L3 | $\lambda_Z > 1, \nu_e^* \rightarrow e W$ |
| > 87 | 95 | | ADRIANI | 93M | L3 | $\lambda_Z > 0.1, \nu^* \rightarrow \nu \gamma$ |
| > 74 | 95 | | ADRIANI | 93M | L3 | $\lambda_Z > 0.1, \nu_e^* \rightarrow e W$ |
| | | 269 | BARDADIN-... | 92 | RVUE | |
| > 91 | 95 | 270 | DECAMP | 92 | ALEP | $\lambda_Z > 1$ |
| > 74 | 95 | 270 | DECAMP | 92 | ALEP | $\lambda_Z > 0.034$ |
| > 91 | 95 | 271,272 | ADEVA | 90O | L3 | $\lambda_Z > 1$ |
| > 83 | 95 | 272 | ADEVA | 90O | L3 | $\lambda_Z > 0.1, \nu^* \rightarrow \nu \gamma$ |
| > 74 | 95 | 272 | ADEVA | 90O | L3 | $\lambda_Z > 0.1, \nu_e^* \rightarrow e W$ |
| > 90 | 95 | 273,274 | DECAMP | 90O | ALEP | $\lambda_Z > 1$ |
| > 74.7 | 95 | 273,274 | DECAMP | 90O | ALEP | $\lambda_Z > 0.06$ |

244 ACHARD 03B result is from $e^+ e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. The quoted limit is for ν_e^* . $f = -f' = \Lambda/m_{\nu^*}$ is assumed. See their Fig. 4 for the exclusion plot in the mass-coupling plane.

245 ADLOFF 02 search for single ν^* production in ep collisions with the decays $\nu^* \rightarrow \nu \gamma, \nu Z, e W$. The quoted limit assumes $f = -f' = \Lambda/m_{\nu^*}$. See their Fig. 1 for the exclusion plots in the mass-coupling plane.

246 CHEKANOV 02D search for single ν^* production in ep collisions with the decays $\nu^* \rightarrow \nu \gamma, \nu Z, e W$. $f = -f' = \Lambda/m_{\nu^*}$ is assumed for the e^* coupling. CHEKANOV 02D also obtain limit for $f = f' = \Lambda/m_{\nu^*}$: $m_{\nu^*} > 135$ GeV. See their Fig. 5c and Fig. 5d for the exclusion plot in the mass-coupling plane.

- 247 ACCIARRI 01D search for $\nu\nu^*$ production in e^+e^- collisions at $\sqrt{s} = 192\text{--}202$ GeV with decays $\nu^* \rightarrow \nu\gamma$, $\nu^* \rightarrow eW$. $f=-f'=1/m_{\nu^*}$ is assumed for the ν^* coupling. See their Fig. 4 for limits in the mass-coupling plane.
- 248 ABBIENDI 00I result is from e^+e^- collisions at $\sqrt{s}=161\text{--}183$ GeV. See their Fig. 7 for limits in mass-coupling plane.
- 249 From e^+e^- collisions at $\sqrt{s}=189$ GeV. ABBIENDI,G 00D obtain limit on $\sigma(e^+e^- \rightarrow \nu^*\nu^*)B(\nu^* \rightarrow \nu\gamma)^2$. See their Fig. 11.
- 250 ACCIARRI 00E result is from e^+e^- collisions at $\sqrt{s}=189$ GeV. See their Fig. 3 for limits in mass-coupling plane.
- 251 ADLOFF 00E search for single ν^* production in ep collisions with the decays $\nu^* \rightarrow \nu\gamma$, νZ , eW . The quoted limit assumes $f=-f'=1/m_{\nu^*}$. See their Fig. 10 for the exclusion plot in the mass-coupling plane.
- 252 From e^+e^- collisions at $\sqrt{s}=130\text{--}183$ GeV, ABBIENDI 99F obtain limit on $\sigma(e^+e^- \rightarrow \nu\nu^*) B(\nu^* \rightarrow \nu\gamma)$. See their Fig. 8.
- 253 ABREU 990 result is from e^+e^- collisions at $\sqrt{s}=183$ GeV. See their Figs. 4 and 5 for the exclusion limit in the mass-coupling plane.
- 254 ACKERSTAFF 98C from e^+e^- collisions at $\sqrt{s}=170\text{--}172$ GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.
- 255 BARATE 98U obtain limits on the $Z\nu\nu^*$ coupling. See their Fig. 13 for limits in mass-coupling plane
- 256 From e^+e^- collisions at $\sqrt{s}=161$ GeV.
- 257 See Fig. 4b and Fig. 5b of ABREU 97B for the exclusion limit in the mass-coupling plane.
- 258 ABREU 97I limit is from $Z \rightarrow \nu\nu^*$. See their Fig. 12 for the exclusion limit in the mass-coupling plane.
- 259 ABREU 97J limit is from $Z \rightarrow \nu\nu^*$. See their Fig. 5 for the exclusion limit in the mass-coupling plane.
- 260 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.
- 261 ACKERSTAFF 97 result is from e^+e^- collisions at $\sqrt{s}=161$ GeV, for homodoublet ν^* . See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 262 ADLOFF 97 search for single e^* production in ep collisions with the decays $e^* \rightarrow e\gamma$, eZ , νW . See their Fig. 4 for the rejection limits on the product of the production cross section and the branching ratio.
- 263 BREITWEG 97C search for single ν^* production in ep collisions with the decay $\nu^* \rightarrow \nu\gamma$. $f=-f'=2/m_{\nu^*}$ is assumed for the ν^* coupling. See their Fig. 10 for the exclusion plot in the mass-coupling plane.
- 264 ACCIARRI 96D result is from e^+e^- collisions at $\sqrt{s}=130\text{--}140$ GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.
- 265 ALEXANDER 96Q result is from e^+e^- collisions at $\sqrt{s}=130\text{--}140$ GeV for homodoublet ν^* . See their Fig. 3b and Fig. 3c for the exclusion limit in the mass-coupling plane.
- 266 BUSKULIC 96W result is from e^+e^- collisions at $\sqrt{s}=130\text{--}140$ GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.
- 267 DERRICK 95B search for single ν^* production via ν^*eW coupling in ep collisions with the decays $\nu^* \rightarrow \nu\gamma$, νZ , eW . See their Fig. 14 for the exclusion plot in the $m_{\nu^*}-\lambda\gamma$ plane.
- 268 ABT 93 search for single ν^* production via ν^*eW coupling in ep collisions with the decays $\nu^* \rightarrow \nu\gamma$, νZ , eW . See their Fig. 4 for exclusion plot in the $m_{\nu^*}-\lambda_W$ plane.
- 269 See Fig. 5 of BARDADIN-OTWINOWSKA 92 for combined limit of ADEVA 900, DECAMP 900, and DECAMP 92.
- 270 DECAMP 92 limit is based on $B(Z \rightarrow \nu^*\bar{\nu}) \times B(\nu^* \rightarrow \nu\gamma) < 2.7 \times 10^{-5}$ (95%CL) assuming Dirac ν^* , $B(\nu^* \rightarrow \nu\gamma) = 1$.
- 271 Limit is either for $\nu^* \rightarrow \nu\gamma$ or $\nu^* \rightarrow eW$.

272 Superseded by ADRIANI 93M.

273 DECAMP 900 limit based on $B(Z \rightarrow \nu\nu^*) \cdot B(\nu^* \rightarrow \nu\gamma) < 6 \times 10^{-5}$ (95%CL), assuming $B(\nu^* \rightarrow \nu\gamma) = 1$.

274 Superseded by DECAMP 92.

MASS LIMITS for Excited q (q^*)

Limits for Excited q (q^*) from Pair Production

These limits are obtained from $e^+ e^- \rightarrow q^* \bar{q}^*$ and thus rely only on the (electroweak) charge of the q^* . Form factor effects are ignored unless noted. Assumptions about the q^* decay are given in the comments and footnotes.

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------|----------|--|
| >45.6 | 95 | 275 ADRIANI | 93M L3 | u or d type, $Z \rightarrow q^* q^*$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| | | 276 BARATE | 98U ALEP | $Z \rightarrow q^* q^*$ |
| | | 277 ADRIANI | 92F L3 | $Z \rightarrow q^* q^*$ |
| >41.7 | 95 | 278 BARDADIN-... | 92 RVUE | u -type, $\Gamma(Z)$ |
| >44.7 | 95 | 278 BARDADIN-... | 92 RVUE | d -type, $\Gamma(Z)$ |
| >40.6 | 95 | 279 DECAMP | 92 ALEP | u -type, $\Gamma(Z)$ |
| >44.2 | 95 | 279 DECAMP | 92 ALEP | d -type, $\Gamma(Z)$ |
| >45 | 95 | 280 DECAMP | 92 ALEP | u or d type, $Z \rightarrow q^* q^*$ |
| >45 | 95 | 279 ABREU | 91F DLPH | u -type, $\Gamma(Z)$ |
| >45 | 95 | 279 ABREU | 91F DLPH | d -type, $\Gamma(Z)$ |
| >21.1 | 95 | 281 BEHREND | 86C CELL | $e(q^*) = -1/3$, $q^* \rightarrow$ qg |
| >22.3 | 95 | 281 BEHREND | 86C CELL | $e(q^*) = 2/3$, $q^* \rightarrow qg$ |
| >22.5 | 95 | 281 BEHREND | 86C CELL | $e(q^*) = -1/3$, $q^* \rightarrow$ $q\gamma$ |
| >23.2 | 95 | 281 BEHREND | 86C CELL | $e(q^*) = 2/3$, $q^* \rightarrow q\gamma$ |

275 ADRIANI 93M limit is valid for $B(q^* \rightarrow qg) > 0.25$ (0.17) for up (down) type.

276 BARATE 98U obtain limits on the form factor. See their Fig. 16 for limits in mass-form factor plane.

277 ADRIANI 92F search for $Z \rightarrow q^* \bar{q}^*$ followed with $q^* \rightarrow q\gamma$ decays and give the limit $\sigma_Z \cdot B(Z \rightarrow q^* \bar{q}^*) \cdot B^2(q^* \rightarrow q\gamma) < 2$ pb at 95%CL. Assuming five flavors of degenerate q^* of homodoublet type, $B(q^* \rightarrow q\gamma) < 4\%$ is obtained for $m_{q^*} < 45$ GeV.

278 BARDADIN-OTWINOWSKA 92 limit based on $\Delta\Gamma(Z) < 36$ MeV.

279 These limits are independent of decay modes.

280 Limit is for $B(q^* \rightarrow qg) + B(q^* \rightarrow q\gamma) = 1$.

281 BEHREND 86C search for $e^+ e^- \rightarrow q^* \bar{q}^*$ for $m_{q^*} > 5$ GeV. But $m < 5$ GeV excluded by total hadronic cross section. The limits are for point-like photon couplings of excited quarks.

Limits for Excited q (q^*) from Single Production

These limits are from $e^+e^- \rightarrow q^*\bar{q}$ or $p\bar{p} \rightarrow q^*X$ and depend on transition magnetic couplings between q and q^* . Assumptions about q^* decay mode are given in the footnotes and comments.

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|--------------|----------|---|
| >775 | 95 | 282 ABAZOV | 04C D0 | $p\bar{p} \rightarrow q^*X, q^* \rightarrow qg$ |
| none 200–520 and 580–760 | 95 | 283 ABE | 97G CDF | $p\bar{p} \rightarrow q^*X, q^* \rightarrow 2$ jets |
| none 80–570 | 95 | 284 ABE | 95N CDF | $p\bar{p} \rightarrow q^*X, q^* \rightarrow qg, q\gamma, qW$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| >205 | 95 | 285 CHEKANOV | 02D ZEUS | $ep \rightarrow q^*X$ |
| >188 | 95 | 286 ADLOFF | 00E H1 | $ep \rightarrow q^*X$ |
| | | 287 ABREU | 990 DLPH | $e^+e^- \rightarrow qq^*$ |
| | | 288 BARATE | 98U ALEP | $Z \rightarrow qq^*$ |
| | | 289 ADLOFF | 97 H1 | Lepton-flavor violation |
| none 40–169 | 95 | 290 BREITWEG | 97C ZEUS | $ep \rightarrow q^*X$ |
| | | 291 DERRICK | 95B ZEUS | $ep \rightarrow q^*X$ |
| none 80–540 | 95 | 292 ABE | 94 CDF | $p\bar{p} \rightarrow q^*X, q^* \rightarrow q\gamma, qW$ |
| > 79 | 95 | 293 ADRIANI | 93M L3 | $\lambda_Z(L3) > 0.06$ |
| >288 | 90 | 294 ALITTI | 93 UA2 | $p\bar{p} \rightarrow q^*X, q^* \rightarrow qg$ |
| | | 295 ABREU | 92D DLPH | $Z \rightarrow qq^*$ |
| | | 296 ADRIANI | 92F L3 | $Z \rightarrow qq^*$ |
| > 75 | 95 | 293 DECAMP | 92 ALEP | $Z \rightarrow qq^*, \lambda_Z > 1$ |
| > 88 | 95 | 297 DECAMP | 92 ALEP | $Z \rightarrow qq^*, \lambda_Z > 1$ |
| > 86 | 95 | 297 AKRAWY | 90J OPAL | $Z \rightarrow qq^*, \lambda_Z > 1.2$ |
| | | 298 ALBAJAR | 89 UA1 | $p\bar{p} \rightarrow q^*X, q^* \rightarrow qW$ |
| > 39 | 95 | 299 BEHREND | 86C CELL | $e^+e^- \rightarrow q^*\bar{q} (q^* \rightarrow qg, q\gamma), \lambda_\gamma=1$ |

282 ABAZOV 04C assume $f_S = f = f' = \Lambda/m_{q^*}$.

283 ABE 97G search for new particle decaying to dijets.

284 ABE 95N assume a degenerate u^* and d^* with $f_S=f=f'=\Lambda/m_{q^*}$. See their Fig. 4 for the excluded region in $m_{q^*} - f$ plane.

285 CHEKANOV 02D search for single q^* production in ep collisions with the decays $q^* \rightarrow q\gamma, qZ, qW$. $f_S = 0$ and $f = f' = \Lambda/m_{q^*}$ is assumed for the q^* coupling. See their Fig. 5b for the exclusion plot in the mass-coupling plane.

286 ADLOFF 00E search for single q^* production in ep collisions with the decays $q^* \rightarrow q\gamma, qZ, qW$. $f_S=0$ and $f=f'=\Lambda/m_{q^*}$ is assumed for the q^* coupling. See their Fig. 11 for the exclusion plot in the mass-coupling plane.

287 ABREU 990 result is from e^+e^- collisions at $\sqrt{s}=183$ GeV. See their Fig. 6 for the exclusion limit in the mass-coupling plane.

288 BARATE 98U obtain limits on the Zqq^* coupling. See their Fig. 16 for limits in mass-coupling plane

289 ADLOFF 97 search for single q^* production in ep collisions with the decay $q^* \rightarrow q\gamma$. See their Fig. 6 for the rejection limits on the product of the production cross section and the branching ratio.

- 290 BREITWEG 97C search for single q^* production in $e p$ collisions with the decays $q^* \rightarrow q\gamma, qW$. $f_S=0$, and $f=-f'=2\Lambda/m_{q^*}$ is assumed for the q^* coupling. See their Fig. 11 for the exclusion plot in the mass-coupling plane.
- 291 DERRICK 95B search for single q^* production via $q^* q\gamma$ coupling in $e p$ collisions with the decays $q^* \rightarrow qW, qZ, qg, q\gamma$. See their Fig. 15 for the exclusion plot in the $m_{q^*}-\lambda\gamma$ plane.
- 292 ABE 94 search for resonances in jet- γ and jet- W invariant mass in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The limit is for $f_S = f = f' = \Lambda/m_{q^*}$ and u^* and d^* are assumed to be degenerate. See their Fig. 4 for the excluded region in $m_{q^*}-f$ plane.
- 293 Assumes $B(q^* \rightarrow qg) = 1$.
- 294 ALITTI 93 search for resonances in the two-jet invariant mass. The limit is for $f_S = f = f' = \Lambda/m_{q^*}$. u^* and d^* are assumed to be degenerate. If not, the limit for u^* (d^*) is 277 (247) GeV if $m_{d^*} \gg m_{u^*}$ ($m_{u^*} \gg m_{d^*}$).
- 295 ABREU 92D give $\sigma(e^+e^- \rightarrow Z \rightarrow q^*\bar{q} \text{ or } q\bar{q}^*) \times B(q^* \rightarrow q\gamma) < 15$ pb (95% CL) for $m_{q^*} < 80$ GeV.
- 296 ADRIANI 92F search for $Z \rightarrow qq^*$ with $q^* \rightarrow q\gamma$ and give the limit $\sigma_Z \cdot B(Z \rightarrow qq^*) \cdot B(q^* \rightarrow q\gamma) < (2-10)$ pb (95%CL) for $m_{q^*} = (46-82)$ GeV.
- 297 Assumes $B(q^* \rightarrow q\gamma) = 0.1$.
- 298 ALBAJAR 89 give $\sigma(q^* \rightarrow W + \text{jet})/\sigma(W) < 0.019$ (90% CL) for $m_{q^*} > 220$ GeV.
- 299 BEHREND 86C has $E_{\text{cm}} = 42.5-46.8$ GeV. See their Fig. 3 for excluded region in the $m_{q^*}-(\lambda_\gamma/m_{q^*})^2$ plane. The limit is for $\lambda_\gamma = 1$ with $\eta_L = \eta_R = 1$.

MASS LIMITS for Color Sextet Quarks (q_6)

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------|-----|-------------|---------|-------------------------------------|
| >84 | 95 | 300 ABE | 89D CDF | $p\bar{p} \rightarrow q_6\bar{q}_6$ |

300 ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color sextet quark is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. A limit of 121 GeV is obtained for a color decuplet.

MASS LIMITS for Color Octet Charged Leptons (l_8)

$$\lambda \equiv m_{l_8}/\Lambda$$

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|--|
| >86 | 95 | 301 ABE | 89D CDF | Stable $l_8: p\bar{p} \rightarrow l_8\bar{l}_8$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| none 3.0-30.3 | 95 | 302 ABT | 93 H1 | $e_8: ep \rightarrow e_8 X$ |
| none 3.5-30.3 | 95 | 303 KIM | 90 AMY | $e_8: e^+e^- \rightarrow ee + \text{jets}$ |
| none 3.5-30.3 | 95 | 303 KIM | 90 AMY | $\mu_8: e^+e^- \rightarrow \mu\mu + \text{jets}$ |
| >19.8 | 95 | 304 KIM | 90 AMY | $e_8: e^+e^- \rightarrow gg; R$ |
| none 5-23.2 | 95 | 305 BARTEL | 87B JADE | $e_8, \mu_8, \tau_8: e^+e^-; R$ |
| none 5-23.2 | 95 | 305 BARTEL | 87B JADE | $\mu_8: e^+e^- \rightarrow \mu\mu + \text{jets}$ |
| | | 306 BARTEL | 85K JADE | $e_8: e^+e^- \rightarrow gg; R$ |

- 301 ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color octet lepton is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. The limit improves to 99 GeV if it always fragments into a unit-charged hadron.
- 302 ABT 93 search for e_8 production via e -gluon fusion in ep collisions with $e_8 \rightarrow eg$. See their Fig. 3 for exclusion plot in the m_{e_8} - Λ plane for $m_{e_8} = 35$ –220 GeV.
- 303 KIM 90 is at $E_{cm} = 50$ –60.8 GeV. The same assumptions as in BARTEL 87B are used.
- 304 KIM 90 result $(m_{e_8} \Lambda_M)^{1/2} > 178.4$ GeV (95%CL, $\alpha_S = 0.16$ used) is subject to the same restriction as for BARTEL 85K.
- 305 BARTEL 87B is at $E_{cm} = 46.3$ –46.78 GeV. The limits assume ℓ_8 pair production cross sections to be eight times larger than those of the corresponding heavy lepton pair production.
- 306 In BARTEL 85K, R can be affected by $e^+e^- \rightarrow gg$ via e_q exchange. Their limit $m_{e_8} > 173$ GeV (CL=95%) at $\lambda = m_{e_8}/\Lambda_M = 1$ ($\eta_L = \eta_R = 1$) is not listed above because the cross section is sensitive to the product $\eta_L \eta_R$, which should be absent in ordinary theory with electronic chiral invariance.

MASS LIMITS for Color Octet Neutrinos (ν_8)

$$\lambda \equiv m_{\ell_8}/\Lambda$$

| VALUE (GeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|--|
| >110 | 90 | 307 BARGER | 89 RVUE | $\nu_8: p\bar{p} \rightarrow \nu_8\bar{\nu}_8$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| none 3.8–29.8 | 95 | 308 KIM | 90 AMY | $\nu_8: e^+e^- \rightarrow$ acoplanar jets |
| none 9–21.9 | 95 | 309 BARTEL | 87B JADE | $\nu_8: e^+e^- \rightarrow$ acoplanar jets |

- 307 BARGER 89 used ABE 89B limit for events with large missing transverse momentum. Two-body decay $\nu_8 \rightarrow \nu g$ is assumed.
- 308 KIM 90 is at $E_{cm} = 50$ –60.8 GeV. The same assumptions as in BARTEL 87B are used.
- 309 BARTEL 87B is at $E_{cm} = 46.3$ –46.78 GeV. The limit assumes the ν_8 pair production cross section to be eight times larger than that of the corresponding heavy neutrino pair production. This assumption is not valid in general for the weak couplings, and the limit can be sensitive to its $SU(2)_L \times U(1)_Y$ quantum numbers.

MASS LIMITS for W_8 (Color Octet W Boson)

| VALUE (GeV) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|--------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| | 310 ALBAJAR | 89 UA1 | $p\bar{p} \rightarrow W_8 X,$ $W_8 \rightarrow W g$ |

- 310 ALBAJAR 89 give $\sigma(W_8 \rightarrow W + \text{jet})/\sigma(W) < 0.019$ (90% CL) for $m_{W_8} > 220$ GeV.

REFERENCES FOR Searches for Quark and Lepton Compositeness

| | | | | |
|---------------|-----|------------------------|------------------------------|----------------------|
| ACOSTA | 05B | PRL 94 101802 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 04C | PR D69 111101R | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABBIENDI | 04G | EPJ C33 173 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 04N | PL B602 167 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABDALLAH | 04N | EPJ C37 405 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| CHEKANOV | 04B | PL B591 23 | S. Chekanov <i>et al.</i> | (ZEUS Collab.) |
| ACHARD | 03B | PL B568 23 | P. Achard <i>et al.</i> | (L3 Collab.) |
| ADLOFF | 03 | PL B568 35 | C. Adloff <i>et al.</i> | (H1 Collab.) |
| BABICH | 03 | EPJ C29 103 | A.A. Babich <i>et al.</i> | |
| ABBIENDI | 02G | PL B544 57 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ACHARD | 02D | PL B531 28 | P. Achard <i>et al.</i> | (L3 Collab.) |
| ACHARD | 02J | PL B549 290 | P. Achard <i>et al.</i> | (L3 Collab.) |
| ADLOFF | 02 | PL B525 9 | C. Adloff <i>et al.</i> | (H1 Collab.) |
| ADLOFF | 02B | PL B548 35 | C. Adloff <i>et al.</i> | (H1 Collab.) |
| CHEKANOV | 02D | PL B549 32 | S. Chekanov <i>et al.</i> | (ZEUS Collab.) |
| ACCIARRI | 01D | PL B502 37 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| AFFOLDER | 01I | PRL 87 231803 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| BOURILKOV | 01 | PR D64 071701 | D. Bourilkov | |
| CHEUNG | 01B | PL B517 167 | K. Cheung | |
| ABBIENDI | 00I | EPJ C14 73 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 00R | EPJ C13 553 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI,G | 00D | EPJ C18 253 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBOTT | 00E | PR D62 031101 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABREU | 00A | PL B491 67 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 00S | PL B485 45 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 00E | PL B473 177 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 00G | PL B475 198 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 00P | PL B489 81 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ADLOFF | 00 | PL B479 358 | C. Adloff <i>et al.</i> | (H1 Collab.) |
| ADLOFF | 00E | EPJ C17 567 | C. Adloff <i>et al.</i> | (H1 Collab.) |
| AFFOLDER | 00I | PR D62 012004 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| BARATE | 00I | EPJ C12 183 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BOURILKOV | 00 | PR D62 076005 | D. Bourilkov | |
| BREITWEG | 00B | EPJ C14 239 | J. Breitweg <i>et al.</i> | (ZEUS Collab.) |
| ABBIENDI | 99 | EPJ C6 1 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 99F | EPJ C8 23 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 99P | PL B465 303 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBOTT | 99C | PRL 82 2457 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABBOTT | 99D | PRL 82 4769 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABREU | 99A | EPJ C11 383 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 99O | EPJ C8 41 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ZARNECKI | 99 | EPJ C11 539 | A.F. Zarnecki | |
| ABBOTT | 98G | PRL 80 666 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABREU | 98J | PL B433 429 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 98J | PL B433 163 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 98T | PL B439 183 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACKERSTAFF | 98 | EPJ C1 21 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 98C | EPJ C1 45 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 98V | EPJ C2 441 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKER...,K... | 98B | PL B438 379 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| BARATE | 98J | PL B429 201 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 98U | EPJ C4 571 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARGER | 98E | PR D57 391 | V. Barger <i>et al.</i> | |
| BERTRAM | 98 | PL B443 347 | I. Bertram, E.H. Simmons | |
| MCFARLAND | 98 | EPJ C1 509 | K.S. McFarland <i>et al.</i> | (CCFR/NuTeV Collab.) |
| MIURA | 98 | PR D57 5345 | M. Miura <i>et al.</i> | (VENUS Collab.) |
| ABE | 97G | PR D55 R5263 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 97T | PRL 79 2198 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 97B | PL B393 245 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 97I | ZPHY C74 57 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| Also | | ZPHY C75 580 (erratum) | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 97J | ZPHY C74 577 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 97G | PL B401 139 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 97W | PL B413 159 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACKERSTAFF | 97 | PL B391 197 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 97C | PL B391 221 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ADLOFF | 97 | NP B483 44 | C. Adloff <i>et al.</i> | (H1 Collab.) |

| | | | | |
|-------------|-----|-------------------------|-------------------------------------|-------------------|
| ARIMA | 97 | PR D55 19 | T. Arima <i>et al.</i> | (VENUS Collab.) |
| BREITWEG | 97C | ZPHY C76 631 | J. Breitweg <i>et al.</i> | (ZEUS Collab.) |
| DEANDREA | 97 | PL B409 277 | A. Deandrea | (MARS) |
| ABE | 96 | PRL 77 438 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96S | PRL 77 5336 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 96K | PL B380 480 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 96D | PL B370 211 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 96L | PL B384 323 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ALEXANDER | 96K | PL B377 222 | G. Alexander <i>et al.</i> | (OPAL Collab.) |
| ALEXANDER | 96Q | PL B386 463 | G. Alexander <i>et al.</i> | (OPAL Collab.) |
| BUSKULIC | 96W | PL B385 445 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 96Z | PL B384 333 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| ABE | 95N | PRL 74 3538 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ACCIARRI | 95G | PL B353 136 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| AID | 95 | PL B353 578 | S. Aid <i>et al.</i> | (H1 Collab.) |
| DERRICK | 95B | ZPHY C65 627 | M. Derrick <i>et al.</i> | (ZEUS Collab.) |
| ABE | 94 | PRL 72 3004 | F. Abe <i>et al.</i> | (CDF Collab.) |
| DIAZCRUZ | 94 | PR D49 R2149 | J.L. Diaz Cruz, O.A. Sampayo | (CINV) |
| VELISSARIS | 94 | PL B331 227 | C. Velissaris <i>et al.</i> | (AMY Collab.) |
| ABE | 93G | PRL 71 2542 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABT | 93 | NP B396 3 | I. Abt <i>et al.</i> | (H1 Collab.) |
| ADRIANI | 93M | PRPL 236 1 | O. Adriani <i>et al.</i> | (L3 Collab.) |
| ALITTI | 93 | NP B400 3 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| BUSKULIC | 93Q | ZPHY C59 215 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| DERRICK | 93B | PL B316 207 | M. Derrick <i>et al.</i> | (ZEUS Collab.) |
| ABE | 92B | PRL 68 1463 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 92D | PRL 68 1104 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 92M | PRL 69 2896 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 92C | ZPHY C53 41 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 92D | ZPHY C53 555 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ADRIANI | 92B | PL B288 404 | O. Adriani <i>et al.</i> | (L3 Collab.) |
| ADRIANI | 92F | PL B292 472 | O. Adriani <i>et al.</i> | (L3 Collab.) |
| BARADIN-... | 92 | ZPHY C55 163 | M. Bardadin-Otwinowska | (CLER) |
| DECAMP | 92 | PRPL 216 253 | D. Decamp <i>et al.</i> | (ALEPH Collab.) |
| HOWELL | 92 | PL B291 206 | B. Howell <i>et al.</i> | (TOPAZ Collab.) |
| KROHA | 92 | PR D46 58 | H. Kroha | (ROCH) |
| PDG | 92 | PR D45, 1 June, Part II | K. Hikasa <i>et al.</i> | (KEK, LBL, BOST+) |
| SHIMOZAWA | 92 | PL B284 144 | K. Shimozawa <i>et al.</i> | (TOPAZ Collab.) |
| ABE | 91D | PRL 67 2418 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 91E | PL B268 296 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 91F | NP B367 511 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ADACHI | 91 | PL B255 613 | I. Adachi <i>et al.</i> | (TOPAZ Collab.) |
| AKRAWY | 91F | PL B257 531 | M.Z. Akrawy <i>et al.</i> | (OPAL Collab.) |
| ALITTI | 91B | PL B257 232 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| BEHREND | 91B | ZPHY C51 143 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| BEHREND | 91C | ZPHY C51 149 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| Also | | ZPHY C51 143 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| ABE | 90I | ZPHY C48 13 | K. Abe <i>et al.</i> | (VENUS Collab.) |
| ADEVA | 90F | PL B247 177 | B. Adeva <i>et al.</i> | (L3 Collab.) |
| ADEVA | 90K | PL B250 199 | B. Adeva <i>et al.</i> | (L3 Collab.) |
| ADEVA | 90L | PL B250 205 | B. Adeva <i>et al.</i> | (L3 Collab.) |
| ADEVA | 90O | PL B252 525 | B. Adeva <i>et al.</i> | (L3 Collab.) |
| AKRAWY | 90F | PL B241 133 | M.Z. Akrawy <i>et al.</i> | (OPAL Collab.) |
| AKRAWY | 90I | PL B244 135 | M.Z. Akrawy <i>et al.</i> | (OPAL Collab.) |
| AKRAWY | 90J | PL B246 285 | M.Z. Akrawy <i>et al.</i> | (OPAL Collab.) |
| DECAMP | 90G | PL B236 501 | D. Decamp <i>et al.</i> | (ALEPH Collab.) |
| DECAMP | 90O | PL B250 172 | D. Decamp <i>et al.</i> | (ALEPH Collab.) |
| KIM | 90 | PL B240 243 | G.N. Kim <i>et al.</i> | (AMY Collab.) |
| ABE | 89 | PRL 62 613 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 89B | PRL 62 1825 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 89D | PRL 63 1447 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 89H | PRL 62 3020 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 89J | ZPHY C45 175 | K. Abe <i>et al.</i> | (VENUS Collab.) |
| ABE | 89L | PL B232 425 | K. Abe <i>et al.</i> | (VENUS Collab.) |
| ADACHI | 89B | PL B228 553 | I. Adachi <i>et al.</i> | (TOPAZ Collab.) |
| ALBAJAR | 89 | ZPHY C44 15 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| BARGER | 89 | PL B220 464 | V. Barger <i>et al.</i> | (WISC, KEK) |
| BEHREND | 89B | PL B222 163 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| BRAUNSCH... | 89C | ZPHY C43 549 | W. Braunschweig <i>et al.</i> | (TASSO Collab.) |
| DORENBOS... | 89 | ZPHY C41 567 | J. Dorenbosch <i>et al.</i> | (CHARM Collab.) |
| HAGIWARA | 89 | PL B219 369 | K. Hagiwara, M. Sakuda, N. Terunuma | (KEK, DURH+) |

| | | | | |
|-------------|-----|----------------------|-------------------------------|-------------------|
| KIM | 89 | PL B223 476 | S.K. Kim <i>et al.</i> | (AMY Collab.) |
| ABE | 88B | PL B213 400 | K. Abe <i>et al.</i> | (VENUS Collab.) |
| BARINGER | 88 | PL B206 551 | P. Baringer <i>et al.</i> | (HRS Collab.) |
| BRAUNSCH... | 88 | ZPHY C37 171 | W. Braunschweig <i>et al.</i> | (TASSO Collab.) |
| BRAUNSCH... | 88D | ZPHY C40 163 | W. Braunschweig <i>et al.</i> | (TASSO Collab.) |
| ANSARI | 87D | PL B195 613 | R. Ansari <i>et al.</i> | (UA2 Collab.) |
| BARTEL | 87B | ZPHY C36 15 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| BEHREND | 87C | PL B191 209 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| FERNANDEZ | 87B | PR D35 10 | E. Fernandez <i>et al.</i> | (MAC Collab.) |
| ARNISON | 86C | PL B172 461 | G.T.J. Arnison <i>et al.</i> | (UA1 Collab.) |
| ARNISON | 86D | PL B177 244 | G.T.J. Arnison <i>et al.</i> | (UA1 Collab.) |
| BARTEL | 86 | ZPHY C31 359 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| BARTEL | 86C | ZPHY C30 371 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| BEHREND | 86 | PL 168B 420 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| BEHREND | 86C | PL B181 178 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| DERRICK | 86 | PL 166B 463 | M. Derrick <i>et al.</i> | (HRS Collab.) |
| Also | | PR D34 3286 | M. Derrick <i>et al.</i> | (HRS Collab.) |
| DERRICK | 86B | PR D34 3286 | M. Derrick <i>et al.</i> | (HRS Collab.) |
| GRIFOLS | 86 | PL 168B 264 | J.A. Grifols, S. Peris | (BARC) |
| JODIDIO | 86 | PR D34 1967 | A. Jodidio <i>et al.</i> | (LBL, NWES, TRIU) |
| Also | | PR D37 237 (erratum) | A. Jodidio <i>et al.</i> | (LBL, NWES, TRIU) |
| APPEL | 85 | PL 160B 349 | J.A. Appel <i>et al.</i> | (UA2 Collab.) |
| BARTEL | 85K | PL 160B 337 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| BERGER | 85 | ZPHY C28 1 | C. Berger <i>et al.</i> | (PLUTO Collab.) |
| BERGER | 85B | ZPHY C27 341 | C. Berger <i>et al.</i> | (PLUTO Collab.) |
| BAGNAIA | 84C | PL 138B 430 | P. Bagnaia <i>et al.</i> | (UA2 Collab.) |
| BARTEL | 84D | PL 146B 437 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| BARTEL | 84E | PL 146B 121 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| EICHTEN | 84 | RMP 56 579 | E. Eichten <i>et al.</i> | (FNAL, LBL, OSU) |
| ALTHOFF | 83C | PL 126B 493 | M. Althoff <i>et al.</i> | (TASSO Collab.) |
| RENARD | 82 | PL 116B 264 | F.M. Renard | (CERN) |