



$$I(J^P) = 0(\frac{1}{2}^+) \text{ Status: } ****$$

The parity of the  $\Lambda_c^+$  is defined to be positive (as are the parities of the proton, neutron, and  $\Lambda$ ). The quark content is  $udc$ . The spin  $J$  has not actually been measured. Results of an analysis of  $pK^-\pi^+$  decays (JEZABEK 92) are consistent with the expected  $J = 1/2$ .

We have omitted some results that have been superseded by later experiments. The omitted results may be found in earlier editions.

### $\Lambda_c^+$ MASS

Our value in 2004,  $2284.9 \pm 0.6$  MeV, was the average of the measurements now filed below as "not used." The BABAR measurement is so much better that we use it alone. Note that it is about 2.6 (old) standard deviations above the 2004 value.

The fit also includes  $\Sigma_c^+ - \Lambda_c^+$  and  $\Lambda_c^{*+} - \Lambda_c^+$  mass-difference measurements, but this doesn't affect the  $\Lambda_c^+$  mass. The new (in 2006)  $\Lambda_c^+$  mass simply pushes all those other masses higher.


| VALUE (MeV)   | EVTS | DOCUMENT ID           | TECN     | COMMENT                                     |
|---|------|-----------------------|----------|---|
| <b>2286.46 ± 0.14 OUR FIT</b>   |      |                       |          |   |
| <b>2286.46 ± 0.14</b>   | 4891 | <sup>1</sup> AUBERT,B | 05S BABR | $\Lambda_c^0 K^+$ and $\Sigma_c^0 K^+$      |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |      |                       |          |   |
| 2284.7 ± 0.6 ± 0.7  | 1134 | AVERY                 | 91 CLEO  | Six modes                                   |
| 2281.7 ± 2.7 ± 2.6  | 29   | ALVAREZ               | 90B NA14 | $pK^-\pi^+$                                 |
| 2285.8 ± 0.6 ± 1.2  | 101  | BARLAG                | 89 NA32  | $pK^-\pi^+$                                 |
| 2284.7 ± 2.3 ± 0.5  | 5    | AGUILAR-...           | 88B LEBC | $pK^-\pi^+$                                 |
| 2283.1 ± 1.7 ± 2.0  | 628  | ALBRECHT              | 88C ARG  | $pK^-\pi^+$ , $p\bar{K}^0$ , $\Lambda 3\pi$ |
| 2286.2 ± 1.7 ± 0.7  | 97   | ANJOS                 | 88B E691 | $pK^-\pi^+$                                 |
| 2281 ± 3  | 2    | JONES                 | 87 HBC   | $pK^-\pi^+$                                 |
| 2283 ± 3  | 3    | BOSETTI               | 82 HBC   | $pK^-\pi^+$                                 |
| 2290 ± 3  | 1    | CALICCHIO             | 80 HYBR  | $pK^-\pi^+$                                 |

<sup>1</sup>AUBERT,B 05S uses low-Q  $\Lambda_c^0 K^+$  and  $\Sigma_c^0 K^+$  decays to minimize systematic errors. The error above includes systematic as well as statistical errors. Many cross checks and adjustments to properties of the BABAR detector, as well as the large number of clean events, make this by far the best measurement of the  $\Lambda_c^+$  mass.

## $\Lambda_c^+$ MEAN LIFE

Measurements with an error  $\geq 100 \times 10^{-15}$  s or with fewer than 20 events have been omitted from the Listings.

| VALUE ( $10^{-15}$ s)   | EVTS               | DOCUMENT ID                         | TECN | COMMENT   |
|---|--------------------|-------------------------------------|------|---|
| <b>200 ± 6</b>  | <b>OUR AVERAGE</b> | Error includes scale factor of 1.6. |      | See the ideogram below.                                     |
| 204.6 ± 3.4 ± 2.5   | 8034               | LINK                                | 02C  | FOCS $pK^- \pi^+$   |
| 198.1 ± 7.0 ± 5.6   | 1630               | KUSHNIR...                          | 01   | SELX $\Lambda_c^+ \rightarrow pK^- \pi^+$                   |
| 179.6 ± 6.9 ± 4.4   | 4749               | MAHMOOD                             | 01   | CLE2 $e^+ e^- \approx \Upsilon(4S)$                         |
| 215 ± 16 ± 8  | 1340               | FRABETTI                            | 93D  | E687 $\gamma \text{Be}, \Lambda_c^+ \rightarrow pK^- \pi^+$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                    |                                     |      |   |
| 180 ± 30 ± 30   | 29                 | ALVAREZ                             | 90   | NA14 $\gamma, \Lambda_c^+ \rightarrow pK^- \pi^+$           |
| 200 ± 30 ± 30   | 90                 | FRABETTI                            | 90   | E687 $\gamma \text{Be}, \Lambda_c^+ \rightarrow pK^- \pi^+$ |
| 196 $^{+23}_{-20}$  | 101                | BARLAG                              | 89   | NA32 $pK^- \pi^+ + \text{c.c.}$                             |
| 220 ± 30 ± 20   | 97                 | ANJOS                               | 88B  | E691 $pK^- \pi^+ + \text{c.c.}$                             |



## $\Lambda_c^+$ DECAY MODES

Nearly all branching fractions of the  $\Lambda_c^+$  are measured relative to the  $pK^- \pi^+$  mode, but there are no model-independent measurements of this branching fraction. We explain how we arrive at our value of  $B(\Lambda_c^+ \rightarrow pK^- \pi^+)$  in a Note at the beginning of the branching-ratio measurements,

below. When this branching fraction is eventually well determined, all the other branching fractions will slide up or down proportionally as the true value differs from the value we use here.

| Mode  | Fraction ( $\Gamma_i/\Gamma$ )                                   | Scale factor/<br>Confidence level      |        |
|---|--|--|--------|
| <b>Hadronic modes with a <math>p</math>: <math>S = -1</math> final states</b> |  |  |        |
| $\Gamma_1$  | $p\bar{K}^0$   | ( 2.3 $\pm$ 0.6 ) %                    |        |
| $\Gamma_2$  | $pK^-\pi^+$  | [a] ( 5.0 $\pm$ 1.3 ) %                |        |
| $\Gamma_3$  | $p\bar{K}^*(892)^0$  | [b] ( 1.6 $\pm$ 0.5 ) %                |        |
| $\Gamma_4$  | $\Delta(1232)^{++}K^-$   | ( 8.6 $\pm$ 3.0 ) $\times 10^{-3}$     |        |
| $\Gamma_5$  | $\Lambda(1520)\pi^+$   | [b] ( 1.8 $\pm$ 0.6 ) %                |        |
| $\Gamma_6$  | $pK^-\pi^+$ nonresonant  | ( 2.8 $\pm$ 0.8 ) %                    |        |
| $\Gamma_7$  | $p\bar{K}^0\pi^0$  | ( 3.3 $\pm$ 1.0 ) %                    |        |
| $\Gamma_8$  | $p\bar{K}^0\eta$   | ( 1.2 $\pm$ 0.4 ) %                    |        |
| $\Gamma_9$  | $p\bar{K}^0\pi^+\pi^-$   | ( 2.6 $\pm$ 0.7 ) %                    |        |
| $\Gamma_{10}$   | $pK^-\pi^+\pi^0$   | ( 3.4 $\pm$ 1.0 ) %                    |        |
| $\Gamma_{11}$   | $pK^*(892)^-\pi^+$   | [b] ( 1.1 $\pm$ 0.5 ) %                |        |
| $\Gamma_{12}$   | $p(K^-\pi^+)_{\text{nonresonant}}\pi^0$                          | ( 3.6 $\pm$ 1.2 ) %                    |        |
| $\Gamma_{13}$   | $\Delta(1232)\bar{K}^*(892)$                                     | seen                                   |        |
| $\Gamma_{14}$   | $pK^-\pi^+\pi^+\pi^-$  | ( 1.1 $\pm$ 0.8 ) $\times 10^{-3}$     |        |
| $\Gamma_{15}$   | $pK^-\pi^+\pi^0\pi^0$  | ( 8 $\pm$ 4 ) $\times 10^{-3}$         |        |
| $\Gamma_{16}$   | $pK^-\pi^+3\pi^0$  |  |        |
| <b>Hadronic modes with a <math>p</math>: <math>S = 0</math> final states</b>  |  |  |        |
| $\Gamma_{17}$   | $p\pi^+\pi^-$  | ( 3.5 $\pm$ 2.0 ) $\times 10^{-3}$     |        |
| $\Gamma_{18}$   | $pf_0(980)$  | [b] ( 2.8 $\pm$ 1.9 ) $\times 10^{-3}$ |        |
| $\Gamma_{19}$   | $p\pi^+\pi^+\pi^-\pi^-$  | ( 1.8 $\pm$ 1.2 ) $\times 10^{-3}$     |        |
| $\Gamma_{20}$   | $pK^+K^-$  | ( 7.7 $\pm$ 3.5 ) $\times 10^{-4}$     |        |
| $\Gamma_{21}$   | $p\phi$  | [b] ( 8.2 $\pm$ 2.7 ) $\times 10^{-4}$ |        |
| $\Gamma_{22}$   | $pK^+K^-$ non- $\phi$  | ( 3.5 $\pm$ 1.7 ) $\times 10^{-4}$     |        |
| <b>Hadronic modes with a hyperon: <math>S = -1</math> final states</b>        |  |  |        |
| $\Gamma_{23}$   | $\Lambda\pi^+$   | ( 1.01 $\pm$ 0.28 ) %                  |        |
| $\Gamma_{24}$   | $\Lambda\pi^+\pi^0$  | ( 3.6 $\pm$ 1.3 ) %                    |        |
| $\Gamma_{25}$   | $\Lambda\rho^+$  | < 5 %                                  | CL=95% |
| $\Gamma_{26}$   | $\Lambda\pi^+\pi^+\pi^-$   | ( 2.6 $\pm$ 0.7 ) %                    |        |
| $\Gamma_{27}$   | $\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+$ | ( 7 $\pm$ 4 ) $\times 10^{-3}$         |        |
| $\Gamma_{28}$   | $\Sigma(1385)^-\pi^+\pi^+, \Sigma^{*-} \rightarrow \Lambda\pi^-$ | ( 5.5 $\pm$ 1.7 ) $\times 10^{-3}$     |        |
| $\Gamma_{29}$   | $\Lambda\pi^+\rho^0$   | ( 1.1 $\pm$ 0.5 ) %                    |        |
| $\Gamma_{30}$   | $\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$     | ( 3.7 $\pm$ 3.1 ) $\times 10^{-3}$     |        |
| $\Gamma_{31}$   | $\Lambda\pi^+\pi^+\pi^-$ nonresonant                             | < 8 $\times 10^{-3}$                   | CL=90% |
| $\Gamma_{32}$   | $\Lambda\pi^+\pi^+\pi^-\pi^0$ total                              | ( 1.8 $\pm$ 0.8 ) %                    |        |
| $\Gamma_{33}$   | $\Lambda\pi^+\eta$   | [b] ( 1.8 $\pm$ 0.6 ) %                |        |

|               |  |     |                                  |        |
|---------------|--|-----|----------------------------------|--------|
| $\Gamma_{34}$ | $\Sigma(1385)^+ \eta$  | [b] | $( 8.5 \pm 3.3 ) \times 10^{-3}$ |        |
| $\Gamma_{35}$ | $\Lambda \pi^+ \omega$                                       | [b] | $( 1.2 \pm 0.5 ) \%$             |        |
| $\Gamma_{36}$ | $\Lambda \pi^+ \pi^+ \pi^- \pi^0$ , no $\eta$ or $\omega$    |     | $< 7 \times 10^{-3}$             | CL=90% |
| $\Gamma_{37}$ | $\Lambda K^+ \bar{K}^0$                                      |     | $( 6.5 \pm 2.0 ) \times 10^{-3}$ |        |
| $\Gamma_{38}$ | $\Xi(1690)^0 K^+$ , $\Xi^{*0} \rightarrow \Lambda \bar{K}^0$ |     | $( 1.9 \pm 0.7 ) \times 10^{-3}$ |        |
| $\Gamma_{39}$ | $\Sigma^0 \pi^+$   |     | $( 1.04 \pm 0.31 ) \%$           |        |
| $\Gamma_{40}$ | $\Sigma^+ \pi^0$   |     | $( 1.00 \pm 0.34 ) \%$           |        |
| $\Gamma_{41}$ | $\Sigma^+ \eta$  |     | $( 5.5 \pm 2.3 ) \times 10^{-3}$ |        |
| $\Gamma_{42}$ | $\Sigma^+ \pi^+ \pi^-$                                       |     | $( 3.6 \pm 1.0 ) \%$             |        |
| $\Gamma_{43}$ | $\Sigma^+ \rho^0$  |     | $< 1.4 \%$                       | CL=95% |
| $\Gamma_{44}$ | $\Sigma^- \pi^+ \pi^+$                                       |     | $( 1.9 \pm 0.8 ) \%$             |        |
| $\Gamma_{45}$ | $\Sigma^0 \pi^+ \pi^0$                                       |     | $( 1.8 \pm 0.8 ) \%$             |        |
| $\Gamma_{46}$ | $\Sigma^0 \pi^+ \pi^+ \pi^-$                                 |     | $( 8.3 \pm 3.1 ) \times 10^{-3}$ |        |
| $\Gamma_{47}$ | $\Sigma^+ \pi^+ \pi^- \pi^0$                                 |     | —                                |        |
| $\Gamma_{48}$ | $\Sigma^+ \omega$  | [b] | $( 2.7 \pm 1.0 ) \%$             |        |
| $\Gamma_{49}$ | $\Sigma^+ K^+ K^-$   |     | $( 2.8 \pm 0.8 ) \times 10^{-3}$ |        |
| $\Gamma_{50}$ | $\Sigma^+ \phi$  | [b] | $( 3.2 \pm 1.0 ) \times 10^{-3}$ |        |
| $\Gamma_{51}$ | $\Xi(1690)^0 K^+$ , $\Xi^{*0} \rightarrow \Sigma^+ K^-$      |     | $( 8.2 \pm 3.1 ) \times 10^{-4}$ |        |
| $\Gamma_{52}$ | $\Sigma^+ K^+ K^-$ nonresonant                               |     | $< 7 \times 10^{-4}$             | CL=90% |
| $\Gamma_{53}$ | $\Xi^0 K^+$  |     | $( 3.9 \pm 1.4 ) \times 10^{-3}$ |        |
| $\Gamma_{54}$ | $\Xi^- K^+ \pi^+$  |     | $( 4.9 \pm 1.7 ) \times 10^{-3}$ |        |
| $\Gamma_{55}$ | $\Xi(1530)^0 K^+$  | [b] | $( 2.6 \pm 1.0 ) \times 10^{-3}$ |        |

#### Hadronic modes with a hyperon: $S = 0$ final states

|               |                       |     |                                  |        |
|---------------|-----------------------|-----|----------------------------------|--------|
| $\Gamma_{56}$ | $\Lambda K^+$         |     | $( 7.5 \pm 2.6 ) \times 10^{-4}$ |        |
| $\Gamma_{57}$ | $\Sigma^0 K^+$        |     | $( 5.8 \pm 2.4 ) \times 10^{-4}$ |        |
| $\Gamma_{58}$ | $\Sigma^+ K^+ \pi^-$  |     | $( 1.7 \pm 0.7 ) \times 10^{-3}$ |        |
| $\Gamma_{59}$ | $\Sigma^+ K^*(892)^0$ | [b] | $( 2.8 \pm 1.1 ) \times 10^{-3}$ |        |
| $\Gamma_{60}$ | $\Sigma^- K^+ \pi^+$  |     | $< 1.0 \times 10^{-3}$           | CL=90% |

#### Doubly Cabibbo-suppressed modes

|               |               |  |                        |        |
|---------------|---------------|--|------------------------|--------|
| $\Gamma_{61}$ | $p K^+ \pi^-$ |  | $< 2.3 \times 10^{-4}$ | CL=90% |
|---------------|---------------|--|------------------------|--------|

#### Semileptonic modes

|               |                           |     |                      |  |
|---------------|---------------------------|-----|----------------------|--|
| $\Gamma_{62}$ | $\Lambda \ell^+ \nu_\ell$ | [c] | $( 2.0 \pm 0.6 ) \%$ |  |
| $\Gamma_{63}$ | $\Lambda e^+ \nu_e$       |     | $( 2.1 \pm 0.6 ) \%$ |  |
| $\Gamma_{64}$ | $\Lambda \mu^+ \nu_\mu$   |     | $( 2.0 \pm 0.7 ) \%$ |  |

#### Inclusive modes

|               |                              |  |                      |  |
|---------------|------------------------------|--|----------------------|--|
| $\Gamma_{65}$ | $e^+$ anything               |  | $( 4.5 \pm 1.7 ) \%$ |  |
| $\Gamma_{66}$ | $p e^+$ anything             |  | $( 1.8 \pm 0.9 ) \%$ |  |
| $\Gamma_{67}$ | $\Lambda e^+$ anything       |  |                      |  |
| $\Gamma_{68}$ | $p$ anything                 |  | $( 50 \pm 16 ) \%$   |  |
| $\Gamma_{69}$ | $p$ anything (no $\Lambda$ ) |  | $( 12 \pm 19 ) \%$   |  |

|               |                              |                 |  |       |  |
|---------------|------------------------------|-----------------|--|-------|--|
| $\Gamma_{70}$ | $p$ hadrons                  |                 |  |       |  |
| $\Gamma_{71}$ | $n$ anything                 | (50 ± 16 ) %    |  |       |  |
| $\Gamma_{72}$ | $n$ anything (no $\Lambda$ ) | (29 ± 17 ) %    |  |       |  |
| $\Gamma_{73}$ | $\Lambda$ anything           | (35 ± 11 ) %    |  | S=1.4 |  |
| $\Gamma_{74}$ | $\Sigma^\pm$ anything        | [d] (10 ± 5 ) % |  |       |  |
| $\Gamma_{75}$ | 3prongs                      | (24 ± 8 ) %     |  |       |  |

**$\Delta C = 1$  weak neutral current (C1) modes, or  
Lepton number (L) violating modes**

|               |                      |      |       |                  |        |
|---------------|----------------------|------|-------|------------------|--------|
| $\Gamma_{76}$ | $p\mu^+\mu^-$        | $CI$ | < 3.4 | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{77}$ | $\Sigma^-\mu^+\mu^+$ | $L$  | < 7.0 | $\times 10^{-4}$ | CL=90% |

[a] See the note on “ $\Lambda_c^+$  Branching Fractions” below.

[b] This branching fraction includes all the decay modes of the final-state resonance.

[c] An  $\ell$  indicates an  $e$  or a  $\mu$  mode, not a sum over these modes.

[d] The value is for the sum of the charge states or particle/antiparticle states indicated.

**CONSTRAINED FIT INFORMATION**

An overall fit to 12 branching ratios uses 22 measurements and one constraint to determine 9 parameters. The overall fit has a  $\chi^2 = 6.6$  for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

|          |       |          |          |          |          |          |          |
|----------|-------|----------|----------|----------|----------|----------|----------|
| $x_{23}$ | 94    |          |          |          |          |          |          |
| $x_{26}$ | 97    | 92       |          |          |          |          |          |
| $x_{39}$ | 87    | 87       | 85       |          |          |          |          |
| $x_{42}$ | 91    | 86       | 89       | 80       |          |          |          |
| $x_{46}$ | 69    | 65       | 70       | 61       | 63       |          |          |
| $x_{49}$ | 87    | 82       | 85       | 76       | 93       | 60       |          |
| $x_{50}$ | 84    | 79       | 81       | 73       | 90       | 58       | 84       |
|          | $x_2$ | $x_{23}$ | $x_{26}$ | $x_{39}$ | $x_{42}$ | $x_{46}$ | $x_{49}$ |

A REVIEW GOES HERE – Check our WWW List of Reviews

## $\Lambda_c^+$ BRANCHING RATIOS

### ————— Hadronic modes with a $p$ : $S = -1$ final states —————

#### $\Gamma(p\bar{K}^0)/\Gamma(pK^-\pi^+)$ $\Gamma_1/\Gamma_2$

| <u>VALUE</u>                 | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                |
|------------------------------|-------------|--------------------|-------------|-------------------------------|
| <b>0.47±0.04 OUR AVERAGE</b> |             |                    |             |                               |
| 0.46±0.02±0.04               | 1025        | ALAM               | 98 CLE2     | $e^+e^- \approx \Upsilon(4S)$ |
| 0.44±0.07±0.05               | 133         | AVERY              | 91 CLEO     | $e^+e^-$ 10.5 GeV             |
| 0.55±0.17±0.14               | 45          | ANJOS              | 90 E691     | $\gamma$ Be 70–260 GeV        |
| 0.62±0.15±0.03               | 73          | ALBRECHT           | 88C ARG     | $e^+e^-$ 10 GeV               |

#### $\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma$

See the note on " $\Lambda_c^+$  Branching Fractions" above.

| <u>VALUE</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u>      | <u>TECN</u> | <u>COMMENT</u>                |
|---|-------------|-------------------------|-------------|-------------------------------|
| <b>0.050±0.013 OUR FIT</b>  |             |                         |             |                               |
| <b>0.050±0.013</b>  |             | PDG                     | 02          | See note at top of ratios     |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |             |                         |             |                               |
| 0.050±0.005±0.012   | 1205        | <sup>2</sup> JAFFE      | 00 CLE2     | $e^+e^-$ 10.52–10.58 GeV      |
| 0.041±0.010   |             | <sup>3,4</sup> ALBRECHT | 920 ARG     | $e^+e^- \approx \Upsilon(4S)$ |
| 0.044±0.012   |             | <sup>3,5</sup> CRAWFORD | 92 CLEO     | $e^+e^-$ 10.5 GeV             |

<sup>2</sup> JAFFE 00 assumes that a  $\bar{D}$  meson and an antiproton in opposite hemispheres tags for a  $\Lambda_c^+$  in the hemisphere of the  $\bar{p}$ . The fraction of such  $\bar{D}\bar{p}$  events with a  $\Lambda_c^+ \rightarrow pK^-\pi^+$  decay then gives the  $pK^-\pi^+$  branching fraction. See the paper for assumptions, caveats, etc.

<sup>3</sup> To extract  $\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$ , we use  $B(\bar{B} \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (0.28 \pm 0.06)\%$ , which is the average of measurements from ARGUS (ALBRECHT 88C) and CLEO (CRAWFORD 92).

<sup>4</sup> ALBRECHT 920 measures  $B(\bar{B} \rightarrow \Lambda_c^+ X) = (6.8 \pm 0.5 \pm 0.3)\%$ .

<sup>5</sup> CRAWFORD 92 measures  $B(\bar{B} \rightarrow \Lambda_c^+ X) = (6.4 \pm 0.8 \pm 0.8)\%$ .

#### $\Gamma(p\bar{K}^*(892)^0)/\Gamma(pK^-\pi^+)$ $\Gamma_3/\Gamma_2$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included.

| <u>VALUE</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u>                     |
|---|-------------|---------------------|-------------|------------------------------------|
| <b>0.31±0.04 OUR AVERAGE</b>  |             |                     |             |                                    |
| 0.29±0.04±0.03  |             | <sup>6</sup> AITALA | 00 E791     | $\pi^- N$ , 500 GeV                |
| 0.35 <sup>+0.06</sup> <sub>-0.07</sub> ±0.03                                  | 39          | BOZEK               | 93 NA32     | $\pi^-$ Cu 230 GeV                 |
| 0.42±0.24   | 12          | BASILE              | 81B CNTR    | $pp \rightarrow \Lambda_c^+ e^- X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |             |                     |             |                                    |
| 0.35±0.11   |             | BARLAG              | 90D NA32    | See BOZEK 93                       |

<sup>6</sup> AITALA 00 makes a coherent 5-dimensional amplitude analysis of  $946 \pm 38 \Lambda_c^+ \rightarrow pK^-\pi^+$  decays.

$\Gamma(\Delta(1232)^{++} K^-)/\Gamma(\rho K^- \pi^+)$   $\Gamma_4/\Gamma_2$

| <u>VALUE</u>                                 | <u>EVTS</u> | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u>                      |
|--|-------------|---------------------|-------------|-------------------------------------|
| <b>0.17±0.04 OUR AVERAGE</b>                 |             |                     |             | Error includes scale factor of 1.1. |
| 0.18±0.03±0.03                               |             | <sup>7</sup> AITALA | 00 E791     | $\pi^- N$ , 500 GeV                 |
| 0.12 <sup>+0.04</sup> <sub>-0.05</sub> ±0.05 | 14          | BOZEK               | 93 NA32     | $\pi^- Cu$ 230 GeV                  |
| 0.40±0.17                                    | 17          | BASILE              | 81B CNTR    | $pp \rightarrow \Lambda_c^+ e^- X$  |

<sup>7</sup> AITALA 00 makes a coherent 5-dimensional amplitude analysis of  $946 \pm 38 \Lambda_c^+ \rightarrow \rho K^- \pi^+$  decays.

$\Gamma(\Lambda(1520)\pi^+)/\Gamma(\rho K^- \pi^+)$   $\Gamma_5/\Gamma_2$

Unseen decay modes of the  $\Lambda(1520)$  are included.

| <u>VALUE</u>                                 | <u>EVTS</u> | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u>      |
|--|-------------|---------------------|-------------|---------------------|
| <b>0.35±0.08 OUR AVERAGE</b>                 |             |                     |             |                     |
| 0.34±0.08±0.05                               |             | <sup>8</sup> AITALA | 00 E791     | $\pi^- N$ , 500 GeV |
| 0.40 <sup>+0.18</sup> <sub>-0.13</sub> ±0.09 | 12          | BOZEK               | 93 NA32     | $\pi^- Cu$ 230 GeV  |

<sup>8</sup> AITALA 00 makes a coherent 5-dimensional amplitude analysis of  $946 \pm 38 \Lambda_c^+ \rightarrow \rho K^- \pi^+$  decays.

$\Gamma(\rho K^- \pi^+ \text{nonresonant})/\Gamma(\rho K^- \pi^+)$   $\Gamma_6/\Gamma_2$

| <u>VALUE</u>                                 | <u>EVTS</u> | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u>      |
|--|-------------|---------------------|-------------|---------------------|
| <b>0.55±0.06 OUR AVERAGE</b>                 |             |                     |             |                     |
| 0.55±0.06±0.04                               |             | <sup>9</sup> AITALA | 00 E791     | $\pi^- N$ , 500 GeV |
| 0.56 <sup>+0.07</sup> <sub>-0.09</sub> ±0.05 | 71          | BOZEK               | 93 NA32     | $\pi^- Cu$ 230 GeV  |

<sup>9</sup> AITALA 00 makes a coherent 5-dimensional amplitude analysis of  $946 \pm 38 \Lambda_c^+ \rightarrow \rho K^- \pi^+$  decays.

$\Gamma(\rho \bar{K}^0 \pi^0)/\Gamma(\rho K^- \pi^+)$   $\Gamma_7/\Gamma_2$

| <u>VALUE</u>          | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                 |
|-----------------------|-------------|--------------------|-------------|--------------------------------|
| <b>0.66±0.05±0.07</b> | 774         | ALAM               | 98 CLE2     | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\rho \bar{K}^0 \eta)/\Gamma(\rho K^- \pi^+)$   $\Gamma_8/\Gamma_2$

Unseen decay modes of the  $\eta$  are included.

| <u>VALUE</u>          | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                 |
|-----------------------|-------------|--------------------|-------------|--------------------------------|
| <b>0.25±0.04±0.04</b> | 57          | AMMAR              | 95 CLE2     | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\rho \bar{K}^0 \pi^+ \pi^-)/\Gamma(\rho K^- \pi^+)$   $\Gamma_9/\Gamma_2$

| <u>VALUE</u>                 | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                 |
|------------------------------|-------------|--------------------|-------------|--------------------------------|
| <b>0.51±0.06 OUR AVERAGE</b> |             |                    |             |                                |
| 0.52±0.04±0.05               | 985         | ALAM               | 98 CLE2     | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.43±0.12±0.04               | 83          | AVERY              | 91 CLEO     | $e^+ e^-$ 10.5 GeV             |
| 0.98±0.36±0.08               | 12          | BARLAG             | 90D NA32    | $\pi^-$ 230 GeV                |

$\Gamma(\rho K^- \pi^+ \pi^0)/\Gamma(\rho K^- \pi^+)$   $\Gamma_{10}/\Gamma_2$

| <u>VALUE</u>          | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                 |
|-----------------------|-------------|--------------------|-------------|--------------------------------|
| <b>0.67±0.04±0.11</b> | 2606        | ALAM               | 98 CLE2     | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\rho K^*(892)^- \pi^+) / \Gamma(\rho \bar{K}^0 \pi^+ \pi^-)$   $\Gamma_{11} / \Gamma_9$

Unseen decay modes of the  $K^*(892)^-$  are included.

| VALUE              | EVTS | DOCUMENT ID | TECN | COMMENT        |
|--------------------|------|-------------|------|----------------|
| <b>0.44 ± 0.14</b> | 17   | ALEEV 94    | BIS2 | $nN$ 20–70 GeV |

$\Gamma(\rho(K^- \pi^+)_{\text{nonresonant}} \pi^0) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{12} / \Gamma_2$

| VALUE                     | EVTS | DOCUMENT ID | TECN | COMMENT            |
|---------------------------|------|-------------|------|--------------------|
| <b>0.73 ± 0.12 ± 0.05</b> | 67   | BOZEK 93    | NA32 | $\pi^-$ Cu 230 GeV |

$\Gamma(\Delta(1232) \bar{K}^*(892)) / \Gamma_{\text{total}}$   $\Gamma_{13} / \Gamma$

| VALUE       | EVTS | DOCUMENT ID  | TECN | COMMENT        |
|-------------|------|--------------|------|----------------|
| <b>seen</b> | 35   | AMENDOLIA 87 | SPEC | $\gamma$ Ge-Si |

$\Gamma(\rho K^- \pi^+ \pi^+ \pi^-) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{14} / \Gamma_2$

| VALUE                | DOCUMENT ID | TECN | COMMENT         |
|----------------------|-------------|------|-----------------|
| <b>0.022 ± 0.015</b> | BARLAG 90D  | NA32 | $\pi^-$ 230 GeV |

$\Gamma(\rho K^- \pi^+ \pi^0 \pi^0) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{15} / \Gamma_2$

| VALUE                     | EVTS | DOCUMENT ID | TECN | COMMENT            |
|---------------------------|------|-------------|------|--------------------|
| <b>0.16 ± 0.07 ± 0.03</b> | 15   | BOZEK 93    | NA32 | $\pi^-$ Cu 230 GeV |

$\Gamma(\rho K^- \pi^+ 3\pi^0) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{16} / \Gamma_2$

| VALUE                     | EVTS | DOCUMENT ID | TECN | COMMENT            |
|---------------------------|------|-------------|------|--------------------|
| <b>0.10 ± 0.06 ± 0.02</b> | 8    | BOZEK 93    | NA32 | $\pi^-$ Cu 230 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

————— **Hadronic modes with a  $\rho$ :  $S = 0$  final states** —————

$\Gamma(\rho \pi^+ \pi^-) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{17} / \Gamma_2$

| VALUE                | DOCUMENT ID | TECN | COMMENT         |
|----------------------|-------------|------|-----------------|
| <b>0.069 ± 0.036</b> | BARLAG 90D  | NA32 | $\pi^-$ 230 GeV |

$\Gamma(\rho f_0(980)) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{18} / \Gamma_2$

Unseen decay modes of the  $f_0(980)$  are included.

| VALUE                | DOCUMENT ID | TECN | COMMENT         |
|----------------------|-------------|------|-----------------|
| <b>0.055 ± 0.036</b> | BARLAG 90D  | NA32 | $\pi^-$ 230 GeV |

$\Gamma(\rho \pi^+ \pi^+ \pi^- \pi^-) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{19} / \Gamma_2$

| VALUE                | DOCUMENT ID | TECN | COMMENT         |
|----------------------|-------------|------|-----------------|
| <b>0.036 ± 0.023</b> | BARLAG 90D  | NA32 | $\pi^-$ 230 GeV |

$\Gamma(\rho K^+ K^-) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{20} / \Gamma_2$

| VALUE   | EVTS | DOCUMENT ID   | TECN | COMMENT                               |
|---|------|---------------|------|---------------------------------------|
| <b>0.015 ± 0.006 OUR AVERAGE</b>  |      |               |      | Error includes scale factor of 2.1.   |
| 0.014 ± 0.002 ± 0.002   | 676  | ABE 02C       | BELL | $e^+ e^- \approx \Upsilon(4S)$        |
| 0.039 ± 0.009 ± 0.007   | 214  | ALEXANDER 96C | CLE2 | $e^+ e^- \approx \Upsilon(4S)$        |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |               |      |                                       |
| 0.096 ± 0.029 ± 0.010   | 30   | FRABETTI 93H  | E687 | $\gamma$ Be, $\bar{E}_\gamma$ 220 GeV |
| 0.048 ± 0.027   |      | BARLAG 90D    | NA32 | $\pi^-$ 230 GeV                       |



$\Gamma(p\phi)/\Gamma(pK^-\pi^+)$   $\Gamma_{21}/\Gamma_2$

Unseen decay modes of the  $\phi$  are included.

| VALUE   | EVTS | DOCUMENT ID | TECN | COMMENT                             |
|---|------|-------------|------|-------------------------------------|
| <b>0.0164 ± 0.0032 OUR AVERAGE</b>  |      |             |      | Error includes scale factor of 1.2. |
| 0.015 ± 0.002 ± 0.002   | 345  | ABE         | 02C  | BELL $e^+e^- \approx \Upsilon(4S)$  |
| 0.024 ± 0.006 ± 0.003   | 54   | ALEXANDER   | 96C  | CLE2 $e^+e^- \approx \Upsilon(4S)$  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |             |      |                                     |
| 0.040 ± 0.027   |      | BARLAG      | 90D  | NA32 $\pi^-$ 230 GeV                |

$\Gamma(pK^+K^-\text{non-}\phi)/\Gamma(pK^-\pi^+)$   $\Gamma_{22}/\Gamma_2$

| VALUE                        | EVTS | DOCUMENT ID | TECN | COMMENT                            |
|------------------------------|------|-------------|------|------------------------------------|
| <b>0.007 ± 0.002 ± 0.002</b> | 344  | ABE         | 02C  | BELL $e^+e^- \approx \Upsilon(4S)$ |

————— Hadronic modes with a hyperon:  $S = -1$  final states —————

$\Gamma(\Lambda\pi^+)/\Gamma(pK^-\pi^+)$   $\Gamma_{23}/\Gamma_2$

| VALUE   | CL% | EVTS | DOCUMENT ID | TECN | COMMENT   |
|---|-----|------|-------------|------|---|
| <b>0.202 ± 0.018 OUR FIT</b>  |     |      |             |      |   |
| <b>0.204 ± 0.019 OUR AVERAGE</b>  |     |      |             |      |   |
| 0.217 ± 0.013 ± 0.020   |     | 750  | LINK        | 05F  | FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.18 ± 0.03 ± 0.04  |     |      | ALBRECHT    | 92   | ARG $e^+e^- \approx 10.4$ GeV                           |
| 0.18 ± 0.03 ± 0.03  |     | 87   | AVERY       | 91   | CLEO $e^+e^-$ 10.5 GeV                                  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |      |             |      |   |
| <0.33   |     | 90   | ANJOS       | 90   | E691 $\gamma$ Be 70–260 GeV                             |
| <0.16   |     | 90   | ALBRECHT    | 88C  | ARG $e^+e^-$ 10 GeV                                     |

$\Gamma(\Lambda\pi^+\pi^0)/\Gamma(pK^-\pi^+)$   $\Gamma_{24}/\Gamma_2$

| VALUE                     | EVTS | DOCUMENT ID | TECN | COMMENT  |
|---------------------------|------|-------------|------|--|
| <b>0.73 ± 0.09 ± 0.16</b> | 464  | AVERY       | 94   | CLE2 $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$ |

$\Gamma(\Lambda\rho^+)/\Gamma(pK^-\pi^+)$   $\Gamma_{25}/\Gamma_2$

| VALUE           | CL% | DOCUMENT ID | TECN | COMMENT  |
|-----------------|-----|-------------|------|--|
| <b>&lt;0.95</b> | 95  | AVERY       | 94   | CLE2 $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$ |

$\Gamma(\Lambda\pi^+\pi^+\pi^-)/\Gamma(pK^-\pi^+)$   $\Gamma_{26}/\Gamma_2$

| VALUE                            | EVTS | DOCUMENT ID | TECN | COMMENT   |
|----------------------------------|------|-------------|------|---|
| <b>0.525 ± 0.032 OUR FIT</b>     |      |             |      |   |
| <b>0.522 ± 0.032 OUR AVERAGE</b> |      |             |      |   |
| 0.508 ± 0.024 ± 0.024            | 1356 | LINK        | 05F  | FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.65 ± 0.11 ± 0.12               | 289  | AVERY       | 91   | CLEO $e^+e^-$ 10.5 GeV                                  |
| 0.82 ± 0.29 ± 0.27               | 44   | ANJOS       | 90   | E691 $\gamma$ Be 70–260 GeV                             |
| 0.94 ± 0.41 ± 0.13               | 10   | BARLAG      | 90D  | NA32 $\pi^-$ 230 GeV                                    |
| 0.61 ± 0.16 ± 0.04               | 105  | ALBRECHT    | 88C  | ARG $e^+e^-$ 10 GeV                                     |

$\Gamma(\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$   $\Gamma_{27}/\Gamma_{26}$

| VALUE                     | DOCUMENT ID | TECN | COMMENT   |
|---------------------------|-------------|------|---|
| <b>0.28 ± 0.10 ± 0.08</b> | LINK        | 05F  | FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma(1385)^- \pi^+ \pi^+, \Sigma^{*-} \rightarrow \Lambda \pi^-) / \Gamma(\Lambda \pi^+ \pi^+ \pi^-)$   $\Gamma_{28} / \Gamma_{26}$

| <u>VALUE</u>          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|-----------------------|--------------------|-------------|---|
| <b>0.21±0.03±0.02</b> | LINK               | 05F         | FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Lambda \pi^+ \rho^0) / \Gamma(\Lambda \pi^+ \pi^+ \pi^-)$   $\Gamma_{29} / \Gamma_{26}$

| <u>VALUE</u>          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|-----------------------|--------------------|-------------|---|
| <b>0.40±0.12±0.12</b> | LINK               | 05F         | FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma(1385)^+ \rho^0, \Sigma^{*+} \rightarrow \Lambda \pi^+) / \Gamma(\Lambda \pi^+ \pi^+ \pi^-)$   $\Gamma_{30} / \Gamma_{26}$

| <u>VALUE</u>          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|-----------------------|--------------------|-------------|---|
| <b>0.14±0.09±0.07</b> | LINK               | 05F         | FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Lambda \pi^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(\Lambda \pi^+ \pi^+ \pi^-)$   $\Gamma_{31} / \Gamma_{26}$

| <u>VALUE</u>   | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|----------------|------------|--------------------|-------------|---|
| <b>&lt;0.3</b> | 90         | LINK               | 05F         | FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\rho \bar{K}^0 \pi^+ \pi^-) / \Gamma(\Lambda \pi^+ \pi^+ \pi^-)$   $\Gamma_9 / \Gamma_{26}$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|----------------|
|--------------|-------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|         |     |       |    |      |                       |
|---------|-----|-------|----|------|-----------------------|
| 2.6±1.2 |     | ALEEV | 96 | SPEC | $n$ nucleus, 50 GeV/c |
| 4.3±1.2 | 130 | ALEEV | 84 | BIS2 | $n$ C 40–70 GeV       |

$\Gamma(\Lambda \pi^+ \pi^+ \pi^- \pi^0 \text{ total}) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{32} / \Gamma_2$

| <u>VALUE</u>          | <u>EVTS</u> | <u>DOCUMENT ID</u>           | <u>TECN</u> | <u>COMMENT</u>                 |
|-----------------------|-------------|------------------------------|-------------|--------------------------------|
| <b>0.36±0.09±0.09</b> | 50          | <sup>10</sup> CRONIN-HEN..03 | CLE3        | $e^+ e^- \approx \Upsilon(4S)$ |

<sup>10</sup> CRONIN-HENNESSY 03 finds this channel to be dominately  $\Lambda \eta \pi^+$  and  $\Lambda \omega \pi^+$ ; see below.

$\Gamma(\Lambda \pi^+ \eta) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{33} / \Gamma_2$

Unseen decay modes of the  $\eta$  are included.

| <u>VALUE</u>                 | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                      |
|------------------------------|-------------|--------------------|-------------|-------------------------------------|
| <b>0.36±0.07 OUR AVERAGE</b> |             |                    |             |                                     |
| 0.41±0.17±0.10               | 11          | CRONIN-HEN..03     | CLE3        | $e^+ e^- \approx \Upsilon(4S)$      |
| 0.35±0.05±0.06               | 116         | AMMAR              | 95          | CLE2 $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma(1385)^+ \eta) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{34} / \Gamma_2$

Unseen decay modes of the  $\Sigma(1385)^+$  and  $\eta$  are included.

| <u>VALUE</u>          | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                      |
|-----------------------|-------------|--------------------|-------------|-------------------------------------|
| <b>0.17±0.04±0.03</b> | 54          | AMMAR              | 95          | CLE2 $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda \pi^+ \omega) / \Gamma(\rho K^- \pi^+)$   $\Gamma_{35} / \Gamma_2$

Unseen decay modes of the  $\omega$  are included.

| <u>VALUE</u>          | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                 |
|-----------------------|-------------|--------------------|-------------|--------------------------------|
| <b>0.24±0.06±0.06</b> | 32          | CRONIN-HEN..03     | CLE3        | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda\pi^+\pi^+\pi^-\pi^0, \text{no } \eta \text{ or } \omega)/\Gamma(\rho K^-\pi^+)$   $\Gamma_{36}/\Gamma_2$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                |
|--------------|------------|--------------------|-------------|-------------------------------|
| <0.13        | 90         | CRONIN-HEN..03     | CLE3        | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda K^+\bar{K}^0)/\Gamma(\rho K^-\pi^+)$   $\Gamma_{37}/\Gamma_2$

| <u>VALUE</u>                   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                     |
|--------------------------------|-------------|--------------------|-------------|--|
| <b>0.131±0.020 OUR AVERAGE</b> |             |                    |             |  |
| 0.142±0.018±0.022              | 251         | LINK               | 05F FOCS    | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.12 ±0.02 ±0.02               | 59          | AMMAR              | 95 CLE2     | $e^+e^- \approx \Upsilon(4S)$                      |

$\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0)/\Gamma(\Lambda K^+\bar{K}^0)$   $\Gamma_{38}/\Gamma_{37}$

| <u>VALUE</u>                 | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                     |
|------------------------------|-------------|--------------------|-------------|--|
| <b>0.28±0.07 OUR AVERAGE</b> |             |                    |             |  |
| 0.32±0.10±0.04               | 84±24       | LINK               | 05F FOCS    | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.26±0.08±0.03               | 93          | ABE                | 02C BELL    | $e^+e^- \approx \Upsilon(4S)$                      |

$\Gamma(\Sigma^0\pi^+)/\Gamma(\rho K^-\pi^+)$   $\Gamma_{39}/\Gamma_2$

| <u>VALUE</u>                  | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                              |
|-------------------------------|-------------|--------------------|-------------|---|
| <b>0.208±0.030 OUR FIT</b>    |             |                    |             |   |
| <b>0.20 ±0.04 OUR AVERAGE</b> |             |                    |             |   |
| 0.21 ±0.02 ±0.04              | 196         | AVERY              | 94 CLE2     | $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$ |
| 0.17 ±0.06 ±0.04              |             | ALBRECHT           | 92 ARG      | $e^+e^- \approx 10.4$ GeV                   |

$\Gamma(\Sigma^0\pi^+)/\Gamma(\Lambda\pi^+)$   $\Gamma_{39}/\Gamma_{23}$

| <u>VALUE</u>             | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                     |
|--------------------------|-------------|--------------------|-------------|--|
| <b>1.03±0.15 OUR FIT</b> |             |                    |             |  |
| <b>1.09±0.11±0.19</b>    |             |                    |             |  |
|                          | 750         | LINK               | 05F FOCS    | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma^+\pi^0)/\Gamma(\rho K^-\pi^+)$   $\Gamma_{40}/\Gamma_2$

| <u>VALUE</u>          | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                |
|-----------------------|-------------|--------------------|-------------|-------------------------------|
| <b>0.20±0.03±0.03</b> |             |                    |             |                               |
|                       | 93          | KUBOTA             | 93 CLE2     | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+\eta)/\Gamma(\rho K^-\pi^+)$   $\Gamma_{41}/\Gamma_2$

Unseen decay modes of the  $\eta$  are included.

| <u>VALUE</u>          | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                |
|-----------------------|-------------|--------------------|-------------|-------------------------------|
| <b>0.11±0.03±0.02</b> |             |                    |             |                               |
|                       | 26          | AMMAR              | 95 CLE2     | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+\pi^+\pi^-)/\Gamma(\rho K^-\pi^+)$   $\Gamma_{42}/\Gamma_2$

| <u>VALUE</u>                           | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                |
|--|-------------|--------------------|-------------|-------------------------------|
| <b>0.73±0.08 OUR FIT</b>               |             |                    |             |                               |
| <b>0.68±0.09 OUR AVERAGE</b>           |             |                    |             |                               |
| 0.74±0.07±0.09                         | 487         | KUBOTA             | 93 CLE2     | $e^+e^- \approx \Upsilon(4S)$ |
| 0.54 <sup>+0.18</sup> <sub>-0.15</sub> | 11          | BARLAG             | 92 NA32     | $\pi^-$ Cu 230 GeV            |

$\Gamma(\Sigma^+\rho^0)/\Gamma(\rho K^-\pi^+)$   $\Gamma_{43}/\Gamma_2$

| <u>VALUE</u>    | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                |
|-----------------|------------|--------------------|-------------|-------------------------------|
| <b>&lt;0.27</b> |            |                    |             |                               |
|                 | 95         | KUBOTA             | 93 CLE2     | $e^+e^- \approx \Upsilon(4S)$ |

| $\Gamma(\Sigma^- \pi^+ \pi^+)/\Gamma(\Sigma^+ \pi^+ \pi^-)$            |             |                    | $\Gamma_{44}/\Gamma_{42}$ |  |  |
|--|-------------|--------------------|---------------------------|--|--|
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.53±0.15±0.07</b>  | 56          | FRABETTI           | 94E E687                  | $\gamma$ Be, $\bar{E}_\gamma$ 220 GeV              |  |
| $\Gamma(\Sigma^0 \pi^+ \pi^0)/\Gamma(pK^- \pi^+)$                      |             |                    | $\Gamma_{45}/\Gamma_2$    |  |  |
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.36±0.09±0.10</b>  | 117         | AVERY              | 94 CLE2                   | $e^+ e^- \approx \Upsilon(3S), \Upsilon(4S)$       |  |
| $\Gamma(\Sigma^0 \pi^+ \pi^+ \pi^-)/\Gamma(pK^- \pi^+)$                |             |                    | $\Gamma_{46}/\Gamma_2$    |  |  |
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.17±0.04 OUR FIT</b>   |             |                    |                           |  |  |
| <b>0.21±0.05±0.05</b>  | 90          | AVERY              | 94 CLE2                   | $e^+ e^- \approx \Upsilon(3S), \Upsilon(4S)$       |  |
| $\Gamma(\Sigma^0 \pi^+ \pi^+ \pi^-)/\Gamma(\Lambda \pi^+ \pi^+ \pi^-)$ |             |                    | $\Gamma_{46}/\Gamma_{26}$ |  |  |
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.31±0.08 OUR FIT</b>   |             |                    |                           |  |  |
| <b>0.26±0.06±0.09</b>  | 480         | LINK               | 05F FOCS                  | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |  |
| $\Gamma(\Sigma^+ \omega)/\Gamma(pK^- \pi^+)$                           |             |                    | $\Gamma_{48}/\Gamma_2$    |  |  |
| Unseen decay modes of the $\omega$ are included.                       |             |                    |                           |  |  |
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.54±0.13±0.06</b>  | 107         | KUBOTA             | 93 CLE2                   | $e^+ e^- \approx \Upsilon(4S)$                     |  |
| $\Gamma(\Sigma^+ K^+ K^-)/\Gamma(pK^- \pi^+)$                          |             |                    | $\Gamma_{49}/\Gamma_2$    |  |  |
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.057±0.008 OUR FIT</b>   |             |                    |                           |  |  |
| <b>0.070±0.011±0.011</b>   | 59          | AVERY              | 93 CLE2                   | $e^+ e^- \approx 10.5$ GeV                         |  |
| $\Gamma(\Sigma^+ K^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$                |             |                    | $\Gamma_{49}/\Gamma_{42}$ |  |  |
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.078±0.009 OUR FIT</b>   |             |                    |                           |  |  |
| <b>0.074±0.009 OUR AVERAGE</b>   |             |                    |                           |  |  |
| 0.076±0.007±0.009  | 246         | ABE                | 02C BELL                  | $e^+ e^- \approx \Upsilon(4S)$                     |  |
| 0.071±0.011±0.011  | 103         | LINK               | 02G FOCS                  | $\gamma$ nucleus, $\approx 180$ GeV                |  |
| $\Gamma(\Sigma^+ \phi)/\Gamma(pK^- \pi^+)$                             |             |                    | $\Gamma_{50}/\Gamma_2$    |  |  |
| Unseen decay modes of the $\phi$ are included.                         |             |                    |                           |  |  |
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.063±0.011 OUR FIT</b>   |             |                    |                           |  |  |
| <b>0.069±0.023±0.016</b>   | 26          | AVERY              | 93 CLE2                   | $e^+ e^- \approx 10.5$ GeV                         |  |
| $\Gamma(\Sigma^+ \phi)/\Gamma(\Sigma^+ \pi^+ \pi^-)$                   |             |                    | $\Gamma_{50}/\Gamma_{42}$ |  |  |
| Unseen decay modes of the $\phi$ are included.                         |             |                    |                           |  |  |
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>               | <u>COMMENT</u>                                     |  |
| <b>0.087±0.012 OUR FIT</b>   |             |                    |                           |  |  |
| <b>0.086±0.012 OUR AVERAGE</b>   |             |                    |                           |  |  |
| 0.085±0.012±0.012  | 129         | ABE                | 02C BELL                  | $e^+ e^- \approx \Upsilon(4S)$                     |  |
| 0.087±0.016±0.006  | 57          | LINK               | 02G FOCS                  | $\gamma$ nucleus, $\approx 180$ GeV                |  |

$\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Sigma^+ K^-) / \Gamma(\Sigma^+ \pi^+ \pi^-)$   $\Gamma_{51} / \Gamma_{42}$

| <u>VALUE</u>                     | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                           |
|----------------------------------|-------------|--------------------|-------------|--|
| <b>0.023 ± 0.005 OUR AVERAGE</b> |             |                    |             |  |
| 0.023 ± 0.005 ± 0.005            | 75          | ABE                | 02C         | BELL $e^+ e^- \approx \Upsilon(4S)$      |
| 0.022 ± 0.006 ± 0.006            | 34          | LINK               | 02G         | FOCS $\gamma$ nucleus, $\approx 180$ GeV |

$\Gamma(\Sigma^+ K^+ K^- \text{ nonresonant}) / \Gamma(\Sigma^+ \pi^+ \pi^-)$   $\Gamma_{52} / \Gamma_{42}$


| <u>VALUE</u>  | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                           |
|---|------------|--------------------|-------------|--|
| <b>&lt;0.018</b>  | 90         | ABE                | 02C         | BELL $e^+ e^- \approx \Upsilon(4S)$      |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |            |                    |             |  |
| <0.028  | 90         | LINK               | 02G         | FOCS $\gamma$ nucleus, $\approx 180$ GeV |

$\Gamma(\Xi^0 K^+) / \Gamma(p K^- \pi^+)$   $\Gamma_{53} / \Gamma_2$

| <u>VALUE</u>                 | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                  |
|------------------------------|-------------|--------------------|-------------|---------------------------------|
| <b>0.078 ± 0.013 ± 0.013</b> | 56          | AVERY              | 93          | CLE2 $e^+ e^- \approx 10.5$ GeV |

$\Gamma(\Xi^- K^+ \pi^+) / \Gamma(p K^- \pi^+)$   $\Gamma_{54} / \Gamma_2$

| <u>VALUE</u>                     | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|----------------------------------|-------------|--------------------|-------------|---|
| <b>0.098 ± 0.021 OUR AVERAGE</b> |             |                    |             | Error includes scale factor of 1.3. See the ideogram below. |
| 0.14 ± 0.03 ± 0.02               | 34          | ALBRECHT           | 95B         | ARG $e^+ e^- \approx 10.4$ GeV                              |
| 0.079 ± 0.013 ± 0.014            | 60          | AVERY              | 93          | CLE2 $e^+ e^- \approx 10.5$ GeV                             |
| 0.15 ± 0.04 ± 0.03               | 30          | AVERY              | 91          | CLEO $e^+ e^- 10.5$ GeV                                     |



$\Gamma(\Xi(1530)^0 K^+)/\Gamma(p K^- \pi^+)$   $\Gamma_{55}/\Gamma_2$

Unseen decay modes of the  $\Xi(1530)^0$  are included.

| <u>VALUE</u>                   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>             |
|--------------------------------|-------------|--------------------|-------------|----------------------------|
| <b>0.052±0.014 OUR AVERAGE</b> |             |                    |             |                            |
| 0.05 ±0.02 ±0.01               | 11          | ALBRECHT           | 95B ARG     | $e^+ e^- \approx 10.4$ GeV |
| 0.053±0.016±0.010              | 24          | AVERY              | 93 CLE2     | $e^+ e^- \approx 10.5$ GeV |

————— **Hadronic modes with a hyperon: S = 0 final states** —————

$\Gamma(\Lambda K^+)/\Gamma(\Lambda \pi^+)$   $\Gamma_{56}/\Gamma_{23}$

| <u>VALUE</u>             | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>               |
|--------------------------|-------------|--------------------|-------------|------------------------------|
| <b>0.074±0.010±0.012</b> | 265         | ABE                | 02C BELL    | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Sigma^0 K^+)/\Gamma(\Sigma^0 \pi^+)$   $\Gamma_{57}/\Gamma_{39}$

| <u>VALUE</u>             | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>               |
|--------------------------|-------------|--------------------|-------------|------------------------------|
| <b>0.056±0.014±0.008</b> | 75          | ABE                | 02C BELL    | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Sigma^+ K^+ \pi^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$   $\Gamma_{58}/\Gamma_{42}$

| <u>VALUE</u>             | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>               |
|--------------------------|-------------|--------------------|-------------|------------------------------|
| <b>0.047±0.011±0.008</b> | 105         | ABE                | 02C BELL    | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Sigma^+ K^*(892)^0)/\Gamma(\Sigma^+ \pi^+ \pi^-)$   $\Gamma_{59}/\Gamma_{42}$

Unseen decay modes of the  $K^*(892)^0$  are included.

| <u>VALUE</u>             | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                      |
|--------------------------|-------------|--------------------|-------------|-------------------------------------|
| <b>0.078±0.018±0.013</b> | 49          | LINK               | 02G FOCS    | $\gamma$ nucleus, $\approx 180$ GeV |

$\Gamma(\Sigma^- K^+ \pi^+)/\Gamma(\Sigma^+ K^*(892)^0)$   $\Gamma_{60}/\Gamma_{59}$

| <u>VALUE</u>    | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                      |
|-----------------|------------|--------------------|-------------|-------------------------------------|
| <b>&lt;0.35</b> | 90         | LINK               | 02G FOCS    | $\gamma$ nucleus, $\approx 180$ GeV |

————— **Doubly Cabibbo-suppressed modes** —————

$\Gamma(p K^+ \pi^-)/\Gamma(p K^- \pi^+)$   $\Gamma_{61}/\Gamma_2$

| <u>VALUE</u>      | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                   |
|-------------------|------------|--------------------|-------------|----------------------------------|
| <b>&lt;0.0046</b> | 90         | LINK               | 05K FOCS    | $R = (0.05 \pm 0.26 \pm 0.02)\%$ |

————— **Semileptonic modes** —————

$\Gamma(\Lambda \ell^+ \nu_\ell)/\Gamma(p K^- \pi^+)$   $\Gamma_{62}/\Gamma_2$

We average here the averages of the next two data blocks.

| <u>VALUE</u>                 | <u>DOCUMENT ID</u> | <u>COMMENT</u>   |
|------------------------------|--------------------|--|
| <b>0.41±0.05 OUR AVERAGE</b> |                    |  |
| 0.42±0.07                    | PDG                | 02 Our $\Gamma(\Lambda e^+ \nu_e)/\Gamma(p K^- \pi^+)$     |
| 0.39±0.08                    | PDG                | 02 Our $\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma(p K^- \pi^+)$ |

### $\Gamma(\Lambda e^+ \nu_e)/\Gamma(p K^- \pi^+)$

$\Gamma_{63}/\Gamma_2$

| <u>VALUE</u>   | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                    |
|--|--------------------|-------------|-----------------------------------|
| <b>0.42±0.07 OUR AVERAGE</b>   |                    |             |                                   |
| 0.43±0.08  | 11,12 BERGFELD 94  | CLE2        | $e^+ e^- \approx \mathcal{Y}(4S)$ |
| 0.38±0.14  | 12,13 ALBRECHT 91G | ARG         | $e^+ e^- \approx 10.4$ GeV        |
| <sup>11</sup> BERGFELD 94 measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.87 \pm 0.28 \pm 0.69)$ pb.   |                    |             |                                   |
| <sup>12</sup> To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e)/\Gamma(\Lambda_c^+ \rightarrow p K^- \pi^+)$ , we use $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (11.2 \pm 1.3)$ pb, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91). |                    |             |                                   |
| <sup>13</sup> ALBRECHT 91G measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.20 \pm 1.28 \pm 0.71)$ pb.  |                    |             |                                   |

### $\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma(p K^- \pi^+)$

$\Gamma_{64}/\Gamma_2$

| <u>VALUE</u>   | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                    |
|--|--------------------|-------------|-----------------------------------|
| <b>0.39±0.08 OUR AVERAGE</b>   |                    |             |                                   |
| 0.40±0.09  | 14,15 BERGFELD 94  | CLE2        | $e^+ e^- \approx \mathcal{Y}(4S)$ |
| 0.35±0.20  | 15,16 ALBRECHT 91G | ARG         | $e^+ e^- \approx 10.4$ GeV        |
| <sup>14</sup> BERGFELD 94 measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu) = (4.43 \pm 0.51 \pm 0.64)$ pb.   |                    |             |                                   |
| <sup>15</sup> To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu)/\Gamma(\Lambda_c^+ \rightarrow p K^- \pi^+)$ , we use $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (11.2 \pm 1.3)$ pb, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91). |                    |             |                                   |
| <sup>16</sup> ALBRECHT 91G measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu) = (3.91 \pm 2.02 \pm 0.90)$ pb.  |                    |             |                                   |

### ———— Inclusive modes ————

### $\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$

$\Gamma_{65}/\Gamma$

| <u>VALUE</u>       | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|--------------------|--------------------|-------------|-----------------------|
| <b>0.045±0.017</b> | VELLA 82           | MRK2        | $e^+ e^-$ 4.5–6.8 GeV |

### $\Gamma(p e^+ \text{ anything})/\Gamma_{\text{total}}$

$\Gamma_{66}/\Gamma$

| <u>VALUE</u>       | <u>DOCUMENT ID</u>     | <u>TECN</u> | <u>COMMENT</u>        |
|--------------------|------------------------|-------------|-----------------------|
| <b>0.018±0.009</b> | <sup>17</sup> VELLA 82 | MRK2        | $e^+ e^-$ 4.5–6.8 GeV |

<sup>17</sup>VELLA 82 includes protons from  $\Lambda$  decay.

### $\Gamma(\Lambda e^+ \text{ anything})/\Gamma_{\text{total}}$

$\Gamma_{67}/\Gamma$

| <u>VALUE</u>  | <u>DOCUMENT ID</u>     | <u>TECN</u> | <u>COMMENT</u>        |
|---|------------------------|-------------|-----------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                        |             |                       |
| 0.011±0.008   | <sup>18</sup> VELLA 82 | MRK2        | $e^+ e^-$ 4.5–6.8 GeV |

<sup>18</sup>VELLA 82 includes  $\Lambda$ 's from  $\Sigma^0$  decay.

### $\Gamma(p \text{ anything})/\Gamma_{\text{total}}$

$\Gamma_{68}/\Gamma$

| <u>VALUE</u>          | <u>DOCUMENT ID</u>        | <u>TECN</u> | <u>COMMENT</u>     |
|-----------------------|---------------------------|-------------|--------------------|
| <b>0.50±0.08±0.14</b> | <sup>19</sup> CRAWFORD 92 | CLEO        | $e^+ e^-$ 10.5 GeV |

<sup>19</sup>This CRAWFORD 92 value includes protons from  $\Lambda$  decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(p \text{ anything (no } \Lambda))/\Gamma_{\text{total}}$

$\Gamma_{69}/\Gamma$

| VALUE                 | DOCUMENT ID | TECN | COMMENT           |
|-----------------------|-------------|------|-------------------|
| <b>0.12±0.10±0.16</b> | CRAWFORD 92 | CLEO | $e^+e^-$ 10.5 GeV |

$\Gamma(n \text{ anything})/\Gamma_{\text{total}}$

$\Gamma_{71}/\Gamma$

| VALUE                 | DOCUMENT ID               | TECN | COMMENT           |
|-----------------------|---------------------------|------|-------------------|
| <b>0.50±0.08±0.14</b> | <sup>20</sup> CRAWFORD 92 | CLEO | $e^+e^-$ 10.5 GeV |

<sup>20</sup>This CRAWFORD 92 value includes neutrons from  $\Lambda$  decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(n \text{ anything (no } \Lambda))/\Gamma_{\text{total}}$

$\Gamma_{72}/\Gamma$

| VALUE                 | DOCUMENT ID | TECN | COMMENT           |
|-----------------------|-------------|------|-------------------|
| <b>0.29±0.09±0.15</b> | CRAWFORD 92 | CLEO | $e^+e^-$ 10.5 GeV |

$\Gamma(p \text{ hadrons})/\Gamma_{\text{total}}$

$\Gamma_{70}/\Gamma$

| VALUE     | DOCUMENT ID  | TECN | COMMENT                |
|-----------|--------------|------|------------------------|
| 0.41±0.24 | ADAMOVICH 87 | EMUL | $\gamma A$ 20–70 GeV/c |


• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\Lambda \text{ anything})/\Gamma_{\text{total}}$

$\Gamma_{73}/\Gamma$

| VALUE                        | EVTS | DOCUMENT ID          | TECN | COMMENT   |
|------------------------------|------|----------------------|------|---|
| <b>0.35±0.11 OUR AVERAGE</b> |      |                      |      | Error includes scale factor of 1.4. See the ideogram below. |
| 0.59±0.10±0.12               |      | CRAWFORD 92          | CLEO | $e^+e^-$ 10.5 GeV   |
| 0.49±0.24                    |      | ADAMOVICH 87         | EMUL | $\gamma A$ 20–70 GeV/c                                      |
| 0.23±0.10                    | 8    | <sup>21</sup> ABE 86 | HYBR | 20 GeV $\gamma p$   |

<sup>21</sup>ABE 86 includes  $\Lambda$ 's from  $\Sigma^0$  decay.





| $\Gamma(\Sigma^\pm \text{ anything})/\Gamma_{\text{total}}$ |      |             |      |         | $\Gamma_{74}/\Gamma$ |
|---|------|-------------|------|---------|----------------------|
| VALUE   | EVTS | DOCUMENT ID | TECN | COMMENT |                      |
| <b>0.1±0.05</b>   | 5    | ABE         | 86   | HYBR    | 20 GeV $\gamma p$    |

| $\Gamma(3\text{prongs})/\Gamma_{\text{total}}$ |      |                |      |                                      | $\Gamma_{75}/\Gamma$ |
|--|------|----------------|------|--------------------------------------|----------------------|
| VALUE  | EVTS | DOCUMENT ID    | TECN | COMMENT                              |                      |
| <b>0.24±0.07±0.04</b>                          |      | KAYIS-TOPAK.03 | CHRS | $\nu_\mu$ emulsion, $\bar{E}=27$ GeV |                      |

————— Rare or forbidden modes —————

| $\Gamma(p\mu^+\mu^-)/\Gamma_{\text{total}}$   |     |      |             |      | $\Gamma_{76}/\Gamma$          |
|---|-----|------|-------------|------|-------------------------------|
| A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions. |     |      |             |      |                               |
| VALUE   | CL% | EVTS | DOCUMENT ID | TECN | COMMENT                       |
| <b>&lt;3.4 × 10<sup>-4</sup></b>  | 90  | 0    | KODAMA      | 95   | E653 $\pi^-$ emulsion 600 GeV |

| $\Gamma(\Sigma^-\mu^+\mu^+)/\Gamma_{\text{total}}$ |     |      |             |      | $\Gamma_{77}/\Gamma$          |
|--|-----|------|-------------|------|-------------------------------|
| A test of lepton-number conservation.              |     |      |             |      |                               |
| VALUE  | CL% | EVTS | DOCUMENT ID | TECN | COMMENT                       |
| <b>&lt;7.0 × 10<sup>-4</sup></b>                   | 90  | 0    | KODAMA      | 95   | E653 $\pi^-$ emulsion 600 GeV |

## $\Lambda_c^+$ DECAY PARAMETERS

See the note on "Baryon Decay Parameters" in the neutron Listings.

### $\alpha$ FOR $\Lambda_c^+ \rightarrow \Lambda\pi^+$

| VALUE                         | EVTS | DOCUMENT ID          | TECN | COMMENT  |
|-------------------------------|------|----------------------|------|--|
| <b>-0.91±0.15 OUR AVERAGE</b> |      |                      |      |  |
| -0.78±0.16±0.19               |      | LINK                 | 06A  | FOCS $\gamma A$ , $\bar{E}_\gamma \approx 180$ GeV |
| -0.94±0.21±0.12               | 414  | <sup>22</sup> BISHAI | 95   | CLE2 $e^+e^- \approx \mathcal{T}(4S)$              |
| -0.96±0.42                    |      | ALBRECHT             | 92   | ARG $e^+e^- \approx 10.4$ GeV                      |
| -1.1 ±0.4                     | 86   | AVERY                | 90B  | CLEO $e^+e^- \approx 10.6$ GeV                     |

<sup>22</sup>BISHAI 95 actually gives  $\alpha = -0.94^{+0.21+0.12}_{-0.06-0.06}$ , chopping the errors at the physical limit  $-1.0$ . However, for  $\alpha \approx -1.0$ , some experiments should *get* unphysical values ( $\alpha < -1.0$ ), and for averaging with other measurements such values (or errors that extend below  $-1.0$ ) should *not* be chopped.

### $\alpha$ FOR $\Lambda_c^+ \rightarrow \Sigma^+\pi^0$

| VALUE                  | EVTS | DOCUMENT ID | TECN | COMMENT                               |
|------------------------|------|-------------|------|---------------------------------------|
| <b>-0.45±0.31±0.06</b> | 89   | BISHAI      | 95   | CLE2 $e^+e^- \approx \mathcal{T}(4S)$ |

### $\alpha$ FOR $\Lambda_c^+ \rightarrow \Lambda\ell^+\nu_\ell$

The experiments don't cover the complete (or same incomplete)  $M(\Lambda\ell^+)$  range, but we average them together anyway.

| VALUE                         | EVTS | DOCUMENT ID            | TECN | COMMENT                               |
|-------------------------------|------|------------------------|------|---------------------------------------|
| <b>-0.86±0.04 OUR AVERAGE</b> |      |                        |      |                                       |
| -0.86±0.03±0.02               | 3201 | <sup>23</sup> HINSON   | 05   | CLEO $e^+e^- \approx \mathcal{T}(4S)$ |
| -0.91±0.42±0.25               |      | <sup>24</sup> ALBRECHT | 94B  | ARG $e^+e^- \approx 10$ GeV           |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                   |     |                        |    |      |                 |
|-----------------------------------|-----|------------------------|----|------|-----------------|
| $-0.82^{+0.09+0.06}_{-0.06-0.03}$ | 700 | <sup>25</sup> CRAWFORD | 95 | CLE2 | See HINSON 05   |
| $-0.89^{+0.17+0.09}_{-0.11-0.05}$ | 350 | <sup>26</sup> BERGFELD | 94 | CLE2 | See CRAWFORD 95 |

<sup>23</sup> HINSON 05 measures the form-factor ratio  $R \equiv f_2/f_1$  for  $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$  events to be  $-0.31 \pm 0.05 \pm 0.04$  and the pole mass to be  $2.21 \pm 0.08 \pm 0.14$  GeV/ $c^2$ , and from these calculates  $\alpha$ , averaged over  $q^2$ , where  $\langle q^2 \rangle = 0.67$  (GeV/ $c$ )<sup>2</sup>.

<sup>24</sup> ALBRECHT 94B uses  $\Lambda e^+$  and  $\Lambda \mu^+$  events in the mass range  $1.85 < M(\Lambda \ell^+) < 2.20$  GeV.

<sup>25</sup> CRAWFORD 95 measures the form-factor ratio  $R \equiv f_2/f_1$  for  $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$  events to be  $-0.25 \pm 0.14 \pm 0.08$  and from this calculates  $\alpha$ , averaged over  $q^2$ , to be the above.

<sup>26</sup> BERGFELD 94 uses  $\Lambda e^+$  events.

## $\Lambda_c^+, \bar{\Lambda}_c^-$ CP-VIOLATING DECAY ASYMMETRIES

### $(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda \pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} \pi^-$

This is zero if CP is conserved.

| VALUE                                       | DOCUMENT ID | TECN | COMMENT   |
|---|-------------|------|---|
| <b><math>-0.07 \pm 0.19 \pm 0.24</math></b> | LINK        | 06A  | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

### $(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e$

This is zero if CP is conserved.

| VALUE                                      | DOCUMENT ID | TECN | COMMENT                             |
|--|-------------|------|-------------------------------------|
| <b><math>0.00 \pm 0.03 \pm 0.02</math></b> | HINSON      | 05   | CLEO $e^+ e^- \approx \Upsilon(4S)$ |

## $\Lambda_c^+$ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1992 edition (Physical Review **D45**, 1 June, Part II) or in earlier editions.

|                |     |                |                                  |                             |
|----------------|-----|----------------|----------------------------------|-----------------------------|
| LINK           | 06A | PL B634 165    | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)        |
| AUBERT,B       | 05S | PR D72 052006  | B. Aubert <i>et al.</i>          | (BABAR Collab.)             |
| HINSON         | 05  | PRL 94 191801  | J.W. Hinson <i>et al.</i>        | (CLEO Collab.)              |
| LINK           | 05F | PL B624 22     | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)        |
| LINK           | 05K | PL B624 166    | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)        |
| CRONIN-HEN...  | 03  | PR D67 012001  | D. Cronin-Hennessy <i>et al.</i> | (CLEO Collab.)              |
| KAYIS-TOPAK... | 03  | PL B555 156    | A. Kayis-Topaksu <i>et al.</i>   | (CERN CHORUS Collab.)       |
| ABE            | 02C | PL B524 33     | K. Abe <i>et al.</i>             | (KEK BELLE Collab.)         |
| LINK           | 02C | PRL 88 161801  | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)        |
| LINK           | 02G | PL B540 25     | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)        |
| PDG            | 02  | PR D66 010001  | K. Hagiwara <i>et al.</i>        |                             |
| KUSHNIR...     | 01  | PRL 86 5243    | A. Kushnirenko <i>et al.</i>     | (FNAL SELEX Collab.)        |
| MAHMOOD        | 01  | PRL 86 2232    | A.H. Mahmood <i>et al.</i>       | (CLEO Collab.)              |
| AITALA         | 00  | PL B471 449    | E.M. Aitala <i>et al.</i>        | (FNAL E791 Collab.)         |
| JAFFE          | 00  | PR D62 072005  | D.E. Jaffe <i>et al.</i>         | (CLEO Collab.)              |
| ALAM           | 98  | PR D57 4467    | M.S. Alam <i>et al.</i>          | (CLEO Collab.)              |
| ALBRECHT       | 96E | PRPL 276 223   | H. Albrecht <i>et al.</i>        | (ARGUS Collab.)             |
| ALEEV          | 96  | JINRRC 3-77 31 | A.N. Aleev <i>et al.</i>         | (Serpukhov EXCHARM Collab.) |
| ALEXANDER      | 96C | PR D53 R1013   | J.P. Alexander <i>et al.</i>     | (CLEO Collab.)              |
| ALBRECHT       | 95B | PL B342 397    | H. Albrecht <i>et al.</i>        | (ARGUS Collab.)             |
| AMMAR          | 95  | PRL 74 3534    | R. Ammar <i>et al.</i>           | (CLEO Collab.)              |
| BISHAI         | 95  | PL B350 256    | M. Bishai <i>et al.</i>          | (CLEO Collab.)              |
| CRAWFORD       | 95  | PRL 75 624     | G. Crawford <i>et al.</i>        | (CLEO Collab.)              |
| KODAMA         | 95  | PL B345 85     | K. Kodama <i>et al.</i>          | (FNAL E653 Collab.)         |
| ALBRECHT       | 94B | PL B326 320    | H. Albrecht <i>et al.</i>        | (ARGUS Collab.)             |
| ALEEV          | 94  | PAN 57 1370    | A.N. Aleev <i>et al.</i>         | (Serpukhov BIS-2 Collab.)   |

Translated from YF 57 1443.

|             |     |                              |                                  |                           |
|-------------|-----|------------------------------|----------------------------------|---------------------------|
| AVERY       | 94  | PL B325 257                  | P. Avery <i>et al.</i>           | (CLEO Collab.)            |
| BERGFELD    | 94  | PL B323 219                  | T. Bergfeld <i>et al.</i>        | (CLEO Collab.)            |
| FRABETTI    | 94E | PL B328 193                  | P.L. Frabetti <i>et al.</i>      | (FNAL E687 Collab.)       |
| AVERY       | 93  | PRL 71 2391                  | P. Avery <i>et al.</i>           | (CLEO Collab.)            |
| BOZEK       | 93  | PL B312 247                  | A. Bozek <i>et al.</i>           | (CERN NA32 Collab.)       |
| FRABETTI    | 93D | PRL 70 1755                  | P.L. Frabetti <i>et al.</i>      | (FNAL E687 Collab.)       |
| FRABETTI    | 93H | PL B314 477                  | P.L. Frabetti <i>et al.</i>      | (FNAL E687 Collab.)       |
| KUBOTA      | 93  | PRL 71 3255                  | Y. Kubota <i>et al.</i>          | (CLEO Collab.)            |
| ALBRECHT    | 92  | PL B274 239                  | H. Albrecht <i>et al.</i>        | (ARGUS Collab.)           |
| ALBRECHT    | 92O | ZPHY C56 1                   | H. Albrecht <i>et al.</i>        | (ARGUS Collab.)           |
| BARLAG      | 92  | PL B283 465                  | S. Barlag <i>et al.</i>          | (ACCMOR Collab.)          |
| CRAWFORD    | 92  | PR D45 752                   | G. Crawford <i>et al.</i>        | (CLEO Collab.)            |
| JEZABEK     | 92  | PL B286 175                  | M. Jezabek, K. Rybicki, R. Rylko | (CRAC)                    |
| ALBRECHT    | 91G | PL B269 234                  | H. Albrecht <i>et al.</i>        | (ARGUS Collab.)           |
| AVERY       | 91  | PR D43 3599                  | P. Avery <i>et al.</i>           | (CLEO Collab.)            |
| ALVAREZ     | 90  | ZPHY C47 539                 | M.P. Alvarez <i>et al.</i>       | (CERN NA14/2 Collab.)     |
| ALVAREZ     | 90B | PL B246 256                  | M.P. Alvarez <i>et al.</i>       | (CERN NA14/2 Collab.)     |
| ANJOS       | 90  | PR D41 801                   | J.C. Anjos <i>et al.</i>         | (FNAL E691 Collab.)       |
| AVERY       | 90B | PRL 65 2842                  | P. Avery <i>et al.</i>           | (CLEO Collab.)            |
| BARLAG      | 90D | ZPHY C48 29                  | S. Barlag <i>et al.</i>          | (ACCMOR Collab.)          |
| FRABETTI    | 90  | PL B251 639                  | P.L. Frabetti <i>et al.</i>      | (FNAL E687 Collab.)       |
| BARLAG      | 89  | PL B218 374                  | S. Barlag <i>et al.</i>          | (ACCMOR Collab.)          |
| AGUILAR-... | 88B | ZPHY C40 321                 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.)        |
| Also        |     | PL B189 254                  | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.)        |
| Also        |     | PL B199 462                  | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.)        |
| Also        |     | SJNP 48 833                  | M. Begalli <i>et al.</i>         | (LEBC-EHS Collab.)        |
|             |     | Translated from YAF 48 1310. |                                  |                           |
| ALBRECHT    | 88C | PL B207 109                  | H. Albrecht <i>et al.</i>        | (ARGUS Collab.)           |
| ANJOS       | 88B | PRL 60 1379                  | J.C. Anjos <i>et al.</i>         | (FNAL E691 Collab.)       |
| ADAMOVICH   | 87  | EPL 4 887                    | M.I. Adamovich <i>et al.</i>     | (Photon Emulsion Collab.) |
| Also        |     | SJNP 46 447                  | F. Viaggi <i>et al.</i>          | (Photon Emulsion Collab.) |
|             |     | Translated from YAF 46 799.  |                                  |                           |
| AMENDOLIA   | 87  | ZPHY C36 513                 | S.R. Amendolia <i>et al.</i>     | (CERN NA1 Collab.)        |
| JONES       | 87  | ZPHY C36 593                 | G.T. Jones <i>et al.</i>         | (CERN WA21 Collab.)       |
| ABE         | 86  | PR D33 1                     | K. Abe <i>et al.</i>             |                           |
| ALEEV       | 84  | ZPHY C23 333                 | A.N. Aleev <i>et al.</i>         | (BIS-2 Collab.)           |
| BOSETTI     | 82  | PL 109B 234                  | P.C. Bosetti <i>et al.</i>       | (AACH3, BONN, CERN+)      |
| VELLA       | 82  | PRL 48 1515                  | E. Vella <i>et al.</i>           | (SLAC, LBL, UCB)          |
| BASILE      | 81B | NC 62A 14                    | M. Basile <i>et al.</i>          | (CERN, BGNA, PGIA, FRAS)  |
| CALICCHIO   | 80  | PL 93B 521                   | M. Calicchio <i>et al.</i>       | (BARI, BIRM, BRUX+)       |

### OTHER RELATED PAPERS

|           |    |               |                           |
|-----------|----|---------------|---------------------------|
| MIGLIOZZI | 99 | PL B462 217   | P. Migliozi <i>et al.</i> |
| DUNIETZ   | 98 | PR D58 094010 | I. Dunietz                |