

CHARMED BARYONS ($C = +1$)

$$\Lambda_c^+ = udc, \quad \Sigma_c^{++} = uuc, \quad \Sigma_c^+ = udc, \quad \Sigma_c^0 = ddc,$$

$$\Xi_c^+ = usc, \quad \Xi_c^0 = dsc, \quad \Omega_c^0 = ssc$$

Λ_c^+

$$I(J^P) = 0(\frac{1}{2}^+)$$

J is not well measured; $\frac{1}{2}$ is the quark-model prediction.

Mass $m = 2286.46 \pm 0.14$ MeV

Mean life $\tau = (200 \pm 6) \times 10^{-15}$ s ($S = 1.6$)

$c\tau = 59.9$ μ m

Decay asymmetry parameters

$$\Lambda\pi^+ \quad \alpha = -0.91 \pm 0.15$$

$$\Sigma^+\pi^0 \quad \alpha = -0.45 \pm 0.32$$

$$\Lambda\ell^+\nu_\ell \quad \alpha = -0.86 \pm 0.04$$

$$[\alpha(\Lambda_c^+) + \alpha(\bar{\Lambda}_c^-)] / [\alpha(\Lambda_c^+) - \alpha(\bar{\Lambda}_c^-)] \text{ in } \Lambda_c^+ \rightarrow \Lambda\pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^- =$$

$$-0.07 \pm 0.31$$

$$[\alpha(\Lambda_c^+) + \alpha(\bar{\Lambda}_c^-)] / [\alpha(\Lambda_c^+) - \alpha(\bar{\Lambda}_c^-)] \text{ in } \Lambda_c^+ \rightarrow \Lambda e^+\nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}e^-\bar{\nu}_e =$$

$$0.00 \pm 0.04$$

Nearly all branching fractions of the Λ_c^+ are measured relative to the $pK^-\pi^+$ mode, but there are no model-independent measurements of this branching fraction. We explain how we arrive at our value of $B(\Lambda_c^+ \rightarrow pK^-\pi^+)$ in a Note at the beginning of the branching-ratio measurements in the Listings. When this branching fraction is eventually well determined, all the other branching fractions will slide up or down proportionally as the true value differs from the value we use here.

Λ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Hadronic modes with a p: $S = -1$ final states			
$p\bar{K}^0$	(2.3 \pm 0.6) %		873
$pK^-\pi^+$	[a] (5.0 \pm 1.3) %		823
$p\bar{K}^*(892)^0$	[b] (1.6 \pm 0.5) %		685
$\Delta(1232)^{++}K^-$	(8.6 \pm 3.0) $\times 10^{-3}$		710
$\Lambda(1520)\pi^+$	[b] (1.8 \pm 0.6) %		627
$pK^-\pi^+$ nonresonant	(2.8 \pm 0.8) %		823
$p\bar{K}^0\pi^0$	(3.3 \pm 1.0) %		823

$\rho \bar{K}^0 \eta$	(1.2 ± 0.4) %	568
$\rho \bar{K}^0 \pi^+ \pi^-$	(2.6 ± 0.7) %	754
$\rho K^- \pi^+ \pi^0$	(3.4 ± 1.0) %	759
$\rho K^*(892)^- \pi^+$	[b] (1.1 ± 0.5) %	580
$\rho (K^- \pi^+)_{\text{nonresonant}} \pi^0$	(3.6 ± 1.2) %	759
$\Delta(1232) \bar{K}^*(892)$	seen	419
$\rho K^- \pi^+ \pi^+ \pi^-$	(1.1 ± 0.8) × 10 ⁻³	671
$\rho K^- \pi^+ \pi^0 \pi^0$	(8 ± 4) × 10 ⁻³	678

Hadronic modes with a ρ : $S = 0$ final states

$\rho \pi^+ \pi^-$	(3.5 ± 2.0) × 10 ⁻³	927
$\rho f_0(980)$	[b] (2.8 ± 1.9) × 10 ⁻³	622
$\rho \pi^+ \pi^+ \pi^- \pi^-$	(1.8 ± 1.2) × 10 ⁻³	852
$\rho K^+ K^-$	(7.7 ± 3.5) × 10 ⁻⁴	616
$\rho \phi$	[b] (8.2 ± 2.7) × 10 ⁻⁴	590
$\rho K^+ K^- \text{ non-}\phi$	(3.5 ± 1.7) × 10 ⁻⁴	616

Hadronic modes with a hyperon: $S = -1$ final states

$\Lambda \pi^+$	(1.01 ± 0.28) %	864
$\Lambda \pi^+ \pi^0$	(3.6 ± 1.3) %	844
$\Lambda \rho^+$	< 5 %	CL=95% 635
$\Lambda \pi^+ \pi^+ \pi^-$	(2.6 ± 0.7) %	807
$\Sigma(1385)^+ \pi^+ \pi^-, \Sigma^{*+} \rightarrow$	(7 ± 4) × 10 ⁻³	688
$\Lambda \pi^+$		
$\Sigma(1385)^- \pi^+ \pi^+, \Sigma^{*-} \rightarrow$	(5.5 ± 1.7) × 10 ⁻³	688
$\Lambda \pi^-$		
$\Lambda \pi^+ \rho^0$	(1.1 ± 0.5) %	523
$\Sigma(1385)^+ \rho^0, \Sigma^{*+} \rightarrow \Lambda \pi^+$	(3.7 ± 3.1) × 10 ⁻³	363
$\Lambda \pi^+ \pi^+ \pi^- \text{ nonresonant}$	< 8 × 10 ⁻³	CL=90% 807
$\Lambda \pi^+ \pi^+ \pi^- \pi^0 \text{ total}$	(1.8 ± 0.8) %	757
$\Lambda \pi^+ \eta$	[b] (1.8 ± 0.6) %	691
$\Sigma(1385)^+ \eta$	[b] (8.5 ± 3.3) × 10 ⁻³	570
$\Lambda \pi^+ \omega$	[b] (1.2 ± 0.5) %	517
$\Lambda \pi^+ \pi^+ \pi^- \pi^0, \text{ no } \eta \text{ or } \omega$	< 7 × 10 ⁻³	CL=90% 757
$\Lambda K^+ \bar{K}^0$	(6.5 ± 2.0) × 10 ⁻³	443
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda \bar{K}^0$	(1.9 ± 0.7) × 10 ⁻³	286
$\Sigma^0 \pi^+$	(1.04 ± 0.31) %	825
$\Sigma^+ \pi^0$	(1.00 ± 0.34) %	827
$\Sigma^+ \eta$	(5.5 ± 2.3) × 10 ⁻³	714
$\Sigma^+ \pi^+ \pi^-$	(3.6 ± 1.0) %	804
$\Sigma^+ \rho^0$	< 1.4 %	CL=95% 575
$\Sigma^- \pi^+ \pi^+$	(1.9 ± 0.8) %	799
$\Sigma^0 \pi^+ \pi^0$	(1.8 ± 0.8) %	803
$\Sigma^0 \pi^+ \pi^+ \pi^-$	(8.3 ± 3.1) × 10 ⁻³	763
$\Sigma^+ \pi^+ \pi^- \pi^0$	—	767

$\Sigma^+ \omega$	[b]	(2.7 ± 1.0) %		569
$\Sigma^+ K^+ K^-$		(2.8 ± 0.8) × 10 ⁻³		349
$\Sigma^+ \phi$	[b]	(3.2 ± 1.0) × 10 ⁻³		295
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow$		(8.2 ± 3.1) × 10 ⁻⁴		286
$\Sigma^+ K^+ K^-$ nonresonant		< 7 × 10 ⁻⁴	CL=90%	349
$\Xi^0 K^+$		(3.9 ± 1.4) × 10 ⁻³		653
$\Xi^- K^+ \pi^+$		(4.9 ± 1.7) × 10 ⁻³		566
$\Xi(1530)^0 K^+$	[b]	(2.6 ± 1.0) × 10 ⁻³		473

Hadronic modes with a hyperon: S = 0 final states

ΛK^+		(7.5 ± 2.6) × 10 ⁻⁴		781
$\Sigma^0 K^+$		(5.8 ± 2.4) × 10 ⁻⁴		735
$\Sigma^+ K^+ \pi^-$		(1.7 ± 0.7) × 10 ⁻³		670
$\Sigma^+ K^*(892)^0$	[b]	(2.8 ± 1.1) × 10 ⁻³		470
$\Sigma^- K^+ \pi^+$		< 1.0 × 10 ⁻³	CL=90%	664

Doubly Cabibbo-suppressed modes

$p K^+ \pi^-$		< 2.3 × 10 ⁻⁴	CL=90%	823
---------------	--	--------------------------	--------	-----

Semileptonic modes

$\Lambda \ell^+ \nu_\ell$	[c]	(2.0 ± 0.6) %		871
$\Lambda e^+ \nu_e$		(2.1 ± 0.6) %		871
$\Lambda \mu^+ \nu_\mu$		(2.0 ± 0.7) %		867

Inclusive modes

e^+ anything		(4.5 ± 1.7) %		—
$p e^+$ anything		(1.8 ± 0.9) %		—
p anything		(50 ± 16) %		—
p anything (no Λ)		(12 ± 19) %		—
n anything		(50 ± 16) %		—
n anything (no Λ)		(29 ± 17) %		—
Λ anything		(35 ± 11) %	S=1.4	—
Σ^\pm anything	[d]	(10 ± 5) %		—
3prongs		(24 ± 8) %		—

$\Delta C = 1$ weak neutral current (C1) modes, or Lepton number (L) violating modes

$p \mu^+ \mu^-$	C1	< 3.4 × 10 ⁻⁴	CL=90%	937
$\Sigma^- \mu^+ \mu^+$	L	< 7.0 × 10 ⁻⁴	CL=90%	812

$\Lambda_c(2593)^+$

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that $\Sigma_c(2455)\pi$ decays, with little available phase space, are dominant. This assumes that $J^P = 1/2^+$ for the $\Sigma_c(2455)$.

$$\text{Mass } m = 2595.4 \pm 0.6 \text{ MeV} \quad (S = 1.1)$$

$$m - m_{\Lambda_c^+} = 308.9 \pm 0.6 \text{ MeV} \quad (S = 1.1)$$

$$\text{Full width } \Gamma = 3.6_{-1.3}^{+2.0} \text{ MeV}$$

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma_c(2455)\pi$ — the latter just barely — are the only strong decays allowed to an excited Λ_c^+ having this mass; and the submode seems to dominate.

$\Lambda_c(2593)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[e] $\approx 67\%$	124
$\Sigma_c(2455)^{++} \pi^-$	$24 \pm 7\%$	28
$\Sigma_c(2455)^0 \pi^+$	$24 \pm 7\%$	28
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	$18 \pm 10\%$	124
$\Lambda_c^+ \pi^0$	[f] not seen	261
$\Lambda_c^+ \gamma$	not seen	291

 $\Lambda_c(2625)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$$\text{Mass } m = 2628.1 \pm 0.6 \text{ MeV} \quad (S = 1.5)$$

$$m - m_{\Lambda_c^+} = 341.7 \pm 0.6 \text{ MeV} \quad (S = 1.6)$$

$$\text{Full width } \Gamma < 1.9 \text{ MeV, CL} = 90\%$$

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma(2455)\pi$ are the only strong decays allowed to an excited Λ_c^+ having this mass.

$\Lambda_c(2625)^+$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[e] $\approx 67\%$		184
$\Sigma_c(2455)^{++} \pi^-$	< 5	90%	102
$\Sigma_c(2455)^0 \pi^+$	< 5	90%	102
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	large		184
$\Lambda_c^+ \pi^0$	[f] not seen		293
$\Lambda_c^+ \gamma$	not seen		319

$\Sigma_c(2455)$

$$I(J^P) = 1(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\Sigma_c(2455)^{++} \text{ mass } m = 2454.02 \pm 0.18 \text{ MeV}$$

$$\Sigma_c(2455)^+ \text{ mass } m = 2452.9 \pm 0.4 \text{ MeV}$$

$$\Sigma_c(2455)^0 \text{ mass } m = 2453.76 \pm 0.18 \text{ MeV}$$

$$m_{\Sigma_c^{++}} - m_{\Lambda_c^+} = 167.56 \pm 0.11 \text{ MeV}$$

$$m_{\Sigma_c^+} - m_{\Lambda_c^+} = 166.4 \pm 0.4 \text{ MeV}$$

$$m_{\Sigma_c^0} - m_{\Lambda_c^+} = 167.30 \pm 0.11 \text{ MeV}$$

$$m_{\Sigma_c^{++}} - m_{\Sigma_c^0} = 0.27 \pm 0.11 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Sigma_c^+} - m_{\Sigma_c^0} = -0.9 \pm 0.4 \text{ MeV}$$

$$\Sigma_c(2455)^{++} \text{ full width } \Gamma = 2.23 \pm 0.30 \text{ MeV}$$

$$\Sigma_c(2455)^+ \text{ full width } \Gamma < 4.6 \text{ MeV, CL} = 90\%$$

$$\Sigma_c(2455)^0 \text{ full width } \Gamma = 2.2 \pm 0.4 \text{ MeV} \quad (S = 1.4)$$

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

 $\Sigma_c(2455)$ DECAY MODESFraction (Γ_i/Γ) p (MeV/c) $\Lambda_c^+ \pi$ $\approx 100\%$

94

 $\Sigma_c(2520)$

$$I(J^P) = 1(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\Sigma_c(2520)^{++} \text{ mass } m = 2518.4 \pm 0.6 \text{ MeV} \quad (S = 1.4)$$

$$\Sigma_c(2520)^+ \text{ mass } m = 2517.5 \pm 2.3 \text{ MeV}$$

$$\Sigma_c(2520)^0 \text{ mass } m = 2518.0 \pm 0.5 \text{ MeV}$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 231.9 \pm 0.6 \text{ MeV} \quad (S = 1.5)$$

$$m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} = 231.0 \pm 2.3 \text{ MeV}$$

$$m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 231.6 \pm 0.5 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 0.3 \pm 0.6 \text{ MeV} \quad (S = 1.2)$$

$$\Sigma_c(2520)^{++} \text{ full width } \Gamma = 14.9 \pm 1.9 \text{ MeV}$$

$$\Sigma_c(2520)^+ \text{ full width } \Gamma < 17 \text{ MeV, CL} = 90\%$$

$$\Sigma_c(2520)^0 \text{ full width } \Gamma = 16.1 \pm 2.1 \text{ MeV}$$

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2520)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	180

$\Sigma_c(2800)$

$$I(J^P) = 1(?^?)$$

$$\begin{aligned} \Sigma_c(2800)^{++} \text{ mass } m &= 2801_{-6}^{+4} \\ \Sigma_c(2800)^+ \text{ mass } m &= 2792_{-5}^{+14} \\ \Sigma_c(2800)^0 \text{ mass } m &= 2802_{-7}^{+4} \\ m_{\Sigma_c(2800)^{++}} - m_{\Lambda_c^+} &= 514_{-6}^{+4} \\ m_{\Sigma_c(2800)^+} - m_{\Lambda_c^+} &= 505_{-5}^{+14} \\ m_{\Sigma_c(2800)^0} - m_{\Lambda_c^+} &= 515_{-7}^{+4} \\ \Sigma_c(2800)^{++} \text{ full width } \Gamma &= 75_{-17}^{+22} \\ \Sigma_c(2800)^+ \text{ full width } \Gamma &= 62_{-40}^{+60} \\ \Sigma_c(2800)^0 \text{ full width } \Gamma &= 61_{-18}^{+28} \end{aligned}$$

$\Sigma_c(2800)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	seen	443

Ξ_c^+

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2467.9 \pm 0.4 \text{ MeV}$$

$$\text{Mean life } \tau = (442 \pm 26) \times 10^{-15} \text{ s} \quad (S = 1.3)$$

$$c\tau = 132 \text{ } \mu\text{m}$$

Ξ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
-----------------------	--------------------------------	------------------	-------------

No absolute branching fractions have been measured.
The following are branching *ratios* relative to $\Xi^- \pi^+ \pi^+$.

Cabibbo-favored ($S = -2$) decays

$p K_S^0 K_S^0$	[g]	0.087 ± 0.022	767
$\Lambda \bar{K}^0 \pi^+$		—	852
$\Sigma(1385)^+ \bar{K}^0$	[b,g]	1.0 ± 0.5	746
$\Lambda K^- \pi^+ \pi^+$	[g]	0.323 ± 0.033	787
$\Lambda \bar{K}^*(892)^0 \pi^+$	[b,g]	< 0.2	90% 608

$\Sigma(1385)^+ K^- \pi^+$	$[b,g] < 0.3$	90%	678
$\Sigma^+ K^- \pi^+$	$[g] 0.94 \pm 0.11$		811
$\Sigma^+ \bar{K}^*(892)^0$	$[b,g] 0.81 \pm 0.15$		658
$\Sigma^0 K^- \pi^+ \pi^+$	$[g] 0.29 \pm 0.16$		735
$\Xi^0 \pi^+$	$[g] 0.55 \pm 0.16$		877
$\Xi^- \pi^+ \pi^+$	$[g] \text{DEFINED AS } 1$		851
$\Xi(1530)^0 \pi^+$	$[b,g] < 0.1$	90%	750
$\Xi^0 \pi^+ \pi^0$	$[g] 2.34 \pm 0.68$		856
$\Xi^0 \pi^+ \pi^+ \pi^-$	$[g] 1.74 \pm 0.50$		818
$\Xi^0 e^+ \nu_e$	$[g] 2.3 \begin{smallmatrix} +0.7 \\ -0.9 \end{smallmatrix}$		884
$\Omega^- K^+ \pi^+$	$[g] 0.07 \pm 0.04$		399

Cabibbo-suppressed decays

$p K^- \pi^+$	$[g] 0.21 \pm 0.03$		944
$p \bar{K}^*(892)^0$	$[b,g] 0.12 \pm 0.02$		828
$\Sigma^+ K^+ K^-$	$[g] 0.15 \pm 0.07$		580
$\Sigma^+ \phi$	$[b,g] < 0.11$	90%	549
$\Xi(1690)^0 K^+, \Xi(1690)^0 \rightarrow \Sigma^+ K^-$	$[g] < 0.05$	90%	501



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2471.0 \pm 0.4$ MeV

$$m_{\Xi_c^0} - m_{\Xi_c^+} = 3.1 \pm 0.5 \text{ MeV}$$

Mean life $\tau = (112_{-10}^{+13}) \times 10^{-15}$ s

$$c\tau = 33.6 \text{ } \mu\text{m}$$


Decay asymmetry parameters

$$\Xi^- \pi^+ \quad \alpha = -0.6 \pm 0.4$$

No absolute branching fractions have been measured. Several measurements of ratios of fractions may be found in the Listings that follow.

Ξ_c^0 DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$p K^- K^- \pi^+$	seen	676
$p K^- \bar{K}^*(892)^0$	seen	413
$p K^- K^- \pi^+$ no $\bar{K}^*(892)^0$	seen	676
ΛK_S^0	seen	906
$\Lambda \bar{K}^0 \pi^+ \pi^-$	seen	787

$\Lambda K^- \pi^+ \pi^+ \pi^-$	seen	703
$\Xi^- \pi^+$	seen	875
$\Xi^- \pi^+ \pi^+ \pi^-$	seen	817
$\Omega^- K^+$	seen	523
$\Xi^- e^+ \nu_e$	seen	882
$\Xi^- \ell^+$ anything	seen	—



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$


J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2575.7 \pm 3.1$ MeV

$$m_{\Xi_c^{'+}} - m_{\Xi_c^+} = 107.8 \pm 3.0 \text{ MeV}$$

The $\Xi_c^{'+} - \Xi_c^+$ mass difference is too small for any strong decay to occur.

$\Xi_c^{'+}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ \gamma$	seen	106



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$


J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2578.0 \pm 2.9$ MeV

$$m_{\Xi_c^{'0}} - m_{\Xi_c^0} = 107.0 \pm 2.9 \text{ MeV}$$

The $\Xi_c^{'0} - \Xi_c^0$ mass difference is too small for any strong decay to occur.

$\Xi_c^{'0}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^0 \gamma$	seen	105



$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$\Xi_c(2645)^+$ mass $m = 2646.6 \pm 1.4$ MeV ($S = 1.6$)

$\Xi_c(2645)^0$ mass $m = 2646.1 \pm 1.2$ MeV

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 175.6 \pm 1.4 \text{ MeV} \quad (S = 1.7)$$

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.2 \pm 1.1 \text{ MeV}$$

$\Xi_c(2645)^+$ full width $\Gamma < 3.1$ MeV, CL = 90%

$\Xi_c(2645)^0$ full width $\Gamma < 5.5$ MeV, CL = 90%

$\Xi_c \pi$ is the only strong decay allowed to a Ξ_c resonance having this mass.

$\Xi_c(2645)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^0 \pi^+$	seen	102
$\Xi_c^+ \pi^-$	seen	107

$\Xi_c(2790)$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

J^P has not been measured; $\frac{1}{2}^-$ is the quark-model prediction.

$\Xi_c(2790)^+$ mass = 2789.2 ± 3.2 MeV

$\Xi_c(2790)^0$ mass = 2791.9 ± 3.3 MeV

$m_{\Xi_c(2790)^+} - m_{\Xi_c^0} = 318.2 \pm 3.2$ MeV

$m_{\Xi_c(2790)^0} - m_{\Xi_c^+} = 324.0 \pm 3.3$ MeV

$\Xi_c(2790)^+$ width < 15 MeV, CL = 90%

$\Xi_c(2790)^0$ width < 12 MeV, CL = 90%

$\Xi_c(2790)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ \pi^-$	seen	159

$\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$\Xi_c(2815)^+$ mass $m = 2816.5 \pm 1.2$ MeV

$\Xi_c(2815)^0$ mass $m = 2818.2 \pm 2.1$ MeV

$m_{\Xi_c(2815)^+} - m_{\Xi_c^+} = 348.6 \pm 1.2$ MeV

$m_{\Xi_c(2815)^0} - m_{\Xi_c^0} = 347.2 \pm 2.1$ MeV

$\Xi_c(2815)^+$ full width $\Gamma < 3.5$ MeV, CL = 90%

$\Xi_c(2815)^0$ full width $\Gamma < 6.5$ MeV, CL = 90%

The $\Xi_c \pi \pi$ modes are consistent with being entirely via $\Xi_c(2645) \pi$.

$\Xi_c(2815)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ \pi^+ \pi^-$	seen	196
$\Xi_c^0 \pi^+ \pi^-$	seen	191



$$I(J^P) = 0(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2697.5 \pm 2.6 \text{ MeV} \quad (S = 1.2)$$

$$\text{Mean life } \tau = (69 \pm 12) \times 10^{-15} \text{ s}$$

$$c\tau = 21 \text{ } \mu\text{m}$$

No absolute branching fractions have been measured.

Ω_c^0 DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Sigma^+ K^- K^- \pi^+$	seen	691
$\Xi^0 K^- \pi^+$	seen	903
$\Xi^- K^- \pi^+ \pi^+$	seen	832
$\Omega^- e^+ \nu_e$	seen	830
$\Omega^- \pi^+$	seen	822
$\Omega^- \pi^+ \pi^0$	seen	798
$\Omega^- \pi^- \pi^+ \pi^+$	seen	754

NOTES

- [a] See the note on " Λ_c^+ Branching Fractions" in the Λ_c^+ Particle Listings.
- [b] This branching fraction includes all the decay modes of the final-state resonance.
- [c] An ℓ indicates an e or a μ mode, not a sum over these modes.
- [d] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [e] Assuming isospin conservation, so that the other third is $\Lambda_c^+ \pi^0 \pi^0$.
- [f] A test that the isospin is indeed 0, so that the particle is indeed a Λ_c^+ .
- [g] No absolute branching fractions have been measured. The value here is the branching *ratio* relative to $\Xi^- \pi^+ \pi^+$.