

N BARYONS

($S = 0, I = 1/2$)

$p, N^+ = uud; \quad n, N^0 = udd$

p

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

- Mass $m = 1.00727646688 \pm 0.00000000013$ u
- Mass $m = 938.27203 \pm 0.00008$ MeV [a]
- $|m_p - m_{\bar{p}}|/m_p < 1.0 \times 10^{-8}$, CL = 90% [b]
- $|\frac{q_{\bar{p}}}{m_{\bar{p}}}|/(\frac{q_p}{m_p}) = 0.99999999991 \pm 0.00000000009$
- $|q_p + q_{\bar{p}}|/e < 1.0 \times 10^{-8}$, CL = 90% [b]
- $|q_p + q_e|/e < 1.0 \times 10^{-21}$ [c]
- Magnetic moment $\mu = 2.792847351 \pm 0.000000028 \mu_N$
- $(\mu_p + \mu_{\bar{p}}) / \mu_p = (-2.6 \pm 2.9) \times 10^{-3}$
- Electric dipole moment $d < 0.54 \times 10^{-23}$ e cm
- Electric polarizability $\alpha = (12.0 \pm 0.6) \times 10^{-4}$ fm³
- Magnetic polarizability $\beta = (1.9 \pm 0.5) \times 10^{-4}$ fm³
- Charge radius = 0.875 ± 0.007 fm
- Mean life $\tau > 2.1 \times 10^{29}$ years, CL = 90% ($p \rightarrow$ invisible mode)
- Mean life $\tau > 10^{31}$ to 10^{33} years [d] (mode dependent)

See the "Note on Nucleon Decay" in our 1994 edition (Phys. Rev. **D50**, 1673) for a short review.

The "partial mean life" limits tabulated here are the limits on τ/B_i , where τ is the total mean life and B_i is the branching fraction for the mode in question. For N decays, p and n indicate proton and neutron partial lifetimes.

| p DECAY MODES | Partial mean life (10 ³⁰ years) | Confidence level | p (MeV/c) |
|----------------------------|---|------------------|----------------|
| Antilepton + meson | | | |
| $N \rightarrow e^+ \pi$ | > 158 (n), > 1600 (p) | 90% | 459 |
| $N \rightarrow \mu^+ \pi$ | > 100 (n), > 473 (p) | 90% | 453 |
| $N \rightarrow \nu \pi$ | > 112 (n), > 25 (p) | 90% | 459 |
| $p \rightarrow e^+ \eta$ | > 313 | 90% | 309 |
| $p \rightarrow \mu^+ \eta$ | > 126 | 90% | 297 |
| $n \rightarrow \nu \eta$ | > 158 | 90% | 310 |
| $N \rightarrow e^+ \rho$ | > 217 (n), > 75 (p) | 90% | 149 |
| $N \rightarrow \mu^+ \rho$ | > 228 (n), > 110 (p) | 90% | 113 |
| $N \rightarrow \nu \rho$ | > 19 (n), > 162 (p) | 90% | 149 |
| $p \rightarrow e^+ \omega$ | > 107 | 90% | 143 |

| | | | |
|--------------------------------|-----------------------------|-----|-----|
| $p \rightarrow \mu^+ \omega$ | > 117 | 90% | 105 |
| $n \rightarrow \nu \omega$ | > 108 | 90% | 144 |
| $N \rightarrow e^+ K$ | > 17 (n), > 150 (p) | 90% | 339 |
| $p \rightarrow e^+ K_S^0$ | > 120 | 90% | 337 |
| $p \rightarrow e^+ K_L^0$ | > 51 | 90% | 337 |
| $N \rightarrow \mu^+ K$ | > 26 (n), > 120 (p) | 90% | 329 |
| $p \rightarrow \mu^+ K_S^0$ | > 150 | 90% | 326 |
| $p \rightarrow \mu^+ K_L^0$ | > 83 | 90% | 326 |
| $N \rightarrow \nu K$ | > 86 (n), > 670 (p) | 90% | 339 |
| $n \rightarrow \nu K_S^0$ | > 51 | 90% | 338 |
| $p \rightarrow e^+ K^*(892)^0$ | > 84 | 90% | 45 |
| $N \rightarrow \nu K^*(892)$ | > 78 (n), > 51 (p) | 90% | 45 |

Antilepton + mesons

| | | | |
|-----------------------------------|-------|-----|-----|
| $p \rightarrow e^+ \pi^+ \pi^-$ | > 82 | 90% | 448 |
| $p \rightarrow e^+ \pi^0 \pi^0$ | > 147 | 90% | 449 |
| $n \rightarrow e^+ \pi^- \pi^0$ | > 52 | 90% | 449 |
| $p \rightarrow \mu^+ \pi^+ \pi^-$ | > 133 | 90% | 425 |
| $p \rightarrow \mu^+ \pi^0 \pi^0$ | > 101 | 90% | 427 |
| $n \rightarrow \mu^+ \pi^- \pi^0$ | > 74 | 90% | 427 |
| $n \rightarrow e^+ K^0 \pi^-$ | > 18 | 90% | 319 |

Lepton + meson

| | | | |
|------------------------------|------|-----|-----|
| $n \rightarrow e^- \pi^+$ | > 65 | 90% | 459 |
| $n \rightarrow \mu^- \pi^+$ | > 49 | 90% | 453 |
| $n \rightarrow e^- \rho^+$ | > 62 | 90% | 150 |
| $n \rightarrow \mu^- \rho^+$ | > 7 | 90% | 114 |
| $n \rightarrow e^- K^+$ | > 32 | 90% | 340 |
| $n \rightarrow \mu^- K^+$ | > 57 | 90% | 330 |

Lepton + mesons

| | | | |
|-----------------------------------|-------|-----|-----|
| $p \rightarrow e^- \pi^+ \pi^+$ | > 30 | 90% | 448 |
| $n \rightarrow e^- \pi^+ \pi^0$ | > 29 | 90% | 449 |
| $p \rightarrow \mu^- \pi^+ \pi^+$ | > 17 | 90% | 425 |
| $n \rightarrow \mu^- \pi^+ \pi^0$ | > 34 | 90% | 427 |
| $p \rightarrow e^- \pi^+ K^+$ | > 75 | 90% | 320 |
| $p \rightarrow \mu^- \pi^+ K^+$ | > 245 | 90% | 279 |

Antilepton + photon(s)

| | | | |
|-----------------------------------|-------|-----|-----|
| $p \rightarrow e^+ \gamma$ | > 670 | 90% | 469 |
| $p \rightarrow \mu^+ \gamma$ | > 478 | 90% | 463 |
| $n \rightarrow \nu \gamma$ | > 28 | 90% | 470 |
| $p \rightarrow e^+ \gamma \gamma$ | > 100 | 90% | 469 |
| $n \rightarrow \nu \gamma \gamma$ | > 219 | 90% | 470 |

Three (or more) leptons

| | | | |
|-----------------------------------|----------|-----|-----|
| $p \rightarrow e^+ e^+ e^-$ | > 793 | 90% | 469 |
| $p \rightarrow e^+ \mu^+ \mu^-$ | > 359 | 90% | 457 |
| $p \rightarrow e^+ \nu \nu$ | > 17 | 90% | 469 |
| $n \rightarrow e^+ e^- \nu$ | > 257 | 90% | 470 |
| $n \rightarrow \mu^+ e^- \nu$ | > 83 | 90% | 464 |
| $n \rightarrow \mu^+ \mu^- \nu$ | > 79 | 90% | 458 |
| $p \rightarrow \mu^+ e^+ e^-$ | > 529 | 90% | 463 |
| $p \rightarrow \mu^+ \mu^+ \mu^-$ | > 675 | 90% | 439 |
| $p \rightarrow \mu^+ \nu \nu$ | > 21 | 90% | 463 |
| $p \rightarrow e^- \mu^+ \mu^+$ | > 6 | 90% | 457 |
| $n \rightarrow 3\nu$ | > 0.0005 | 90% | 470 |

Inclusive modes

| | | | |
|------------------------------------|------------------|-----|---|
| $N \rightarrow e^+$ anything | > 0.6 (n, p) | 90% | — |
| $N \rightarrow \mu^+$ anything | > 12 (n, p) | 90% | — |
| $N \rightarrow e^+ \pi^0$ anything | > 0.6 (n, p) | 90% | — |

$\Delta B = 2$ dinucleon modes

The following are lifetime limits per iron nucleus.

| | | | |
|------------------------------------|------------------------|-----|---|
| $pp \rightarrow \pi^+ \pi^+$ | > 0.7 | 90% | — |
| $pn \rightarrow \pi^+ \pi^0$ | > 2 | 90% | — |
| $nn \rightarrow \pi^+ \pi^-$ | > 0.7 | 90% | — |
| $nn \rightarrow \pi^0 \pi^0$ | > 3.4 | 90% | — |
| $pp \rightarrow e^+ e^+$ | > 5.8 | 90% | — |
| $pp \rightarrow e^+ \mu^+$ | > 3.6 | 90% | — |
| $pp \rightarrow \mu^+ \mu^+$ | > 1.7 | 90% | — |
| $pn \rightarrow e^+ \bar{\nu}$ | > 2.8 | 90% | — |
| $pn \rightarrow \mu^+ \bar{\nu}$ | > 1.6 | 90% | — |
| $nn \rightarrow \nu_e \bar{\nu}_e$ | > 0.000049 | 90% | — |
| $pn \rightarrow$ invisible | > 2.1×10^{-5} | 90% | — |
| $pp \rightarrow$ invisible | > 0.00005 | 90% | — |

\bar{p} DECAY MODES

| \bar{p} DECAY MODES | Partial mean life (years) | Confidence level | p (MeV/c) |
|------------------------------------|------------------------------|------------------|----------------|
| $\bar{p} \rightarrow e^- \gamma$ | > 7×10^5 | 90% | 469 |
| $\bar{p} \rightarrow \mu^- \gamma$ | > 5×10^4 | 90% | 463 |
| $\bar{p} \rightarrow e^- \pi^0$ | > 4×10^5 | 90% | 459 |
| $\bar{p} \rightarrow \mu^- \pi^0$ | > 5×10^4 | 90% | 453 |
| $\bar{p} \rightarrow e^- \eta$ | > 2×10^4 | 90% | 309 |
| $\bar{p} \rightarrow \mu^- \eta$ | > 8×10^3 | 90% | 297 |

| | | | |
|--|-------------------|-----|-----|
| $\bar{p} \rightarrow e^- K_S^0$ | > 900 | 90% | 337 |
| $\bar{p} \rightarrow \mu^- K_S^0$ | > 4×10^3 | 90% | 326 |
| $\bar{p} \rightarrow e^- K_L^0$ | > 9×10^3 | 90% | 337 |
| $\bar{p} \rightarrow \mu^- K_L^0$ | > 7×10^3 | 90% | 326 |
| $\bar{p} \rightarrow e^- \gamma\gamma$ | > 2×10^4 | 90% | 469 |
| $\bar{p} \rightarrow \mu^- \gamma\gamma$ | > 2×10^4 | 90% | 463 |
| $\bar{p} \rightarrow e^- \omega$ | > 200 | 90% | 143 |

n $I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$

Mass $m = 1.0086649156 \pm 0.0000000006$ u

Mass $m = 939.56536 \pm 0.00008$ MeV [a]

$m_n - m_p = 1.2933317 \pm 0.0000005$ MeV
 $= 0.0013884487 \pm 0.0000000006$ u

Mean life $\tau = 885.7 \pm 0.8$ s

$c\tau = 2.655 \times 10^8$ km

Magnetic moment $\mu = -1.9130427 \pm 0.0000005 \mu_N$

Electric dipole moment $d < 0.63 \times 10^{-25}$ e cm, CL = 90%

Mean-square charge radius $\langle r_n^2 \rangle = -0.1161 \pm 0.0022$
 fm^2 (S = 1.3)

Electric polarizability $\alpha = (11.6 \pm 1.5) \times 10^{-4} \text{ fm}^3$

Magnetic polarizability $\beta = (3.7 \pm 2.0) \times 10^{-4} \text{ fm}^3$

Charge $q = (-0.4 \pm 1.1) \times 10^{-21} e$

Mean $n\bar{n}$ -oscillation time > 8.6×10^7 s, CL = 90% (free n)

Mean $n\bar{n}$ -oscillation time > 1.3×10^8 s, CL = 90% [e] (bound n)

Decay parameters [f]

$p e^- \bar{\nu}_e$ $\lambda \equiv g_A / g_V = -1.2695 \pm 0.0029$ (S = 2.0)

" $A = -0.1173 \pm 0.0013$ (S = 2.3)

" $B = 0.981 \pm 0.004$ (S = 1.1)

" $a = -0.103 \pm 0.004$

" $\phi_{AV} = (180.06 \pm 0.07)^\circ$ [g]

" $D = (-4 \pm 6) \times 10^{-4}$

| n DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | $\frac{P}{\text{MeV}/c}$ |
|---|--------------------------------|------------------|--------------------------|
| $p e^- \bar{\nu}_e$ | 100 % | | 1 |
| $p e^- \bar{\nu}_e \gamma$ | [h] < 6.9×10^{-3} | 90% | 1 |
| Charge conservation (Q) violating mode | | | |
| $p \nu_e \bar{\nu}_e$ | Q < 8×10^{-27} | 68% | 1 |

$N(1440) P_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Breit-Wigner mass = 1420 to 1470 (≈ 1440) MeVBreit-Wigner full width = 200 to 450 (≈ 300) MeV $p_{\text{beam}} = 0.61 \text{ GeV}/c$ $4\pi\lambda^2 = 31.0 \text{ mb}$ Re(pole position) = 1350 to 1380 (≈ 1365) MeV $-2\text{Im}(\text{pole position}) = 160 \text{ to } 220$ (≈ 190) MeV

| $N(1440)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $N\pi$ | 0.55 to 0.75 | 398 |
| $N\pi\pi$ | 30–40 % | 347 |
| $\Delta\pi$ | 20–30 % | 147 |
| $N\rho$ | <8 % | † |
| $N(\pi\pi)_{S\text{-wave}}^{I=0}$ | 5–10 % | – |
| $p\gamma$ | 0.035–0.048 % | 414 |
| $p\gamma$, helicity=1/2 | 0.035–0.048 % | 414 |
| $n\gamma$ | 0.009–0.032 % | 413 |
| $n\gamma$, helicity=1/2 | 0.009–0.032 % | 413 |

 $N(1520) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Breit-Wigner mass = 1515 to 1525 (≈ 1520) MeVBreit-Wigner full width = 100 to 125 (≈ 115) MeV $p_{\text{beam}} = 0.74 \text{ GeV}/c$ $4\pi\lambda^2 = 23.5 \text{ mb}$ Re(pole position) = 1505 to 1515 (≈ 1510) MeV $-2\text{Im}(\text{pole position}) = 105 \text{ to } 120$ (≈ 110) MeV

| $N(1520)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $N\pi$ | 0.55 to 0.65 | 457 |
| $N\eta$ | $(2.3 \pm 0.4) \times 10^{-3}$ | 155 |
| $N\pi\pi$ | 40–50 % | 414 |
| $\Delta\pi$ | 15–25 % | 230 |
| $N\rho$ | 15–25 % | † |
| $N(\pi\pi)_{S\text{-wave}}^{I=0}$ | <8 % | – |
| $p\gamma$ | 0.46–0.56 % | 470 |
| $p\gamma$, helicity=1/2 | 0.001–0.034 % | 470 |
| $p\gamma$, helicity=3/2 | 0.44–0.53 % | 470 |
| $n\gamma$ | 0.30–0.53 % | 470 |
| $n\gamma$, helicity=1/2 | 0.04–0.10 % | 470 |
| $n\gamma$, helicity=3/2 | 0.25–0.45 % | 470 |

$N(1535) S_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

Breit-Wigner mass = 1525 to 1545 (≈ 1535) MeV
 Breit-Wigner full width = 125 to 175 (≈ 150) MeV
 $p_{\text{beam}} = 0.76 \text{ GeV}/c$ $4\pi\lambda^2 = 22.5 \text{ mb}$
 Re(pole position) = 1490 to 1530 (≈ 1510) MeV
 $-2\text{Im}(\text{pole position}) = 90 \text{ to } 250$ (≈ 170) MeV

| $N(1535)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $N\pi$ | 35–55 % | 468 |
| $N\eta$ | 45–60 % | 186 |
| $N\pi\pi$ | 1–10 % | 426 |
| $\Delta\pi$ | <1 % | 244 |
| $N\rho$ | <4 % | † |
| $N(\pi\pi)_{S\text{-wave}}^{I=0}$ | <3 % | – |
| $N(1440)\pi$ | <7 % | † |
| $p\gamma$ | 0.15–0.35 % | 481 |
| $p\gamma$, helicity=1/2 | 0.15–0.35 % | 481 |
| $n\gamma$ | 0.004–0.29 % | 480 |
| $n\gamma$, helicity=1/2 | 0.004–0.29 % | 480 |

 $N(1650) S_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

Breit-Wigner mass = 1645 to 1670 (≈ 1655) MeV
 Breit-Wigner full width = 145 to 185 (≈ 165) MeV
 $p_{\text{beam}} = 0.97 \text{ GeV}/c$ $4\pi\lambda^2 = 16.2 \text{ mb}$
 Re(pole position) = 1640 to 1670 (≈ 1655) MeV
 $-2\text{Im}(\text{pole position}) = 150 \text{ to } 180$ (≈ 165) MeV

| $N(1650)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $N\pi$ | 0.60 to 0.95 | 551 |
| $N\eta$ | 3–10 % | 354 |
| ΛK | 3–11 % | 179 |
| $N\pi\pi$ | 10–20 % | 517 |
| $\Delta\pi$ | 1–7 % | 349 |
| $N\rho$ | 4–12 % | † |
| $N(\pi\pi)_{S\text{-wave}}^{I=0}$ | <4 % | – |
| $N(1440)\pi$ | <5 % | 156 |
| $p\gamma$ | 0.04–0.18 % | 562 |
| $p\gamma$, helicity=1/2 | 0.04–0.18 % | 562 |
| $n\gamma$ | 0.003–0.17 % | 561 |
| $n\gamma$, helicity=1/2 | 0.003–0.17 % | 561 |

$N(1675) D_{15}$

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^-)$$

Breit-Wigner mass = 1670 to 1680 (≈ 1675) MeVBreit-Wigner full width = 130 to 165 (≈ 150) MeV

$$p_{\text{beam}} = 1.01 \text{ GeV}/c \quad 4\pi\lambda^2 = 15.4 \text{ mb}$$

Re(pole position) = 1655 to 1665 (≈ 1660) MeV–2Im(pole position) = 125 to 150 (≈ 135) MeV

| $N(1675)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $N\pi$ | 0.35 to 0.45 | 564 |
| $N\eta$ | (0.0 \pm 1.0) % | 376 |
| ΛK | <1 % | 216 |
| $N\pi\pi$ | 50–60 % | 532 |
| $\Delta\pi$ | 50–60 % | 366 |
| $N\rho$ | < 1–3 % | † |
| $p\gamma$ | 0.004–0.023 % | 575 |
| $p\gamma$, helicity=1/2 | 0.0–0.015 % | 575 |
| $p\gamma$, helicity=3/2 | 0.0–0.011 % | 575 |
| $n\gamma$ | 0.02–0.12 % | 574 |
| $n\gamma$, helicity=1/2 | 0.006–0.046 % | 574 |
| $n\gamma$, helicity=3/2 | 0.01–0.08 % | 574 |

 $N(1680) F_{15}$

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^+)$$

Breit-Wigner mass = 1680 to 1690 (≈ 1685) MeVBreit-Wigner full width = 120 to 140 (≈ 130) MeV

$$p_{\text{beam}} = 1.02 \text{ GeV}/c \quad 4\pi\lambda^2 = 15.0 \text{ mb}$$

Re(pole position) = 1665 to 1680 (≈ 1675) MeV–2Im(pole position) = 110 to 135 (≈ 120) MeV

| $N(1680)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $N\pi$ | 0.65 to 0.70 | 571 |
| $N\eta$ | (0.0 \pm 1.0) % | 387 |
| $N\pi\pi$ | 30–40 % | 539 |
| $\Delta\pi$ | 5–15 % | 374 |
| $N\rho$ | 3–15 % | † |
| $N(\pi\pi)_{S\text{-wave}}^{I=0}$ | 5–20 % | – |
| $p\gamma$ | 0.21–0.32 % | 581 |

| | | |
|--------------------------|---------------|-----|
| $p\gamma$, helicity=1/2 | 0.001–0.011 % | 581 |
| $p\gamma$, helicity=3/2 | 0.20–0.32 % | 581 |
| $n\gamma$ | 0.021–0.046 % | 581 |
| $n\gamma$, helicity=1/2 | 0.004–0.029 % | 581 |
| $n\gamma$, helicity=3/2 | 0.01–0.024 % | 581 |

$N(1700) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Breit-Wigner mass = 1650 to 1750 (≈ 1700) MeV

Breit-Wigner full width = 50 to 150 (≈ 100) MeV

$$p_{\text{beam}} = 1.05 \text{ GeV}/c \quad 4\pi\chi^2 = 14.5 \text{ mb}$$

Re(pole position) = 1630 to 1730 (≈ 1680) MeV

$-2\text{Im}(\text{pole position}) = 50$ to 150 (≈ 100) MeV

| $N(1700)$ DECAY MODES | Fraction (Γ_i/Γ) | ρ (MeV/c) |
|---|--------------------------------|----------------|
| $N\pi$ | 5–15 % | 581 |
| $N\eta$ | (0.0 ± 1.0) % | 402 |
| ΛK | <3 % | 255 |
| $N\pi\pi$ | 85–95 % | 550 |
| $N\rho$ | <35 % | † |
| $p\gamma$ | 0.01–0.05 % | 591 |
| $p\gamma$, helicity=1/2 | 0.0–0.024 % | 591 |
| $p\gamma$, helicity=3/2 | 0.002–0.026 % | 591 |
| $n\gamma$ | 0.01–0.13 % | 590 |
| $n\gamma$, helicity=1/2 | 0.0–0.09 % | 590 |
| $n\gamma$, helicity=3/2 | 0.01–0.05 % | 590 |

$N(1710) P_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Breit-Wigner mass = 1680 to 1740 (≈ 1710) MeV

Breit-Wigner full width = 50 to 250 (≈ 100) MeV

$$p_{\text{beam}} = 1.07 \text{ GeV}/c \quad 4\pi\chi^2 = 14.2 \text{ mb}$$

Re(pole position) = 1670 to 1770 (≈ 1720) MeV

$-2\text{Im}(\text{pole position}) = 80$ to 380 (≈ 230) MeV

| N(1710) DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|-----------------------------------|--------------------------------|-------------|
| $N\pi$ | 10–20 % | 588 |
| $N\eta$ | (6.2 ± 1.0) % | 412 |
| $N\omega$ | (13.0 ± 2.0) % | † |
| ΛK | 5–25 % | 269 |
| $N\pi\pi$ | 40–90 % | 557 |
| $\Delta\pi$ | 15–40 % | 394 |
| $N\rho$ | 5–25 % | † |
| $N(\pi\pi)_{S\text{-wave}}^{I=0}$ | 10–40 % | – |
| $p\gamma$ | 0.002–0.05% | 598 |
| $p\gamma$, helicity=1/2 | 0.002–0.05% | 598 |
| $n\gamma$ | 0.0–0.02% | 597 |
| $n\gamma$, helicity=1/2 | 0.0–0.02% | 597 |

N(1720) P_{13}

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

Breit-Wigner mass = 1700 to 1750 (≈ 1720) MeV
 Breit-Wigner full width = 150 to 300 (≈ 200) MeV
 $p_{\text{beam}} = 1.09$ GeV/c $4\pi\lambda^2 = 13.9$ mb
 Re(pole position) = 1660 to 1690 (≈ 1675) MeV
 $-2\text{Im}(\text{pole position}) = 115$ to 275 MeV

| N(1720) DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$ | 10–20 % | 594 |
| $N\eta$ | (4.0 ± 1.0) % | 422 |
| ΛK | 1–15 % | 283 |
| $N\pi\pi$ | >70 % | 564 |
| $N\rho$ | 70–85 % | 73 |
| $p\gamma$ | 0.003–0.10 % | 604 |
| $p\gamma$, helicity=1/2 | 0.003–0.08 % | 604 |
| $p\gamma$, helicity=3/2 | 0.001–0.03 % | 604 |
| $n\gamma$ | 0.002–0.39 % | 603 |
| $n\gamma$, helicity=1/2 | 0.0–0.002 % | 603 |
| $n\gamma$, helicity=3/2 | 0.001–0.39 % | 603 |

N(2190) G_{17}

$$I(J^P) = \frac{1}{2}(\frac{7}{2}^-)$$

Breit-Wigner mass = 2100 to 2200 (≈ 2190) MeV
 Breit-Wigner full width = 300 to 700 (≈ 500) MeV
 $p_{\text{beam}} = 2.07$ GeV/c $4\pi\lambda^2 = 6.21$ mb
 Re(pole position) = 2050 to 2100 (≈ 2075) MeV
 $-2\text{Im}(\text{pole position}) = 400$ to 520 (≈ 450) MeV

| N(2190) DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$ | 10–20 % | 888 |
| $N\eta$ | (0.0 ± 1.0) % | 792 |

N(2220) H_{19}

$$I(J^P) = \frac{1}{2}(\frac{9}{2}^+)$$

Breit-Wigner mass = 2200 to 2300 (≈ 2250) MeVBreit-Wigner full width = 350 to 500 (≈ 400) MeV

$$p_{\text{beam}} = 2.21 \text{ GeV}/c \quad 4\pi\lambda^2 = 5.74 \text{ mb}$$

Re(pole position) = 2130 to 2200 (≈ 2170) MeV–2Im(pole position) = 400 to 560 (≈ 480) MeV

| N(2220) DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$ | 10–20 % | 924 |

N(2250) G_{19}

$$I(J^P) = \frac{1}{2}(\frac{9}{2}^-)$$

Breit-Wigner mass = 2200 to 2350 (≈ 2275) MeVBreit-Wigner full width = 230 to 800 (≈ 500) MeV

$$p_{\text{beam}} = 2.27 \text{ GeV}/c \quad 4\pi\lambda^2 = 5.56 \text{ mb}$$

Re(pole position) = 2150 to 2250 (≈ 2200) MeV–2Im(pole position) = 350 to 550 (≈ 450) MeV

| N(2250) DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$ | 5–15 % | 938 |

N(2600) $I_{1,11}$

$$I(J^P) = \frac{1}{2}(\frac{11}{2}^-)$$

Breit-Wigner mass = 2550 to 2750 (≈ 2600) MeVBreit-Wigner full width = 500 to 800 (≈ 650) MeV

$$p_{\text{beam}} = 3.12 \text{ GeV}/c \quad 4\pi\lambda^2 = 3.86 \text{ mb}$$

| N(2600) DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$ | 5–10 % | 1126 |

NOTES

- [a] The masses of the p and n are most precisely known in u (unified atomic mass units). The conversion factor to MeV, $1 u = 931.494043 \pm 0.000080$ MeV, is less well known than are the masses in u .
- [b] These two results are not independent, and both use the more precise measurement of $|q_{\bar{p}}/m_{\bar{p}}|/(q_p/m_p)$.
- [c] The limit is from neutrality-of-matter experiments; it assumes $q_n = q_p + q_e$. See also the charge of the neutron.
- [d] The first limit is for $p \rightarrow$ anything or "disappearance" modes of a bound proton. The second entry, a rough range of limits, assumes the dominant decay modes are among those investigated. For antiprotons the best limit, inferred from the observation of cosmic ray \bar{p} 's is $\tau_{\bar{p}} > 10^7$ yr, the cosmic-ray storage time, but this limit depends on a number of assumptions. The best direct observation of stored antiprotons gives $\tau_{\bar{p}}/B(\bar{p} \rightarrow e^- \gamma) > 7 \times 10^5$ yr.
- [e] There is some controversy about whether nuclear physics and model dependence complicate the analysis for bound neutrons (from which the best limit comes). The first limit here is from reactor experiments with free neutrons.
- [f] The parameters g_A , g_V , and g_{WM} for semileptonic modes are defined by $\bar{B}_f[\gamma_\lambda(g_V + g_A\gamma_5) + i(g_{WM}/m_{B_i}) \sigma_{\lambda\nu} q^\nu]B_i$, and ϕ_{AV} is defined by $g_A/g_V = |g_A/g_V|e^{i\phi_{AV}}$. See the "Note on Baryon Decay Parameters" in the neutron Particle Listings.
- [g] Time-reversal invariance requires this to be 0° or 180° .
- [h] This limit is for γ energies between 35 and 100 keV.