

$\Xi(1820) D_{13}$

$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$ Status: ***

The clearest evidence is an 8-standard-deviation peak in ΛK^- seen by GAY 76C. TEODORO 78 favors $J=3/2$, but cannot make a parity discrimination. BIAGI 87C is consistent with $J=3/2$ and favors negative parity for this J value.

$\Xi(1820)$ MASS


We only average the measurements that appear to us to be most significant and best determined.

| <u>VALUE (MeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> |
|---|-------------|-----------------------|-------------|------------|---|
| 1823 ± 5 | | | | | OUR ESTIMATE |
| 1823.4 ± 1.4 | | | | | OUR AVERAGE |
| 1819.4 ± 3.1 ± 2.0 | 280 | ¹ BIAGI | 87 | SPEC | 0 $\Xi^- \text{Be} \rightarrow (\Lambda K^-) X$ |
| 1826 ± 3 ± 1 | 54 | BIAGI | 87C | SPEC | 0 $\Xi^- \text{Be} \rightarrow (\Lambda \bar{K}^0) X$ |
| 1822 ± 6 | | JENKINS | 83 | MPS | - $K^- p \rightarrow K^+$ (MM) |
| 1830 ± 6 | 300 | BIAGI | 81 | SPEC | - SPS hyperon beam |
| 1823 ± 2 | 130 | GAY | 76C | HBC | - $K^- p$ 4.2 GeV/c |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| 1817 ± 3 | | ADAMOVICH | 99B | WA89 | Σ^- nucleus, 345 GeV |
| 1797 ± 19 | 74 | BRIEFEL | 77 | HBC | 0 $K^- p$ 2.87 GeV/c |
| 1829 ± 9 | 68 | BRIEFEL | 77 | HBC | -0 $\Xi(1530) \pi$ |
| 1860 ± 14 | 39 | BRIEFEL | 77 | HBC | - $\Sigma^- \bar{K}^0$ |
| 1870 ± 9 | 44 | BRIEFEL | 77 | HBC | 0 $\Lambda \bar{K}^0$ |
| 1813 ± 4 | 57 | BRIEFEL | 77 | HBC | - ΛK^- |
| 1807 ± 27 | | DIBIANCA | 75 | DBC | -0 $\Xi \pi \pi, \Xi^* \pi$ |
| 1762 ± 8 | 28 | ² BADIER | 72 | HBC | -0 $\Xi \pi, \Xi \pi \pi, Y K$ |
| 1838 ± 5 | 38 | ² BADIER | 72 | HBC | -0 $\Xi \pi, \Xi \pi \pi, Y K$ |
| 1830 ± 10 | 25 | ³ CRENNELL | 70B | DBC | -0 3.6, 3.9 GeV/c |
| 1826 ± 12 | | ⁴ CRENNELL | 70B | DBC | -0 3.6, 3.9 GeV/c |
| 1830 ± 10 | 40 | ALITTI | 69 | HBC | - $\Lambda, \Sigma \bar{K}$ |
| 1814 ± 4 | 30 | BADIER | 65 | HBC | 0 $\Lambda \bar{K}^0$ |
| 1817 ± 7 | 29 | SMITH | 65C | HBC | -0 $\Lambda \bar{K}^0, \Lambda K^-$ |
| 1770 | | HALSTEINSLID | 63 | FBC | -0 K^- freon 3.5 GeV/c |

$\Xi(1820)$ WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|---|------|-----------------------|------|------|--|
| 24 $\begin{smallmatrix} +15 \\ -10 \end{smallmatrix}$ | | | | | OUR ESTIMATE |
| 24 ± 6 | | | | | OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below. |
| 24.6 \pm 5.3 | 280 | ¹ BIAGI | 87 | SPEC | 0 $\Xi^- \text{Be} \rightarrow (\Lambda K^-) X$ |
| 12 \pm 14 \pm 1.7 | 54 | BIAGI | 87C | SPEC | 0 $\Xi^- \text{Be} \rightarrow (\Lambda \bar{K}^0) X$ |
| 72 \pm 20 | 300 | BIAGI | 81 | SPEC | – SPS hyperon beam |
| 21 \pm 7 | 130 | GAY | 76C | HBC | – $K^- p$ 4.2 GeV/c |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| 23 \pm 13 | | ADAMOVICH | 99B | WA89 | Σ^- nucleus, 345 GeV |
| 99 \pm 57 | 74 | BRIEFEL | 77 | HBC | 0 $K^- p$ 2.87 GeV/c |
| 52 \pm 34 | 68 | BRIEFEL | 77 | HBC | –0 $\Xi(1530)\pi$ |
| 72 \pm 17 | 39 | BRIEFEL | 77 | HBC | – $\Sigma^- \bar{K}^0$ |
| 44 \pm 11 | 44 | BRIEFEL | 77 | HBC | 0 $\Lambda \bar{K}^0$ |
| 26 \pm 11 | 57 | BRIEFEL | 77 | HBC | – ΛK^- |
| 85 \pm 58 | | DIBIANCA | 75 | DBC | –0 $\Xi \pi \pi, \Xi^* \pi$ |
| 51 \pm 13 | | ² BADIER | 72 | HBC | –0 Lower mass |
| 58 \pm 13 | | ² BADIER | 72 | HBC | –0 Higher mass |
| 103 $\begin{smallmatrix} +38 \\ -24 \end{smallmatrix}$ | | ³ CRENNELL | 70B | DBC | –0 3.6, 3.9 GeV/c |
| 48 $\begin{smallmatrix} +36 \\ -19 \end{smallmatrix}$ | | ⁴ CRENNELL | 70B | DBC | –0 3.6, 3.9 GeV/c |
| 55 $\begin{smallmatrix} +40 \\ -20 \end{smallmatrix}$ | | ALITTI | 69 | HBC | – $\Lambda, \Sigma \bar{K}$ |
| 12 \pm 4 | | BADIER | 65 | HBC | 0 $\Lambda \bar{K}^0$ |
| 30 \pm 7 | | SMITH | 65B | HBC | –0 $\Lambda \bar{K}$ |
| < 80 | | HALSTEINSLID63 | FBC | –0 | K^- freon 3.5 GeV/c |

WEIGHTED AVERAGE
 24 ± 6 (Error scaled by 1.5)



$\Xi(1820)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| Γ_1 $\Lambda \bar{K}$ | large |
| Γ_2 $\Sigma \bar{K}$ | small |
| Γ_3 $\Xi \pi$ | small |
| Γ_4 $\Xi(1530)\pi$ | small |
| Γ_5 $\Xi \pi \pi$ (not $\Xi(1530)\pi$) | |

$\Xi(1820)$ BRANCHING RATIOS

The dominant modes seem to be $\Lambda \bar{K}$ and (perhaps) $\Xi(1530)\pi$, but the branching fractions are very poorly determined.

| $\Gamma(\Lambda \bar{K})/\Gamma_{\text{total}}$ | | | | | Γ_1/Γ |
|---|-------------|------|-----|---------|---------------------|
| VALUE | DOCUMENT ID | TECN | CHG | COMMENT | |
| 0.30 ± 0.15 | ALITTI | 69 | HBC | — | $K^- p$ 3.9–5 GeV/c |

| $\Gamma(\Xi \pi)/\Gamma_{\text{total}}$ | | | | | Γ_3/Γ |
|---|-------------|------|-----|---------|---------------------|
| VALUE | DOCUMENT ID | TECN | CHG | COMMENT | |
| 0.10 ± 0.10 | ALITTI | 69 | HBC | — | $K^- p$ 3.9–5 GeV/c |

| $\Gamma(\Xi\pi)/\Gamma(\Lambda\bar{K})$ | | | | | | | Γ_3/Γ_1 |
|---|------------|--------------------|-------------|------------|----------------|-------------------|---------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | | |
| <0.36 | 95 | GAY | 76C | HBC | – | $K^- p$ 4.2 GeV/c | |
| 0.20±0.20 | | BADIER | 65 | HBC | 0 | $K^- p$ 3 GeV/c | |

| $\Gamma(\Xi\pi)/\Gamma(\Xi(1530)\pi)$ | | | | | | | Γ_3/Γ_4 |
|--|--|--------------------|-------------|------------|----------------|--------------------|---------------------|
| <u>VALUE</u> | | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | | |
| 1.5^{+0.6}_{-0.4} | | APSELL | 70 | HBC | 0 | $K^- p$ 2.87 GeV/c | |

| $\Gamma(\Sigma\bar{K})/\Gamma_{total}$ | | | | | | | Γ_2/Γ |
|--|--|--------------------|-------------|------------|----------------|---------------------|-------------------|
| <u>VALUE</u> | | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | | |
| 0.30±0.15 | | ALITTI | 69 | HBC | – | $K^- p$ 3.9–5 GeV/c | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|-------|--|-------|----|------|--|---------------|
| <0.02 | | TRIPP | 67 | RVUE | | Use SMITH 65C |
|-------|--|-------|----|------|--|---------------|

| $\Gamma(\Sigma\bar{K})/\Gamma(\Lambda\bar{K})$ | | | | | | | Γ_2/Γ_1 |
|--|--|--------------------|-------------|------------|----------------|-------------------|---------------------|
| <u>VALUE</u> | | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | | |
| 0.24±0.10 | | GAY | 76C | HBC | – | $K^- p$ 4.2 GeV/c | |

| $\Gamma(\Xi(1530)\pi)/\Gamma_{total}$ | | | | | | | Γ_4/Γ |
|---------------------------------------|--|--------------------|-------------|------------|----------------|---------------------|-------------------|
| <u>VALUE</u> | | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | | |
| 0.30±0.15 | | ALITTI | 69 | HBC | – | $K^- p$ 3.9–5 GeV/c | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|----------|--|----------------------|-----|------|--|-------------------|
| seen | | ASTON | 85B | LASS | | $K^- p$ 11 GeV/c |
| not seen | | ⁵ HASSALL | 81 | HBC | | $K^- p$ 6.5 GeV/c |
| <0.25 | | ⁶ DAUBER | 69 | HBC | | $K^- p$ 2.7 GeV/c |

| $\Gamma(\Xi(1530)\pi)/\Gamma(\Lambda\bar{K})$ | | | | | | | Γ_4/Γ_1 |
|---|--|-------------------------------------|-------------|------------|----------------|------------------------|---------------------|
| <u>VALUE</u> | | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | | |
| 0.38±0.27 OUR AVERAGE | | Error includes scale factor of 2.3. | | | | | |
| 1.0 ±0.3 | | GAY | 76C | HBC | – | $K^- p$ 4.2 GeV/c | |
| 0.26±0.13 | | SMITH | 65C | HBC | –0 | $K^- p$ 2.45–2.7 GeV/c | |

| $\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(\Lambda\bar{K})$ | | | | | | | Γ_5/Γ_1 |
|--|--|--------------------|-------------|------------|----------------|--------------------|---------------------|
| <u>VALUE</u> | | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | | |
| 0.30±0.20 | | BIAGI | 87 | SPEC | – | Ξ^- Be 116 GeV | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|-------|--|---------------------|-----|-----|----|------------------------|
| <0.14 | | ⁷ BADIER | 65 | HBC | 0 | 1 st. dev. limit |
| >0.1 | | SMITH | 65C | HBC | –0 | $K^- p$ 2.45–2.7 GeV/c |

| $\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(\Xi(1530)\pi)$ | | | | | Γ_5/Γ_4 |
|---|---------------------|------|-----|---------|---------------------|
| VALUE | DOCUMENT ID | TECN | CHG | COMMENT | |
| consistent with zero | GAY | 76C | HBC | — | $K^- p$ 4.2 GeV/c |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.3 ± 0.5 | ⁸ APSELL | 70 | HBC | 0 | $K^- p$ 2.87 GeV/c |

$\Xi(1820)$ FOOTNOTES

- ¹ BIAGI 87 also sees weak signals in the in the $\Xi^- \pi^+ \pi^-$ channel at 1782.6 ± 1.4 MeV ($\Gamma = 6.0 \pm 1.5$ MeV) and 1831.9 ± 2.8 MeV ($\Gamma = 9.6 \pm 9.9$ MeV).
- ² BADIER 72 adds all channels and divides the peak into lower and higher mass regions. The data can also be fitted with a single Breit-Wigner of mass 1800 MeV and width 150 MeV.
- ³ From a fit to inclusive $\Xi\pi$, $\Xi\pi\pi$, and ΛK^- spectra.
- ⁴ From a fit to inclusive $\Xi\pi$ and $\Xi\pi\pi$ spectra only.
- ⁵ Including $\Xi\pi\pi$.
- ⁶ DAUBER 69 uses in part the same data as SMITH 65C.
- ⁷ For the decay mode $\Xi^- \pi^+ \pi^0$ only. This limit includes $\Xi(1530)\pi$.
- ⁸ Or less. Upper limit for the 3-body decay.

$\Xi(1820)$ REFERENCES

| | | | | |
|--------------|-----|------------------|-------------------------------|--------------------------|
| ADAMOVICH | 99B | EPJ C11 271 | M.I. Adamovich <i>et al.</i> | (CERN WA89 Collab.) |
| BIAGI | 87 | ZPHY C34 15 | S.F. Biagi <i>et al.</i> | (BRIS, CERN, GEVA+) |
| BIAGI | 87C | ZPHY C34 175 | S.F. Biagi <i>et al.</i> | (BRIS, CERN, GEVA+) JP |
| ASTON | 85B | PR D32 2270 | D. Aston <i>et al.</i> | (SLAC, CARL, CNRC, CINC) |
| JENKINS | 83 | PRL 51 951 | C.M. Jenkins <i>et al.</i> | (FSU, BRAN, LBL+) |
| BIAGI | 81 | ZPHY C9 305 | S.F. Biagi <i>et al.</i> | (BRIS, CAVE, GEVA+) |
| HASSALL | 81 | NP B189 397 | J.K. Hassall <i>et al.</i> | (CAVE, MSU) |
| TEODORO | 78 | PL 77B 451 | D. Teodoro <i>et al.</i> | (AMST, CERN, NIJM+) JP |
| BRIEFEL | 77 | PR D16 2706 | E. Briefel <i>et al.</i> | (BRAN, UMD, SYRA+) |
| Also | | PRL 23 884 | S.P. Apseil <i>et al.</i> | (BRAN, UMD, SYRA+) |
| GAY | 76C | PL 62B 477 | J.B. Gay <i>et al.</i> | (AMST, CERN, NIJM) IJ |
| DIBIANCA | 75 | NP B98 137 | F.A. Dibianca, R.J. Endorf | (CMU) |
| BADIER | 72 | NP B37 429 | J. Badier <i>et al.</i> | (EPOL) |
| APSELL | 70 | PRL 24 777 | S.P. Apseil <i>et al.</i> | (BRAN, UMD, SYRA+) I |
| CRENNELL | 70B | PR D1 847 | D.J. Crennell <i>et al.</i> | (BNL) |
| ALITTI | 69 | PRL 22 79 | J. Alitti <i>et al.</i> | (BNL, SYRA) I |
| DAUBER | 69 | PR 179 1262 | P.M. Dauber <i>et al.</i> | (LRL) |
| TRIPP | 67 | NP B3 10 | R.D. Tripp <i>et al.</i> | (LRL, SLAC, CERN+) |
| BADIER | 65 | PL 16 171 | J. Badier <i>et al.</i> | (EPOL, SACL, AMST) I |
| SMITH | 65B | Athens Conf. 251 | G.A. Smith, J.S. Lindsey | (LRL) |
| SMITH | 65C | PRL 14 25 | G.A. Smith <i>et al.</i> | (LRL) IJP |
| HALSTEINSLID | 63 | Siena Conf. 1 73 | A. Halsteinslid <i>et al.</i> | (BERG, CERN, EPOL+) I |

OTHER RELATED PAPERS

| | | | | |
|---------|----|------------------|--------------------------------|------------------------|
| TEODORO | 78 | PL 77B 451 | D. Teodoro <i>et al.</i> | (AMST, CERN, NIJM+) JP |
| BRIEFEL | 75 | PR D12 1859 | E. Briefel <i>et al.</i> | (BRAN, UMD, SYRA+) |
| SCHMIDT | 73 | Purdue Conf. 363 | P.E. Schmidt | (BRAN) |
| MERRILL | 68 | PR 167 1202 | D.W. Merrill, J. Button-Shafer | (LRL) |
| SMITH | 64 | PRL 13 61 | G.A. Smith <i>et al.</i> | (LRL) IJP |