

# $\eta_c(2S)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Quantum numbers are quark model predictions.


## $\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3637 ±4</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.7. See the ideogram below.		
3626 ±5 ±6	311	<sup>1</sup> ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
3645.0 ± 5.5 <sup>+4.9</sup> <sub>-7.8</sub>	121 ± 27	AUBERT	05C BABR	$e^+e^- \rightarrow J/\psi c\bar{c}$
3642.9 ± 3.1 ± 1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
3630.8 ± 3.4 ± 1.0	112 ± 24	AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
3654 ±6 ±8	39 ± 11	CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3639 ±7	98 ± 52	<sup>2</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
3594 ±5		<sup>3</sup> EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

<sup>1</sup> From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

<sup>2</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>3</sup> Assuming mass of  $\psi(2S) = 3686$  MeV.



## $\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>14 ± 7 OUR AVERAGE</b>					
6.3 ± 12.4 ± 4.0		61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
17.0 ± 8.3 ± 2.5		112 ± 24	AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>					
<23	90	98 ± 52	<sup>4</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
22 ± 14		121 ± 27	AUBERT	05C BABR	$e^+e^- \rightarrow J/\psi c\bar{c}$
<55	90	39 ± 11	<sup>5</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
<8.0	95		<sup>6</sup> EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$
<p><sup>4</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.  <sup>5</sup> For a mass value of <math>3654 \pm 6</math> MeV  <sup>6</sup> For a mass value of <math>3594 \pm 5</math> MeV</p>					

## $\eta_c(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons	not seen	
$\Gamma_2$ $K\bar{K}\pi$	seen	
$\Gamma_3$ $2\pi^+2\pi^-$	not seen	
$\Gamma_4$ $K^+K^-\pi^+\pi^-$	not seen	
$\Gamma_5$ $2K^+2K^-$	not seen	
$\Gamma_6$ $\rho\bar{\rho}$	not seen	
$\Gamma_7$ $\gamma\gamma$	$<5 \times 10^{-4}$	90%

## $\eta_c(2S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$					$\Gamma_7$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>					
1.3 ± 0.6	<sup>7</sup> ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$		
<p><sup>7</sup> They measure <math>\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)</math>. The value for <math>\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)</math> is derived assuming that the branching fractions for <math>\eta_c(2S)</math> and <math>\eta_c(1S)</math> decays to <math>K_S K\pi</math> are equal and using <math>\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3</math> keV.</p>					

## $\eta_c(2S)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_3\Gamma_7/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<6.5	90	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$	

$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_4\Gamma_7/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

$\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_5\Gamma_7/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+K^-)$

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}^2$   $\Gamma_6\Gamma_7/\Gamma^2$

VALUE (units $10^{-8}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90	<sup>8,9,10</sup> AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8.0	90	<sup>8,9,11</sup> AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
<12.0	90	<sup>9,11</sup> AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$

<sup>8</sup> Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

<sup>9</sup> For a total width  $\Gamma=5$  MeV.

<sup>10</sup> For the resonance mass region 3589–3599 MeV/ $c^2$ .

<sup>11</sup> For the resonance mass region 3575–3660 MeV/ $c^2$ .

$\eta_c(2S)$  BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABREU 98o	DLPH	$e^+e^- \rightarrow e^+e^- + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	<sup>12</sup> EDWARDS 82c	CBAL	$e^+e^- \rightarrow \gamma X$

<sup>12</sup> For a mass value of  $3594 \pm 5$  MeV

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	$39 \pm 11$	<sup>13</sup> CHOI 02	BELL	$B \rightarrow K K_S K^- \pi^+$

<sup>13</sup> For a mass value of  $3654 \pm 6$  MeV

$\Gamma(2\pi^+2\pi^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(2K^+2K^-)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$

**$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{\gamma}/\Gamma$**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.0005</b>	90	<sup>14</sup> WICHT 08	BELL	$B^{\pm} \rightarrow K^{\pm} \gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.01                      90                      LEE                      85                      CBAL                       $\psi' \rightarrow \text{photons}$

<sup>14</sup> WICHT 08 reports  $[B(\eta_c(2S) \rightarrow \gamma\gamma)] \times [B(B^+ \rightarrow \eta_c(2S) K^+)] < 0.18 \times 10^{-6}$ . We divide by our best value  $B(B^+ \rightarrow \eta_c(2S) K^+) = 0.00034$ .

**$\eta_c(2S)$  REFERENCES**

UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

**OTHER RELATED PAPERS**

PENNINGTON	07A	PR D76 077502	M.R. Pennington, DJ. Wilson	(PITT)
SWANSON	06	PRPL 429 243	E.S. Swanson	(PITT)
BADALIAN	03	PR D67 071901	A.M. Badalian, B.L.G. Bakker	
EICHTEN	02	PRL 89 162002	E.J. Eichten, K. Lane, C. Quigg	
ACCIARRI	99T	PL B461 155	M. Acciarri <i>et al.</i>	(L3 Collab.)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
PORTER	81	SLAC Summer Inst. 355	F.C. Porter <i>et al.</i>	(CIT, HARV, PRIN+)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)