

# Technicolor

A REVIEW GOES HERE – Check our WWW List of Reviews

## MASS LIMITS for Resonances in Models of Dynamical Electroweak Symmetry Breaking

<u>VALUE (GeV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>280	95	1 ABAZOV 07I	D0	$\rho\bar{p} \rightarrow \rho_T / \omega_T \rightarrow W\pi_T$
		2 ABULENCIA 05A	CDF	$\rho_T \rightarrow e^+e^-, \mu^+\mu^-$
		3 CHEKANOV 02B	ZEUS	color octet techni- $\pi$
>207	95	4 ABAZOV 01B	D0	$\rho_T \rightarrow e^+e^-$
		5 ABDALLAH 01	DLPH	$e^+e^- \rightarrow \rho_T$
none 90–206.7	95	6 AFFOLDER 00F	CDF	color-singlet techni- $\rho$ , $\rho_T \rightarrow W\pi_T, 2\pi_T$
		7 AFFOLDER 00K	CDF	color-octet techni- $\rho$ , $\rho_{T8} \rightarrow 2\pi_{LQ}$
>480	95	8 AFFOLDER 00L	CDF	top-color $Z'$
		9 ABE 99F	CDF	color-octet techni- $\rho$ , $\rho_{T8} \rightarrow \bar{b}b$
none 350–440	95	10 ABE 99H	CDF	color-octet techni- $\rho$ , $\rho_{T8} \rightarrow 2\pi_{LQ}$
		11 ABE 99N	CDF	techni- $\omega$ , $\omega_T \rightarrow \gamma\bar{b}b$
>465	95	12 ABE 97G	CDF	color-octet techni- $\rho$ , $\rho_{T8} \rightarrow 2\text{jets}$

<sup>1</sup> ABAZOV 07I search for the vector techni-resonances ( $\rho_T, \omega_T$ ) decaying into  $W\pi_T$  with  $W \rightarrow e\nu$  and  $\pi_T \rightarrow b\bar{b}$  or  $b\bar{c}$ . See their Fig. 2 for the exclusion plot in  $M_{\pi_T}$ - $M_{\rho_T}$  plane.

<sup>2</sup> ABULENCIA 05A search for resonances decaying to electron or muon pairs in  $p\bar{p}$  collisions. at  $\sqrt{s} = 1.96$  TeV. The limit assumes Technicolor-scale mass parameters  $M_V = M_A = 500$  GeV.

<sup>3</sup> CHEKANOV 02B search for color octet techni- $\pi$   $P$  decaying into dijets in  $e p$  collisions. See their Fig. 5 for the limit on  $\sigma(ep \rightarrow ePX) \cdot B(P \rightarrow 2j)$ .

<sup>4</sup> ABAZOV 01B searches for vector techni-resonances ( $\rho_T, \omega_T$ ) decaying to  $e^+e^-$ . The limit assumes  $M_{\rho_T} = M_{\omega_T} < M_{\pi_T} + M_W$ .

<sup>5</sup> The limit is independent of the  $\pi_T$  mass. See their Fig. 9 and Fig. 10 for the exclusion plot in the  $M_{\rho_T}$ - $M_{\pi_T}$  plane. ABDALLAH 01 limit on the techni-pion mass is  $M_{\pi_T} > 79.8$  GeV for  $N_D=2$ , assuming its point-like coupling to gauge bosons.

<sup>6</sup> AFFOLDER 00F search for  $\rho_T$  decaying into  $W\pi_T$  or  $\pi_T\pi_T$  with  $W \rightarrow \ell\nu$  and  $\pi_T \rightarrow \bar{b}b, \bar{b}c$ . See Fig. 1 in the above Note on “Dynamical Electroweak Symmetry Breaking” for the exclusion plot in the  $M_{\rho_T}$ - $M_{\pi_T}$  plane.

<sup>7</sup> AFFOLDER 00K search for the  $\rho_{T8}$  decaying into  $\pi_{LQ}\pi_{LQ}$  with  $\pi_{LQ} \rightarrow b\nu$ . For  $\pi_{LQ} \rightarrow c\nu$ , the limit is  $M_{\rho_{T8}} > 510$  GeV. See their Fig. 2 and Fig. 3 for the exclusion plot in the  $M_{\rho_{T8}}$ - $M_{\pi_{LQ}}$  plane.

<sup>8</sup> AFFOLDER 00L search for top-color  $Z'_{\text{top}}$  decaying into  $\bar{t}t$ . The quoted limit is for  $Z'_{\text{top}}$  with decay width  $\Gamma=0.012 M_{Z'}$ . For  $\Gamma=0.04 M_{Z'}$ , the limit becomes 780 GeV.

<sup>9</sup> ABE 99F search for a new particle  $X$  decaying into  $b\bar{b}$  in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.8$  TeV. See Fig. 7 in the above Note on “Dynamical Electroweak Symmetry Breaking” for the

- upper limit on  $\sigma(p\bar{p} \rightarrow X) \times B(X \rightarrow b\bar{b})$ . ABE 99F also exclude top gluons of width  $\Gamma=0.3M$  in the mass interval  $280 < M < 670$  GeV, of width  $\Gamma=0.5M$  in the mass interval  $340 < M < 640$  GeV, and of width  $\Gamma=0.7M$  in the mass interval  $375 < M < 560$  GeV.
- <sup>10</sup> ABE 99H search for the color-octet techni- $\rho$  decaying into a pair of color-triplet technipions which subsequently decay into  $\tau + \text{jet}$ . See Fig. 6 in the above Note on “Dynamical Electroweak Symmetry Breaking” for the exclusion plot in the  $M_{\rho_{T8}} - M_{\pi_{LQ}}$  plane.
- <sup>11</sup> ABE 99N search for the techni- $\omega$  decaying into  $\gamma\pi_T$ . The technipion is assumed to decay  $\pi_T \rightarrow b\bar{b}$ . See Fig. 2 in the above Note on “Dynamical Electroweak Symmetry Breaking” for the exclusion plot in the  $M_{\omega_T} - M_{\pi_T}$  plane.
- <sup>12</sup> ABE 97G search for a new particle  $X$  decaying into dijets in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.8$  TeV. See Fig. 5 in the above Note on “Dynamical Electroweak Symmetry Breaking” for the upper limit on  $\sigma(p\bar{p} \rightarrow X) \times B(X \rightarrow 2j)$ .

---

## REFERENCES FOR Technicolor

ABAZOV	07I	PRL 98 221801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABULENCIA	05A	PRL 95 252001	A. Abulencia <i>et al.</i>	(CDF Collab.)
CHEKANOV	02B	PL B531 9	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABAZOV	01B	PRL 87 061802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	01	EPJ C22 17	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
AFFOLDER	00F	PRL 84 1110	T. Affolder <i>et al.</i>	(CDF Collab.)
AFFOLDER	00K	PRL 85 2056	T. Affolder <i>et al.</i>	(CDF Collab.)
AFFOLDER	00L	PRL 85 2062	T. Affolder <i>et al.</i>	(CDF Collab.)
ABE	99F	PRL 82 2038	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	99H	PRL 82 3206	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	99N	PRL 83 3124	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	97G	PR D55 R5263	F. Abe <i>et al.</i>	(CDF Collab.)

---